
**Numerical
implementation and
oceanographic
application**

R. Feistel

Numerical implementation and oceanographic application of the Gibbs thermodynamic potential of seawater

R. Feistel

Baltic Sea Research Institute, Seestraße 15, D-18119 Warnemünde, Germany

Received: 11 November 2004 – Accepted: 16 November 2004 – Published: 17 November 2004

Correspondence to: R. Feistel (rainer.feistel@io-warnemuende.de)

© 2004 Author(s). This work is licensed under a Creative Commons License.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

Abstract

The 2003 Gibbs thermodynamic potential function represents a very accurate, compact, consistent and comprehensive formulation of equilibrium properties of seawater. It is expressed in the International Temperature Scale ITS-90 and is fully consistent with the current scientific pure water standard, IAPWS-95. Source code examples in FORTRAN, C++ and Visual Basic are presented for the numerical implementation of the potential function and its partial derivatives, as well as for potential temperature. A collection of thermodynamic formulas and relations is given for possible applications in oceanography, from density and chemical potential over entropy and potential density to mixing heat and entropy production. For colligative properties like vapour pressure, freezing points, and for a Gibbs potential of sea ice, the equations relating the Gibbs function of seawater to those of vapour and ice are presented.

1. Introduction

Thermodynamic potential functions (also called fundamental equations of state) offer a very compact and consistent way of representing equilibrium properties of a given substance, both theoretically and numerically (Alberty, 2001). This was very successfully demonstrated by subsequent standard formulations for water and steam (Wagner and Pruß, 2002). For seawater, this method was first studied by Fofonoff (1962) and later applied numerically in three subsequently improved versions by Feistel (1993), Feistel and Hagen (1995), and Feistel (2003), expressing free enthalpy (also called Gibbs energy) as function of pressure, temperature and practical salinity. Their mathematical structures are polynomial-like and have remained identical throughout these versions with only slight modifications of their sets of coefficients. The structure was chosen for its simplicity in analytical partial derivatives and its numerical implementation, as discussed in Feistel (1993).

This paper provides code examples for the numerical computation in FORTRAN,

Numerical implementation and oceanographic application

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Numerical
implementation and
oceanographic
application**R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

C++ and Visual Basic 6, and describes their algorithms for the latter case. This code is neither very compact, nor very fast, nor definitely error-free; it is just intended as functioning example and guide for the development of individual implementations into custom program environments. Users are free to use, modify and distribute the code at their own responsibility.

The recent Gibbs potential formulation of seawater thermodynamics has a number of advantages compared to the classical “EOS80”, the 1980 Equation of State (Fofonoff and Millard, 1983), as explained in detail by Feistel (2003). One important reason is that it is fully consistent with the current international scientific standard formulation of liquid and gaseous pure water, IAPWS-95 (Wagner and Pruß, 2002), and with a new comprehensive description of ice (Feistel and Wagner, 2005). It is valid for pressures from the triple point to 100 MPa (10 000 dbar), temperatures from -2°C to 40°C , for practical salinities up to 42 psu and up to 50 psu at normal pressure.

For faster computation, as e.g. required in circulation models, modified equations of state derived from the 1995 and 2003 Gibbs potential functions have recently been constructed by McDougall et al. (2003) and Jackett et al. (2004)¹.

A significant advantage compared to the usual EOS80 formulation of seawater properties is the new availability of quantities like energy, enthalpy, entropy, or chemical potential. We present in section 3 a collection of important thermodynamic and oceanographic relations with brief explanations, for which the new potential function can be applied. Such formulas are often only found scattered over various articles and textbooks. In chapter 4, the Gibbs function of seawater is used in conjunction with numerically available thermodynamic formulations for water vapour and water ice, consistent with the current one (Tillner-Roth, 1998; Wagner and Pruß, 2002; Feistel, 2003; Feistel and Wagner, 2005). This way colligative properties like vapour pressure or freezing points can be computed, as well as various properties of sea ice.

¹Jackett, D. R., McDougall, T. J., Feistel, R., Wright, D. G., and Griffies, S. M.: Updated algorithms for density, potential temperature, conservative temperature and freezing temperature of seawater, J. Atm. Ocean Technol., submitted, 2004.

2. Gibbs potential and its derivatives

Specific free enthalpy (also called Gibbs function, Gibbs energy, Gibbs free energy, or free energy in the literature) of seawater, $g(S, t, p)$, is assumed to be a polynomial-like function of the independent variables practical salinity, $S = x^2 \cdot S_U$, temperature, $t = y \cdot t_U$, and applied pressure, $p = z \cdot p_U$, as,

$$\frac{g(S, t, p)}{g_U} = (g_{100} + g_{110}y) x^2 \ln x + \sum_{j,k} \left(g_{0jk} + \sum_{i>1} g_{ijk} x^i \right) y^j z^k. \quad (1)$$

The unit specific free enthalpy is $g_U = 1 \text{ J kg}^{-1}$. The reference values are defined arbitrarily as $S_U = 40 \text{ psu}$ for salinity (PSS-78) (Lewis and Perkin, 1981; Unesco, 1981), $t_U = 40^\circ\text{C}$ for temperature (ITS-90) (Blanke, 1989; Preston-Thomas, 1990), and $p_U = 100 \text{ MPa} = 10\,000 \text{ dbar}$ for pressure. The dimensionless variables x, y, z for salinity, temperature and pressure are not to be confused with spatial coordinates. We follow Fofonoff's (1992) proposal here and write for clarity "psu" as unit expressing practical salinity, even though this notion is formally not recommended (Siedler, 1998). We shall use capital symbols $T = T_0 + t$ for absolute temperatures, with Celsius zero point $T_0 = 273.15 \text{ K}$, and $P = P_0 + p$ for absolute pressures, with normal pressure $P_0 = 0.101325 \text{ MPa}$, in the following. The polynomial coefficients g_{ijk} are listed in Table 1. The specific dependence on salinity results from Planck's theory of ideal solutions and the Debye-Hückel theory of electrolytes (Landau and Lifschitz, 1966; Falkenhagen et al., 1971), providing a thermodynamically correct low-salinity limit of the equation.

There are three first derivatives of g with respect to its independent variables p, t , and S .

Density, ρ , and specific volume, v :

$$\frac{1}{\rho} = v = \left(\frac{\partial g}{\partial p} \right)_{S,t} \quad (2)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

$$\text{with } \left(\frac{\partial g}{\partial p}\right)_{S,t} = \frac{g_U}{p_U} \sum_{j,k>0} \left(g_{0jk} + \sum_{i>1} g_{ijk} x^i \right) \cdot k \cdot y^j z^{k-1}.$$

Specific entropy, σ :

$$\sigma = - \left(\frac{\partial g}{\partial t}\right)_{S,p}, \quad (3)$$

$$\text{with } \left(\frac{\partial g}{\partial t}\right)_{S,p} = \frac{g_U}{t_U} \left[g_{110} x^2 \ln x + \sum_{j>0,k} \left(g_{0jk} + \sum_{i>1} g_{ijk} x^i \right) \cdot j \cdot y^{j-1} z^k \right].$$

5 Relative chemical potential, μ :

$$\mu = \left(\frac{\partial g}{\partial S}\right)_{t,p} \quad (4)$$

$$\text{with } \left(\frac{\partial g}{\partial S}\right)_{t,p} = \frac{g_U}{2S_U} \left[(g_{100} + g_{110} y) (2 \ln x + 1) + \sum_{i>1,j,k} g_{ijk} \cdot i \cdot x^{i-2} y^j z^k \right].$$

Several thermodynamic coefficients require second derivatives of g .

Isothermal compressibility, K :

$$K = -\frac{1}{v} \left(\frac{\partial v}{\partial p}\right)_{S,t} = -\frac{\left(\partial^2 g / \partial p^2\right)_{S,t}}{\left(\partial g / \partial p\right)_{S,t}} \quad (5)$$

$$\text{with } \left(\frac{\partial^2 g}{\partial p^2}\right)_{S,t} = \frac{g_U}{p_U^2} \sum_{j,k>1} \left(g_{0jk} + \sum_{i>1} g_{ijk} x^i \right) \cdot k(k-1) \cdot y^j z^{k-2}.$$

Isobaric thermal expansion coefficient, α :

$$\alpha = \frac{1}{v} \left(\frac{\partial v}{\partial t}\right)_{S,p} = \frac{\left(\partial^2 g / \partial t \partial p\right)_S}{\left(\partial g / \partial p\right)_{S,t}} \quad (6)$$

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

Numerical implementation and oceanographic application

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

$$\text{with } \left(\frac{\partial^2 g}{\partial p \partial t} \right)_S = \frac{g_U}{\rho_U t_U} \sum_{j>0, k>0} \left(g_{0jk} + \sum_{i>1} g_{ijk} x^i \right) \cdot j \cdot k \cdot y^{j-1} z^{k-1}$$

Isobaric specific heat capacity, c_p :

$$c_p = T \left(\frac{\partial \sigma}{\partial t} \right)_{S,p} = \left(\frac{\partial h}{\partial t} \right)_{S,p} = -T \left(\frac{\partial^2 g}{\partial t^2} \right)_{S,p} \quad (7)$$

$$\text{with } \left(\frac{\partial^2 g}{\partial t^2} \right)_{S,p} = \frac{g_U}{t_U^2} \sum_{j>1, k} \left(g_{0jk} + \sum_{i>1} g_{ijk} x^i \right) \cdot j(j-1) \cdot y^{j-2} z^k.$$

5 h is specific enthalpy, as defined below in Eq. (10).

Isothermal haline contraction coefficient, β :

$$\beta = -\frac{1}{v} \left(\frac{\partial v}{\partial S} \right)_{t,p} = -\frac{\left(\partial^2 g / \partial p \partial S \right)_t}{\left(\partial g / \partial p \right)_{S,t}} \quad (8)$$

$$\text{with } \left(\frac{\partial^2 g}{\partial p \partial S} \right)_t = \frac{g_U}{2\rho_U S_U} \sum_{i>1, j, k>0} g_{ijk} \cdot i \cdot k \cdot x^{i-2} y^j z^{k-1}.$$

10 The Gibbs potential and its partial derivatives as given by Eqs. (1)–(8) are available in the sample code by a function call of **GSTP03(nS, nT, nP, S, Tabs, Pabs)**. Input parameters **nS**, **nP** and **nT** are the orders of partial derivatives to be carried out with respect to S , T and P . Input parameters **S**, **TabS**, **Pabs** are the arguments for salinity S in psu, for absolute temperature T in K, and for absolute pressure P in Pa. Only lowest salinity derivatives are supported by the code, $\mathbf{nS} \leq 2$ for $\mathbf{S} > 0$ and $\mathbf{nS} \leq 1$ for $\mathbf{S} \geq 0$. Higher S -derivatives are hardly required in practical applications. A prior call of the procedure **COEFFS03** is mandatory to initialise the array of coefficients g_{ijk} before calling **GSTP03** the first time.

15 The function **GSTP03** is a wrapper for the function **Gxyz(nx, ny, nz, x, y, z)** which represents the right-hand side of Eq. (1) without the leading logarithm term. Input parameters **nx**, **ny** and **nz** are the orders of derivatives with respect to the dimensionless

variables **x**, **y** and **z**. A call of the procedure **COEFFS03** is required to initialise the array of coefficients g_{ijk} before calling **Gxyz** the first time.

When the code is compiled and started, a procedure **F03demo(psu, degC, dbar)** is executed automatically. It creates a sample output with the input values **psu=35**,
5 **degC=20**, **dbar=2000**. The corresponding piece of code in VB looks like,

```
Sub F03demo(ByVal S_psu As Double, _  
            ByVal t_degC As Double, _  
            ByVal p_dbar As Double)  
  
10 Dim S As Double, T As Double, P As Double  
  
Call COEFFS03  
  
15 S = S_psu           'psu -> psu  
T = t_degC + 273.15   'degC -> K  
P = p_dbar * 10000# + 101325# 'dBar -> Pa  
  
Debug.Print "S=", S_psu; "psu"  
20 Debug.Print "T=", t_degC; "°C"  
Debug.Print "P=", p_dbar; "dbar"  
Debug.Print ""  
Debug.Print "free enthalpy", GSTP03(0, 0, 0, S, T, P); "J/kg"  
Debug.Print "chem. pot.", GSTP03(1, 0, 0, S, T, P); "J/kg psu"  
25 Debug.Print "entropy", -GSTP03(0, 1, 0, S, T, P); "J/kgK"  
Debug.Print "density", 1#/GSTP03(0, 0, 1, S, T, P); "kg/m^3"  
Debug.Print "heat capacity", -T * GSTP03(0, 2, 0, S, T, P); "J/kgK"  
Debug.Print "Ch. pot. H2O", GSTP03(0, 0, 0, S, T, P)-S*GSTP03(1, 0, 0, S, T, P);  
    "J/kg"
```

**Numerical
implementation and
oceanographic
application**

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

```
Debug.Print "therm. exp.", GSTP03(0, 1, 1, S, T, P)/GSTP03(0, 0, 1, S, T, P); "1/K"  
Debug.Print "compressib.", -GSTP03(0, 0, 2, S, T, P)/GSTP03(0, 0, 1, S, T, P); "1/Pa"  
Debug.Print "lapse rate", -GSTP03(0, 1, 1, S, T, P)/GSTP03(0, 2, 0, S, T, P); "K/Pa"  
Debug.Print "pot. temp.", PotTemp(S_psu, t_degC, p_dbar); "°C"
```

5

End Sub

Only 8-byte floating points should be used ("**Double**"), indicated here by "#" in VB.
The produced data should look like the following check value printout:

10

```
S=35 psu  
T=20°C  
P=2000 dbar
```

15

```
free enthalpy 16583.1806714797 J/kg  
chem. pot. 60.0099366692805 J/kg psu  
entropy 276.780886190056 J/kgK  
density 1033.32930433584 kg/m3  
heat capacity 3951.77837149032 J/kgK  
20 Ch. pot. H2O 14482.8328880549 J/kg  
therm. exp. 2.78522499678412E-04 1/K  
compressib. 4.06129773355324E-10 1/Pa  
lapse rate 1.99948825300137E-08 K/Pa  
25 pot. temp. 19.617987328589 °C
```

The reader may modify the input values as desired in the startup routine. In the case of Visual Basic, this procedure is **Form.Load**, and the output goes to the immediate (**Debug**) window of the VB developer environment.

Numerical implementation and oceanographic application

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

3. Related oceanographic quantities

Many additional properties of seawater can be computed by combinations of the derivatives given in the former section. A first group is that of thermodynamic functions available from g by mathematical so-called Legendre transforms (Alberty, 2001).

5 Specific free energy (also called Helmholtz energy or Helmholtz free energy), f :

$$f = g - Pv = g - P \cdot \left(\frac{\partial g}{\partial p} \right)_{S,t} \quad (9)$$

Specific enthalpy, h :

$$h = g + T\sigma = g - T \cdot \left(\frac{\partial g}{\partial t} \right)_{S,p} \quad (10)$$

Specific internal energy, e :

$$10 \quad e = g + T\sigma - Pv = g - T \cdot \left(\frac{\partial g}{\partial t} \right)_{S,p} - P \cdot \left(\frac{\partial g}{\partial p} \right)_{S,t} \quad (11)$$

Chemical potential of water in seawater, μ^W :

$$\mu^W = g - S\mu = g - S \cdot \left(\frac{\partial g}{\partial S} \right)_{t,p} \quad (12)$$

A second group is that of adiabatic quantities, describing isentropic processes, i.e. without heat exchange.

15 Adiabatic lapse rate, Γ :

$$\Gamma = \left(\frac{\partial t}{\partial p} \right)_{S,\sigma} = - \frac{\left(\partial^2 g / \partial t \partial p \right)_S}{\left(\partial^2 g / \partial t^2 \right)_{S,p}} = \frac{\alpha T v}{c_p} \quad (13)$$

Adiabatic compressibility, κ , and sound speed, U :

$$\kappa = -\frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_{S,\sigma} = \frac{\left(\frac{\partial^2 g}{\partial t \partial p} \right)_S^2 - \left(\frac{\partial^2 g}{\partial t^2} \right)_{S,p} \left(\frac{\partial^2 g}{\partial p^2} \right)_{S,t}}{\left(\frac{\partial g}{\partial p} \right)_{S,t} \left(\frac{\partial^2 g}{\partial t^2} \right)_{S,p}}. \quad (14)$$

$$\kappa = \frac{v}{U^2} = K - \frac{\alpha^2 T v}{c_p} = K - \alpha \Gamma$$

Adiabatic haline contraction coefficient, β_σ :

$$\beta_\sigma = -\frac{1}{v} \left(\frac{\partial v}{\partial S} \right)_{\sigma,p} = \frac{\left(\frac{\partial^2 g}{\partial S \partial t} \right)_p \left(\frac{\partial^2 g}{\partial t \partial p} \right)_S - \left(\frac{\partial^2 g}{\partial t^2} \right)_{S,p} \left(\frac{\partial^2 g}{\partial S \partial p} \right)_t}{\left(\frac{\partial g}{\partial p} \right)_{S,t} \left(\frac{\partial^2 g}{\partial t^2} \right)_{S,p}}. \quad (15)$$

Closely related to the adiabatic quantities are the so-called ‘potential’ ones, which can be directly computed from entropy (Bradshaw, 1978; Feistel, 1993; Feistel and Hagen, 1994; McDougall et al., 2003; McDougall and Feistel, 2003). They are obtained by formally replacing in-situ temperature t and in-situ pressure p by potential temperature θ and reference pressure p_r . They describe the property a water parcel would take if moved from in-situ pressure p to reference pressure p_r without exchange of matter and heat. By definition of θ , specific entropy is equal to “potential” specific entropy.

Potential temperature, $\theta(S, t, p, p_r)$, is implicitly given by

$$\sigma(S, t, p) = \sigma(S, \theta, p_r). \quad (16)$$

This equation can be solved numerically by Newton iteration and avoids Runge-Kutta integration (Fofonoff, 1985). An example for the algorithm is provided in the accompanying code by the function **PotTemp(Spsu, tdegC, pdBar, prefdBar)**, which uses the input parameters S , t , p , p_r and returns potential temperature θ in °C as solution of Eq. (16). Once potential temperature $\theta(S, t, p, p_r)$ is known, other related “potential” quantities can be computed straight forward.

Potential density, ρ_θ , is defined as

$$\rho_\theta(S, t, p, p_r) = \rho(S, \theta, p_r). \quad (17)$$

For a given profile at geographic position (x, y) , the vertical derivative of potential density provides Brunt-Väisälä frequency, N , describing vertical stability of the water column (z pointing in direction of gravity acceleration, G)

$$N^2 = \frac{G}{\rho} \left(\frac{\partial \rho_\theta}{\partial z} \right)_{x,y} = \frac{G}{\rho} \left(\frac{\partial \rho}{\partial z} \right)_{x,y} - \frac{G^2}{U^2}. \quad (18)$$

Potential enthalpy, h_θ ,

$$h_\theta(S, t, p, p_r) = h(S, \theta, p_r) \quad (19)$$

is supposed to benefit from the combination of conservative behaviour of potential temperature during adiabatic excursions, Eq. (16), and conservation of enthalpy during isobaric mixing, Eq. (28). For a more detailed discussion of potential enthalpy, see McDougall (2003), McDougall and Feistel (2003).

The hydrostatic equilibrium pressure stratification in an external gravity field with acceleration G in z -direction is given by the solution of the differential equation

$$\frac{1}{\rho} \frac{\partial p}{\partial z} = G. \quad (20)$$

If G is a constant, and the vertical profiles $S(p)$ of salinity and $t(p)$ of temperature are known e.g. from a CTD cast, the solution of Eq. (20) is usually obtained by separation of variables and numerical integration over p , as,

$$\int_{p_0}^p v \{S(p), t(p), p\} dp = G \cdot (z - z_0). \quad (21)$$

For important special cases, however, Eq. (20) can be solved analytically in compact form. If both temperature t and salinity S are constant over the water column, we can use Eq. (2) and obtain upon integration

$$g \{S, t, \rho(z)\} - g \{S, t, \rho_0\} = G \cdot (z - z_0), \quad (22)$$

5 i.e. free enthalpy g is a linear function of depth.

If, however, salinity S and entropy σ (or potential temperature) are constant over the water column, as e.g. in the cases of winter convection or wind mixing, we can use the relation $v = (\partial h / \partial p)_{S, \sigma}$ to find

$$h \{S, \sigma, \rho(z)\} - h \{S, \sigma, \rho_0\} = G \cdot (z - z_0). \quad (23)$$

10 In this case, enthalpy h grows linear with depth, Eq. (10). These equations implicitly define pressure as function of depth, $\rho(z)$, and can be solved numerically by Newton iteration at any given z without integrating over the entire column as in case of Eq. (21).

The total energy per mass of a water parcel moving with advection speed \mathbf{u} is specific internal energy e , Eq. (11), plus kinetic plus potential energy in the gravity field:

$$15 e_{tot} = e + u^2/2 - Gz. \quad (24)$$

If dissipative processes are neglected, energy conservation in the ocean is expressed locally by the continuity equation (τ is used for time here)

$$\frac{\partial}{\partial \tau} (\rho e_{tot}) + \text{div} \left\{ \rho \mathbf{u} \left(h + u^2/2 - Gz \right) \right\} = 0. \quad (25)$$

In a stationary ocean, this equation reduces to

$$20 \mathbf{u} \nabla b = 0, \quad (26)$$

i.e. the Bernoulli function $b \equiv h + u^2/2 - Gz = e_{tot} + p\nu$ is always conserved along the advection trajectories in time-independent flows (Landau and Lifschitz, 1974; Gill,

Numerical implementation and oceanographic application

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Numerical
implementation and
oceanographic
application**

R. Feistel

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

1982; Feistel, 1993; Saunders, 1995). In the practically interesting approximation of geostrophic currents, this equation is expressed as

$$\frac{\partial p}{\partial x} \cdot \frac{\partial b}{\partial y} - \frac{\partial p}{\partial y} \cdot \frac{\partial b}{\partial x} = \frac{\partial(p, b)}{\partial(x, y)} = 0. \quad (27)$$

This vanishing Jacobian implies that the Bernoulli function b is a function of pressure alone on any given depth horizon z , i.e. $b(x, y, z) = b(p, z)$. In other words, in the geostrophic special case b is conserved along the horizontal isobars, which of course are identical with the trajectories.

If a given parcel of seawater is initially inhomogeneous in salinity and/or temperature, and during the progressing mixing process it remains at constant pressure p and without exchange of heat or salt with its surrounding, then its total enthalpy is conserved (Fofonoff, 1962, 1992). Denoting by brackets $\langle \dots \rangle$ the average over the parcel's mass elements, we thus find, comparing the inhomogeneous initial with the final homogeneous state,

$$h(\langle S \rangle, \langle t \rangle + \Delta t, p) = \langle h(S, t, p) \rangle. \quad (28)$$

This equation permits the numerical calculation of excess temperature Δt upon isobaric mixing, e.g. by Newton iteration. The corresponding excess of specific volume, Δv , is then given by

$$v(\langle S \rangle, \langle t \rangle + \Delta t, p) + \Delta v = \langle v(S, t, p) \rangle. \quad (29)$$

If heat is assumed to be exchanged as necessary to keep the parcel's average temperature constant, we can get released mixing heat Δh from

$$h(\langle S \rangle, \langle t \rangle, p) + \Delta h = \langle h(S, t, p) \rangle. \quad (30)$$

This formula is commonly applied to mixing processes with isothermal initial state, $t = \langle t \rangle$. The values of Δt and Δh do not have definite signs for seawater mixing, i.e. either cooling or warming can occur depending on the values of S, t, p .

Internal energy is conserved due to the First Law if the mixing process is conducted isochorically (isopycnically), and without exchange of heat and salt. The corresponding balance

$$e(\langle S \rangle, \langle t \rangle + \Delta t, p + \Delta p) = \langle e(S, t, p) \rangle. \quad (31)$$

5 together with supposed constant specific volume

$$v(\langle S \rangle, \langle t \rangle + \Delta t, p + \Delta p) = \langle v(S, t, p) \rangle \quad (32)$$

represent the two equations required to determine the changes caused in both pressure and temperature, Δp and Δt . From these, nonnegative entropy production $\Delta\sigma$ of the mixing process can be computed, obeying the Second Law,

$$10 \Delta\sigma = \sigma(\langle S \rangle, \langle t \rangle + \Delta t, p + \Delta p) - \langle \sigma(S, t, p) \rangle \geq 0. \quad (33)$$

4. Phase equilibria

Equilibria between seawater and other aqueous phases are controlled by equal chemical potentials of water in both. It is important to use for these computations only formulae with mutually consistent reference points, which for the IAPWS-95 pure water standard is zero entropy and zero internal energy of liquid water at the triple point
15 (Wagner and Pruß, 2002). The same reference point is valid for the 2003 seawater formulation, but not for the earlier ones.

Osmotic pressure of seawater, $\pi(S, t, p)$ is the excess pressure of seawater in equilibrium with pure water behind a membrane impenetrable for salt. It is implicitly given
20 by

$$\mu^W(S, t, p + \pi) = \mu^W(0, t, p). \quad (34)$$

Vapour pressure, $p_V(S, t, p)$, above seawater under pressure p , is implicitly given by

$$\mu^W(S, t, p) = g^{\text{vapour}}(t, p_V). \quad (35)$$

Numerical implementation and oceanographic application

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Additionally to the chemical potential of water in seawater, μ^W , Eq. (12), the chemical potential of water vapour, g^{vapour} , is required here. It is available from the IAPWS-95 formulation for the fluid water phases (Wagner and Pruß, 2002).

Freezing point temperature of seawater, $t_f(S, \rho)$, is implicitly given by

$$5 \quad \mu^W(S, t_f, \rho) = g^{\text{lce}}(t_f, \rho). \quad (36)$$

The chemical potential of ice, g^{lce} , is required here additionally. A low-pressure Gibbs potential of ice is given by Feistel (2003), and a high-pressure formulation by Tillner-Roth (1998). However, the new and significantly improved high-pressure version by Feistel and Wagner (2005) is recommended for use here instead.

10 Sea ice is considered a mixture of ice and seawater, which is usually called brine then, at thermodynamic equilibrium. Its Gibbs potential function, g^{SI} , is given as function of temperature t , pressure p , and bulk salinity s by,

$$g^{\text{SI}}(s, t, p) = w \cdot g(S, t, p) + (1 - w) \cdot g^{\text{lce}}(T, P) \quad (37)$$

(Feistel and Hagen, 1998). The salt of sea ice is entirely contained in the liquid brine phase, so bulk salinity s is related to brine salinity S by the mass fraction w of brine, $s = w \cdot S$. Brine salinity, and therefore the mass ratio of the liquid and solid fractions, follows from the equilibrium condition that the chemical potentials of water in brine, μ^W , and of ice, μ^{lce} , must be equal,

$$\mu^{\text{lce}}(T, P) \equiv g^{\text{lce}}(T, P) = \mu^W(S, t, p) = g(S, t, p) - S \cdot \left(\frac{\partial g}{\partial S} \right)_{t,p}. \quad (38)$$

20 Depending on the pair of independent variables chosen, this equation implicitly defines either brine salinity $S_B(t, p)$ of sea ice, or its melting pressure $P_m(S, t)$, or freezing temperature of seawater $t_f(S, p)$. Assuming brine salinity to be known this way, we can express Eq. (37) in the form

$$g^{\text{SI}}(s, t, p) = g^{\text{lce}}(T, P) + s \cdot \mu^B(t, p), \quad (39)$$

Numerical implementation and oceanographic application

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

where the relative chemical potential of brine, $\mu^B = \left(\frac{\partial g}{\partial S}\right)_{t,p}$ has to be taken at brine salinity $S = S_B(t, p)$. Thus, the Gibbs function of sea ice is linear in bulk salinity with coefficients being functions of pressure and temperature, describing both its separate components, ice and brine.

Thermodynamic properties of sea ice can be obtained from Eq. (39) by partial derivatives in the usual way, e.g. specific entropy $\sigma^{SI} = -\left(\frac{\partial g^{SI}}{\partial t}\right)_{s,p}$, specific enthalpy

$h^{SI} = g^{SI} + T\sigma^{SI}$, or specific volume $v^{SI} = \left(\frac{\partial g^{SI}}{\partial p}\right)_{s,t}$. While density, enthalpy or

entropy are strictly additive in the contributions of ice and brine, coefficients like heat capacity or compressibility include significant additional parts due to phase equilibrium shifts, like latent heat, dilution heat, or haline contraction, which make the properties of sea ice so distinct from those of either ice or seawater alone. Vapour pressure over sea ice equals the one over pure ice at same temperature and pressure, as follows from Eq. (39). Thermodynamic functions for sea ice obtained by the Gibbs function formalism are discussed in more detail by Feistel and Hagen (1998).

Acknowledgements. The author is grateful to S. Feistel, A. Feistel and D. Webb for writing, correcting or improving parts of the source code examples.

References

- Alberty, R. A.: Use of Legendre transforms in chemical thermodynamics, *Pure Appl. Chem.*, 73, 1349–1380, 2001.
- Blanck, W.: Eine neue Temperaturskala. Die Internationale Temperaturskala von 1990 (ITS-90), *PTB-Mitteilungen*, 99, 411–418, 1989.
- Bradshaw, A.: Calculation of the potential temperature of seawater from the effect of pressure on entropy, *Deep-Sea Res.*, 25, 1253–1257, 1978.
- Falkenhagen, H. W., Ebeling, W., and Hertz, H. G.: *Theorie der Elektrolyte*, S. Hirzel Verlag, Leipzig, 1971.

Numerical implementation and oceanographic application

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Numerical
implementation and
oceanographic
application**

R. Feistel

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

- Feistel, R.: Equilibrium thermodynamics of seawater revisited, *Progr. Oceanogr.*, 31, 101–179, 1993.
- Feistel, R.: A new extended Gibbs thermodynamic potential of seawater, *Progr. Oceanogr.*, 58, 43–115, 2003, Corrigendum, 61, 99, 2004.
- 5 Feistel, R. and Hagen, E.: Thermodynamic Quantities in Oceanography, in: *The Oceans: Physical-Chemical Dynamics and Human Impact*, edited by Majumdar, S. K., Miller, E. W., Forbes, G. S., Schmalz, R. F., and Panah, A. A., The Pennsylvania Academy of Science, Easton, 1–16, 1994.
- Feistel, R. and Hagen, E.: On the GIBBS thermodynamic potential of seawater, *Progr. Oceanogr.*, 36, 249–327, 1995.
- 10 Feistel, R. and Hagen, E.: A Gibbs thermodynamic potential of sea ice, *Cold Reg. Sci. Technol.*, 28, 83–142, 1998, Corrigendum, 29, 173–176, 1999.
- Feistel, R. and Wagner, W.: High-pressure thermodynamic Gibbs functions of ice and sea ice, *J. Mar. Res.*, in press, 2005.
- 15 Fofonoff, N. P.: Physical properties of sea-water, in: *The Sea*, edited by Hill, M. N., J. Wiley and Sons, 3–30, 1962.
- Fofonoff, N. P. and Millard, R. C.: Algorithms for the computation of fundamental properties of seawater, *Unesco Tech. Pap. Mar. Sci.*, 44, 1–53, 1983.
- Fofonoff, N. P.: Physical Properties of Seawater: A New Salinity Scale and Equation of State of Seawater, *J. Geophys. Res.*, 90, 3332–3342, 1985.
- 20 Fofonoff, N. P.: *Physical Oceanography*, Lecture Notes EPP-226, Harvard University, 1992.
- Gill, A. E.: *Atmosphere-Ocean Dynamics*, Academic Press, Inc., 1982.
- Lewis, E. L. and Perkin, R. G.: The practical salinity scale 1978: Conversion of existing data, *Deep-Sea Res.*, 28A, 307–328, 1981.
- 25 Landau, L. D. and Lifschitz E. M.: *Statistische Physik*, Akademie-Verlag Berlin, 1966.
- Landau, L. D. and Lifschitz E. M.: *Hydrodynamik*, Akademie-Verlag Berlin, 1974.
- McDougall, T. J.: Potential enthalpy: A conservative oceanic variable for evaluating heat content and heat fluxes, *J. Phys. Oceanogr.*, 33, 945–963, 2003.
- McDougall, T. J. and Feistel, R.: What causes the adiabatic lapse rate?, *Deep-Sea Res. I*, 50, 1523–1535, 2003.
- 30 McDougall, T. J., Jackett, D. R., Wright, D. G., and Feistel, R.: Accurate and Computationally Efficient Algorithms for Potential Temperature and Density of Seawater, *J. Atm. Oceanic Technol.*, 20, 730–741, 2003.

Preston-Thomas, H.: The international temperature scale of 1990 (ITS-90), Metrologia, 27, 3–10, 1990.

Saunders, P. M.: The Bernoulli Function and Flux of Energy in the Ocean, J. Geophys. Res., 100, 22 647–22 648, 1995.

5 Siedler, G.: SI-Einheiten in der Ozeanographie, SI Units in Oceanography, Ber. Inst. Meeresk., Christian-Albrechts-Univ. Kiel, 101, 1–18, 1998.

Tillner-Roth, R.: Fundamental Equations of State, Shaker Verlag, Aachen, 1998.

Unesco: Background papers and supporting data on the Practical Salinity Scale 1978, Unesco Techn. Pap. Mar. Sci., 37, 1–144, 1981.

10 Wagner, W. and Pruß, A.: The IAPWS Formulation 1995 for the Thermodynamic Properties of Ordinary Water Substance for General and Scientific Use, J. Phys. Chem. Ref. Data, 31, 387–535, 2002.

**Numerical
implementation and
oceanographic
application**

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Table 1. Coefficients g_{ijk} of specific free enthalpy $g(S, t, \rho)$, Eq. (1).

i	j	k	g_{ijk}	i	j	k	g_{ijk}	i	j	k	g_{ijk}
0	0	0	101.342743139672	0	5	4	6.48190668077221	2	4	2	74.726141138756
0	0	1	100015.695367145	0	6	0	-18.9843846514172	2	4	3	-36.4872919001588
0	0	2	-2544.5765420363	0	6	1	63.5113936641785	2	5	0	-17.43743842213
0	0	3	284.517778446287	0	6	2	-22.2897317140459	3	0	0	-2432.0947227047
0	0	4	-33.3146754253611	0	6	3	8.17060541818112	3	0	1	199.459603073901
0	0	5	4.20263108803084	0	7	0	3.05081646487967	3	0	2	-52.2940909281335
0	0	6	-0.546428511471039	0	7	1	-9.63108119393062	3	0	3	68.0444942726459
0	1	0	5.90578348518236	1	0	0	5813.28667992895	3	0	4	-3.41251932441282
0	1	1	-270.983805184062	1	1	0	851.295871122672	3	1	0	-493.512590658728
0	1	2	776.153611613101	2	0	0	1376.28030233939	3	1	1	-175.292041186547
0	1	3	-196.51255088122	2	0	1	-3310.49154044839	3	1	2	83.1923927801819
0	1	4	28.9796526294175	2	0	2	384.794152978599	3	1	3	-29.483064349429
0	1	5	-2.13290083518327	2	0	3	-96.5324320107458	3	2	0	-158.720177628421
0	2	0	-12357.785933039	2	0	4	15.8408172766824	3	2	1	383.058066002476
0	2	1	1455.0364540468	2	0	5	-2.62480156590992	3	2	2	-54.1917262517112
0	2	2	-756.558385769359	2	1	0	140.576997717291	3	2	3	25.6398487389914
0	2	3	273.479662323528	2	1	1	729.116529735046	3	3	0	67.5232147262047
0	2	4	-55.5604063817218	2	1	2	-343.956902961561	3	3	1	-460.319931801257
0	2	5	4.34420671917197	2	1	3	124.687671116248	3	4	0	-16.8901274896506
0	3	0	736.741204151612	2	1	4	-31.656964386073	3	4	1	234.565187611355
0	3	1	-672.507778314507	2	1	5	7.04658803315449	4	0	0	2630.93863474177
0	3	2	499.360390819152	2	2	0	929.460016974089	4	0	1	-54.7919133532887
0	3	3	-239.545330654412	2	2	1	-860.764303783977	4	0	2	-4.08193978912261
0	3	4	48.8012518593872	2	2	2	337.409530269367	4	0	3	-30.1755111971161
0	3	5	-1.66307106208905	2	2	3	-178.314556207638	4	1	0	845.15825213234
0	4	0	-148.185936433658	2	2	4	44.2040358308	4	1	1	-22.6683558512829
0	4	1	397.968445406972	2	2	5	-7.92001547211682	5	0	0	-2559.89065469719
0	4	2	-301.815380621876	2	3	0	-260.427286048143	5	0	1	36.0284195611086
0	4	3	152.196371733841	2	3	1	694.244814133268	5	1	0	-810.552561548477
0	4	4	-26.3748377232802	2	3	2	-204.889641964903	6	0	0	1695.91780114244
0	5	0	58.0259125842571	2	3	3	113.561697840594	6	1	0	506.103588839417
0	5	1	-194.618310617595	2	3	4	-11.1282734326413	7	0	0	-466.680815621115
0	5	2	120.520654902025	2	4	0	97.1562727658403	7	1	0	-129.049444012372
0	5	3	-55.2723052340152	2	4	1	-297.728741987187				

**Numerical
implementation and
oceanographic
application**

R. Feistel

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion