



Role of deep water formation in multidecadal variability

K. Lohmann et al.

The role of subpolar deep water formation and Nordic Seas overflows in simulated multidecadal variability of the Atlantic overturning

K. Lohmann¹, J. H. Jungclauss¹, D. Matei¹, J. Mignot^{2,3}, M. Menary⁴,
H. R. Langehaug^{5,6}, J. Ba⁷, Y. Gao^{5,6}, O. H. Otterå^{6,8}, W. Park⁷, and S. Lorenz¹

¹Max Planck Institute for Meteorology, Hamburg, Germany

²LOCEAN, Institute Pierre Simon Laplace, University Pierre and Marie Curie, Paris, France

³Climate and Environmental Physics, and Oeschger Centre of Climate Change Research, University of Bern, Switzerland

⁴Met Office Hadley Center, Exeter, UK

⁵Nansen Environmental and Remote Sensing Center, Bergen, Norway

⁶Bjerknes Centre for Climate Research, Bergen, Norway

⁷Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

⁸Uni Research, Bergen, Norway

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Received: 18 September 2013 – Accepted: 11 October 2013 – Published: 30 October 2013

Correspondence to: K. Lohmann (katja.lohmann@mpimet.mpg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper

OSD

10, 1895–1931, 2013

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

We investigate the respective role of variations in subpolar deep water formation and Nordic Seas overflows for the decadal to multidecadal variability of the Atlantic meridional overturning circulation (AMOC). This is done by analysing long (order of 1000 yr) control simulations with five coupled climate models as well as sensitivity experiments performed with one of the models, in which we suppress the variability of either subpolar deep water formation or Nordic Seas overflows. For all models, the maximum influence of variations in subpolar deep water formation is found at about 45° N, while the maximum influence of variations in Nordic Seas overflows is rather found at 55° N to 60° N. Regarding the two overflow branches, the influence of variations in the Denmark Strait overflow is, for all models, substantially larger than that of variations in the overflow across the Iceland–Scotland–Ridge. The influence of variations in subpolar deep water formation is, on multi-model average, larger than that of variations in the Denmark Strait overflow. This is true both at 45° N, where the maximum standard deviation of decadal to multidecadal AMOC variability is located for all but one model, and at the more classical latitude of 30° N. At 30° N, variations in subpolar deep water formation and Denmark Strait overflow explain, on multi-model average, about half and one third respectively of the decadal to multidecadal AMOC variance.

1 Introduction

The strength of the Atlantic meridional overturning circulation (AMOC) significantly influences the climate in north-western Europe through the associated northward heat transport (e.g. Hall and Bryden, 1982). A detailed understanding of the processes that govern the AMOC variability is therefore a topic of much interest. In particular, the respective role of deep water formation variability north and south of the Greenland–Scotland–Ridge (GSR) in modulating the AMOC strength is still not fully understood.

OSD

10, 1895–1931, 2013

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Several studies based on multi-century long control simulations with coupled climate models (e.g. Bentsen et al., 2004; Jungclauss et al., 2005; Msadek and Frankignoul, 2009; Medhaug et al., 2012; Escudier et al., 2013) suggest that AMOC variability is closely linked to convective mixing in the subpolar North Atlantic, but not strongly related to convective mixing in the Nordic Seas. Using multi-century long control simulations with ten coupled climate models, Ba et al. (2013), on the other hand, find in some of the models significant correlation between AMOC variability and convective mixing in the Nordic Seas. Since the source waters of the overflow water masses at the GSR are relatively complex (e.g. Tanhua et al., 2008; Våge et al., 2013), variations in Nordic Seas convection do not necessarily match with variations in the overflows across the GSR. Using an ocean reanalysis for the last 50 yr, Köhl and Stammer (2008) find a high correlation of the AMOC variability at 48° N with the Denmark Strait overflow as well as with the density at intermediate depth south of the Denmark Strait, but, in contrast to above mentioned modelling studies, a low correlation with the convective mixing in the Labrador Sea.

Based on a 50 yr long simulation with a coupled model of the Arctic/North Atlantic Ocean, Bailey et al. (2005) find that the AMOC variability is well correlated with the integrated surface heat flux over the Labrador Sea. Their model simulation, however, has a shortage of buoyant surface water, leading to an overly strong convection. Using a regional model for the Atlantic Ocean forced with atmospheric reanalysis fields, Böning et al. (2006) show that convection events in the Labrador Sea are followed by a strengthening of the AMOC. Their model configuration, however, excludes changes in the Nordic Seas overflows by imposing climatological conditions at the northern boundary. Based on observations, Latif et al. (2006) suggest that the Atlantic dipole sea surface temperature anomaly index over the last century, which is a fingerprint for AMOC variability (Latif et al., 2004), is a response to the North Atlantic Oscillation and associated changes in Labrador Sea convection. Based on relations derived from ocean-only simulations, they estimate, however, that the observed changes in the density of the Denmark Strait overflow over the last decades translate into AMOC changes

that are of comparable order as the AMOC variability associated with Labrador Sea convection during the last century.

Here we investigate the respective role of deep water formation variability north and south of the GSR for the low-frequency AMOC variability at different latitudes in the North Atlantic in a multi-model framework by analysing long (order of 1000 yr) control simulations with five coupled climate models. Different from some of the previous studies mentioned above (e.g. Bentsen et al., 2004; Jungclauss et al., 2005; Msadek and Frankignoul, 2009; Escudier et al., 2013; Ba et al., 2013), we use the thickness of Labrador Sea Water and the strength of the Nordic Seas overflows rather than convective mixing as indicators for deep water formation south and north of the GSR respectively. Labrador Sea Water thickness and Nordic Seas overflow strength have a more direct effect on the AMOC. Regarding the Nordic Seas overflows, we consider the transport both through the Denmark Strait and across the Iceland–Scotland–Ridge. The influence of the latter on the AMOC variability has so far received little attention in the literature. Based on control simulations the influence of subpolar deep water formation and Nordic Seas overflow variations on the AMOC variability can only be estimated by means of statistical analysis. In order to more directly estimate the influence of these two processes, we perform and analyse two sensitivity experiments with one of the models, in which we artificially suppress the variations in either subpolar deep water formation or Nordic Seas overflows.

In this study, we focus on the decadal to multidecadal AMOC variability. Variability on other time scales, such as centennial or multicentennial, can be found in previous studies: Fischer (2011) discusses centennial variability of AMOC, subpolar deep water formation and Nordic Seas overflows in a 6000 yr long transient Holocene simulation with the MPI coupled climate model applying the same grid configuration as MPI-ESM-CR. Park and Latif (2008) suggest that multidecadal AMOC variability in the KCM control simulation originates in the North Atlantic, whereas multicentennial variability is driven by the Southern Ocean. Menary et al. (2012) link centennial AMOC variability in the HadCM3, KCM and MPI-ESM-CR control simulation to a meridional

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

shift in the intertropical convergence zone and the advection of the associated salinity anomalies into the deep water formation regions. Regarding the seasonal to interannual AMOC variability, according to the literature, it appears to be mainly wind-driven through local Ekman transport (e.g. Köhl, 2005; Biastoch et al., 2008).

The paper is organized as follows: the coupled climate models and the simulated state in the North Atlantic are described in Sect. 2. In Sect. 3, we present the analysis of the multi-model control simulations. In Sect. 4, the sensitivity experiments with one of the models are presented. We discuss the results and summarize the main conclusions in Sect. 5.

2 Model description and simulated state in the North Atlantic Ocean

We analyse millennial scale control simulations with five global coupled climate models, namely the Max Planck Institute for Meteorology Earth System Model (MPI-ESM-CR; Jungclaus et al., 2010), the Kiel Climate Model (KCM; Park and Latif, 2008; Park et al., 2009), the coupled climate model developed at the Institute Pierre-Simon Laplace (IPSLCM4 v2, hereafter IPSLCM4; Marti et al., 2010; Servonnat et al., 2010), the Hadley Center Climate Model (HadCM3; Gordon et al., 2000; Pope et al., 2000) and the Bergen Climate Model (BCM; Furevik et al., 2003; Otterå et al., 2009). These simulations were available within the EU-project THOR (ThermoHaline Overturning – at Risk?). We limit our study to simulations from the project partners, as non-standard model output, such as the overflow transports across the GSR, is needed. The models use relatively coarse-resolution grid configurations and are sufficiently computationally efficient to perform long simulations (1000 yr, BCM 700 yr), allowing robust results on decadal to multidecadal time scales. The horizontal resolution in the North Atlantic is of the order of 100 km. We also include a control simulation with a higher-resolution grid configuration in this study: a 500 yr long control simulation with a previous version of the MPI coupled climate model (MPI-AO-LR; Jungclaus et al., 2006) with an approxi-

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

mately double-resolution grid configuration compared to MPI-ESM-CR. An overview of the model configurations used in this study is given in Table 1.

Regarding the simulated AMOC and North Atlantic climate variability, recent multi-model control simulation studies (including many of the simulations used here) discuss differences among the various coupled climate models in both the representation of the low-frequency North Atlantic climate variability as well as in the mechanisms and feedbacks involved (e.g. Menary et al., 2012; Langehaug et al., 2012b; Gastineau and Frankignoul, 2012; Ba et al., 2013).

The overall maximum AMOC strength in the North Atlantic in the different control simulations (Fig. 1) varies between about 10 and 21 Sv (1 Sverdrup = $10^6 \text{ m}^3 \text{ s}^{-1}$). It is located at a depth of about 1000 m between 30° N and 35° N, with the exception of IPSLCM4, where it is located at 45° N. Compared to the observational estimate at 26.5° N (Cunningham et al., 2007; Kanzow et al., 2010), suggesting a mean AMOC strength of about 18.5 Sv with an interannual variability of about ± 2 Sv, the simulated AMOC strength lies within the observed range at this latitude in all models except for KCM (13 Sv) and IPSLCM4 (9 Sv). For IPSLCM4, we note that the AMOC strength is drifting throughout the simulation due to a relatively short spin-up phase. Most of the models show a too shallow meridional overturning cell. This is indeed a common problem in z-coordinate models due to unrealistically large diapycnal mixing associated with the Nordic Seas overflows (e.g. Willebrand et al., 2001; Danabasoglu et al., 2010; Yeager and Danabasoglu, 2012). The excessive mixing results in a density of the simulated overflow water masses which is too low compared to observations. The simulated lower branch of the AMOC extends below 3000 m only in BCM – the only model using an isopycnic vertical coordinate.

We have determined the dominant periodicity of decadal to multidecadal AMOC variability at different latitudes in the North Atlantic in the various control simulations based on peaks in the power spectra exceeding the 95 % confidence level when compared to a red noise process. The AMOC strength at a particular latitude, we define as the maximum value of the Atlantic meridional overturning streamfunction over all depth

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

levels at that latitude. The dominant periodicity varies among models and latitudes and provides a rather wide range of variability from decadal to multidecadal time scales (20 to 75 yr). Furthermore, Zanchettin et al. (2010, their Fig. 2), using the MPI-ESM-CR control simulation, and Menary et al. (2012, their Fig. 1), using the HadCM3, KCM and MPI-ESM-CR control simulations, show that the dominant periodicity of AMOC variability at 30° N is non-stationary in time. To focus on the decadal to multidecadal time scales, we apply a 10th order Butterworth bandpass filter for periods ranging from 15 to 80 yr to all model data. In the following, we refer to this variability as multidecadal variability.

The mean mixed layer depth (MLD) in March, which indicates the location of deep water formation, differs among the various control simulations (Fig. 2). Deep convection in or off the Labrador Sea is found in MPI-ESM-CR, MPI-AO-LR and BCM. Compared to the observed winter convection depth in the Labrador Sea, which varies from about 800 m to 2300 m, MPI-AO-LR and KCM show a relatively large convection depth exceeding 2500 m in the mean. Apart from deep convection in the Labrador Sea, observations show relatively shallow convection in the Irminger Sea, with a winter convection depth generally not exceeding 500 m. Shallow convection in the Irminger Sea is also found in all models except for KCM and IPSLCM4. In the latter two models, deep convection is shifted south of Iceland. Regarding BCM, where an isopycnic vertical coordinate is used, MLD is not the best indicator for deep water formation (Medhaug et al., 2012). We note, however, that in our definition of subpolar deep water formation variability in BCM, the MLD is not taken into account.

For the definition of a subpolar deep water formation index, we follow Zhu and Jungclauss (2008), using the thickness of Labrador Sea Water. Nevertheless, due to the varying location of subpolar deep water formation in the different models (Fig. 2), we use the expression subpolar deep water, rather than Labrador Sea Water, in the present study. A description of how we determine the thickness of subpolar deep water in the different models is given in Appendix A. As an index for subpolar deep water formation (SDWI), we average the annual thickness of subpolar deep water over the subpolar

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

region (60° W to 15° W and 45° N to 65° N). The entire subpolar region is chosen to account for the different location of the subpolar deep water formation in the various models. Based on the MPI-ESM-CR control simulation, we have also tested defining the SDWI by using the subpolar deep water thickness in March or by averaging the subpolar deep water thickness over the convection region (defined by the grid cells where the MLD in March exceeds 1000 m in at least 100 integration years). The multidecadal variability of the different indices is, however, rather similar, with correlation coefficients exceeding 0.9 (not shown). We are therefore confident that our results do not critically depend on the definition of the SDWI.

Apart from the subpolar deep convection, deep convection is found in the Nordic Seas (Fig. 2). Also in the Nordic Seas, mean location and depth of the deep water formation vary among the different control simulations. As mentioned earlier, we do not directly consider deep water formation in the Nordic Seas, but rather use the overflow transports across the GSR. The overflow transports through Denmark Strait (DSO) and across the Iceland–Scotland–Ridge (ISRO) are defined based on density (and depth) criteria according to the respective water mass properties at the GSR in the different models. In general, the total transport out of the Nordic Seas with a density threshold of $\sigma > 27.8 \text{ kg m}^{-3}$ is used. A description of the criteria used in the different models is given in Appendix B.

Mean simulated values of DSO and ISRO lie between about 2 and 4 Sv (Table 2), in reasonable agreement with observational estimates (e.g. Hansen et al., 2008). An exception is the DSO in KCM, IPSLCM4 and HadCM3, where mean values of the order of 10 Sv are found. The reason for the unrealistically large DSO in the three models is not clear, but we note that all models have realistic sill depths at the GSR. In contrast to observations, where an overflow transport of about 1 Sv is found between Iceland and the Faroes (e.g. Østerhus et al., 2008), the simulated ISRO is (mostly) restricted to the Faroe–Shetland–Channel in all models. South of the Iceland–Scotland–Ridge, one major bias in all models except for KCM is that (a large part of) the Iceland–Scotland overflow water spreads southward in the eastern part of the North Atlantic basin (e.g.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Langehaug et al., 2012a; their Fig. 5), rather than flowing around the Reykjanes Ridge (through fracture zones in the Mid Atlantic Ridge) and joining the Denmark Strait overflow water and the deep western boundary current.

3 Analysis of multi-model control simulations

In this section, we present the results from a correlation analysis between the maximum AMOC strength at different latitudes in the North Atlantic and respectively SDWI, DSO and ISRO in the multi-model control simulations.

Maximum correlation coefficients between SDWI and AMOC strength on multi-decadal time scales (Fig. 3) are found at 45° N in all models, with the exception of MPI-ESM-CR, where they are located at about 40° N. In IPSLCM4, similarly high correlation coefficients are found between about 45° N and 55° N. Maximum correlation coefficients amount to 0.85 to 0.9, with the exception of HadCM3, where they reach a value of only 0.55. We note that HadCM3 also exhibits substantially weaker regression coefficients (per standard deviation of SDWI) compared to the other models (not shown). Maximum regression coefficients amount to 0.35 Sv (per standard deviation of SDWI) in HadCM3 compared to 0.5–0.8 Sv (per standard deviation of SDWI) in the other models. The negative correlation coefficients alternating with the positive ones (e.g. Fig. 3b) indicate the dominant periodicity of AMOC variability and are not further discussed here.

Significant correlation coefficients between SDWI and AMOC strength are found, except for MPI-ESM-CR, at all latitudes in the North Atlantic (latitude range 0° to 15° N not shown). In MPI-ESM-CR, correlation coefficients at zero lag or with the AMOC strength lagging are significant only between about 25° N and 50° N. In HadCM3, correlation coefficients are relatively low compared to the other models (except for partly MPI-ESM-CR) at all latitudes. The same is true, if regression coefficients (per standard deviation of SDWI) rather than correlation coefficients are considered (not shown). We

speculate that this is due to the fact that HadCM3 exhibits a rather shallow subpolar deep water formation (Fig. 2e).

Maximum correlation coefficients between DSO and AMOC strength on multidecadal time scales (Fig. 4) are found at about 60° N in all models. They amount to 0.85 to 0.9, with the exception of BCM, where they reach a value of 0.75. In addition, strong correlation coefficients are found at about 45° N in MPI-AO-LR and KCM as well as for all subpolar latitudes in IPSLCM4. We note that KCM and IPSLCM4 exhibit a much larger variability of the DSO than the other models (standard deviation of the order of 1.5 Sv compared to less than 0.5 Sv in the other models).

Significant correlation coefficients between DSO and AMOC strength are found at all latitudes in the North Atlantic (latitude range 0° to 15° N not shown), with the exception of BCM. In the latter, significant correlation coefficients are only found close to the GSR. We note that BCM is the only model with an isopycnic vertical coordinate. In another multi-model study using partly the same climate models as here, Swingedouw et al. (2013) underline the limited grid resolution of this model in the Denmark Strait region. In KCM and IPSLCM4, correlation coefficients between AMOC strength and respectively SDWI and DSO are rather similar (Fig. 3c and d compared to Fig. 4c and d). In both models, however, SDWI and DSO are highly correlated with each other (correlation coefficient of about 0.8). We note that this could be due to a specific feature of the oceanic component as these two coupled climate models use the same one.

Regarding the correlation between ISRO and AMOC strength on multidecadal time scales (Fig. 5), correlation coefficients are generally rather low compared to the correlation between DSO and AMOC strength. The same is true, if regression coefficients (per standard deviation of ISRO and DSO respectively) rather than correlation coefficients are considered (not shown). In IPSLCM4, HadCM3 and BCM, an out-of-phase relation of DSO and ISRO is found on multidecadal time scales. Such an out-of-phase relation on time scales of 3 to 10 yr is described by Serra et al. (2010), based on an ocean-only simulation with the MITgcm model forced with atmospheric reanalysis fields. Consistent with the out-of-phase relation of the two overflow branches, positive correlation coeffi-

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

cients are found when the AMOC strength leads the ISRO by a few years in IPSLCM4 and HadCM3. BCM shows almost no significant correlation coefficients between ISRO and AMOC strength.

In MPI-ESM-CR and MPI-AO-LR, significant correlation coefficients between ISRO and AMOC strength are found for the subpolar latitudes. In MPI-ESM-CR, correlation coefficients lie between 0.5 and 0.7, while in MPI-AO-LR they are just above the significance level. KCM is the only model, where significant correlation coefficients are found at all latitudes between 10° N and 60° N (latitude range 10° N to 15° N not shown), with correlation coefficients of 0.4 to 0.6. We note that the simulated influence of the ISRO on the multidecadal AMOC variability might, however, be underestimated, as the models (except for KCM) do not realistically simulate the flow path of the Iceland–Scotland overflow water south of the Iceland–Scotland–Ridge.

The latitude where the maximum correlation coefficients between SDWI and AMOC strength are found (about 45° N), coincides, with the exception of HadCM3, with the latitude where the maximum standard deviation of multidecadal AMOC variability is located. The maximum standard deviation is of the order of 0.5 to 1 Sv and is, with the exception of HadCM3, located at about 45° N at a depth of 1500 m (Fig. 6), below the depth at which the overall maximum AMOC strength in the North Atlantic is found. In HadCM3, the maximum standard deviation of multidecadal AMOC variability is found at 60° N, coinciding with the latitude where the maximum correlation coefficients between DSO and AMOC strength are found. We note that HadCM3 shows relatively low correlation coefficients between SDWI and AMOC strength compared to the other models (except for partly MPI-ESM-CR, Fig. 3). Whether the multidecadal AMOC variability at 45° N and 60° N is, to a large extent, indeed driven by variations in subpolar deep water formation and Denmark Strait overflow respectively, can, however, not definitely be determined from the statistical analysis presented in this section.

Based on the correlation analysis for the individual models, we have deduced the multi-model average of the multidecadal AMOC variance explained by SDWI and DSO respectively (Table 3). The ISRO explains substantially less variance than the DSO.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Furthermore, in some models (IPSLCM4, HadCM3 and BCM), it is out-of-phase with the AMOC variability. Thus we do not comment further its influence here. At 45° N, where the maximum standard deviation of multidecadal AMOC variability is found, the variance explained by SDWI and DSO amounts, on multi-model average, to 68 % and 57 % respectively. The sum of the explained variances by SDWI and DSO amounts to more than 100 %, indicating that these two processes are not completely independent from each other. At the more classical latitude of 30° N, the variance explained by SDWI and DSO amounts, on multi-model average, to 50 % and 30 % respectively, suggesting that the influence of both subpolar deep water formation and Nordic Seas overflow on the multidecadal AMOC variability is weaker at this latitude.

4 Sensitivity experiments with MPI-ESM-CR

4.1 Design of sensitivity experiments

The statistical analysis of the multi-model control simulations presented in the previous section could be biased by the fact that subpolar deep water formation and Nordic Seas overflows are somehow coupled. In order to more directly estimate the respective role of these two processes, we have performed two 1000 yr long sensitivity experiments with the MPI-ESM-CR model set-up (Table 1) by suppressing the variations in either subpolar deep water formation (in the following referred to as SDW experiment) or Nordic Seas overflows (in the following referred to as overflow experiment). Initial conditions are the same as for the control simulation.

In the SDW experiment, we relax temperature and salinity in the western subpolar region (60° W to 30° W, 48° N to 64° N; see box in Fig. 2a) towards monthly climatological values obtained from the control simulation. The relaxation time scale is one day. The region where relaxation is applied corresponds to the subpolar deep water formation site. We also include the Irminger Sea in this area, as previous modelling studies suggest a significant correlation between convective mixing in the Irminger Sea and

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

decadal AMOC variability (e.g. Bentsen et al., 2004; Ba et al., 2013). Ba et al. (2013) show that this is true also for the MPI-ESM-CR control simulation. In the Irminger Sea region (45° W to 30° W, 60° N to 64° N; see small box in Fig. 2a), where the convection is relatively shallow, relaxation is applied down to a depth of 500 m, otherwise down to a depth of 900 m. The deeper levels as well as the upper four levels (approximately the upper 40 m of the water column) are excluded from relaxation to keep the variability of the overflow water masses as well as the coupling between ocean and atmosphere unaffected.

In the overflow experiment, we relax temperature and salinity in the Nordic Seas and along the GSR (boundaries indicated by the black lines in Fig. 2a) towards monthly climatological values obtained from the control simulation. The relaxation time scale is one day. The relaxation is applied in the entire water column, excluding the upper four levels to keep the coupling between ocean and atmosphere unaffected. The relaxation of Nordic Seas temperature and salinity towards climatological values also leads to a density of the overflow water masses at the GSR, which is almost constant with time.

Since temperature and salinity are relaxed towards climatological mean values taken from the control simulation, the simulated mean state in the North Atlantic in the two sensitivity experiments remains approximately unchanged compared to the simulated mean state in the control simulation (as described in Sect. 2). SDWI, DSO and ISRO in the sensitivity experiments are defined in the same way as described in Sect. 2 for the control simulation.

4.2 Results

In the SDW experiment, the multidecadal variability of the SDWI is reduced by 82 % compared to the control simulation (Fig. 7a and Table). A reduction of 29 % is, however, also found in the multidecadal variability of the ISRO (Fig. 7c and Table). One possible explanation of this phenomenon is as follows: Reduced AMOC variability (due to reduced subpolar deep water formation variability) leads to reduced SST variability in the eastern part of the Nordic Seas, as indicated by the correlation pattern between

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the maximum AMOC strength and North Atlantic SST in the control simulation (not shown). The variability of the overflow across the Iceland–Scotland–Ridge in ocean-only and coupled simulations using the ocean component of MPI-ESM-CR is significantly correlated with the SST variability in the eastern part of the Nordic Seas and related changes in surface density, convective mixing and sea surface height (Olsen et al., 2008; Lohmann et al., 2013).

In the overflow experiment, the multidecadal variability of both DSO and ISRO is reduced by about 90 % compared to the control simulation (Fig. 7b and c and Table). A reduction of 38 % is, however, also found in the multidecadal variability of the SDWI (Fig. 7a and Table). This might be due to reduced AMOC variability (due to reduced overflow variability) inducing reduced subpolar deep water formation variability caused by ocean internal modes such as the one described in Zhu and Jungclaus (2008). Variations in Nordic Seas overflow and subpolar deep water formation might thus not be completely independent, at least in this model. We note, however, that in the control simulation, SDWI and DSO/ISRO are not significantly correlated with each other. Another possible explanation is that the relaxation of Nordic Seas temperature and salinity towards climatological values reduces, via advection along the East Greenland Current, upper-ocean temperature and salinity variability in the subpolar region, especially the Irminger Sea, and consequently probably affects variability of subpolar deep water formation.

Regarding the multidecadal AMOC variability, the amplitude of variability is reduced in the sensitivity experiments compared to the control simulation (Fig. 8). We quantify the reduction in the standard deviation of the AMOC strength in the sensitivity experiments compared to the control simulation by $100 \cdot [1 - (\sigma(\text{AMOC}_{\text{sensitivity experiment}})/\sigma(\text{AMOC}_{\text{control simulation}}))]$. Note that positive values correspond to a reduction in the standard deviation of the AMOC strength in the sensitivity experiments.

In the SDW experiment (Fig. 9a), the maximum reduction in the standard deviation of multidecadal AMOC variability amounts to about 60 % and is found at about 45° N.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

This latitude is similar to the latitude where the maximum correlation coefficients between SDWI and AMOC strength are found in the control simulation (Fig. 3a). At 30° N, the reduction in the standard deviation of multidecadal AMOC variability in the SDW experiment amounts to about 50 %.

In the overflow experiment (Fig. 9b), the maximum reduction in the standard deviation of multidecadal AMOC variability amounts to about 80 % and is found at 55° N to 60° N, again similarly to the latitude of the maximum correlation coefficients between both DSO and AMOC strength and ISRO and AMOC strength in the control simulation (Figs. 4a and 5a). At 45° N and south of this latitude, the reduction in the standard deviation of multidecadal AMOC variability is similar to what is found in the SDW experiment. We remind the reader, however, that in the overflow experiment also the variability of the SDWI is reduced by almost 40 % (Fig. 7a and Table). It is interesting to note that the maximum reduction in the standard deviation in the overflow experiment is found below the depth at which the overall maximum AMOC strength in the North Atlantic is located (solid line in Fig. 9b). This indicates that the overflow water masses, as in the real ocean, contribute to the lower North Atlantic Deep Water.

In summary, we conclude from the sensitivity experiments that the multidecadal AMOC variability north of about 50° N is mainly related to variations in Nordic Seas overflows. At 45° N and south of this latitude, variations in both subpolar deep water formation and Nordic Seas overflows contribute to the multidecadal AMOC variability, with neither of the processes being very dominant compared to the other.

5 Discussion and conclusions

In this study, we have analysed long control simulations with five coupled climate models as well as sensitivity experiments performed with one of the models in order to add to the discussion in the literature about the respective role of variations in subpolar deep water formation and Nordic Seas overflows for the multidecadal AMOC variability.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Regarding the multi-model control simulations (Table 3), the multidecadal AMOC variance explained by subpolar deep water formation is higher than or of similar order as the one explained by the Denmark Strait overflow. This is true for all models at both 45° N, where the maximum standard deviation of multidecadal AMOC variability is found, and at the more classical latitude of 30° N, with the exception of 45° N in HadCM3. In the latter, the Denmark Strait overflow seems to have a relatively strong influence on the AMOC variability. We note that HadCM3 indeed shows a rather shallow subpolar deep water formation compared to the other models (Fig. 2). We also note that HadCM3 is the only model, in which the maximum standard deviation of multidecadal AMOC variability is found at 60° N, rather than at 45° N. The multidecadal AMOC variance explained by the Denmark Strait overflow is, for all models, substantially larger than the one explained by the overflow across the Iceland–Scotland–Ridge. The latter might, however, be underestimated, as the models in general do not realistically simulate the flow path of the Iceland–Scotland overflow water south of the Iceland–Scotland–Ridge.

The respective role of subpolar deep water formation and Nordic Seas overflows on the multidecadal AMOC variability suggested from the control simulations is supported by the sensitivity experiments. The latter indicate that the maximum influence of Nordic Seas overflow variations is found below the depth at which the overall maximum AMOC strength in the North Atlantic is located. The conclusions of our study do, however, not change, if the correlation analysis based on the multi-model control simulations is repeated for depth levels located in the lower branch of the AMOC.

Our results are in contrast with previous studies, which find the major influence from the Denmark Strait overflow (e.g. Köhl and Stammer, 2008), and rather in line with the ones suggesting a larger influence of subpolar deep water formation on the AMOC variability (e.g. Bailey et al., 2005; Latif et al., 2006; Böning et al., 2006). We note, however, that in only one of the models (BCM) the multidecadal AMOC variance explained by subpolar deep water formation is substantially larger than the one explained by the Denmark Strait overflow. We also note that one possible caveat of the model simula-

tions used here is the generally biased representation of the overflow water masses in z-coordinate models (e.g. Willebrand et al., 2001; Danabasoglu et al., 2010; Yeager and Danabasoglu, 2012).

In some previous modelling studies (e.g. Jungclaus et al., 2005; Msadek and Frankignoul, 2009; Escudier et al., 2013), the AMOC variability was found not to be strongly related to convective mixing in the Nordic Seas. Variations in Nordic Seas convection do, however, not necessarily match with variations in the overflows across the GSR, since the source waters of the overflow water masses at the GSR are relatively complex (e.g. Tanhua et al., 2008; Våge et al., 2013). Our study suggests that the Denmark Strait overflow, though its influence is, on multi-model average, smaller than that of subpolar deep water formation, still explains a substantial part of multidecadal AMOC variance. At the classical latitude of 30° N, variations in Denmark Strait overflow and subpolar deep water formation explain, on multi-model average, about one third and half of multidecadal AMOC variance respectively.

Appendix A

Definition of subpolar deep water thickness

In all models except for BCM, the density range of subpolar deep water is defined based on mean profiles of temperature, salinity and σ_2 (potential density with reference depth 2000 m). σ_2 rather than σ values are chosen, as Labrador Sea Water in the North Atlantic spreads at a depth of about 2000 m. In MPI-ESM-CR, MPI-AO-LR, KCM and IPSLCM4, profiles are averaged over all integration years and the subpolar (60° W to 10° W, 45° N to 65° N) grid cells where the MLD in March exceeds 1000 m in at least 10 % of the integration years. These grid cells correspond approximately to the grid cells where the mean MLD in March exceeds 500 m (Fig. 2).

In HadCM3, no subpolar grid cells are found, where the MLD in March exceeds 1000 m in at least 10 % of the integration years. The same is true, if other winter

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



months (December, January, February) rather than March are considered. If all grid cells, where the MLD in March exceeds 1000 m in at least one integration year, are taken into account, subpolar deep convection is found in the Labrador and Irminger Sea. In HadCM3, profiles of temperature, salinity and σ_2 are therefore averaged only over those integration years and subpolar grid cells, when and where the MLD in March exceeds 1000 m.

The upper isopycnal defining subpolar deep water is chosen below the temperature and salinity maximum representing water from the subtropical Atlantic, at a depth of about 800 m. The lower isopycnal is chosen at the beginning of the deep pycnocline, at a depth of about 2500 to 3000 m. The resulting density range of subpolar deep water for the various models is as follows: MPI-ESM-CR: $\sigma_2 = 36.62\text{--}36.76 \text{ kg m}^{-3}$, MPI-AO-LR: $\sigma_2 = 36.65\text{--}36.77 \text{ kg m}^{-3}$, KCM: $\sigma_2 = 36.90\text{--}37.01 \text{ kg m}^{-3}$, IPSLCM4: $\sigma_2 = 36.76\text{--}36.87 \text{ kg m}^{-3}$, HadCM3: $\sigma_2 = 36.74\text{--}37.03 \text{ kg m}^{-3}$. The thickness of subpolar deep water we define as the depth interval between the upper and lower isopycnal specifying the density range of subpolar deep water.

For BCM, where an isopycnic vertical coordinate is used, we follow Langehaug et al. (2012a, analysing the same simulation as used in our study), defining the layers with $\sigma_2 = 36.846 \text{ kg m}^{-3}$ and $\sigma_2 = 36.946 \text{ kg m}^{-3}$ as (upper) Labrador Sea Water. These density layers occupy approximately the depth range between 600 m and 1800 m in the Labrador Sea (Langehaug et al., 2012a; their Fig. 9). The thickness of subpolar deep water we define as the sum of the layer thickness of these two density layers.

Appendix B

Definition of Nordic Seas overflow transports

The following criteria are used to define the overflow transport through Denmark Strait and across the Iceland–Scotland–Ridge in the various models: MPI-ESM-CR: transport out of Nordic Seas for $\sigma > 27.76 \text{ kg m}^{-3}$ and depth $> 300 \text{ m}$ (through Den-

mark Strait) and for $\sigma > 27.8 \text{ kg m}^{-3}$ (across the Iceland–Scotland–Ridge), MPI-AO-LR: transport out of Nordic Seas for $\sigma > 27.8 \text{ kg m}^{-3}$, KCM: transport out of Nordic Seas for $\sigma > 27.8 \text{ kg m}^{-3}$, IPSLCM4: transport out of Nordic Seas for $\sigma > 27.75 \text{ kg m}^{-3}$, HadCM3: Transport out of Nordic Seas for depth $> 300 \text{ m}$, BCM: net transport for $\sigma_2 > 36.946 \text{ kg m}^{-3}$ (corresponding to about $\sigma > 27.83 \text{ kg m}^{-3}$). The transport is restricted to depth levels greater than 300 m in some of the models to ensure a separation between the Denmark Strait overflow and the lower East Greenland Current. In BCM, σ_2 isopycnal layers are used as the vertical coordinate in the ocean component. Therefore, the density threshold is specified as σ_2 value. In this model, also the net transport rather than the transport out of the Nordic Seas is used. We note, however, that this difference is negligible, as a transport into the Nordic Seas with the given density threshold generally does not exist.

Acknowledgements. This work was supported by the European Community's 7th framework programme (FP7/2007-2013) under grant agreement No. GA212643 (THOR: Thermohaline circulation – at risk?, 2008–2012). KL also received funding through the Cluster of Excellence “CLISAP”, funded by the German Science Foundation (DFG). DM was supported by the Federal Ministry for Education and Research (BMBF) NORTH ATLANTIC and RACE projects. The authors wish to thank Michael Botzet and Helmuth Haak for their help with setting up the sensitivity experiments, Manfred Bersch for his help with the definition of the subpolar deep water formation index and Xihua Zhu for a critical review of the manuscript prior to submission. The MPI-ESM-CR and MPI-AO-LR simulations were conducted at the German Climate Computing Center (DKRZ). This study is a contribution to the Centre for Climate Dynamics at the Bjerknes Centre.

The service charges for this open access publication have been covered by the Max Planck Society.

OSD

10, 1895–1931, 2013

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

References

- Ba, J., Keenlyside, N., Park, W., Latif, M., Hawkins, E., and Ding, H.: A mechanism for Atlantic multidecadal variability in the Kiel Climate Model, *Clim. Dynam.*, 41, 2133–2144, doi:10.1007/s00382-012-1633-4, 2013.
- 5 Ba, J., Keenlyside, N., Latif, M., Park, W., Ding, H., Lohmann, K., Mignot, J., Menary, M., Otterå, O. H., Wouters, B., Salas, D., Oka, A., Belluci, A., and Volodin, E.: A multi-model comparison for Atlantic multidecadal variability, *Clim. Dynam.*, in review, 2013.
- Bailey, D., Rhines, P., and Häkkinen, S.: Formation and pathways of north atlantic deep water in a coupled ice–ocean model of the arctic-north atlantic oceans, *Clim. Dynam.*, 25, 497–516, 10 2005.
- Bentsen, M., Drange, H., Furevik, T., and Zhou, T.: Simulated variability of the Atlantic meridional overturning circulation, *Clim. Dynam.*, 22, 701–720, 2004.
- Bjastoch, A., Böning, C., Getzlaff, J., Molines, J.-M., and Madec, G.: Causes of interannual–decadal variability in the meridional overturning circulation of the midlatitude North Atlantic Ocean, *J. Climate*, 21, 6599–6615, 2008.
- 15 Böning, C., Scheinert, M., Dengg, J., Bjastoch, A., and Funk, A.: Decadal variability of subpolar gyre transport and its reverberation in the North Atlantic overturning, *Geophys. Res. Lett.*, 33, L21S01, doi:10.1029/2006GL026906, 2006.
- Cunningham, S., Kanzow, T., Rayner, D., Baringe, M. O., Johns, W. E., Marotzke, J., Longworth, H. R., Grant, E. M., Hirschi, J. J.-M., Beal, L. M., Meinen, C. S., and Bryden, H. L.: Temporal variability of the atlantic meridional overturning circulation at 26.5° N, *Science*, 317, 935–938, 2007.
- 20 Danabasoglu, G., Large, W. G., and Briegleb, B. P.: Climate impacts of parameterized Nordic Sea overflows, *J. Geophys. Res.*, 115, C11005, doi:10.1029/2010JC006243, 2010.
- Escudier, R., Mignot, J., and Swingedouw, D.: A 20-yr coupled ocean-sea ice-atmosphere variability mode in the North Atlantic in an AOGCM, *Clim. Dynam.*, 40, 619–636, doi:10.1007/s00382-012-1402-4, 2013.
- 25 Fischer, N.: Holocene and Eemian climate variability, Ph.D. thesis, Berichte zur Erdsystemforschung No. 91, Max Planck Institute for Meteorology, Hamburg, Germany, 106 pp., 2011.
- 30 Furevik, T., Bentsen, M., Drange, H., Kindem, I., Kvamstø, N., and Sorteberg, A.: Description and evaluation of the Bergen climate model: ARPEGE coupled with MICOM, *Clim. Dynam.*, 21, 27–51, 2003.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Gastineau, G. and Frankignoul, C.: Cold-season atmospheric response to the natural variability of the Atlantic meridional overturning circulation, *Clim. Dynam.*, 39, 37–57, 2012.
- Gordon, C., Cooper, C., Senior, C., Banks, H., Gregory, J., Johns, T., Mitchell, J., and Wood, R.: The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments, *Clim. Dynam.*, 16, 147–168, 2000.
- Hall, M. and Bryden, H.: Direct estimates and mechanisms of ocean heat transport, *Deep-Sea Res.*, 29, 339–359, 1982.
- Hansen, B., Østerhus, S., Turrell, B., Jónsson, S., Valdimarsson, H., Hátún, H., and Olsen, S.: The inflow of Atlantic water, heat, and salt to the Nordic Seas across the Greenland–Scotland Ridge, in: *Arctic–Subarctic Ocean Fluxes*, edited by: Dickson, R., Meincke, J., and Rhines, P., Springer Verlag, 15–43, 2008.
- Jungclaus, J. H., Haak, H., Latif, M., and Mikolajewicz, U.: Arctic–North Atlantic interactions and multidecadal variability of the meridional overturning circulation, *J. Climate*, 18, 4013–4031, 2005.
- Jungclaus, J. H., Keenlyside, N., Botzet, M., Haak, H., Luo, J.-J., Latif, M., Marotzke, J., Mikolajewicz, U., and Roeckner, E.: Ocean circulation and tropical variability in the coupled model ECHAM5/MPI-OM, *J. Climate*, 19, 3952–3972, 2006.
- Jungclaus, J. H., Lorenz, S. J., Timmreck, C., Reick, C. H., Brovkin, V., Six, K., Segschneider, J., Giorgetta, M. A., Crowley, T. J., Pongratz, J., Krivova, N. A., Vieira, L. E., Solanki, S. K., Klocke, D., Botzet, M., Esch, M., Gayler, V., Haak, H., Raddatz, T. J., Roeckner, E., Schnur, R., Widmann, H., Claussen, M., Stevens, B., and Marotzke, J.: Climate and carbon-cycle variability over the last millennium, *Clim. Past*, 6, 723–737, doi:10.5194/cp-6-723-2010, 2010.
- Kanzow, T., Cunningham, S. A., Johns, W. E., Hirschi, J. J.-M., Marotzke, J., Baringer, M. O., Meinen, C. S., Chidichimo, M. P., Atkinson, C., Beal, L. M., Bryden, H. L., and Collins, J.: Seasonal variability of the Atlantic meridional overturning circulation at 26.5°N, *J. Climate*, 23, 5678–5698, 2010.
- Köhl, A.: Anomalies of meridional overturning: mechanisms in the North Atlantic, *J. Phys. Oceanogr.*, 35, 1455–1472, 2005.
- Köhl, A. and Stammer, D.: Variability of the meridional overturning in the North Atlantic from the 50-year GECCO state estimate, *J. Phys. Oceanogr.*, 38, 1913–1930, 2008.
- Langehaug, H. R., Medhaug, I., Eldevik, T., and Ottera, O. H.: Arctic/Atlantic exchanges via the subpolar gyre, *J. Climate*, 25, 2421–2439, 2012a.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Langehaug, H. R., Rhines, P. B., Eldevik, T., Mignot, J., and Lohmann, K.: Water mass transformation and the North Atlantic Current in three multicentury climate model simulations, *J. Geophys. Res.*, 117, C11001, doi:10.1029/2012JC008021, 2012b.
- Latif, M. et al.: Reconstructing, monitoring and predicting decadal-scale changes in the North Atlantic thermohaline circulation with sea surface temperature, *J. Climate*, 17, 1605–1614, 2004.
- Latif, M., Böning, C., Willebrand, J., Biastoch, A., Dengg, J., Keenlyside, N., Schweckendiek, U., and Madec, G.: Is the thermohaline circulation changing?, *J. Climate*, 19, 4631–4637, 2006.
- Lohmann, K. et al.: Using simulations of the last millennium to understand climate variability seen in paleo-observations: similar variation of Iceland–Scotland overflow strength and Atlantic Multidecadal Oscillation, *Clim. Past.*, accepted, 2013.
- Marti, O. et al.: Key features of the IPSL ocean atmosphere model and its sensitivity to atmospheric resolution, *Clim. Dynam.*, 34, 1–26, 2010.
- Medhaug, I., Langehaug, H. R., Eldevik, T., Furevik, T., and Bentsen, M.: Mechanisms for decadal scale variability in a simulated Atlantic meridional overturning circulation, *Clim. Dynam.*, 39, 77–93, 2012.
- Menary, M., Park, W., Lohmann, K., Vellinga, M., Palmer, M., Latif, M., and Jungclauss, J.: A multimodel comparison of centennial Atlantic meridional overturning circulation variability, *Clim. Dynam.*, 38, 2377–2388, 2012.
- Msadek, R. and Frankignoul, C.: Atlantic multidecadal oceanic variability and its influence on the atmosphere in a climate model, *Clim. Dynam.*, 33, 45–62, 2009.
- Olsen, S. M., Hansen, B., Quadfasel, D., and Østerhus, S.: Observed and modeled stability of overflow across the Greenland–Scotland ridge, *Nature*, 455, 519–523, 2008.
- Østerhus, S., Sherwin, T., Quadfasel, D., and Hansen, B.: The overflow transport east of Iceland, in: *Arctic–Subarctic Ocean Fluxes*, edited by: Dickson, R., Meincke, J., and Rhines, P., Springer Verlag, 427–441, 2008.
- Otterå, O. H., Bentsen, M., Bethke, I., and Kvamstø, N. G.: Simulated pre-industrial climate in Bergen Climate Model (version 2): model description and large-scale circulation features, *Geosci. Model Dev.*, 2, 197–212, doi:10.5194/gmd-2-197-2009, 2009.
- Park, W. and Latif, M.: Multidecadal and multicentennial variability of the meridional overturning circulation, *Geophys. Res. Lett.*, 35, L22703, doi:10.1029/2008GL035779, 2008.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Park, W., Keenlyside, N., Latif, M., Ströh, A., Redler, R., Röckner, E., and Madec, G.: Tropical Pacific climate and its response to global warming in the Kiel Climate Model, *J. Climate*, 22, 71–92, 2009.
- Pope, V., Gallani, M., Rowntree, P., and Stratton, R.: The impact of new physical parametrizations in the Hadley Centre climate model – HadCM3, *Clim. Dynam.*, 16, 123–146, 2000.
- Serra, N., Käse, R., Köhl, A., Stammer, D., and Quadfasel, D.: On the low-frequency phase relation between the Denmark Strait and the Faroe-Bank Channel overflows, *Tellus A*, 62, 530–550, 2010.
- Servonnat, J., Yiou, P., Khodri, M., Swingedouw, D., and Denvil, S.: Influence of solar variability, CO₂ and orbital forcing between 1000 and 1850 AD in the IPSLCM4 model, *Clim. Past*, 6, 445–460, doi:10.5194/cp-6-445-2010, 2010.
- Swingedouw, D., Rodehacke, C., Behrens, E., Menary, M., Olsen, S., Gao, Y., Mikolajewicz, U., Mignot, J., and Biastoch, A.: Decadal fingerprints of fresh water discharge around Greenland in a multi-model ensemble, *Clim. Dynam.*, 41, 695–720, doi:10.1007/s00382-012-1479-9, 2013.
- Tanhua, T., Olsson, K., and Jeansson, E.: Tracer evidence of the origin and variability of Denmark Strait Overflow Water, in: *Arctic–Subarctic Ocean Fluxes*, edited by: Dickson, R., Meincke, J., and Rhines, P., Springer Verlag, 475–503, 2008.
- Våge, K., Pickart, R., Spall, M., Moore, G., Valdimarsson, H., Torres, D., Erofeeva, S., and Nilsen, J. E.: Revised circulation scheme north of the Denmark Strait, *Deep-Sea Res. Pt. I*, 79, 20–39, 2013.
- Willebrand, J. et al.: Circulation characteristics in three eddy-permitting models of the North Atlantic, *Prog. Oceanogr.*, 48, 123–161, 2001.
- Yeager, S. and Danabasoglu, G.: Sensitivity of Atlantic meridional overturning circulation variability to parameterized Nordic Sea overflows in CCSM4, *J. Climate*, 25, 2077–2103, 2012.
- Zanchettin, D., Rubino, A., and Jungclaus, J. H.: Intermittent multidecadal-to-centennial fluctuations dominate global temperature evolution over the last millennium, *Geophys. Res. Lett.*, 37, L14702, doi:10.1029/2010GL043717, 2010.
- Zhu, X. and Jungclaus, J. H.: Interdecadal variability of the meridional overturning circulation as an ocean internal mode, *Clim. Dynam.*, 31, 731–741, 2008.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Table 1. Overview of the coupled climate models used in this study.

Model	Ocean component	Atmosphere component	Ocean grid	Resolution in North Atlantic	Atmosphere grid	Ocean levels	Atmosphere levels	Length of control simulation
MPI-ESM-CR	MPI-OM	ECHAM5	conformal mapping, 3°, northern pole over Greenland	50–200 km	spectral, T31 ^a	40 z-levels	19	1000 yr
MPI-AO-LR	MPI-OM	ECHAM5	conformal mapping, 1.5°, northern pole over Greenland	25–100 km	spectral, T63 ^a	40 z-levels	31	500 yr
KCM	NEMO ^b	ECHAM5	conformal mapping, 2°, tripolar (ORCA2 grid)	100–200 km	spectral, T31 ^a	31 z-levels	19	1000 yr
IPSLCM4	OPA8.2	LMDz4	conformal mapping, 2°, tripolar (ORCA2 grid)	100–200 km	3.75° (lon) × 2.535° (lat)	31 z-levels	19	1000 yr
HadCM3	No specific name	HadAM3	1.25° × 1.25°	~ 100 km	3.25° (lon) × 2.5° (lat)	20 z-levels	19	1000 yr
BCM	MICOM	ARPEGE	conformal mapping, 2.4°, northern pole over Siberia	150–200 km	spectral, T63 ^a	34 isopycnic layers, non-isopycnic surface mixed layer	31	700 yr

^aCorresponding to about 3.75° (1.875°) on a Gaussian grid for T31 (T63).

^bNEMO (Nucleus for European Modelling of the Ocean) is based on the OPA model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Table 2. Mean value of the overflow transport through Denmark Strait (DSO) and across the Iceland–Scotland–Ridge (ISRO) in the different control simulations.

Model	Mean DSO	Mean ISRO
MPI-ESM-CR	2.8 Sv	2.3 Sv
MPI-AO-LR	2.8 Sv	3.5 Sv
KCM	12.7 Sv	2.3 Sv
IPSLCM4	10.8 Sv	2.4 Sv
HadCM3	8.0 Sv	3.4 Sv
BCM	1.9 Sv	3.8 Sv

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Role of deep water formation in multidecadal variability

K. Lohmann et al.

Table 3. Summary of the maximum variance of the bandpass filtered (15 to 80 yr) Atlantic meridional overturning streamfunction at 45° N and 30° N explained by subpolar deep water formation (SDWI) and Denmark Strait overflow (DSO) in the different control simulations, deduced from Figs. 3 and 4 (correlation coefficient²).

Model	SDWI – AMOC 45°N	DSO – AMOC 45°N	SDWI – AMOC 30°N	DSO – AMOC 30°N
MPI-ESM-CR	72 %	49 %	49 %	25 %
MPI-AO-LR	81 %	72 %	64 %	36 %
KCM	72 %	72 %	42 %	49 %
IPSLCM4	81 %	72 %	56 %	49 %
HadCM3	30 %	56 %	25 %	20 %
BCM	72 %	20 %	64 %	not significant
Multi-model average	68 %	57 %	50 %	30 %

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Role of deep water formation in multidecadal variability

K. Lohmann et al.




Fig. 1. Mean Atlantic meridional overturning streamfunction (in Sv) in the control simulation with **(a)** MPI-ESM-CR, **(b)** MPI-AO-LR, **(c)** KCM, **(d)** IPSLCM4, **(e)** HadCM3 and **(f)** BCM. Solid (dashed) lines indicate positive (negative) values. Contour interval is 2 Sv, starting from ± 1 Sv. An additional contour line is added to show the respective location of the overall maximum strength. The respective value of the overall maximum strength is given in the lower right corner.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Role of deep water formation in multidecadal variability

K. Lohmann et al.





Fig. 2. Mean mixed layer depth in Mar (in meter) in the control simulation with (a) MPI-ESM-CR, (b) MPI-AO-LR, (c) KCM, (d) IPSLCM4, (e) HadCM3 and (f) BCM. The boxes in (a) indicate the area, where in the sensitivity experiments temperature and salinity are relaxed towards climatological values.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Role of deep water formation in multidecadal variability

K. Lohmann et al.

**Fig. 3.** Caption on next page.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Fig. 3. Lag correlation coefficients between the bandpass filtered (15 to 80 yr) subpolar deep water formation index (SDWI) and maximum Atlantic meridional overturning streamfunction at different latitudes in the North Atlantic in the control simulation with **(a)** MPI-ESM-CR (depth level 1020 m), **(b)** MPI-AO-LR (depth level 1020 m), **(c)** KCM (depth level 870 m), **(d)** IPSLCM4 (depth level 1210 m), **(e)** HadCM3 (depth level 800 m) and **(f)** BCM (depth level 1000 m). Only correlation coefficients significant at the 95 % level for a two-tailed t test (taking into account both length and decorrelation time scale of the SDWI and the overall maximum strength of the Atlantic meridional overturning streamfunction) are plotted (significance level: 0.31 for MPI-AO-LR, 0.27 for BCM and 0.23 for the other models). Positive (negative) lags indicate that the SDWI is leading (lagging).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Role of deep water formation in multidecadal variability

K. Lohmann et al.




Fig. 4. Same as Fig. 3, but for the correlation between Denmark Strait overflow transport (DSO) and Atlantic meridional overturning streamfunction.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Role of deep water formation in multidecadal variability

K. Lohmann et al.




Fig. 5. Same as Fig. 3, but for the correlation between overflow transport across the Iceland–Scotland–Ridge (ISRO) and Atlantic meridional overturning streamfunction.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Role of deep water formation in multidecadal variability

K. Lohmann et al.





Fig. 6. Standard deviation of the bandpass filtered (15 to 80 yr) Atlantic meridional overturning streamfunction (in Sv) in the control simulation with **(a)** MPI-ESM-CR, **(b)** MPI-AO-LR, **(c)** KCM, **(d)** IPSLCM4, **(e)** HadCM3 and **(f)** BCM. Contour interval is 0.1 Sv, starting from 0.1 Sv.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



	Control simulation	SDW experiment	Overflow experiment
SDWI	60 meter	11 meter (82 %)	37 meter (38 %)
DSO	0.45 Sv	0.42 Sv (7 %)	0.05 Sv (89 %)
ISRO	0.21 Sv	0.15 Sv (29 %)	0.02 Sv (90 %)

Fig. 7. Anomalous bandpass filtered (15 to 80 yr) **(a)** subpolar deep water formation index (SDWI, in meter), **(b)** Denmark Strait overflow transport (DSO, in Sv) and **(c)** overflow transport across the Iceland–Scotland–Ridge (ISRO, in Sv) in the control simulation (black lines), the SDW experiment (red lines) and the overflow experiment (green lines) with MPI-ESM-CR. The respective standard deviations are given in the table. Numbers in brackets in the table give the reduction (in %) in the standard deviation in the respective sensitivity experiment compared to the control simulation.

Role of deep water formation in multidecadal variability

K. Lohmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 8. Anomalous bandpass filtered (15 to 80 yr) maximum Atlantic meridional overturning streamfunction (in Sv) at different latitudes in the North Atlantic (depth level 1020 m) in **(a)** the control simulation, **(b)** the SDW experiment and **(c)** the overflow experiment with MPI-ESM-CR.




Fig. 9. Change (in %, for definition see text) in the standard deviation of the bandpass filtered (15 to 80 yr) Atlantic meridional overturning streamfunction in **(a)** the SDW experiment and **(b)** the overflow experiment compared to the control simulation with MPI-ESM-CR. Note that positive (negative) values correspond to a reduction (increase) in the standard deviation in the sensitivity experiments. The colour scale is chosen to match the explained variance, i.e. (correlation coefficient)², in Figs. 3 to 5. The depth level presented in Figs. 3 to 5 is indicated by the solid line.