# Design and validation of MEDRYS, a Mediterranean Sea

# 2 reanalysis over 1992-2013

3

M. Hamon<sup>1</sup>, J. Beuvier<sup>1,2</sup>, S. Somot<sup>2</sup>, J.M. Lellouche<sup>1</sup>, E. Greiner<sup>3</sup>, G. Jordà<sup>4</sup>,
M.N. Bouin<sup>2</sup>, T. Arsouze<sup>5</sup>, K. Béranger<sup>6</sup>, F. Sevault<sup>2</sup>, C. Dubois<sup>1</sup>, M. Drevillon<sup>1</sup>
and Y. Drillet<sup>1</sup>.

7 [1]{Mercator Océan, 10 rue Hermès, 31520 Ramonville-Saint-Agne, France}

8 [2]{Météo France, 42 av. Gaspard Coriolis, 31057 Toulouse cedex, France}

9 [3] {CLS, 11 rue Hermès, 31520 Ramonville-Saint-Agne, France}

10 [4] {Department of Ecology and Marine Resources, IMEDEA (CSIC-UIB), Institut

11 Mediterrani d'Estudis Avançats, Esporles (Illes Balears), Spain}

12 [5] {LMD, école Polytechnique, 91128 Palaiseau cedex, France}

13 [6] {LTHE, rue de la Piscine, 38400 Saint-Martin d'Hère, France}

14 Correspondence to: J. Beuvier (jonathan.beuvier@mercator-ocean.fr)

15

#### 16 Abstract

17

18 The French research community on the Mediterranean Sea modelling and the French 19 operational ocean forecasting center Mercator Océan have gathered their skill and expertise in physical oceanography, ocean modelling, atmospheric forcings and data assimilation, to carry 20 21 out a MEDiterranean sea ReanalYsiS (MEDRYS) at high resolution for the period 1992-22 2013. The ocean model used is NEMOMED12, a Mediterranean configuration of NEMO with a 1/12° (~ 7 km) horizontal resolution and 75 vertical z-levels with partial steps. At the 23 24 surface, it is forced by a new atmospheric forcing dataset (ALDERA), coming from a 25 dynamical downscaling of the ERA-Interim atmospheric reanalysis by the regional climate 26 model ALADIN-Climate with a 12-km horizontal and 3-hour temporal resolutions. This 27 configuration is used to carry a 34-year free simulationhindcast simulation over the period 1979-2013 (NM12-FREE) which is the initial state of the reanalysis in October 1992. The 28

version of MEDRYS uses the existing Mercator Océan data assimilation system SAM2 1 first that is based on a reduced-order Kalman filter with a 3D multivariate modal decomposition of 2 the forecast error. Altimeter data, satellite SST and temperature and salinity vertical profiles 3 are jointly assimilated. This paper describes the configuration we used to perform the 4 MEDRYS-simulation. We then first validate the skills of the data assimilation system. It is 5 shown that the data assimilation restores a good averaged temperature and salinity in-at 6 7 intermediate layers compared to the free simulationhindcast. No particular biases are 8 identified in the bottom layers. However, the reanalysis show slight positive biases of 0.02 9 psu and 0.15°C above 150m depth. In the validation stage, it is also shown that the 10 assimilation allows to better reproduce water, heat and salt transports through the Strait of 11 Gibraltar. Finally, the ability of the reanalysis to represent the sea surface high frequency 12 variability is pointed out.

#### 13

#### 14 1. Introduction

15

16 The Mediterranean Sea is a semi-enclosed sea located between 5.5°W and 36°E and 17 between 30°N and 46°N. It is connected to the Atlantic Ocean through the Strait of Gibraltar 18 and to the Black Sea through the Dardanelles and the Bosphorus Straits. The surrounding 19 orography tends to generate cold and dry regional northern winds over the Mediterranean Sea. 20 This leads to strong heat and freshwater losses by evaporation and latent heat transfer. The 21 heat loss is estimated around 5 W/m2 (MacDonald et al., 1994) while the freshwater loss is 22 about 0.6 m/yr (Mariotti et al., 2008). The main part of the heat and water atmospheric losses 23 are balanced by warm Atlantic Waters (AW) entering through the Strait of Gibraltar while it is estimated that only about 10% of the net water flux is balanced with river runoff (Struglia 24 25 et al., 2004).

26

In a climate change context, the Mediterranean area is considered as a hot spot and shows an increase in the temperature and precipitation interannual variability, and a strong warming and drying (Giorgi et al., 2006). The vulnerability of the population is likely to increase with a higher probability of occurrence of events leading to floods and droughts, which are among the most devastating natural hazards. In this context, it is necessary to simulate the water

cycle over the Mediterranean basin (Drobinski et al., 2013) and to understand how it will 1 2 impact water resources. We must improve our understanding of the variability of the water 3 cycle, from extreme events to the seasonal and interannual scales. In addition to the socio-4 economic motivations and from a strictly physical point of view, the specific configuration of 5 the basin also permits the study of a wide variety of dynamical oceanic processes. For example, the Mediterranean Sea has been found to have a dominant mesoscale circulation 6 7 component (Robinson et al., 1987; Ayoub et al., 1998; Hamad et al., 2005; Fernandez et al., 8 2005) in addition to a thermohaline circulation similar to the world ocean (Wüst, 1961; 9 Robinson et al., 2001). The Mediterranean eddy field also shows semi-permanent structures 10 (Rhodes and South Adriatic gyres for example) which that define the general circulation in 11 the basin. The time persistency of this small scale field is still an unresolved problem, especially in ocean modelling because for example of approximations and uncertainties on 12 13 nonlinear dynamical balance, atmospheric forcing or the bathymetry (Sorgente et al., 2011; Pinardi et al., 2013). Modeling the different time and spatial scales of this circulation is still 14 challenging because for example of approximations and uncertainties on non-linear dynamical 15 balance, atmospheric forcing or the bathymetry (Sorgente et al., 2011; Pinardi et al., 2013). 16

17

18 The ocean reanalysis is a reconstruction technique that allows the production of a consistent 19 four-dimensional estimate of a physical field from observations and numerical modeling 20 simulation. Observations are used to constrain the model trajectory to be near as close as 21 possible from to the "real" state of the ocean. Ocean reanalyses are thus reference products 22 which help to improve our knowledge of the ocean variability at various space and time 23 scales. Several techniques have been used in the past to produce large-scale reanalysis but 24 regional reanalysis are challenging because observational datasets are scarcer and the use of 25 high resolution model requires to adequately represent fluxes through the air/sea interface. 26 This is even more important in the Mediterranean Sea due to the complex orography.-for Mediterranean Sea models in a region surrounded by a various and complex topography. 27 Many small-size islands and a particularly complex coastline limit the low-level air flow, 28 29 channeling potentially strong and recurring regional winds (Mistral, Tramontane, Bora, 30 Etesian, Sirocco; Herrmann et al., 2011). The role of the spatial resolution of the forcing has 31 been highlighted as a key aspect of the representation of Mediterranean Sea phenomena such 32 as local winds (Sotillo et al., 2005; Ruti et al., 2007; Herrmann et al., 2011; Lebeaupin

Brossier et al., 2012), open-sea deep convection (Herrmann and Somot, 2008; Béranger et al., 1 2 2010), shelf-cascading (Dufau-Julliand et al., 2004; Langlais et al., 2009), coastal upwelling 3 (Estournel et al., 2009; Casella et al., 2011), permanent circulation features (Estournel et al., 4 2003; Ourmières et al., 2011) or intermittent eddies (Marullo et al., 2003; Ciappa, 2009; 5 Rubio et al., 2009). The infra-diurnal temporal resolution of the forcing has also been identified as necessary to represent key phenomena such as large salinity anomalies following 6 7 intense rainfall events (Lebeaupin Brossier et al., 2012) or the SST diurnal cycle (Lebeaupin 8 Brossier et al., 2011, 2014). Other studies demonstrated the importance of the good 9 representation of the atmospheric synoptic chronology linked with the so-called weather 10 patterns or weather regimes (Josey et al., 2011 ; Papadopoulos et al., 2012 ; Durrieu de 11 Madron et al., 2013) or with the passage of Mediterranean storms associated with strong airsea exchanges (Herrmann and Somot, 2008; Herrmann et al., 2010). At a longer time scale, 12 13 interannual to decadal variability of the atmospheric forcings (water or heat fluxes) is known 14 to dominate the climate variability of the deep water mass formation in both basins of the Mediterranean Sea (Beuvier et al. 2010; Herrmann et al. 2010; L'Heveder et al., 2013) leading 15 16 sometimes to exceptionnal decadal events such as the Eastern Mediterranean Transient 17 (Roether et al., 2007) or the Western Mediterranean Transition (Schroeder et al., 2008).

18

19 The first regional Mediterranean reanalyses, initiated by Oddo et al. (2009), have been 20 recently produced over the 1985-2007 period by Adani et al. (2011), using a reduced-order 21 optimal interpolation and a three-dimensional variational scheme. Their OPA ocean model 22 (Madec et al., 1997) on a 1/16° regular horizontal grid (Tonani et al., 2008) is forced by daily 23 atmospheric fields from the European Center Medium-Range Weather Forecast (ECMWF) 24 with bulk parameterizations and a monthly precipitation climatology. They used the 25 reanalysis ERA-15 for the 1985-1992 period and then the operational analyses for the 1993-26 2007 period. We note thus several successive changes in the atmospheric forcing, in particular 27 during the 1993-2007 period, for which the resolution of the ECMWF analyses has 28 progressively increased in several steps from about 100km to 25km., Such changes suggesting 29 then that temporal continuity and coherence in atmospheric forcing were are not guaranteed. 30 However, Tthe first results of these reanalyses pointed out for example that such products 31 allow to better simulate the AW salinity inflow, the sea surface height variability, and current-32 jet pathways.

2	In the same way of these previous studies and in order to enhance the diversity of the
3	Mediterranean Sea reanalyses, we present in this study another reanalysis of the
4	Mediterranean circulation, MEDRYS, performed with different tools and covering the
5	altimetry 1992-2013 period. Our ocean model used is NEMOMED12 (Beuvier et al., 2012a),
6	a Mediterranean configuration of NEMO (Madec and the NEMO team, 2008; an update
7	version of the OPA code) with the ORCA standard NEMO grid, giving a close horizontal
8	resolution of NEMOMED16. Our ocean model used is NEMOMED12 (Beuvier et al., 2012a),
9	a Mediterranean configuration of NEMO (Madec and the NEMO team, 2008; an update
10	version of the OPA code) with the ORCA12 standard grid. The ORCA12 grid shows a
11	varying resolution around 1/12° over the world ocean. Within our numerical domain, the
12	ORCA grid has a horizontal resolution ranging between 6 and 7.5km. Note that this spatial
13	resolution is similar is to the 1/16° regular horizontal grid used in Adani et al. (2011). This
14	MEDRYS reanalysis differs also by the use of a reduced-order Kalman filter in the
15	assimilation scheme from the French operational oceanography center Mercator Océan and
16	the long-term 12-km high-resolution fields of the atmospheric forcing called ALDERA. We
17	pay a special attention to the temporal homogeneity of the atmospheric forcing (same
18	resolution, same model physics) in order to allow robust studies of the interannual to decadal
19	variability of the Mediterranean circulation and trends. Even if we cannot overcome other
20	homogeneity issues resulting from the coverage of the observing network (applying in both
21	MEDRYS and ALDERA), we pay a special attention to the consistency of the atmospheric
22	forcing (same resolution, same model physics) in order to reduce as most as possible the
23	sources of inhomogeneity in MEDRYS. This reanalysis then contributes to better describe the
24	interannual to decadal variability of the Mediterranean circulation and trends

1

In the current paper, Wwe first present the configuration of the reanalysis MEDRYS and the twin hindcast NM12-FREE in section 2, with the experimental set up, detailing the specifications of the regional configuration of the ocean model, the ALDERA atmospheric forcing and the assimilated dataset. Then, section 3 presents validation diagnostics and some scientific assessments. Finally, a-discussions and conclusion are is conducted in section 4.on main results and on improvements to be included in further versions of the reanalysis.

#### 2. Experimental set up

2

1

Two twin simulations have been produced: MEDRYS, a Mediterranean reanalysis
covering the 1992-2013 period with data assimilation and its associated free run NM12FREE, a 34-year hindcast simulation covering the 1979-2013 period without assimilation.
Both simulations use the same ocean model configuration, NEMOMED12, described in
sections 2.1 and the high resolution atmospheric forcing ALDERA, presented in section 2.3.
Specific set up concerning data assimilation in the reanalysis are then presented in sections
2.4 and 2.5.

#### 10

11

#### 2.1 Ocean model configuration : NM12-FREE NEMOMED12

12

13 We use the ocean general circulation model NEMO (Madec and the NEMO team, 2008) 14 in a regional configuration of the Mediterranean Sea called NEMOMED12 (Lebeaupin 15 Brossier et al., 2011, 2012, Beuvier et al., 2012a and 2012b ; hereafter NM12). The 16 development of NEMOMED12 is made in the continuity of the evolution of the French 17 modeling of the Mediterranean Sea, following OPAMED16 (Beranger et al., 2005), OPAMED8 (Somot et al., 2006) and NEMOMED8 (Beuvier et al., 2010). We describe the 18 19 NM12 FREE configuration used for the reference 34 year simulation covering the 1979 2013 20 period and performed with NEMOMED12 without assimilation. More details concerning the 21 physical parametrizations and the boundary conditions in the ocean model-NM12 are can be 22 found in Beuvier et al. (2012a).

23

The NM12 configuration covers the whole Mediterranean Sea and a buffer zone including a 24 part of the Atlantic basin, but not the Black Sea. The horizontal resolution is  $1/12^{\circ}$  and 25 26 corresponds to a varying grid cell size between 6 and 7.5km (the distance between two points 27 varying with the cosine of the latitude). NM12 has 75 vertical stretched z-levels (from  $\Delta z$ =1m at the surface to  $\Delta z$  = 135m at the bottom, with 43 levels in the first 1000m) in a partial 28 29 step configuration. The bottom layer thickness is varying to fit the bathymetry (Mercator-30 LEGOS version 10 bathymetry at 1/120° resolution). The no-slip boundary condition is used 31 and the conservation of the model volume is assumed. The mean tidal effect of the quadratic

1	bottom friction formulation computed from a tidal model (Lyard et al., 2006) has been taken
2	into account leading to significant additional bottom friction in the Strait of Gibraltar,
3	Channel of Sicily, Gulf of Gabes and the northern Adriatic sub-basin. As a lateral boundary
4	conditions and in order to represent the exchanges with the Atlantic ocean, a buffer zone is
5	used: from 11° to 7.5°W, 3D temperature and salinity, as well as the Sea Surface Height
6	(SSH) fields are relaxed toward ORAS4 global ocean reanalysis monthly fields (Balmaseda et
7	al. 2013), produced by the European Centre for Medium Range Weather Forecast (ECMWF).
8	For temperature and salinity, the restoring term in the buffer zone is weak west of Cadiz and
9	Gibraltar areas and increases westwards. As the Mediterranean Sea is an evaporation basin,
10	the model volume is conserved through the damping of the SSH in the buffer zone toward
11	prescribed SSH anomalies with a very strong restoring. The SSH from ORAS4 is set in the
12	Atlantic according to a strong damping with a very small characteristic time-scale ( $\tau = 2$ s).

14 We use the climatological averages of the interannual dataset of Ludwig et al. (2009) to 15 compute monthly runoff values, split in two parts (Beuvier et al., 2012a). The 33 main rivers of the NM12 domain are added as precipitation at mouth points (29 in the Mediterranean Sea 16 17 and 4 in the buffer zone). As the Ludwig et al. (2009) dataset consists in 239 mouth points, 18 the inputs of the 210 other rivers in the Mediterranean basin are gathered as a coastal runoff in 19 each subbasin (following the same dividing as in Ludwig et al. 2009). Until 2000, we use the 20 interannual values from Ludwig et al. (2009) and then the climatological average representing 21 the 1960-2000 period. The Black Sea, not included NM12, is taken into account with a 22 monthly average one layer net flow across the Marmara Sea and the Dardanelles Strait. We 23 assume that the flow is a freshwater flux (Beuvier et al., 2012a). Until 1997, we use the 24 interannual values from Stanev et Peneva (2002) and then the climatological average 25 representing the 1960-1997 period.

13

26

The free simulation NM12-FREE starts in October 1979 and ends in June 2013. Initial
conditions are provided by the monthly mean potential temperature and salinity 3D fields
from the state of the MEDATLAS 1979 climatology (Rixen et al., 2005) in the Mediterranean
side and from ORAS4 monthly anomalies (Balmaseda et al., 2013) in the Atlantic side. The
exchanges with the Atlantic basin are performed through a buffer zone. From 11°W to 7.5°W,
3D temperature and salinity are relaxed toward ORAS4 fields. The restoring term is weak

1 west of Cadiz and Gibraltar areas and increases westwards. As the Mediterranean Sea is an 2 evaporation basin, the model volume is conserved through a damping of the Sea Surface 3 Height (SSH) in the buffer zone toward prescribed SSH anomalies with a very strong 4 restoring. The SSH from ORAS4 is set in the Atlantic according to a strong damping with a 5 very small characteristic time scale ( $\tau = 2$  s).

6 7

8 9

10

11 12 We use the climatological averages of the interannual dataset of Ludwig et al. (2009) to compute monthly runoff values, split in two parts (Beuvier et al., 2012a). Runoffs from the 33 main rivers are added as precipitation at mouth points whereas the averaged values of the inputs of the other rivers are gathered in each subbasin and put as a coastal runoff. The Black Sea is not included in the NM12 FREE configuration but taken into account with a monthly average one layer net flow across the Marmara Sea and the Dardanelles Strait. We assume that the flow is a freshwater flux (Beuvier et al., 2012a).

# 14 15

16

13

#### 2.2 Simulations: NM12-FREE and MEDRYS

17 The hindcast NM12-FREE starts in October 1979 and ends in June 2013. In the 18 Mediterranean side, initial conditions are provided by a monthly mean potential temperature 19 and salinity 3-D fields based on the MedAtlas interannual dataset (Rixen et al., 2005). A field representing the state of the Mediterranean Sea in October 1979 has been produced 20 21 combining the MedAtlas monthly climatology (MEDAR/MEDATLAS Group, 2002) to the 3year filtered interannual fields from Rixen et al. (2005). Following Rixen et al. (2005), the 22 23 filtered interannual product is used in order to reduce the impact of large spatio-temporal gaps 24 in the data distribution. In the buffer zone, potential temperature and salinity are initialized 25 from ORAS4 global ocean reanalysis fields in order to maintain consistency with the relaxation. In the initial condition fields, a linear transition between 7.5°W and 6°W is applied 26 27 between the ORAS4 and the MedAtlas fields. MEDRYS starts from the state of NM12-FREE 28 in October 1992 and ends in June 2013.

29

30

2.22.3 Atmospheric forcing: ALDERA

2 The most recent long-term hindcast simulations using the NEMOMED12 ocean model (Beuvier et al. 2012B; Soto-Navarro et al. 2014; Palmiéri et al. 2015) were driven by the 3 4 ARPERA2 dataset (Herrmann et al. 2010). This forcing was obtained by a dynamical 5 downscaling using the stretched-grid Regional Climate Model (RCM) ARPEGE-Climate and a spectral nudging technique. ARPERA2 covers the period 1958-2013 with a daily temporal 6 7 resolution and a 50-km spatial resolution over the Mediterranean Sea. It may include temporal 8 inhomogeneities in the 1970s, a period known for ERA 40 humidity deficiency and in 2001 9 when the large scale driving fields changes from ERA 40 to ECMWF analysis. It may include 10 temporal inhomogeneity especially in 2001 when the large-scale driving fields changes from 11 ERA-40 to ECMWF analysis.

12

1

13 In order to overcome the main deficiencies of the ARPERA2 dataset (relatively coarse spatial and temporal resolution, temporal homogeneity issue), we have developed a new forcing 14 dataset for the MEDRYS reanalysis. This dataset called ALDERA is based on a dynamical 15 16 downscaling of the ERA Interim reanalysis (Dee et al., 2011) over the period 1979 2013 by 17 the RCM ALADIN Climate (Radu et al., 2008; Colin et al., 2010; Herrmann et al., 2011). We use here the version 5 of ALADIN-Climate firstly described in Colin et al. (2010). In order to 18 19 overcome the main deficiencies of the ARPERA2 dataset (relatively coarse spatial and 20 temporal resolution, temporal homogeneity issue), we are using a new forcing dataset for 21 MEDRYS and NM12-FREE. This dataset (called hereafter ALDERA) is based on a 22 dynamical downscaling of the ERA-Interim reanalysis (Dee et al., 2011) over the period 23 1979-2013 by the RCM ALADIN-Climate (Radu et al., 2008; Colin et al., 2010; Herrmann et 24 al., 2011). The dynamical downscaling technique is commonly used to overcome the lack of 25 atmospheric regional reanalysis over sea and to improve locally the resolution of the air-sea 26 forcing in areas dominated by small-scale atmospheric pattern as the Mediterranean Sea (Sotillo et al. 2005, Herrmann and Somot 2008, Beuvier et al. 2010, Herrmann et al. 2010, 27 28 Herrmann et al. 2011, Josey et al. 2011, Beuvier et al. 2012a, Lebeaupin-Brossier et al. 2012, 29 Solé et al. 2012, Vervatis et al. 2013, Auger et al. 2014, Harzallah et al. 2014). In ALDERA, we use the version 5 of ALADIN-Climate firstly described in Colin et al. (2010). For the 30 31 model definition, we used a Lambert conformal projection for pan-Mediterranean area at the 32 horizontal resolution of 12 km centred at 14°E, 43°N with 432405 grid points in longitude

and 288-261 grid points in latitude including the bi-periodization (11) and the relaxation (2 x 1 8) zones excluding the coupling zone. The model version has 31 vertical levels. The time step 2 3 used is 600 seconds. This geographical set-up allows the Med-CORDEX official area (Ruti et al. 2015 in revision, www.medcordex.eu) to be fully included in the model central zone. In 4 this configuration, the RCM is driven at its lateral boundary conditions-by the ERA-Interim 5 reanalysis 80-km full resolution. Dee al.. 6 (T255, at its et 2011. 7 http://www.ecmwf.int/research/era/do/get/era-interim) which are updated every 6 hours. The 8 ERA-Interim data assimilation system uses a 2006 release of the Integrated Forecasting 9 System, which contains many improvements both in the forecasting model and analysis 10 methodology relative to ERA-40. The period simulated is 1979-2013. Before starting this 11 simulation, a two-year long spin-up is carried out allowing the land water content to reach its 12 equilibrium. Land surface parameters and aerosols concentration are updated every month 13 following a climatological seasonal cycle coming from observations. The sea surface 14 temperatures and the sea ice limit (Black Sea) are updated every month with a seasonal and 15 interannual variability following ERA Interim SST and sea ice analysis. The sea surface 16 temperatures and the sea ice limit are updated every month with a seasonal and interannual 17 variability using the same SST and sea ice analyses as the one used to drive the ERA-Interim 18 reanalysis (Dee et al 2011). As ERA Interim constitutes the best knowledge of the 4D 19 dynamic of the atmosphere available over the last decades, such a simulation is often called "perfect boundary simulation" or "poor man regional reanalysis". As atmospheric reanalyses 20 21 constitutes today the best knowledge of the 4-D dynamic of the atmosphere available over the 22 last decades, such a simulation is often called "perfect-boundary simulation" or "poor-man 23 regional reanalysis".

24

25 ALDERA is available at a 12-km spatial resolution and a 3-hour temporal resolution over the 26 whole Mediterranean Sea, Black Sea and near-Atlantic Ocean. It includes a representation of 27 the effect of the aerosols on the long-wave and short-wave radiations and uses the same bulk formula as in ARPERA2 (Louis, 1979) to compute the turbulent fluxes (sensible heat, latent 28 29 heat and momentum fluxes). To our knowledge, ALDERA is the longest and finest 30 homogeneous atmospheric forcing available for the Mediterranean Sea. All variables required 31 to drive regional ocean models using bulk formula or flux formulation are available. For the 32 MEDRYS-NEMOMED12 configuration (both NM12-FREE and MEDRYS), the various

1	fluxes have been interpolated every 3-hour on the NEMOMED12 grid using a conservative
2	interpolation scheme. NEMOMED12 receives heat fluxes (total and solar for the light
3	penetration), net freshwater fluxes (evaporation and precipitation) and wind stresses every 3
4	hours. A retroaction term towards the same SST fields as the one seen by ALADIN-Climate is
5	added in the heat flux, following the method of Barnier et al. (1995), with a retroaction
6	coefficient of -40 W.m <sup>-2</sup> .K <sup>-1</sup> . The total heat flux, including the retroaction term, has been
7	stored when running the hindcast NM12-FREE and is used to force MEDRYS, ensuring thus
8	that both simulations have exactly the same atmospheric forcing.
9	
1.0	
10	No SSS damping is used but a 2D-smoothed monthly climatological freshwater flux
11	correction is added, following the same method as in Beuvier et al. (2012a), but with the 2D
12	spatial variability kept : these monthly 2D fields have been computed by averaging the SSS
13	relaxation term through a previous companion simulation with NEMOMED12 and the same
14	atmospheric forcing, and then filtered at the resolution of 1° by a spatial averaging. The
15	surface freshwater budget is thus balanced without altering the spatial and temporal variations
16	of the freshwater flux and so of the SSS. This correction term is added to the water fluxes
17	coming from the atmospheric fields and from the rivers and Black Sea runoff. All the
18	ALDERA outputs are openly available through the Med CORDEX database
19	(www.medcordex.eu).
20	
21	Within the frame of the Med-CORDEX initiative, the RCM ALADIN-Climate is also run at
22	lower spatial resolutions (150km, and 50km) with exactly the same setting as ALDERA in
23	order to prove the 12 km added value as illustrated below in Table 1, 2 and Figures 1 and 2.
24	to illustrate the small-scale features of the 12km resolution model with respect to lower
25	resolution models (see later comments for Tables 1 and 2 and Figs. 1 and 2).
26	

# 27 <u>2.2.12.3.1</u> Long-term Mediterranean Sea surface heat and water budgets

28

The various terms of the spatially averaged Mediterranean Sea water and heat surface budget
 are given in Table 1 and 2 (flux are positive downward in W.m-2 and mm.day-1). References

# come from Sevault et al. (2014) for the Mediterranean Sea heat budget terms over the 1985 2004 period and from Sanchez Gomez et al. (2011) for the water budget terms.

Table 1 and 2 compares the spatially and temporally averaged values of the Mediterranean 3 Sea surface heat and water budget terms of the ALDERA forcing with past studies and 4 observed-based references (flux are positive downward in W/m2 and mm/day). ALDERA 5 shows values within the range of the references for the net heat and water surface fluxes, 6 respectively with -3 W.m<sup>-2</sup> over the 1985-2004 period (-4 W.m<sup>-2</sup> over the 1979-2012 period) 7 and -1.69 mm.day<sup>-1</sup> (1979-2011). Over the 20-year period considered, ALDERA shows 8 compensating errors between an overestimated shortwave and an overestimated latent heat 9 loss when compared to the observation-based estimates (Sevault et al. 2014). Both values are 10 11 in equilibrium with the heat and water transports at the Strait of Gibraltar (see section 3.2.6). However some individual terms show biases. This is especially true for the shortwave 12 13 radiation, the latent heat flux (and consequently the evaporation) and the precipitation 14 averaged over the sea surface. Note that ALDERA and ARPERA2 show very similar results, 15 what is expected as they share most of their physical parameterizations. This also means that 16 increasing the spatial resolution in the RCMs does not fundamentally change the mean biases 17 at least from 50km to 12km. This is confirmed when comparing ALDERA to the ALADIN-18 Climate simulation at 50km resolution. ALADIN-Climate ran at 150 km is however closer to 19 ERA-Interim with a weaker latent heat loss. Note that Pettenuzzo et al. (2010) dataset also 20 achieves the Mediterranean sea heat budget balance but with lower values both for the shortwave radiation and the latent heat loss. When compared to the ENSEMBLES RCMs 21 22 used in the last published multi-model intercomparison study with Atmosphere RCM (Sanchez-Gomez et al. 2011), ALDERA always fits inside the uncertainty range. 23

24

## 25

# 26 27

## 2.2.22.3.2 Variability : from the synoptic scale to multi-decadal trendsInterannual variability and trends

- Sanchez-Gomez et al. (2009) proved that RCMs (including ALADIN-Climate in its
  version 4) forced by reanalysis over an extended European domain are able to reproduce very
  well the spatial pattern of the large scale weather regimes of the driving model as well as their
  seasonal and interannual variability. However the day-to-day chronology is less well
  reproduced by the RCMs without spectral nudging. Even if the analysis has not been repeated
  - 12

with the version of ALADIN-Climate used in ALDERA, we are confident that their results are applicable here too as the ALDERA domain is smaller than the ENSEMBLES domain used in their study. Note in addition that the links between the weather pattern and the Mediterranean air sea fluxes are very strong and now better understood (see Josev et al. 2011,

- 5 Papadopoulos et al. 2012, Durrieu de Madron et al. 2013).
- 6

1

2

3

4

7 At the basin scale, the interannual variability of the various terms of the Mediterranean Sea heat budget heat fluxes can also be evaluated overfor the period 1985-2004 of the reference 8 dataset of Table 1 (1985-2004, Sevault et al. 2014). For example, for the basin-averaged net 9 shortwave radiation flux, the interannual standard deviation in ALDERA (1.6 Wm<sup>-2</sup>) is 10 underestimated with respect to ISCCP observations (2.8 W.m<sup>-2</sup>) whereas the interannual 11 12 temporal correlation is equal to 0.84. For the latent heat loss, the 1985-2004 interannual standard deviation is equal to 5.6 W.m<sup>-2</sup> in ALDERA within the range of the observations (4.7 13  $W.m^{-2}$  for NOCS and 6.7  $W.m^{-2}$  for OAFLUX) and the interannual temporal correlation is 14 good (0.83 with NOCS and 0.81 with OAFLUX). Interannual standard deviation are lower for 15 the net longwave radiation flux  $(1.2 \text{ W.m}^{-2} \text{ in ALDERA})$  and for the sensible heat loss  $(1.3 \text{ m}^{-2})$ 16 W.m<sup>-2</sup> in ALDERA) and the various observation-based –estimates disagree (not shown). 17 Concerning the trends in air sea fluxes, ALDERA does not include yet a trend in the 18 19 anthropogenic aerosols and therefore does not reproduce the shortwave trend identified in 20 Nabat et al. (2014). However the SST used to drive the RCM does include the observed trend and leads to a realistic positive trend of the latent heat flux though underestimated with 21 respect to Mariotti (2008). Concerning the surface heat flux terms in ALDERA, only the trend 22 in latent heat flux is significant with an increase of the heat loss by the sea equal to + 4.1 23 W.m-2/decade over the 1979-2012 period. 24

25

Trends in the surface forcing are relevant in long-term simulations as they can induce longterm trends in the water mass characteristics. Concerning the surface heat flux terms in
ALDERA, only the trend in latent heat flux is significant with an increase in the heat loss by
the sea equal to +4.1W/m<sup>2</sup>/decade over the 1979–2012 period. This trend is similar to the one
obtained in Mariotti et al. (2008) and is mostly driven by the SST trends (Sevault et al. 2014).
Note that ALDERA does not include the observed trend in European anthropogenic aerosols
and therefore does not reproduce the shortwave trend identified in Nabat et al. (2014).

#### 2.2.3 Added-value of the 12-km resolution of ALDERA

### 3

4

### 2.3.3 Illustration of the small-scalle features in the ALDERA forcing

- Concerning the sea wind representation and the air sea fluxes, the added value of the model 5 spatial high resolution has been already demonstrated by various studies (Ruti et al. 2007, 6 7 Herrmann and Somot 2008, Béranger et al. 2010, Herrmann et al. 2011, Lebeaupin Brossier 8 et al. 2012) especially when reaching the 50 km resolution. Herrmann et al. (2011) show that 9 the 12-km resolution does not bring automatically added value everywhere, and in particular, in the open ocean, but clearly improves the sea wind representation closer to the coast. Over 10 11 the Mediterranean Sea, the added-value of high-resolution models has been shown in 12 particular concerning the representation of the heat and water budget terms (Elguindi et al. 2011, Josey et al. 2011), of wind field especially close to the coast and islands (Sotillo et al. 13 14 2005, Ruti et al. 2007, Herrmann and Somot 2008, Langlais et al. 2009, Herrmann et al. 2011, 15 Vrac et al. 2012) and of the events of strong air-sea fluxes (Herrmann and Somot 2008, Béranger et al. 2010, Lebeaupin-Brossier et al. 2012). Dynamical downscaling of reanalyses 16 17 have therefore been used to force long-term hindcast simulations (Beuvier et al. 2010, 2012b, 18 Herrmann et al. 2010, Solé et al. 2012, Vervatis et al. 2013, Auger et al. 2014, Harzallah et al. 19 2014). Figure 1 illustrates the role of the atmospheric resolution in the representation of the 20 wind and the latent heat flux on March 14th 2013 in the Gulf of Lions by comparing 21 ALDERA at 12 km with ALADIN-Climate runs at lower-resolution. This particular date has 22 been selected because of the strong Mistral event in the Gulf of Lions. Increasing the 23 resolution allows ALDERA to create small-scale features of the wind near the coast as well as 24 the associated pattern of latent heat flux during the Mistral event. The comparison of latent 25 heat flux at 42°N, 5°E also indicates that the maximum of latent heat flux is resolutiondependent. In ALADIN-12km (the so-called ALDERA), the maximum of latent heat loss is 26 about 900W.m-2 whereas in ALADIN-150km, it barely reaches 500W.-m<sup>-2</sup> with ALADIN-27 28 50km being intermediate.
- 29

Figure 2 also illustrates the resolution dependency of the surface wind field but over the
Eastern Mediterranean basin during a Meltem (or Etesian) event (August 16th 2012). This

case shows the clear shadowing effect of the Greek islands. The wind channeling at 12 km 1 2 leads locally to increased wind speed, changes in wind direction and increased vorticity inputs 3 for the ocean due to strong horizontal gradients. All these effects are visible at the South-4 Eastern part of Crete, an area where the Ierapetra anticyclone is formed regularly (see below). Note that the goal here is not to prove the added value of the 12 km with respect to lower 5 resolution as in-situ observations and regridded would be required for this purpose but to 6 7 illustrate differences between the 3 resolutions (150, 50 and 12 km) and to show ALDERA 8 small-scale features with potential impacts on local to regional Mediterranean Sea circulation.

- 9
- 10

#### 2.32.4 Data assimilation scheme

11

12 The data assimilation system used in MEDRYS is SAM2 (Système d'Assimilation 13 Mercator 2nd version), which is used at Mercator Océan for operational oceanography 14 purposes. The Mercator Océan monitoring and forecasting system has especially 15 demonstrated its skills within the MyOcean project for the global ocean forecast (Lellouche et 16 al., 2013) and we used it in a regional configuration. As the main part of the assimilation 17 scheme used in this paper is already described by Lellouche et al. (2013), we will sum up 18 summarize the assimilation methodology and focus on the specifications inherent to the 19 Mediterranean configuration.

20

21 The SAM2 data assimilation method relies on a reduced-order Kalman filter based on the 22 singular evolutive extended Kalman filter (SEEK) with a 7-day assimilation window 23 (hereafter referred as the assimilation cycle). For each assimilation cycle in MEDRYS, SAM2 24 produces increments of SSH, temperature, salinity and velocity (zonal and meridional 25 components) from the model and the observations, weighted by the forecast error covariance 26 and the specified observation error. Increments are then applied as a tendency term in the 27 model prognostic equations. The forecast error covariance is based on the statistics of a 28 collection of 3D ocean state anomalies. It relies on a fixed basis seasonally variable ensemble of anomalies. For the Mediterranean configuration, we computed about 900 anomaly fields 29 from the NM12 FREE free simulation for a given assimilation cycle. The forecast error 30 covariance is based on the statistics of a collection of 3D ocean state anomalies. For a given 31

cycle centred on the Nth day of a given year, ocean state anomalies computed from NM12-1 2 FREE within the window [N - 60 days; N + 60 days] of each year are gathered and define the 3 covariance of the model forecast error. For the Mediterranean configuration, we computed about 900 anomaly fields from NM12-FREE for a given assimilation cycle. Compared to a 4 5 global configuration, the moderate size of the domain allows us to use such a number of anomaly fields (about 300 in a global configuration) in order to statically compute an accurate 6 7 error covariance field. Moreover, as the analysis increment is a linear combination of the 8 anomalies, a large amount of anomalies is desirable in order to better reproduce span the 9 oceanic variability.

#### 10

In Lellouche et al. (2013), increments of SSH are computed as the sum of barotropic and 11 dynamic height increments (computed from temperature and salinity) in their global 12 configuration. This assumption is only valid far from the coast and in open seas, where the 13 14 local SSH variations due to the remote wind are negligible. In the Mediterranean Sea, strong 15 regional winds occur in areas with low bathymetry and near important straits like Gibraltar 16 and Sicily. A significant part of SSH is then driven non locally by the wind. Shelf surge and 17 hydraulic control effects are typically 10 times larger in the Mediterranean Sea than in the middle of the ocean. Take into account these wind effects in MEDRYS, we no more use the 18 19 barotropic plus baroclinic approximation employed in global sytems. As  $\eta$  is a prognostic 20 variable of the model, the SSH increment is computed straight from the n anomaly modes.

21 In the original formulation of SAM2, SSH increments are analytically computed from 22 temperature and salinity increments through barotropic/dynamic height balances (Lellouche et 23 al., 2013). This assumption is only valid far from the coast and in open seas, where the local 24 SSH variations due to the remote wind are negligible. In the Mediterranean Sea, strong 25 regional winds occur in areas with low bathymetry and near important straits like Gibraltar 26 and Sicily. A significant part of SSH is then driven non locally by the wind. Shelf surge and 27 hydraulic control effects are typically 10 times larger in the Mediterranean Sea than in the 28 middle of the ocean. In our regional configuration, SSH increments are purely statistical and 29 derived by the covariances between SSH (the prognostic variable of the model), temperature 30 and salinity implied by the ensemble of anomalies.

31

32

2.42.5 Observational datasets

2 The assimilated observations in MEDRYS consist in-of Sea Surface Temperature (SST) maps, along track Sea Level Anomaly (SLA) data and in situ temperature and salinity 3 profiles. For each cycle, we assimilate the associated centered SST map coming from the 4 daily NOAA Reynolds 0.25° AVHRR-AMSR product (Reynolds et al., 2007). In MEDRYS, 5 **W**We assimilate SST only each 1 degree to avoid correlation problem between observations. 6 7 Moreover, we noted a negative average bias of 0.2°c between AVHRR-AMSR product and 8 the ERA-Interim reanalysis SST that has been used for fluxes computation. For the sake of 9 consistency between fluxes and assimilated SST in MEDRYS, we decided to add 0.2°C to the 10 AVHRR-AMSR maps as a constant offset.

11

The along track SLA is provided by AVISO (SSALTO/DUACS handbook, 2009) and comes 12 from different satellite datasets. Names and acronyms used in this paper as well as the 13 measurement period of each satellite are summarized in Table 3. Small scales signals are 14 removed by filtering the data and a sub-sampling is applied. The filtering and sub-sampling is 15 16 adapted to each region and product as a function of the characteristics of the area and of the assimilation needs. In the Mediterranean Sea, only one point over two is retained to avoid 17 18 redundant information (Dufau et al., 2013). Along-track SLA delayed-time products, specifically reprocessed for Mediterranean Sea, and distributed by AVISO 19 (http://www.aviso.altimetry.fr) in April 2014 in the framework of MyOcean project, are 20 21 assimilated in MEDRYS. These products include along-track filtering (low pass filtered with 22 a cut-off wavelength of 65km for the whole domain) and along-track sub-sampling (only one 23 point over two is retained to avoid taking into account redundant information). For these 24 products, the reference period of the SLA is based on a 20-year [1993-2012] period. Names 25 and acronyms used in this paper as well as the measurement period of each satellite are 26 summarized in Table 3. The assimilation of SLA observations requires the knowledge of a 27 Mean Dynamic Topography (MDT). The mean surface reference used is a hybrid product 28 between the CNES-CLS09 MDT (Rio et al., 2011) adjusted with GOCE and reanalysis data 29 (Lellouche et al., 2013). The assimilation of SLA observation requires the knowledge the observation error and of a Mean Dynamic Topography (MDT). As the simulated 30 31 Mediterranean Sea has a constant volume in the NM12 configuration, a volume correction 32 term is also needed for the computation of the observation operator in MEDRYS. Concerning

the observation error, we choose to not trust observations near the coastal areas. The 1 2 observation error is then artificially increased within 50km of the whole Mediterranean coast. 3 The mean surface reference used is a hybrid product between the CNES-CLS09 MDT (Rio et al., 2011) adjusted with the data from the Gravity field and steady-state Ocean Circulation 4 5 Explorer (GOCE) and from the Mercator-Ocean 1/4° Reanalysis GLORYS2V1(Lellouche et al., 2013) representing the 1993-2012 period. In MEDRYS, the volume correction consists in 6 7 adding a term in the SLA observation operator, representing the effect of the Glacial Isostatic 8 Adjustment (GIA) and the barystatic effect due to the mass intake of continental ice melting. 9 the effects of the Glacial Isostatic Adjustment (GIA) is taken into account with a correction is 10 about 0.3mm/year. The spatial fluctuations of the GIA are also applied on the MDT to 11 compensate for the local deformation of the geoid due to the ongoing deformation of the solid 12 Earth (Peltier et al., 2008). For the global ocean oin average, the correction is about -13 0.3mm/year. In addition we also apply a correction to compensate the mass intake of 14 continental ice melting in the Mediterranean basin. On average, the mass intake corresponds 15 to a rise of 0.85 mm/year.

16

17 In situ temperature and salinity profiles come from the CORA4 (Cabanes et al., 2013) in situ 18 database provided by CORIOLIS data center from the start of the reanalysis up to December 19 2012. For the last 6 months we used the real-time database. A check through objective quality 20 control and a data thinning have been done on the dataset in CORA4. Indeed, for each 21 instrument, only one profile per day and within a  $0.1^{\circ}$  distance is selected. The best profile is 22 identified thanks to a set of objective criteria on measurement resolution and number of 23 validated measurements flagged as good along the profile. In addition to the quality check 24 done by CORIOLIS, SAM2 carries out a supplementary quality control on in situ 25 observations. In order to minimize the risk of erroneous data being assimilated, the system 26 automatically removes, through different criteria, the data too far from a seasonal climatology 27 (Lellouche et al., 2013). On average over the whole period, 79 observations of temperature 28 per year and 16 observations of salinity per year are rejected by this supplementary quality 29 control performed by SAM2.

30

As for SLA, we choose to not assimilate surface salinity observations near coastal areas. Due
 to how we model the continental freshwater intake along the coast (section 2.1), we apply a

<u>coastal surface mask within which the salinity observations are artificially replace by the</u>
 <u>hindcast value. This concept of pseudo observation near the coast has already been used in</u>
 <u>Lellouche et al. (2013) to overcome the deficiencies of the background error, in particular for</u>
 <u>poorly observed variables.</u>

#### 3. Validation methodology and scientific assessment

#### 3.1 Validation methodology

5

6

7

8

9

10

11 12

13

14

15

During the MyOcean project, scientists have defined validation metrics by region and type of product, including observational products. Many efforts were made to synthesize and homogenise quality information in order to provide quality summaries and accuracy numbers. All these rely on the same basis of metrics that can be divided into four main categories derived from Crosnier and Le Provost (2007).

The consistency between two-system solutions or between a system and observations can be 16 17 checked by "eyeball" verification. This consists in comparing subjectively two instantaneous 18 or time mean spatial maps of a given parameter. Coherent spatial structures or oceanic 19 processes such as main currents, fronts and eddies are evaluated. This process is referred to as 20 CLASS1 metrics. The consistency over time is checked using CLASS2 metrics which include 21 comparisons of moorings time series, and statistics between time series. Space and/or time integrated values such as volume and heat transports, heat content and eddy kinetic energy are 22 23 referred to as CLASS3. Their values are generally compared with literature values or values 24 obtained with past time observations such as climatologies or reanalyses. Finally, CLASS4 metrics give a measure of the real time accuracy of systems, by calculating various statistics 25 26 of the differences between all available oceanic observations (in situ or satellite datasets 27 before data thinning and online quality check) and their model equivalent at the time and 28 location of the observation. The validation procedure thus involves all classes of metrics. It 29 checks improvements between versions of a system, and ensures that a version is robust and 30 its performance stable over time.

2	Firstly, we present assimilation statistics directly coming from SAM2 and then results from
3	both NM12-FREE and MEDRYS (daily outputs for all variables and additional hourly
4	outputs for sea surface variables) are presented. As CLASS1 diagnostic, we thus focus on the
5	impact of the assimilation of SLA data on surface circulation. As CLASS3, the assessment of
6	the interannual variability is made using integrated heat and salt contents. Then a CLASS4
7	diagnostic is made using the entire CORA4 database (without data thinning/quality check)
8	and the high frequency surface variability is presented through a comparison to a fixed
9	mooring in the Gulf of Lions (CLASS2). Even if the assimilation process corrects a part of
10	the distance between the model and the observation, the fluxes play a major role in
11	determining the water masses in the Mediterranean Sea and are thereby a good indicator
12	regarding the quality of an experiment. That is why, as CLASS3, we point out in the last
13	section, the benefit of the assimilation in terms of transport through the Strait of Gibraltar.

15 The validation of MEDRYS has been done in compliance with a recognized 16 methodology used for global and regional configurations performed within Mercator Océan. 17 As the assimilation system gives us a large amount of information, we first of all check some 18 assimilation statistics. Among the information given, we especially show here statistics on 19 "observation minus forecast" (called innovation) for temperature and salinity profiles, SST 20 and SLA. Then a large number of diagnostics derived from Crosnier and Le Provost (2007) 21 have been produced in order to assess the improvements between MEDRYS and the 22 associated free simulation NM12 FREE. In the following sections, results from the final 23 products (daily outputs for all variables and additional hourly outputs for sea surface variables) are then presented. We first focus on the assimilation of SLA data and its impact on 24 surface circulation. Then, the assessment of the interannual variability is made using 25 26 integrated heat and salt content. The high frequency surface variability is assessed through a comparison to a fixed mooring in the Gulf of Lions. Finally, we show the effect of the assimilation in terms of transports through the Strait of Gibraltar.

27 28 29

1

14

30

3.2 Scientific assessment

31

#### 3.2.1 Assimilation Statistics

2

3

4

5

1

We present here assimilation diagnostics to highlight that the reanalysis system is stable and well constrained by the assimilated observations. In this section, the evolution of the mean and the RMS innovation for all SLA, SST and in situ profiles are shown.

6

7 The mean and the RMS of SLA innovation are presented in Figure 3. The mean SLA 8 innovation has a slight linear decrease of 0.65mm/year. This suggests that the volume 9 correction (effect of the GIA and ice melting, see section 2.54) we applied is not accurate 10 enough. On average over the whole period, the mean SLA innovation shows then a slight 11 negative anomaly of -8mm. We also note a seasonal cycle. This is probably due to 12 inconsistency between ORAS4 interannual SSH fields in the Atlantic part and the assimilated 13 data but a part of this problem could also come from runoff forcing. If the seasonal variations 14 represented in the runoff climatological values are not realistic enough, the error in the intake 15 of water mass through the Mediterranean basin is directly transferred to the SLA innovation. 16 The RMS of the innovation is steady all along the reanalysis and close to 6.5cm. This result is 17 quite good, knowing that the RMS standard deviation of observations over time is 8cm (not 18 shown here).

#### 19

The main constraint on the SST consists in the assimilation of in situ surface data and gridded 20 21 maps derived from satellite measurements. Thus, for each cycle, we assimilate at least 243 22 values uniformly distributed every spatial degree and a variable amount of in situ surface data 23 from CORA (Figure 4). Before 2004, we note that the main part of assimilated data comes 24 from the satellite data. The mean satellite SST innovation is close to 0°C during the whole period of the reanalysis. The RMS of innovation is about 0.7°C all along the time period and 25 26 exhibits a seasonal signal with 0.25°C amplitude whose maximum is reached at the end of 27 summer. The same diagnostic using in situ profile observation at the surface exhibits some similar features but we note a weak positive bias between in situ and satellite data of about 28 29 0.12°C at the end of the period (the RMS and the mean values from in situ measurements are 30 only significant between 2005 and 2012).

Finally, we present data assimilation diagnostics for temperature and salinity profiles function 1 2 of the depth (Fig. 5 and 6). Diagnostics on the amount of assimilated data show that before 3 the Argo era, i.e. before about 2005, there are few profiles deployed in the Mediterranean Sea, 4 and most of them only reach 1000m depth. This being so, the mean innovation is close to zero 5 in average between the surface and 2000m depth for temperature and salinity. From 2005 to the end of the experiment, we note a positive anomaly (of observation minus model) of about 6 7 0.2°C and 0.03 psu around 400m depth. According to Figure 5 and 6, this seems to result 8 from a propagation of anomalies from surface layers started in 2003. Those positive 9 anomalies at intermediate depths suggest that the Levantine Intermediate Water (LIW) in the 10 model is too cold and too fresh compared to assimilated data in this layer. Conversely, the 11 innovation in surface and deep layers shows a slight negative anomaly. On average, the RMS 12 of the innovation shows reasonable values compared to the mean innovation and the specified 13 observation errors but we note a clear seasonal variation, especially for temperature profiles. 14 During summer, the surface layers become more stratified. Due to the strong gradients, a 15 small variation in the forecast-trajectory of the ocean model is then more likely to drift from 16 observations and the RMS naturally increases. Moreover, the cold bias in surface associated 17 to a warm bias in subsurface illustrates that there is a lack of stratification in MEDRYS during 18 summer. 19

#### 20 3.2 Scientific assessment results

21

22

23

#### 3.2.24 Mean Sea Surface Height and surface circulation

24 The mean Eddy Kinetic Energy (EKE) and the mean currents of MEDRYS and NM12-FREE over the 1993-2012 period are shown in Fig. 7. The Mean Sea Surface Height 25 (MSSH) of the reanalysis and of the associated free simulation NM12 FREE over the period 26 27 1993-2012 are compared to the mean surface reference CNES-CLS09 product in Figures 7. On average over the whole basin, the assimilation has little impact ( 2mm). The impact is 28 29 strong in term of features. The most important difference concerns the western part of the basin. In NM12 FREE, a significant positive anomaly of MSSH is detected in the Algero-30 Provençal and the Alboran sub basin. A quick comparison between NM12-FREE and 31

MEDRYS mean EKE reveals that the assimilation process has a strong impact in the western 1 2 Mediterranean sub-basin. In NM12-FREE, aThe strong positive mean EKE value-anomaly 3 has been located North of Majorca Island. It corresponds to the fingerprint of a too permanent anticyclonic eddy in NM12 FREE (Figure 8). Thanks to altimetric data, Pascual et al. (2002) 4 identified such an intense eddy in 1998 in the Balearic sub-basin but described it as a 5 temporary event. Actually, in 1998, this anticyclonic eddy develops in September due to 6 7 circumstantial atmospheric and oceanic conditions and disappears during cold seasons. The 8 quasi-permanent occurrence of this eddy in NM12-FREE experiment suggests that the model 9 and its high resolution atmospheric forcing ALDERA are able to produce it but not to 10 dissipate it afterward. This results in a large perturbation in the general circulation in western 11 Mediterranean in NM12-FREE. Indeed, aAccording to figure 87, the Liguro-Provencal current in NM12-FREE is deflected at the southern limit of the Gulf of Lions and a significant 12 13 part of the Atlantic waters is driven along the Spanish coast. This influences the circulation in 14 the Algero-Provençal and the Alboran sub-basins. In MEDRYS, the assimilation process restores realistic SSH and surface circulation. The Atlantic Water (AW) migrates into the 15 16 western Mediterranean trough the Strait of Gibraltar and reaches the Sicily channel through 17 the Algerian current remaining close to the African coast.

18

19 In the reanalysis, the mean kinetic energyEKE especially increases in the Ionian sub-basin 20 compared to the free simulation hindcast (Fig. 8). This is partially due to the characteristic of the observation error we used in the assimilation process (section 2.5). The choice has been 21 22 made not to trust SLA observations within 50 km of the coast and to increase observation 23 errors in these coastal areas. Around the center of the Ionian sub-basin the observation error is 24 not increased, compared to coastal areas, and Mmore energy and features are thus injected by 25 the assimilation process. We also notice that the Levantine sub-basin, and more specifically 26 both the Ierapetra and Pelops anticyclonic eddies, are more energetic suggesting that the 27 mesoscale circulation component have been increased thanks to the assimilation of 28 observational data.

- 29
- 30

#### 3.2.32 Integrated temperature and salinity

1 Integrated temperature and salinity from the EN3 climatological gridded products 2 (Ingleby and Huddleston, 2007) and the IMEDEA (Mediterranean Institute for Advanced 3 Studies) reconstruction (Jordà et al. 2014) are compared below with MEDRYS and NM12-FREE. Basin integrals of the various products are compared whatever real data is present or 4 5 Monthly evolution over three different layers representing surface (0m 150m), 6 intermediate (150m-600m) and deep (600m-bottom) waters are shown in Figure 9 and Figure 7 10. In this section, observational gridded products will serve as the reference for the heat and 8 salt content. In this validation exercise, we will consider the observational gridded products as 9 the reference. Integrated temperature and salinity from two hydrographic products are 10 compared with MEDRYS and NM12-FREE. The two products are EN3 (Ingleby and 11 Huddleston, 2007) and IMEDEA (Jordà et al. 2016, submitted paper; the reconstruction methodology has been described in Llasses et al., 2015). Both products differ in the details of 12 13 the mapping algorithm and the quality control applied to the observations. The difference between them can be viewed as a first estimate of the uncertainties linked to the observational 14 15 products, which cannot be neglected (Jordà and Gomis, 2013 ; Llasses et al., 2015). Basin integrals of the various products are compared whatever real data is present or not. Monthly 16 evolution over three different layers representing surface (0-150m), intermediate (150-600m) 17 and deep (600m-bottom) waters are shown in Fig. 8 and 9. 18

19

20 The time series of the averaged temperature between the surface and 150m depth in Fig. 8 21 point out the good representation of the seasonal cycle in both NM12-FREE and MEDRYS. 22 The phase and the magnitude of the seasonal cycle are consistent with the EN3 and IMEDEA 23 gridded products. In terms of mean value, the two experiments are very close and present a 24 positive bias compared to the gridded products. Indeed, in the 0-150m layer, the difference between the simulations and EN3 is about 0.15°C and twice more compared to the IMEDEA 25 26 reconstruction. This is also consistent with the assimilation statistics of in situ profiles shown 27 in section 3.2.1. In the upper layer, the averaged salinity in MEDRYS and NM12-FREE is comparable with that in EN3 and the-IMEDEA-reconstruction. However, between 1992 and 28 29 2013, MEDRYS show a slight positive bias of about 0.02 psu whereas NM12-FREE show a 30 slight negative bias of -0.03 psu compared to the reference products. Before 1993, the free 31 simulationhindcast presents a clear negative bias of -0.07 psu. In 1993, the data assimilation 32 corrects this surface salinity bias. The interannual variability of the atmospheric water fluxes

(Evaporation-Precipitation-Runoffs, not shown) present a less evaporative period followed by
 a stronger one in the late 90's and early 2000's. This leads to similar variability in the surface
 salt content in both MEDRYS and NM12-FREE. As there are few in situ data, especially for
 salinity, the stronger evaporation <u>combined to a weak salinity constraint</u> during the early
 2000's leads to high surface salinization in MEDRYS.

6

7 Concerning the intermediate waters, one clearly sees on Fig. 8 and 9, the drift of the free simulation in NM12-FREE. The model in a free configuration tends to warm and salinize 8 9 intermediate waters. The assimilation of data restores good average values and realistic variability. It is interesting to notice that despite poor data coverage in the early 90's, the 10 11 assimilation system is able to restore a realistic averaged salinity. As we noted in the previous 12 section 3.2.1, we note a spurious positive anomaly in the MEDRYS salinity in the early 13 2000's. Those too salty and too dense waters have been formed in the surface layers and have 14 been advected toward the bottom layers. This bias is probably explained by a bad adjustment 15 of the volume correction term of the SLA model equivalent (section 2.5). In section 3.2.1, we noted that the mean SLA innovation (obs-model) was decreasing, meaning that the simulated 16 17 sea level trends to rise too quickly compared to the observations. In response thereto, the system tends to compensate by densifying surface waters. As the assimilation system is more 18 19 constrained on temperature (due to better data coverage) it has a strong effect on salinity. The 20 resulting bias is also detected in the bottom layer until 2005. Considering the small number of 21 assimilated data below 600m depth, the model is only slightly constrained beyond this depth, 22 especially before 2005. Thus, the reanalysis is quite close to the free simulation hindcast in 23 terms of tendency and mean value for both temperature and salinity.

24

25 According to Fig. 8, Concerning the temperature in the deepest layers, it is difficult to 26 establish whether, both the free simulation hindcast and the reanalysis, are able to represent a 27 realistic signaltemperature in the deepest layer. Actually, we cannot clearly distinguish any 28 reference values as the two gridded products show different signals. However, the two 29 experiments present a linear trend of warming of about 4\*10-3 °C/year comparable to EN3 for the 1993-2012 period. The-IMEDEA reconstruction-presents a lower warming of about 30 31 1.5\*10-3 °C/year. In the deepest layer, EN3 and IMEDEA reconstruction show similar mean salinity (respectively 38.63 psu and 38.64 psu between 1979 and 2010) and a similar 32

interannual variability. NM12-FREE presents a linear salinization over the whole period of
 the experiment in agreement with the gridded product (1.2\*10-3psu/year). With a limited
 number of data to assimilate, MEDRYS show an episode of high salinization from 1997 to
 2004. Thanks to better data coverage after 2005, the reanalysis becomes more constrained and
 show a more realistic average salinity, in accordance to our reference products.

6

24

7 Following Adani et al. 2011, the vertical distribution of the temperature and salinity anomalies is then presented in Fig. 10 and 11. Temperature and salinity anomalies have been 8 9 computed with respect to the monthly cycle of the MEDATLAS-1979 climatology, from 10 which the October month has been taken to initialise NM12-FREE (see section 2.2). These 11 figures complete the vertical view given by Fig. 5 and 6 which were computed only at 12 observation locations, and the integrated view given by Fig. 8 and 9. Moreover, this kind of 13 diagnostics is presented in Adani et al. (2011) allowing thus a qualitative comparison of two 14 available reanalyses. For temperature, both NM12-FREE and MEDRYS show a similar 15 behaviour in the surface layer (above 100m depth); we can thus attribute these anomalies to the model configuration (for instance issues with the vertical mixing) and to interannual 16 17 variations, both simulations being forced by the same realistic atmospheric forcings in 18 surface. In the intermediate layer, NM12-FREE becomes slowly warmer and warmer, starting 19 with a cold anomaly of about -0.1°C in 1993 and ending with a warm anomaly of about 20 +0.2°C in 2013, in the core of the LIW layer. For MEDRYS, this core is too cold of about -21  $0.2^{\circ}$ C to  $-0.1^{\circ}$ C, this anomaly becoming smaller at the end of the period. In the bottom layer, 22 NM12-FREE remains slightly colder than its initial state, around -0.1 °C, whereas MEDRYS shows a slight warming during the 20 years, in agreement with Fig.8. 23

25 For salinity, again the anomalies above 100m depth are similar in both simulations; the 26 succession of positive and negative anomalies can be related to interannual variability. 27 Nevertheless, the surface layer is more salty in MEDRYS than in NM12-FREE, especially during the last years. In the intermediate layer, around the core of the LIW layer, NM12-28 29 FREE becomes saltier and saltier during the 20 years, from +0.05 psu in 1993 up to +0.15 psu at the end of the period. In MEDRYS the intermediate anomalies are negative, around -0.0530 31 psu, and located deeper than in NM12-FREE, around 650m depth, thus at the base of the LIW 32 layer. In the bottom layer (below 1200m), NM12-FREE has small salinity anomalies around 0

psu, become slightly negative below 2000m between 2003 and 2007, and slightly positive 1 2 between 1200m and 2000m at the end of the period, displaying interannual variability. In 3 MEDRYS, the deep layer is slightly saltier, with a small trend during the period, starting with anomalies around 0 psu in 1993 and ending with anomalies up to +0.1 psu. Moreover, the 4 5 positive anomalies in the surface layer in MEDRYS around year 2000 seems to propagate downwards (as seen in Fig.9), leading to the end of the negative anomaly in the intermediate 6 7 layer between 2001 and 2005 and to a stronger positive anomaly in the bottom layer between 8 2002 and 2006.

We can qualitatively compare Fig.10 and 11 to a similar diagnostic performed by Adani et al. 10 (2011) (their figures 8 and 9); the common period is 1993-2007. One can notice similar 11 12 patterns in both reanalyses: high variability in surface layer, a slightly too cold intermediate 13 layer, and a deep layer becoming warmer and saltier during the simulated period, the amplitude of the anomalies being smaller in MEDRYS. As these reanalyses are performed 14 15 with different numerical modelling choices, different atmospheric forcing and different assimilation schemes, these common features could be related to realistic physical processes, 16 17 which could be interesting to assess in a common dedicated work.

# 18 19

9

#### 3.2.43 Temperature and salinity vertical profiles

21

20

22 The model equivalent at the time and spatial location of the observations has been 23 computed from daily averaged outputs. Mean and RMS differences over the whole 24 Mediterranean basin were computed for- 3 layers (0-150m, 150-600m, 600-4000m) for 25 temperature and salinity profiles (CLASS4-validation; Lellouche et al. 2013) and are presented in Figures 12 and 13. In order to evaluate the improvement with respect to a 26 27 constant state, Www applied the same process with the profiles from MEDATLAS-1998. The 28 MEDATLAS-1998 temperature and salinity fields are the initial states of short simulations 29 used for process studies such as in Beuvier at al. (2012a). Those fields have been obtained 30 pondering by a low pass filtering with a time-window of three years, the MEDATLAS data 31 covering the 1997-1999 period. The choice of centering the climatology on the late 90's

1 corresponds to a compromise between a recent year (before 2002, the last field in 2 MEDATLAS) and a sufficient data coverage in both temperature and salinity, knowing that 3 the uncertainty associated with the MEDATLAS fields increases after 2000. Only a daily 4 dataset, checked through objective quality control, have been assimilated in MEDRYS. Large 5 differences may appear locally in the CLASS4 scores with spurious observations. CLASS4 6 results complements here the statistics made against one week forecasts in section 3.2.1.

7

8 We first assess the mean and RMS temperature differences between the analysis and the 9 observations in Figure 124. Concerning the layer-averaged mean differences, results are not 10 fully consistent with comparisons made with integrated content in section 3.2.32. Indeed, those statistics show that, on average, MEDRYS is very close to the observations (at the 11 12 location of the observations). We only note a significant negative bias of 0.03°C in the layer 13 150/600m on average over the period 1993-2012. The mean temperature difference in the two 14 first layers of the reanalysis reproduces the interannual variability present in the observations. 15 As MEDATLAS-1998 is a climatology, the magnitude of the oceanic interannual variability is then represented by the blue curve. We also point out that, in average, no particular 16 temperature bias occurred in the deepest layer in MEDRYS. This highlights that the system is 17 18 well constrained and efficiently responds to the assimilation of in situ profiles. As in average 19 MEDRYS remains close to temperature measurements, that also confirms that the reference 20 products shown in the section 3.2.32 are subject to uncertainties, especially in the deepest 21 layers where the estimated mean temperature may vary widely from a product to another. In 22 term of mean salinity (Fig. 132), MEDRYS is also close to the observations in the deepest 23 layers but, as expected, presents a slight positive bias of about 0.02psu between the surface 24 and 150m depth. When we compared integrated salinity of the reanalysis with other gridded 25 products, we noted a spurious salinization in MEDRYS in the early 2000s that propagated 26 toward deeper layers. In average, the CLASS4 mean difference in salinity is only about 27 0.1psu between the surface and 150m depth and is not noticeable below. Assuming that the major part of the salinity observations are used in both MEDRYS and the reference gridded 28 29 products, Fthis suggests that the signal of the deeper salinization is not in the observations but 30 is a consequence of the propagation of the simulated surface anomaly through the ocean 31 model. However, as the uncertainties in the salinity products are large (Llasses et al., 2015), 32 it cannot be discarded that the observationnal products missed that change.

2 The RMS of the difference is quite good both in temperature and salinity considering the 3 variability in the different layers. However, we note that the RMS of the difference in salinity 4 increases in the waters deeper than 600m, meaning that, despite a realistic estimation of the mean value, the spatial variability is not robust. This can be explained by the lack of salinity 5 measurements and the poor data coverage in Mediterranean Sea under 1000m depth, 6 7 especially before 2005. In average, MEDRYS presents a lower RMS of the difference of 8 temperature and salinity than MEDATLAS-1998. It is not surprising considering that 9 MEDATLAS-1998 is composed of climatological monthly fields and does not represent the 10 variability of the Mediterranean Sea along the whole period of 21 years. In the first 150m, the 11 RMS of the difference in MEDRYS increases with the summertime stratification.

12

1

13

#### 3.2.5 High frequency variability: validation comparison at LION buoy

14

15 We assess show here the ability of NM12-FREE and MEDRYS to reproduce the high 16 frequency variability at the surface in the Mediterranean basin. In Figure 143, we compare the high frequency measurements of SST and SSS at the LION buoy (first level of CTD 17 18 measurements) during HyMeX SOP2 (Special Observation Period 2 from 27/01/201301/27/2013 to 15/03/201303/15/2013) to the hourly outputs of the two numerical 19 20 experiments at the same location. As we noted in paragraph 2.54, the real-time database have 21 been assimilated in 2013. Data from LION buoy were not yet available in real-time and were 22 not assimilated. Note, that this kind of punctual comparison don't allow to assess the high 23 frequency variability over the whole domain of the simulations, but only give an overview of 24 their own abilities.

25

For both SST and SSS comparisons, MEDRYS is slightly closer to the independent observations than the free simulation<u>hindcast</u>, in terms of mean values and variability. Indeed, the mean surface water of MEDRYS shows a positive bias of 0.07°C and 0.03psu while NM12-FREE shows negative biases which are larger in magnitude (0.13°C and 0.06psu). The major part of the mean bias in SSS between MEDRYS and the observations can be explained by the large difference during January (+0.1psu in average) because the mean bias afterward is very weak (less than 0.01psu). Indeed, we notice a strong jump in the observed SSS the
30th of January (+0.04psu) corresponding to a salinity sensor repair (personal communication
from M.N. Bouin). The water-pump was defective and affected the conductivity
measurement. Assuming that a constant negative bias of 0,04psu contaminated the
observation during January, MEDRYS finally presents very good results in SSS during SOP2
at the LION buoy.

7

MEDRYS has a better correlation with LION buoy for both SST (75.8%) and SSS (78.5%) 8 9 than NM12-FREE (respectively 31.1% and 78.3%). Regarding the SST, MEDRYS has a 10 better correlation with LION buoy than NM12-FREE (respectively 76% and 31%). However, MEDRYS and NM12-FREE show a similar correlation for SSS of 78%. For all that, the free 11 12 simulation hindcast is very similar to MEDRYS in the second half of SOP2. This is not 13 surprising since the variability at the surface is controlled by fluxes (identical for both 14 experiments) during the mixed phase of the convection. We especially note the good 15 representation in phase and amplitude of the diurnal variations of SST. This is especially obvious around the 20th February and during many days in March during a temporary 16 17 restratification period, when the diurnal cycle of ALDERA heat fluxes have a higher daily 18 amplitude (beginning of spring season).

19

#### 20 **3.2.6 Transport through the Strait of Gibraltar**

21

22 The Mediterranean Sea is a semi-enclosed basin showing a negative heat and fresh 23 water loss through the air/sea interface. The main part of this loss is balanced by entering heat 24 and water from the Strait of Gibraltar (while only 10% of the net water flux is due to river 25 runoff). That is why it is necessary to ensure that the fluxes through Gibraltar are realistic. 26 Even if the assimilation scheme corrects a part of the distance between the model and the 27 observation in the Mediterranean Sea, the fluxes play a major role in determining the water 28 mass characteristics and are thereby a good indicator regarding the quality of an experiment 29 over the whole basin. We present here water, heat and salt transport through the Strait of 30 Gibraltar at 5.5°W in Figure 154. Heat and salt fluxes are computed from temperature (T) and salinity (S) using equations 1 and 2. Ux represents the zonal component of the current at 31

1  $5.5^{\circ}W, \rho_0$  is the reference sea water density (1020 Kg.m<sup>-3</sup>), S<sub>med</sub> and V<sub>med</sub> are respectively the 2 surface and the volume of the simulated Mediterranean Sea and N<sub>sec</sub> is the number of seconds 3 in a year. Characteristics of the inflow (surface layers) and the outflow (deep layers) and the 4 difference between the two (net flow) are presented. The interface between inflow and 5 outflow has been determined using the horizontal velocity through the strait at daily time-6 scale.

8 Eq. 1: 
$$HeatFlux_{gib} = \frac{\rho_0 c_p}{s_{med}} \iint T(y, z) U_x(y, z) dy dz$$

9

10 Eq. 2: 
$$SaltFlux_{gib} = \frac{N_{sec}}{V_{med}} \iint S(y,z) U_x(y,z) dy dz$$

11

12 Although the characteristics of the ocean are the same in the buffer zone in the two 13 experiments, the amplitude of both inflow and outflow has been improved thanks to data 14 assimilation in MEDRYS (Fig. 154). Despite the realistic value of the net flow through the 15 Strait of Gibraltar, outflow and inflow are underestimated in NM12-FREE in comparison with recent results published (Soto-Navarro et al., 2010, 2014). According to those studies, the 16 17 acceptable range for inflow and outflow at Gibraltar Strait are respectively [+0.76; +0.86]Sy 18 and [-0.84 ; -0.72]Sv. Despite the realistic value of the net flow through the Strait of Gibraltar, 19 outflow and inflow are underestimated in NM12 FREE in comparison with results published 20 in the last twenty years (Bryden and Kinder, 1991 ; Bryden et al., 1994 ; Tsimplis and 21 Bryden, 2000 ; Candela, 2001 ; Baschek et al., 2001 ; Lafuente et al., 2002, Soto Navarro, 22  $\frac{2010}{100}$ . The reason of having a more accurate exchange at Gibraltar in MEDRYS is that the 23 density difference between the inflowing and outflowing waters is larger (-2.34 kg/m3 in 24 MEDRYS and -2.30 kg/m3 in NM12-FREE). In terms of net heat transport, the reanalysis and 25 the free simulation hindcast (respectively 6.6±0.4 W/m2 and 5.5±0.4 W/m2) are consistent with MacDonald et al. (1994). Both averaged temperature and salinity of the inflow in the 26 27 simulations are also consistent with the characteristics of the ORAS4 Atlantic Water in the 28 buffer zone. We also compare the properties of the inflow in MEDRYS and NM12-FREE 29 with results from Soto-Navarro et al. (2014) at the sill of Espartel. They used, inter alia, the 30 experiment NM12-ARPERA. This simulation show similar results with an interface around 150m depth. At this particular depth, we also report similar results with AW at 15.4°C and
 2 36.7psu in MEDRYS and at 15.5°C and 36,5psu in NM12-FREE.

3

4 The net salt transport through the Strait of Gibraltar at 5.5°W is 1.8±2.8 10-3 psu/year in 5 MEDRYS and 3.0±2.6 10-3psu/year in NM12-FREE (Fig. 14).-It directly impacts the salinity of the basin. Assuming that the Mediterranean volume is constant, the evolution of 6 7 Mediterranean salinity is directly linked to the net transport of salt through the Strait of 8 Gibraltar. The trend in salinity ( $\Delta$ sref) of a the reference hydrographic gridded products (EN3 9 and IMEDEA) over the whole basin serves as a way to estimate a reference net salt transport 10 entering at Gibraltar (SaltFlux<sub>gib</sub> from Eq.2), using SaltFlux<sub>gib</sub> =  $\Delta$ sref. Using EN3 and the IMEDEA reconstruction, From the hydrographic products, we estimate the a reference net salt 11 12 intake at approximately 1.7\*10-3psu/year between 1993 and 2012. In MEDRYS, the averaged 13 net salt transport through the Strait of Gibraltar is very close to this reference value but this is 14 not representative of the evolution of the salinity over the whole basin because of the addition 15 of salinity increments coming from the assimilation scheme. Indeed, NM12-FREE and 16 MEDRYS have a similar trend in salinity in spite of a different net salt transport at Gibraltar.

17

#### 18

4. Discussion and conclusion

#### 19

20 This study describes the configuration and the quality of the high resolution reanalysis 21 MEDRYS and its companion free simulationhindcast NM12-FREE, for the Mediterranean 22 Sea over the period 1992-2013. Both simulations have a common configuration: a high-23 resolution oceanic model NEMOMED12, relaxed in the Atlantic buffer zone to ORAS4 24 interannual fields and forced at the surface with the homogeneous and high-resolution ALDERA atmospheric fluxes. The common element to both simulations is the ocean model, 25 NEMOMED12, a high resolution regional configuration of the ocean general circulation 26 27 model NEMO. The model is relaxed to ORAS4 interannual fields in the Atlantic buffer zone 28 and forced at the surface with the homogeneous and high resolution ALDERA atmospheric 29 fluxes. The 21 years of the reanalysis have been produced using all available in situ profiles in 30 from the CORA4 database, SST maps from the daily NOAA AVHRR-AMSR product and 31 along-track SLA from SSALTO/DUACS associated to SAM2 the assimilation scheme from Mercator Océan. The 12-km and 3-hour spatio-temporal resolution of ALDERA fields allows MEDRYS to explicitly reproduce diurnal cycle, and thus SST, and to simulate the impact of local winds on coastal oceanic areas. As we pay a special attention in reducing sources of inhomogeneity in the atmospheric forcing <u>The consistency of</u> ALDERA dataset along the whole 1979-2013, <u>allows us this suggests</u> to trust in the consistency of the interannual variability of processes known to be driven by air-sea interactions (mixed layer variability, surface circulation variability, etc.) in MEDRYS.

8

9 The validation process has highlighted the good results of the reanalysis in terms of mean 10 circulation and integrated heat and salt contents. The data assimilation has a positive impact, 11 especially in the western basin, where it restores a correct circulation of the Liguro-Provençal 12 current and of the Algerian current. The assimilation process leads to stronger mesoscale 13 variability in the Ionian and Levantine sub-basin, especially at the location of Ierapetra and 14 Pelops eddies. Looking at in situ profiles, the reanalysis shows a realistic water masses at 15 intermediate depths, unlike in the free simulationhindcast. In this layer, the simulation without assimilation NM12-FREE drifts from the observations and show a strong positive trend in 16 both temperature and salinity. Transports through the Strait of Gibraltar have also been 17 18 improved in the reanalysis. Despite the same forcing in the Atlantic buffer zone, both inflow 19 and outflow in MEDRYS have been increased compared to NM12-FREE and are now 20 comparable to historical values. The net heat and salt budgets through the strait are also 21 consistent with independent products. The improvement of the Atlantic/Mediterranean fluxes 22 at Gibraltar ensures a better budget in the Mediterranean Ssea.

23

We showed that surface waters in MEDRYS were in average too salty (about 0.02psu). This 24 25 problem probably comes from the adjustment of the volume correction during the computation of SLA model equivalent.-Indeed, we noted that the mean SLA innovation 26 (observation minus model forecast) was decreasing of 0.65mm/yr, meaning that the simulated 27 28 sea level tends to rise too quickly compared to the observations. In response thereto, the 29 system tends to compensate by densifying surface waters. As the assimilation system is more constrained on temperature (due to better data coverage) it has a strong effect on salinity. We 30 31 also point out that it had inconsistencies between ORAS4 interannual fields in the buffer zone 32 and the assimilated data. To correct for those inconsistencies, it will be necessary to apply a

correction to the ORAS4 SSH fields in order to better represent the seasonal variations of sea 1 2 level in the Mediterranean. In further version of MEDRYS, we simply propose to correct the 3 seasonal cycle and the trends of sea level anomalies in ORAS4 in order to match with altimetry observations in the buffer zone. According to additional works (not shown in this 4 5 study), we realized that SLA innovations were strongly correlated with the mean wind 6 patterns (Mistral-Tramontane, Aegean winds), suggesting that the hydraulic constraint 7 component is not negligible in the Mediterranean Sea. Knowing that, the configuration of 8 SAM2 has should been adjusted in order to take into account the wind component in SSH. 9 Moreover, as the effect of the wind at high frequency has been filtered from the 10 SSALTO/DUACS database (SSALTO/DUACS User Handbook, 2014), it will-would be also 11 necessary to filter it in the model.

12

#### 13 Acknowledgements

14

We acknowledge the two anonymous reviewers for their comments and suggestions which helped to improve this article. This work is a contribution to the HyMeX program (HYdrological cycle in The Mediterranean EXperiment) through INSU-MISTRALS support and the Med-CORDEX program (COordinated Regional climate Downscaling EXperiment – Mediterranean region). This research has received funding from the French National Research Agency (ANR) project REMEMBER (contract ANR-12-SENV-001). All the ALDERA outputs are openly available through the Med-CORDEX database (www.medcordex.eu).

#### 1 References

- 2 Adani, M., Dobricic, S., Pinardi, N., 2011. Quality assessment of a 1985–2007 Mediterranean
- 3 Sea reanalysis. Journal of Atmospheric and Oceanic Technology 28 (4), 569–589.
  4 http://dx.doi.org/10.1175/2010JTECHO798.1.
- 5 Auger P.A., Ulses C., Estournel C., Stemman L., Somot S., Diaz F. (2014) Interannual control
- 6 of plankton ecosystem in a deep convection area as inferred from a 30-year 3D modeling
- 7 <u>study: winter mixing and prey/predator in the NW Mediterranean. Progress in Oceanography:</u>
- 8 <u>12-27, DOI: 10.1016/j.pocean.2014.04.004</u>
- 9 Ayoub, N., Le Traon, P.-Y., De Mey, P., 1998. A description of the Mediterranean surface
- 10 variable circulation from combined ers-1 and topex/poseidon altimetric data. Journal of
- 11 Marine Systems 18 (1-3),3-40,
- 12 http://www.sciencedirect.com/science/article/pii/S0924796398800043"http://www.sciencedir
- 13 ect.com/science/article/pii/S0924796398800043.
- Balmaseda, M. A., K. E. Trenberth, and E. Källén (2013), Distinctive climate signals in
  reanalysis of global ocean heat content, Geophys. Res. Lett., 40, 1754–1759,
  doi:10.1002/grl.50382.
- 17 Barnier B., L. Siefridt, P. Marchesiello, 1995. Thermal forcing for a global ocean circulation
- 18 model using a three-year climatology of ECMWF analyses. Journal of Marine Systems,
- 19 Volume 6, Issue 4, June 1995, Pages 363–380.
- Baschek, B., U. Send, J. Garcia Lafuente, and J. Candela, 2001. Transport estimates in the
  Strait of Gibraltar with a tidal inverse model, Journal of Geophysical Research, 112,
  31,03331,044.
- Béranger, K., L. Mortier, and M. Crépon, 2005. Seasonal variability of water transport
  through the Straits of Gibraltar, Sicily and Corsica, derived from a high-resolution model of
  the Mediterranean circulation, Progress in Oceanography, 66, 341–364.
- 26 Béranger, K., Y. Drillet, M-N. Houssais, P. Testor, R. Bourdallé-Badie, B. Alhammoud, A.
- Bozec, L. Mortier, P. Bouruet-Aubertot, and M. Crépon (2010), Impact of the spatial
  distribution of the atmospheric forcing on water mass formation in the Mediterranean Sea,
- 29 Journal of Geophysical Research, 115(C12041). doi:10.1029/2009JC005648.

Mis en forme : Anglais (États Unis)

- 1 Beuvier, J., F. Sevault, M. Herrmann, H. Kontoyiannis, W. Ludwig, M. Rixen, E. Stanev, K.
- 2 Béranger, and S. Somot, 2010. Modeling the Mediterranean Sea interannual variability during
- 3 1961-2000 : Focus on the Eastern Mediterranean Transient, Journal of Geophysical Research,
- 4 115(C08017), doi :10.1029/2009JC005950.
- Beuvier J., Béranger K., Lebeaupin-Brossier C., Somot S., Sevault F., Drillet Y., BourdalleBadie R., Ferry N., Lyard F., 2012a. Spreading of the Western Mediterranean Deep Water
  after winter 2005: time scales and deep cyclone transport. J Geophys Res-Oceans.
- 8 doi:10.1029/2011JC007679.
- 9 Beuvier, J., C. Lebeaupin-Brossier, K. Béranger, T. Arsouze, R. Bourdallé-Badie, C. Deltel,
- 10 Y. Drillet, P. Drobinski, N. Ferry, F. Lyard, F. Sevault, and S. Somot, 2012b. MED12,
- 11 oceanic component for the modeling of the regional Mediterranean Earth System, Mercator
- 12 Ocean Quaterly Newsletter, 46, Octobre 2012, 60-66.
- Bloom, S. C., Takas, L. L., Da Silva, A. M., and Ledvina, D., 1996. Data assimilation using
   incremental analysis updates. Mon. Weather Rev., 124, 1256–1271.
- Bryden, H. L., and T. H. Kinder, 1991. Steady two layer exchange through the Strait of
   Gibraltar, Deep-Sea Research, 38, 445–463.
- Bryden, H. L., J. Candela, and T. H. Kinder, 1994. Exchange through the Strait of Gibraltar,
   Progress in Oceanography, 33, 201–248.
- 19 Cabanes, C., Grouazel, A., von Schuckmann, K., Hamon, M., Turpin, V., Coatanoan, C.,
- 20 Paris, F., Guinehut, S., Boone, C., Ferry, N., de Boyer Montégut, C., Carval, T., Reverdin, G.,
- 21 Pouliquen, S., and Le Traon, P.-Y., 2013. The CORA dataset: validation and diagnostics of
- in-situ ocean temperature and salinity measurements, Ocean Sci., 9, 1-18, doi:10.5194/os-9-12013.
- 1
- 24 Candela, J., 2001. The Mediterranean water and the global circulation, in Observing and
- Modelling the Global Ocean, pp. 419–429, Edited by G. Siedler, J. Church, J. Gould,
   Academic, San Diego, California.
- 27 Casella E., A. Molcard, A. Provenzale (2011) Mesoscale vortices in the Ligurian Sea and
- 28 their effect on coastal upwelling processes, Journal of Marine Systems, 88, 12-19,
- 29 doi:10.1016/j.jmarsys.2011.02.019.

- 1 Ciappa A. C. (2009) Surface circulation patterns in the Sicily Channel and Ionian Sea as
- 2 revealed by MODIS chlorophyll images from 2003 to 2007, Continental Shelf Research 29,
- 3 2099–2109, doi:10.1016/j.csr.2009.08.002.
- Colin, J., M. Déqué, R. Radu and S. Somot, 2010. Sensitivity study of heavy precipitation in
  Limited Area Model climate simulations: influence of the size of the domain and the use of
  the spectral nudging technique. Tellus A, 62: 591–604. doi: 10.1111/j.16000870.2010.00467.x.
- 8 Crosnier, L. and Le Provost, C., 2007. Inter-comparing five forecast operational systems in
  9 the North Atlantic and Mediterranean basins: the MERSEA-strand1 methodology, J. Mar.
  10 Syst., 65, 354–375.
- Dee, D. P., S. M. Uppala, A. J. Simmons, P. Berrisford, P. Poli, S. Kobayashi, U. Andrae, M. 11 12 A. Balmaseda, G. Balsamo, P. Bauer, P. Bechtold, A. C. M. Beljaars, L. van de Berg, J. 13 Bidlot, N. Bormann, C. Delsol, R. Dragani, M. Fuentes, A. J. Geer, L. Haimberger, S. B. 14 Healy, H. Hersbach, E. V. Hólm, L. Isaksen, P. Kållberg, M. Köhler, M. Matricardi, A. P. 15 McNally, B. M. Monge-Sanz, J.-J. Morcrette, B.-K. Park, C. Peubey, P. de Rosnay, C. Tavolato, J.-N. Thépaut, F. Vitart, 2011. The ERA-Interim reanalysis: configuration and 16 17 performance of the data assimilation system. Q.J.R. Meteorol. Soc., 137: 553-597. doi: 18 10.1002/qj.828.
- Drobinski P. Ducrocq V., Alpert P., Anagnostou E., Béranger K., Borga M., Braud I., Chanzy
  A., Davolio S., Delrieu G., Estournel C., Filali Boubrahmi N., Font J., Grubisic V., Gualdi S.,
- 21 Homar V., Ivancan-Picek B., Kottmeier C., Kotroni V., Lagouvardos K., Lionello P., Llasat
- 22 M.C., Ludwig W., Lutoff C., Mariotti A., Richard E., Romero R., Rotunno R., Roussot O.,
- 23 Ruin I., Somot S., Taupier-Letage I., Tintoré J., Uijlenhoet R., Wernli H. (2013) HyMeX, a
- 24 10-year multidisciplinary program on the Mediterranean water cycle, Bulletin of the
- 25 American Meteorological Society, doi: http://dx.doi.org/10.1175/BAMS-D-12-00242.1.
- Dufau-Julliand C., P. Marsaleix, A. Petrenko, I. Dekeyser (2004) Three-dimensional
  modeling of the Gulf of Lion's hydrodynamics (northwest Mediterranean) during January
  1999 (MOOGLI3 Experiment) and late winter 1999: Western Mediterranean Intermediate
  Water's (WIW's) formation and its cascading over the shelf break , jOURNAL OF
  GEOPHYSICAL RESEARCH, VOL. 109, C11002, doi:10.1029/2003JC002019, 2004

- 1 Durrieu de Madron, X., et al., 2013. Interaction of dense shelf water cascading and open-sea
- 2 convection in the northwestern Mediterranean during winter 2012. Geophys. Res. Lett., 40,
- 3 1379–1385, doi:10.1002/grl.50331.
  4 <u>Elguindi N., Somot S., Déqué M., Ludwig W. (2011) "Climate change evolution of the</u>
  5 hydrological balance of the Mediterranean, Black and Caspian Seas: impact of climate model
- 6 resolution" Clim. Dyn., 36 (1-2), 205-228, doi: 10.1007/s00382-009-0715-4
- 7 Estournel C., X. Durrieu de Madron, P. Marsaleix, F. Auclair, C. Julliand, and R. Vehil
  (2003) Observation and modeling of the winter coastal oceanic circulation in the Gulf of Lion
  9 under wind conditions influenced by the continental orography (FETCH experiment)
  10 OURNAL OF GEOPHYSICAL RESEARCH, VOL. 108, NO. C3, 8059,
  11 doi:10.1029/2001JC000825, 2003.
- Estournel C., F. Auclair, M. Lux, C. Nguyen, and P. Marsaleix (2009) "Scale oriented"
  embedded modeling of the North-Western Mediterranean in the frame of MFSTEP, Ocean
  E. J. 5, 72, 00, 2000
- 14 Sci., 5, 73–90, 2009, www.ocean-sci.net/5/73/2009/.
- 15 Fernandez V. D. E. Dietrich, R. L. Haney, J. Tintore (2005) Mesoscale, seasonal and
- interannual variability in the Mediterranean Sea using a numerical ocean model, Progress in
  Oceanography, doi:10.1016/j.pocean.2004.07.010.
- Ferry, N., Parent, L., Garrie, G., Bricaud, C., Testut, C. E., Le Galloudee, O., Lellouche, J.
  M., Drevillon, M., Greiner, E., Barnier, B., Molines, J. M., Jourdain, N. C., Guinehut, S.,
  Cabanes, C., and Zawadzki, L., 2012. GLORYS2V1 global ocean reanalysis of the altimetric
  era (1992–2009) at mesoscale, Mercator Newsletter 44, 29–39.
- Giorgi, F., 2006. Climate change hot-spots, Geophys. Res. Lett., 33, L08707, doi:
  http://dx.doi.org/10.1029/2006GL025734"10.1029/2006GL025734.
- Hamad, N, Millot, C , Taupier-Letage, I (2005) A new hypothesis about the surface
  circulation in the eastern basin of the Mediterranean Sea, Progress in Oceanography, 66, 28726 | 298.
- 27 Harzallah, A., Alioua, M., & Li, L. (2014). Mass exchange at the Strait of Gibraltar in
- 28 response to tidal and lower frequency forcing as simulated by a Mediterranean Sea model.
- 29 <u>Tellus A, 66.</u>

- 1 Herrmann, M. J., and S. Somot, 2008. Relevance of ERA40 dynamical downscaling for
- 2 modeling deep convection in the Mediterranean Sea, Geophysical Research Letters,
- 3 35(L04607), doi:10.1029/2007GL03244.
- Herrmann, M., F. Sevault, J. Beuvier, and S. Somot, 2010. What induced the exceptional
  2005 convection event in the northwestern Mediterranean basin ? Answers from a modeling
  study. Journal of Geophysical Research, 115(C12051), doi:10.1029/2010JC006162.
- Herrmann, M., S. Somot, S. Calmanti, C. Dubois, and F. Sevault, 2011. Representation of
  spatial and temporal variability of daily wind speed and of intense wind events over the
  Mediterranean Sea using dynamical downscaling : impact of the regional climate model
  configuration, Natural Hazards and Earth System Sciences, 11, 1983–2001, doi:
  10.5194/nhess- 11-1983-2011.
- 12 Ingleby, B. and M. Huddleston, 2007. Quality control of ocean temperature and salinity 13 profiles - historical and real-time data. Journal of Marine Systems, 65, 158–175.
- 14 doi:10.1016/j.jmarsys.2005.11.019.
- 15 Jordà G. and D. Gomis, 2013. On the interpretation of the steric and mass components of sea
- level variability: The case of the Mediterranean basin. Journal of Geophysical Research, Vol.
   <u>118, Issue 2, 953-963. 10.1002/jgrc.20060</u>
- Josey, S. A., S. Somot, and M. Tsimplis, 2011. Impacts of atmospheric modes of variability
  on Mediterranean Sea surface heat exchange, Journal of Geophysical Research, 116(C02032),
  doi:10.1029/2010JC006685.
- Lafuente, J. G., J. Delgado, J. M. Vargas, M. Vargas, F. Plaza, and T. Sarhan, 2002. Low
   frequency variability of the exchanged flows through the Strait of Gibraltar during CANIGO,
   Deep Sea Research II, 49, 4051–4067.
- Langlais, C., B. Barnier, J.M. Molines, P. Fraunié, D. Jacob, S. Kotlarski, 2009. Evaluation of
  a dynamically downscaled atmospheric reanalyse in the prospect of forcing long term
  simulations of the ocean circulation in the Gulf of Lions, Ocean Modelling, Volume 30, Issue
  4, Pages 270-286, ISSN 1463-5003. http://dx.doi.org/10.1016/j.ocemod.2009.07.004.
- <u>Llasses, J, G. Jordà, D. Gomis 2015. Skills of different hydrographic networks in capturing</u>
  changes in the Mediterranean Sea at climate scales. Climate Research. Vol. 63: 1–18, 2015
- 23 <u>changes in the Medhemanean Sea at chinate scales. Chinate Research. Vol. 05. 1–18,</u>
- 30 <u>doi: 10.3354/cr01270</u>

- 1 Lebeaupin Brossier, C., K. Béranger, C. Deltel, and P. Drobinski, 2011. The Mediterranean
- 2 response to different space-time resolution atmospheric forcings using perpetual mode
- 3 sensitivity simulations, Ocean Modelling, 36, 1–25, doi :10.1016/j.ocemod.2010.10.008.
- 4 Lebeaupin Brossier, C., K. Béranger and P. Drobinski, 2012. Sensitivity of the northwestern
- 5 Mediterranean Sea coastal and thermohaline circulations simulated by the 1/12° -resolution
- 6 ocean model NEMO-MED12 to the spatial and temporal resolution of atmospheric forcing,
- 7 Ocean Modelling, 43-44, 94–107, doi :10.1016/j.ocemod.2011.12.007.
- 8 Lebeaupin Brossier C., Arsouze T, Béranger K, Bouin M-N., Bresson E., Ducrocq V.,
- 9 Giordani H., Nureta M., Rainaud R., Taupier-Letage I. (2014) Ocean mixed layer responses to
- 10 intense meteorological events during HyMeX-SOP1 from a high-resolution ocean simulation,
- 11 Ocean Modelling, Vol 84, 84-103.
- 12 Lellouche J.M., O. Le Galloudec, M. Drévillon, C. Régnier, E. Greiner, G. Garric, N. Ferry,
- 13 C. Desportes, C.-E. Testut, C. Bricaud, R. Bourdallé-Badie, B. Tranchant, M. Benkiran, Y.
- 14 Drillet, A. Daudin and C. De Nicola, 2013. Evaluation of global monitoring and forecasting
- 15 systems at Mercator Océan. Ocean Sci., 9, 57–81. doi:10.5194/os-9-57-2013.
- Le Traon, P. Y., Dibarboure, G., and Ducet, N., 2001. Use of a High Resolution Model to
   Analyse the Mapping Capabilities of Multiple-Altimeter Missions, J. Atmos. Ocean. Tech.,
- 18 <del>18, 1277 1288</del>.
- L'Hévéder, B., L. Li, F. Sevault and S. Somot, 2013. Interannual variability of deep
  convection in the Northwestern Mediterranean simulated with a coupled AORCM. Clim.
  Dyn., Vol. 41, Issue 3-4, pp937-960.
- -Louis, J.-F. 1979. A parametric model of vertical eddy fluxes in the atmosphere. Boundary
   Layer Meteorol. 17, 187202.
- Ludwig, W., E. Dumont, M. Meybeck, and S. Heussner, 2009. River discharges of water and
  nutrients to the Mediterranean and Black Sea : Major drivers for ecosystem changes during
  past and future decades ?, Progress in Oceanography, 80, 199–217.
- 27 Lyard, F., F. Lefevre, T. Letellier, and O. Francis, 2006. Modelling the global ocean tides :
- 28 modern insights from FES2004, Ocean Dynamics, 56(5-6), doi :10.1007/s10236-006-0086-x.
- 29 Macdonald, A. M., J. Candela, and H. L. Bryden 1994. An estimate of the net heat transport
- 30 through the Strait of Gibraltar, in Seasonal and Interannual Variability of the Western

Mis en forme : Anglais (États Unis)

- 1 Mediterranean Sea Coastal and Estuarine Studies, Coastal Estuarine Stud., vol. 46, edited by
- 2 P. E. La Violette et al., pp. 13–32, AGU, Washingtion, D. C.
- Madec, G., Delecluse, P., Imbard, M., Levy, C., 1997. OPA, release 8, Ocean general
  circulation reference manual. Technical Report 96/xx, LODYC/IPSL, France.
- 5 Madec, G. and The-NEMO-Team (2008) NEMO, Technical report (IPSL), France.
- 6 Mariotti, A., N. Zeng, J. Yoon, V. Artale, A. Navarra, P. Alpert and L. Li, 2008.
- 7 Mediterranean water cycle changes: transition to drier 21st century conditions in observations
- 8 and CMIP3 simulations. Res. Lett. 3 044001 doi:10.1088/1748-9326/3/4/044001.
- 9 Marullo, S., E. Napolitano, R. Santoleri, B. Manca, and R. Evans, 2003. Variability of Rhodes
- and Ierapetra Gyres during Levantine Intermediate Water Experiment: Observations and
  model results, J. Geophys. Res., 108, 8119, doi:10.1029/2002JC001393, C9.
- 12 Millot C. and Taupier-Letage (2005) Circulation in the Mediterranean Sea, The handbook of
- 13 environmental chemistry, Vol.1, 29-66, Springer-Verlag.
- 14 Nabat, P., S. Somot, M. Mallet, A. Sanchez-Lorenzo, and M. Wild, 2014. Contribution of
- 15 anthropogenic sulfate aerosols to the changing Euro-Mediterranean climate since 1980,
- 16 Geophys. Res. Lett., 41, 5605–5611, doi:10.1002/2014GL060798.
- 17 Ourmières Y, Zakardjian B, Béranger K, Langlais C (2011) Assessment of a NEMO-based
- 18 downscaling experiment for the North-Western Mediterranean region: impacts on the
- 19 Northern Current and comparison with ADCP data and altimetry products. Ocean Modelling
- 20 39, 386-404.
- 21 Palmiéri, J., Orr, J. C., Dutay, J.-C., Béranger, K., Schneider, A., Beuvier, J., and Somot, S.,
- 22 2015. Simulated anthropogenic CO2 storage and acidification of the Mediterranean Sea,
- 23 Biogeosciences, 12, 781-802, doi:10.5194/bg-12-781-2015.
- 24 Papadopoulos Vassilis P., Simon A. Josey, Aristides Bartzokas, Samuel Somot, Simon Ruiz,
- and Paraskevi Drakopoulou, 2012. Large-Scale Atmospheric Circulation Favoring Deep- and
- 26 Intermediate-Water Formation in the Mediterranean Sea. J. Climate, 25, 6079–6091. doi:
- 27 http://dx.doi.org/10.1175/JCLI-D-11-00657.1.
- 28 Pascual, A., B. Buongiorno Nardelli, G. Larnicol, M. Emelianov, and D. Gomis, 2002. A case
- 29 of an intense anticyclonic eddy in the Balearic Sea (western Mediterranean), J. Geophys. Res.,
- 30 107(C11), 3183, doi:10.1029/2001JC000913.

- 1 Peltier, W. R., and R. Drummond, 2008. Rheological stratification of the lithosphere: A direct
- 2 inference based upon the geodetically observed pattern of the glacial isostatic adjustment of
- 3 the North American continent, Geophys. Res. Lett., 35, L16314, doi:10.1029/2008GL034586.
- Pettenuzzo, D., W. G. Large, and N. Pinardi, 2010. On the corrections of ERA-40 surface flux
  products consistent with the Mediterranean heat and water budgets and the connection
  between basin surface total heat flux and NAO, J. Geophys. Res., 115, C06022,
  doi:10.1029/2009JC005631.
- 8 Pinardi, N., Zavatarelli, M., Adani, M., Coppini, G., Fratianni, C., Oddo, P., Simoncelli, S.,
- 9 Tonani, M., Lyubartsev, V., Dobricic, S., Bonaduce, A., 2013. Mediterranean Sea large-scale
- 10 low-frequency ocean variability and water mass formation rates from 1987 to 2007: A
- 11 retrospective analysis. Prog. Oceanogr. (2013),
- 12 http://dx.doi.org/10.1016/j.pocean.2013.11.003.
- Radu, R., Déqué, M. and S. Somot, 2008. Spectral nudging in a spectral regional climate
  model. Tellus A, 60: 898–910. doi: 10.1111/j.1600-0870.2008.00341.x.
- Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., and Schlax, M. G.,
  2007. Daily high-resolution blended analyses for sea surface temperature, J. Climate, 20,
  5473–5496.
- Rio, M. H., Guinehut, S., and Larnicol, G., 2011. New CNES-CLS09 global mean dynamic
  topography computed from the combination of GRACE data, altimetry, and in situ
  measurements, J. Geophys. Res., 116, C07018, doi:10.1029/2010JC006505.
- 21 Rixen, M., et al., 2005. The Western Mediterranean Deep Water : A proxy for climate change,
- 22 Geophysical Research Letters, 32(L12608), 1–4.
- 23 Robinson, A.R., Hecht, A., Pinardi, N., Bishop, J., Leslie, W.G., Rosentroub, Z., Mariano,
- A.J., Brenner, S., 1987. Small synoptic/mesoscale eddies and energetic variability of the
- 25 eastern levantine basin. Nature 327 (6118), 131–134.
- 26 http://dx.doi.org/"http://dx.doi.org/10.1038/327131a0.
- 27 Robinson, A. R., W. G. Leslie, A. Theocharis, and A. Laskaratos, 2001. Mediterranean Sea
- 28 circulation. Encyclopedia of Ocean Sciences, J. H. Steele, S. A. Thorpe, and K. K. Turekian,
- 29 Eds., Academic Press, 1689–1706.

- 1 Roether W., B. Klein, B. B. Manca, A. Theocharis, S. Kioroglou (2007) Transient Eastern
- 2 Mediterranean deep waters in response to the massive dense-water output of the Aegean Sea
- 3 in the 1990s , Progress in Oceanography 74 (2007) 540–571,
- 4 doi:10.1016/j.pocean.2007.03.001.
- 5 Rubio, A., B. Barnier, G. Jorda`, M. Espino, and P. Marsaleix (2009), Origin and dynamics of
- 6 mesoscale eddies in the Catalan Sea (NW Mediterranean): Insight from a numerical model
- 7 study, J. Geophys. Res., 114, C06009, doi:10.1029/2007JC004245.
- 8 Ruti, P., S. Marullo, F. D'Ortenzio, M. Tremant, 2007. Comparison of analyzed and measured
- 9 wind speeds in the perspective of oceanic simulations over the Mediterranean basin:
- 10 Analyses, QuikSCAT and buoy data. Journal of Marine Systems, Vol. 70, Issues 1-2, P. 33-
- 11 48. doi:10.1016/j.jmarsys.2007.02.026.
- 12 Ruti PM, Somot S, Giorgi F, Dubois C, Flaounas E, Obermann A, Dell'Aquila A, Pisacane
- 13 G, Harzallah A, Lombardi E, Ahrens B, Akhtar N, Alias A, Arsouze T, Aznar R, Bastin S,
- 14 Bartholy J, Béranger K, Beuvier J, Bouffies-Cloché S, Brauch J, Cabos W, Calmanti S,
- 15 Calvet J-C, Carillo A, Conte D, Coppola E, Djurdjevic V, Drobinski P, Elizalde-Arellano A,
- 16 Gaertner M, Galàn P, Gallardo C, Gualdi S, Goncalves M, Jorba O, Jordà G, L'Heveder B,
- 17 Lebeaupin-Brossier C, Li L, Liguori G, Lionello P, Maciàs D, Nabat P, Onol B, Raikovic B,
- 18 Ramage K, Sevault F, Sannino G, MV Struglia, Sanna A, Torma C, Vervatis V., 2015. MED-
- 19 CORDEX initiative for Mediterranean Climate studies (BAMS, in revision).
- Sanchez Gomez, E., Somot, S., Déqué, M., 2009. Ability of an ensemble of regional climate
   models to reproduce weather regimes over Europe Atlantic during the period 1961–2000.
   Climate dynamics, 33(5), 723–736.
- 23 Sanchez-Gomez, E., S. Somot, S. A. Josey, C. Dubois, N. Elguindi and M. Déqué, 2011.
- 24 Evaluation of Mediterranean Sea water and heat budgets simulated by an ensemble of high
- 25 resolution regional climate models. Clim. Dyn., Vol. 37, Issue 9-10, pp 2067-2086.
- 26 Schroeder K., A. Ribotti, M. Borghini, R. Sorgente, A. Perilli, and G. P. Gasparini (2008) An
- 27 extensive western Mediterranean deep water renewal between 2004 and 2006 ,
- 28 GEOPHYSICAL RESEARCH LETTERS, VOL. 35, L18605, doi:10.1029/2008GL035146.
- 29 Sevault, F., S. Somot, A. Alias, C. Dubois, C. Lebeaupin-Brossier, P. Nabat, F. Adloff, M.
- 30 Déqué and B. Decharme, 2014. A fully coupled Mediterranean regional climate system

model : design and evaluation of the ocean component for the 1980-2012 period. Tellus A
 2 2014, 66, 23967, http:/dx.doi.org/10.3402/tellusa.v66.23967.

Solé J., Ruiz S., Pascual A., Somot S., Tintoré J. (2012) Ocean color response to wind forcing
in the Alboran Sea : a new forecasting method. J. Mar. Syst. Vol 98-99:1-8. Doi:
10.1016/j.jmarsys.2012.02.007.

6 Somot, S., F. Sevault, and M. Deque, 2006. Transient climate change scenario simulation of

the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation
model, Climate Dynamics, 27, 851–879.

Sorgente R, A. Olita, P. Oddo, L. Fazioli, A. Ribotti (2011) Numerical simulation and
decomposition of kinetic energy in the Central Mediterranean: insight on mesoscale
circulation and energy conversion, Ocean Sci., 7, 503–519, 2011, www.oceansci.net/7/503/2011/, doi:10.5194/os-7-503-2011.

Sotillo, M. G., A. W. Ratsimandresy, J. C. Carretero, A. Bentamy, F. Valero, and F.
Gonzalez-Rouco, 2005. A high-resolution 44-year atmospheric hindcast for the
Mediterranean Basin : contribution to the regional improvement of global reanalysis, Climate
Dynamics, 25, 219–236.

Soto-Navarro, J., F. Criado-Aldeanueva, J. Garcia-Lafuente, and A. Sanchez-Roman, 2010.
Estimation of the Atlantic inflow through the Strait of Gibraltar from climatological and in
situ data, Journal of Geophysical Research, 115(C10023), doi:10.1029/2010JC006302.

20 Soto-Navarro, J., S. Somot, F. Sevault, J. Beuvier, F. Criado-Aldeanueva, J. Garcia-Lafuente

and K. Béranger, 2014. Evaluation of regional ocean circulation models for the Mediterranean

22 Sea at the Strait of Gibraltar: volume transport and thermohaline properties of the outflow.

23 Clim. Dyn. doi :10.1007/s00382-014-2179-4.

24 <u>Stanev, E. V., and E. L. Peneva, 2002. Regional sea level response to global climatic change:</u>

25 <u>Black Sea examples, Global Planet. Changes, 32, 33–47.</u>

26 Struglia, M.V., Mariotti, A., Filograsso, A., 2004. River discharge into the Mediterranean

27 Sea: climatology and aspects of the observed variability. Journal of Climate 17 (24), 4740–

- 28 4751. <u>http://dx.doi.org/10.1175/JCLI-3225.1"http://dx.doi.org/10.1175/JCLI-3225.1</u>.
- 29 Tonani, M., Pinardi, N., Dobricic, S., Pujol, I., and Fratianni, C.: A high-resolution free30 surface model of the Mediterranean Sea, Ocean Sci., 4, 1-14, doi:10.5194/os-4-1-2008, 2008.

1	Tsimplis, M. N., and H. L. Bryden, 2000. Estimation of the transport through the Strait of
2	Gibraltar, Deep Sea Research I, 47, 2219-2242.
3	Vervatis, V. D., S. S. Sofianos, N. Skliris, S. Somot, A. Lascaratos, and M. Rixen (2013),
4	Mechanisms controlling the thermohaline circulation pattern variability in the Aegean-
5	Levantine region. A hindcast simulation (1960-2000) with an eddy resolving model, Deep-
6	Sea Research Part I, 74:82-97, doi:10.1016/j.dsr.2012.12.011
7	Vrac M., P. Drobinski, A. Merlo, M. Herrmann, C. Lavaysse, L. Li, S. Somot, 2012.
8	Dynamical and statistical downscaling of the French Mediterranean climate: uncertainty
9	assessment. Natural Hazards and Earth System Sciences, European Geosciences Union, 2012,
10	<u>12, pp.2769-2784. &lt;10.5194/nhess-12-2769-2012&gt;</u>
11	Wüst, G., 1961. On the vertical circulation of the Mediterranean Sea, J. Geophys. Res.,
12	66(10), 3261-3271, doi:10.1029/JZ066i010p03261.
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	

Dataset	Shortwave	Longwave	Latent heat	Sensible heat	Net surface heat flux
Reference	[183, 185]	[-84 , -75]	[-90, -88]	[-14,-6]	[-5,-1]
ALDERA	204	-85	-112	-10	-3
ARPERA2	187	-79	-111	-12	-15
ERA-Int (1989-2004)	198	-83	-97	-12	+6
Pettenuzzo et al. 2010 (1958-2001)	180	-80	-91	-14	-5
ALADIN at 50km	196	-81	-111	-11	-7
ALADIN at 150km	200	-82	-94	-10	+14
ENSEMBLES RCMs	[154, 214]	[-100, -70]	[-128 , -85]	[-22,-8]	[-40, +21]

Table 1: Mediterranean Sea averaged and temporal averaged values of the various terms of the sea surface heat budget (W/m2). Values are computed over the 1985-2004 period except for when indicated. The reference comes from Sevault et al. (2014). The so-called ENSEMBLES RCMs is an ensemble of 15 runs carried out with state-of-the-art RCMs during the EU project ENSEMBLES at 25km, driven by the ERA-40 reanalysis over the 1958-2001

7 period (see Sanchez-Gomez et al. 2011).

Dataset	Evaporation	Precipitation	River runoff	Black Sea freshwater inputs	Net surface water flux
Reference 1	-3.1	0.7	0.4	0.3	-1.7
Reference 2	[-3.3 , -2.9]	[0.6 , 0.8]	[0.3, 0.5]	[0.2, 0.4]	[-2.0, -1.4]
ALDERA (1979-2011)	-4.0	1.6	0.4*	0.3*	-1.7
ARPERA2 (1958-2008)	-3.9	1.8	0.2**	0.3**	-1.6**
ERA-Int (1989-2004)	-3.2	1.4	-	-	
Pettenuzzo et al. 2010 (1958-2001)	-3.2	1.4	-	-	
ALADIN at 50km (1979-2011)	-4.0	1.4	0.4*	0.3*	-1.9
ALADIN at 150km (1979-2011)	-3.3	1.1	0.4*	0.3*	-1.5
ENSEMBLES RCMs	[-4.4 , -2.9]	[1.0, 1.7]	[0.2, 0.6]	[0.1, 0.5]	[-2.0, -1.2]

3 Table 2: Same as Table 1 but for the Mediterranean Sea surface water budget terms 4 (mm/day). The reference 1 comes from Sanchez-Gomez et al. (2011) and the reference 2 from 5 Dubois et al. (2010). The reference values do not always cover a common period. \* : the 6 ALDERA atmospheric forcing is here completed by the river runoff and Black Sea freshwater inputs coming respectively from Ludwig et al. (2009) and Stanev et al. (2008) as used in 7 Beuvier et al. (2012b) and in the MEDRYS simulation. \*\* : the ARPERA2 atmosphere 8 9 forcing is here completed by the river runoff and Black Sea freshwater inputs coming respectively from Ludwig et al. (2009) and Stanev et al. (2008) as used in the Herrmann et al. 10 11 (2010) paper.



Figure 1: Daily average wind direction (arrows) and latent heat flux (color in W.m<sup>-2</sup>) on March 14th 2013 in (a) ALADIN-150km, (b) ALADIN-50km and (c) ALADIN-12km (the so-called ALDERA).



Figure 2: Daily average wind direction (arrows) and wind speed (color in m.s<sup>-1</sup>) on August
16th 2012 in (a) ALADIN-150km, (b) ALADIN-50km and (c) ALADIN-12km (the so-called
ALDERA).

l Setellite second	<b>A</b>	De eter	E., J
Satemite name	Acronym	Begin	Ena
ERS2	e2	15/05/1995	09/04/2003
Topex/Poseidon	tp	25/09/1992	24/04/2002
Topex/Poseidon (interleaved)	tpn	16/09/2002	08/10/2005
Geosat Follow-On	g2	07/01/2000	07/09/2008
Jason 1	j1	24/04/2002	19/10/2008
Envisat	en	09/10/2002	22/10/2010
Jason 2	j2	19/10/2008	now
Jason 1 (interleaved)	jln	14/02/2009	now
Envisat (interleaved)	enn	22/10/2010	now
Cryosat 2	c2	19/02/2012	now
Jason 1 Geodetic	jlg	14/05/2012	now

4 assimilation process.

<sup>3</sup> Table 3: Name, acronym and period of SLA measurement for all satellite used by the



3 Figure 3: Time series of weekly sea level anomaly (SLA, m) data assimilation statistics

- 4 averaged over the whole Mediterranean basin : mean innovation (top) and RMS of innovation
- 5 (bottom). The colors stand for different satellites (please refer to Tab. 3).





3 Figure 4: Time series of weekly sea surface temperature (SST, °C) data assimilation statistics

from in situ (blue) and satellite SST AVHRR-AMSR (black), averaged over the whole 4

5 Mediterranean basin : number of data (top), mean innovation (middle) and RMS of innovation

6 (bottom).



Figure 5: Evolution of weekly temperature data assimilation statistics from in situ profiles,
function of the depth averaged over the whole Mediterranean basin : number of profiles
observations (top), mean innovation (middle) and RMS of the innovation (bottom).



Figure 6: Evolution of weekly salinity data assimilation statistics from in situ profiles, function of the depth averaged over the whole Mediterranean basin : number of profiles observations (top), mean innovation (middle) and RMS of the innovation (bottom). 



Figure 7: Mean Eddy Kinetic Energy (EKE in cm<sup>2</sup>.s<sup>-2</sup>) at 40m depth over the period 19922013 for NM12-FREE (top) and MEDRYS<del>1V1</del> (bottom). Arrows represent the mean currents
(in cm<sup>2</sup>.s<sup>-1</sup>) over the same period and at the same depth.



Figure <u>89</u>: Evolution of the monthly integrated heat content (expressed as mean temperature
in °C) over the Mediterranean basin for the layers 0m-150m (top), 150m-600m (middle) and
600m-bottom (bottom) from MEDRYS<del>1V1</del> (red line), NM12-FREE (black line), EN3 (dotted
green line) and the IMEDEA (blue dotted line) hydrographic gridded products. The blue
shaded area indicates the uncertainty ranges around the values of the IMEDEA-reconstruction.



Figure <u>910</u>: Same as Figure <u>89</u> but for integrated salt content (expressed as mean salinity in psu).



Figure 10: NM12-FREE basin mean temperature (°C, above) and salinity (psu, below) anomalies with respect to MedAtlas-1979.





4

5

7

8

9

Figure 121: Temperature (°C) mean (upper row) and RMS (bottom row) differences between analysis and minus observation (black), and between MEDATLAS-1998 and minus observation (blue). For these diagnostics, all available T/S observations from the CORIOLIS 6 database and MEDRYS<sup>1V1</sup> daily average analysis, collocated (temporally and spatially) with observations, are used. The number of observations is shown with gray bars. Averages are performed in the 0-150m (left), 150-600m (middle) and 600m-4000m (right) layers in the whole Mediterranean basin.





Figure 1<u>3</u><del>2</del>: Same as Fig. 1<u>2</u><del>1</del> but for salinity (psu).



Figure 1<u>4</u>3: Evolution of the hourly Sea Surface Temperature (SST, top) and Sea Surface Salinity (SSS, bottom) at the LION buoy location <u>(red dot on the map)</u> between 01/01/2013 and <u>31/03/201303/31/2013</u>. The observation is shown with the green lines, NM12-FREE with the black lines and MEDRYS<del>1V1</del> with the red lines.

	IN	OUT	NET
Water (Sv)			
NM12-FREE	+0.70 ± 0.03	-0.65 ± 0.03	+0.047 ± 0.009
MEDRYS	+0.81 ± 0.03	-0.77 ± 0.03	+0.048 ± 0.009
Heat (W/m <sup>2</sup> )			
NM12-FREE (averaged temperature)	+19.6 ± 0.9 17.18 °C	-14.1 ± 0.6 13.62 °C	+5.5 ± 0.4 (diff = +3.56 °C)
MEDRYS (averaged temperature)	+22.8 ± 0.9 17.13 °C	-16.2 ± 0.6 13.31 °C	+6.6 ± 0.4 (diff = +3.82 °C)
Salt (10 <sup>-3</sup> psu/year)			
NM12-FREE (averaged salinity) MEDRYS (averaged salinity)	+208 ± 8 36.49 psu +243 ± 9 36.54 psu	-205 ± 9 <sup>38.44 psu</sup> -241 ± 10 <sup>38.45 psu</sup>	+3.0 ± 2.6 (diff = -1.95 psu) +1.8 ± 2.8 (diff = -1.91 psu)

Figure 1<u>5</u>4: Average flow, heat and salt transport of the inflow and the outflow through the
Strait of Gibraltar at 5.5°W between 1992 and 2013 for NMED12-FREE and MEDRYS1V1.
The uncertainty corresponds to the annual standard deviation. For heat and salt transport, the
associated mean temperature and salinity in the layer are specified. The green color represents
values consistent with literature or/and reference products and the red color, those that are not
consistent.