

1 **The flow field of the upper hypoxic Eastern Tropical North Atlantic**
2 **oxygen minimum zone**

3
4 **L. Stramma¹, R. Czeschel¹, T. Tanhua¹, P. Brandt^{1,2}, M. Visbeck^{1,2}, and B.S.**
5 **Giese³**

6 [1]{GEOMAR Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20, 24105
7 Kiel, Germany}

8 [2]{Kiel University, Christian-Albrechts-Platz 4, 24118 Kiel, Germany}

9 [3]{Department of Oceanography, Texas A&M University, College Station, Texas, USA}

10
11
12
13
14
15 Status: 02 December 2015

16 Received: 13 August 2015 - Accepted: 31 August 2015 –Published:

17 Correspondence to: L. Stramma (lstramma@geomar.de)

18 Published by Copernicus Publications on behalf of the European Geosciences Union.

19
20
21 Short title: Upper OMZ flow field in the eastern North Atlantic

1

2 **Abstract**

3 A subsurface low oxygen zone is located in the eastern tropical North Atlantic Ocean (ETNA)
4 in the upper ocean with the core of the hypoxic ($O_2 \leq 60 \mu\text{mol kg}^{-1}$) oxygen minimum zone
5 (OMZ) at 400 to 500 m depth. The subsurface circulation in the OMZ region is derived from
6 observations and data assimilation results. Measurements in the ETNA of velocity, oxygen
7 and of a tracer (CF_3SF_5) that was released in April 2008 at $\sim 8^\circ\text{N}$, 23°W (at ~ 330 m depth) in
8 November/December 2008, in November/December 2009 and October/November 2010 show
9 the circulation in the upper part of the OMZ with spreading to the east in the North Equatorial
10 Countercurrent (NECC) region and northwestward around the Guinea Dome. Three floats
11 equipped with oxygen sensors deployed at $\sim 8^\circ\text{N}$, 23°W with parking depths at 330, 350 and
12 400 m depths were used to estimate velocity along the float trajectory at the surface and at the
13 parking depth. At the 350 m park depth north of 9°N a cyclonic northwestward flow across
14 the OMZ was observed. The northward drift of a float into the upper OMZ and a stronger
15 cyclonic flow around the Guinea Dome seem to be connected to a strong Atlantic Meridional
16 Mode (AMM) event in 2009. A near-surface cyclonic circulation cell east of the Cape Verde
17 Islands reaches down into the OMZ layer. The circulation of the upper OMZ mirrors the near
18 surface circulation. Oxygen measurements from the cruises used here, as well as from other
19 recent cruises up to the year 2014 confirm the continuous deoxygenation trend in the upper
20 OMZ since the 1960's near the Guinea Dome. The three floats deployed with the tracer show
21 spreading paths consistent with the overall observed tracer spreading. Oxygen sensors on the
22 floats remained well calibrated for more than 20 months and so the oxygen profiles can be
23 used to investigate mesoscale eddy signatures. Mesoscale eddies may modify the oxygen
24 distribution in OMZs. However, in general eddies are less energetic in the ETNA south of the
25 Cape Verde Islands compared to similar latitudes in the Eastern Tropical South Pacific.

26

27 **1 Introduction**

28 In the Eastern Tropical North Atlantic (ETNA) a subsurface low-oxygen zone exists with a
29 pronounced minimum in oxygen at about 400 to 500 m depth. South and east of the Cape
30 Verde Islands the oxygen minimum is strongest and is referred to as an oxygen minimum
31 zone (OMZ). This OMZ is hypoxic (oxygen concentrations drop below ~ 60 to $120 \mu\text{mol kg}^{-1}$;

1 e.g. Stramma et al., 2008a) while OMZs in the Eastern Tropical South Pacific and northern
2 Indian Ocean are suboxic (oxygen concentrations below about 4.5-10.0 $\mu\text{mol kg}^{-1}$; e.g.
3 Karstensen et al., 2008; Stramma et al., 2008a). Under hypoxic conditions key mobile macro
4 organisms, such as tuna and marlin, are stressed (Stramma et al., 2012) while in suboxic
5 regions dramatically different ecosystems exist and under extreme circumstances nitrate
6 becomes involved in respiration (e.g. Kalvelage et al., 2013). A vertical expansion of the
7 OMZ and a decrease of OMZ core oxygen concentrations are detected in the tropical Atlantic
8 and Pacific Oceans (Stramma et al., 2008a). Since 2009 record low dissolved oxygen values
9 with less than 40 $\mu\text{mol kg}^{-1}$ were observed in the core of the ETNA OMZ (Stramma et al.,
10 2009). If such a trend continues the ETNA region might become suboxic in the future.

11 The structure of the OMZ is a consequence of the balance between the supply of oxygen
12 through ventilation and circulation, oxygen production by photosynthesis and oxygen
13 consumption by remineralization of sinking organic matter. Brandt et al. (2015) summarize
14 the oxygen supply by energetic equatorial current bands as well as the present knowledge on
15 vertical and lateral mixing, advection and consumption within the OMZ. Oxygen variability
16 might be related to changes in the strength of latitudinally stacked zonal jets as derived by
17 Brandt et al. (2010) which result in changes in the advective pathways to the ETNA OMZ,
18 with likely the strongest impact in the upper 300-400 m of the water column (Hahn et al.,
19 2014). The importance of the equatorial current system for the mean oxygen distribution is
20 also revealed by high-resolution model simulations showing the oxygen supply at the equator
21 from a case with a more realistic representation of the circulation, as compared with coarser
22 resolution models (Duteil et al. 2014).

23 In terms of water masses, the core of the North Atlantic OMZ is comprised of Atlantic
24 Central Water and Antarctic Intermediate Water (AAIW) layers. The Central Water is
25 bounded by the isopycnals $\sigma_{\theta} = 25.8$ and 27.1 kg m^{-3} (Stramma et al., 2008b). Two types of
26 Central Water are found in the eastern tropical Atlantic Ocean. North of the Cape Verde
27 Islands North Atlantic Central Water (NACW) is found while south of the Cape Verde Islands
28 South Atlantic Central Water (SACW) dominates. There is an inclined boundary between
29 NACW and SACW rising from south to north, i.e. SACW lying on top of NACW (Tomczak,
30 1984), hence near the Cape Verde Islands the lower OMZ is more influenced by NACW than
31 in the upper OMZ layers. Based on this Central Water distribution Peña-Izquierdo et al.
32 (2015) proposed different flow regimes for the upper and intermediate central water layer

1 separated by the isopycnal $\sigma_{\theta} = 26.8 \text{ kg m}^{-3}$ at about 300 m depth. The AAIW signature is
2 most prominent below the OMZ core and spreads northward near the African continent
3 reaching as far north as 32.5°N (Machin and Pelegrí, 2009).

4 To better understand the existence of the OMZ, the oxygen supply paths and the oxygen
5 changes it is necessary to know the flow field in this region. Some flow field schematics exist
6 for the upper ocean circulation, however knowledge of the circulation in the depth range of
7 the OMZ is very limited. The major supply path of oxygen above and in the uppermost
8 reaches of the OMZ in the eastern tropical North Atlantic are from the eastward flowing and
9 latitudinally stacked zonal jets at and near the equator (Brandt et al., 2015) and from the North
10 Equatorial Countercurrent (NECC). Sometimes the NECC is obscured by the westward
11 Ekman flow component at the surface. Two current bands of the NECC were briefly
12 mentioned by Richardson and Reverdin (1987) and “rediscovered” in observations (Stramma
13 et al., 2005) and modelling efforts (Urbano et al., 2006), the northern branch named the
14 nNECC. The two NECC branches (Fig. 1) exist between $\sim 3^{\circ}\text{N}$ and $\sim 10^{\circ}\text{N}$. The NECC
15 originates in the North Brazil Current Retroflexion (e.g. Schott et al., 1998) and carries water
16 from the South Atlantic. Below the Ekman layer recirculation from the Northern Hemisphere
17 subtropical gyre contributes to the flow within the NECC (Lumpkin and Garzoli, 2005). The
18 property distribution within the ETNA OMZ shows the lowest oxygen concentrations north of
19 the NECC at about 400 to 500 m depth just above the boundary between Central Water and
20 Antarctic Intermediate Water. The NECC is seasonally connected to the North Equatorial
21 Undercurrent (NEUC) at 4 to 6°N and together with its northern branch at 8 to 10°N supplies
22 oxygen-rich water to the OMZ most pronounced in summer and fall (Stramma et al., 2008b).
23 The NECC velocities are strongest in summer and fall but a weak eastward NECC also exists
24 in winter and spring with lower oxygen content compared to summer and fall (Stramma et al.,
25 2008b). Radon transform analysis (Deans, 1983) indicates that the zonal propagation
26 characteristics of the NECC are consistent with long Rossby waves (Hormann et al., 2012).
27 Along-shore wind fluctuations and equatorially forced coastal Kelvin waves are found to be
28 responsible for the excitation of annual- and semiannual-propagating Rossby waves in the
29 eastern sub-basin (Chu et al., 2007). The NECC splits in the eastern basin (e.g. Stramma and
30 Schott, 1999) with one current band continuing eastward into the Guinea Basin while the
31 other one turns northward and flows off the African coast and is called the Mauretania
32 Current (MC; e.g. Peña-Izquierdo et al., 2012) off Mauretania. South of the NECC a northern
33 band of the South Equatorial Current (nSEC) flows westward north of the equator, while the

1 North Equatorial Current (NEC) limits the OMZ region to the north of the Cape Verde Islands
2 (Fig. 1). Between the Cape Verde Islands and the African shelf a permanent cyclonic feature
3 exists year-round (Mittelstaedt, 1983).

4 Ventilation of the OMZ at 300 to 600 m depth is weaker compared to the strong oxygen
5 supply at 150 to 300 m depth (e.g. Fischer et al., 2013; Hahn et al., 2014). South of the Cape
6 Verde Islands there is a large-scale cyclonic circulation feature with an upward displacement
7 of isotherms at a depth of more than 300 m called the Guinea Dome (Siedler et al., 1992).
8 The Guinea Dome (GD) exists all year-round, although it is weaker in winter. The upper
9 thermocline center of the dome is found at 9°N, 25°W in summer and at 10.5°N, 22°W in
10 winter (Siedler et al., 1992). Westward drainage of oxygen-poor water takes place north of the
11 Guinea Dome center, i.e. north of 10°N, and is most pronounced at a depth of 400 to 600 m
12 (Stramma et al., 2008b). The flow field at 400 m depth (Stramma et al. 2008b; their Fig. 12b)
13 is similar to the flow field described for the upper ocean, including the cyclonic feature
14 described for the region between the Cape Verde Islands and Africa.

15 Mesoscale eddies are the main source of lateral oxygen supply to OMZs. Hahn et al. (2014)
16 found that the eddy-driven meridional oxygen supply is the dominant term in the oxygen
17 balance at the core depth of the OMZ. Moreover, energetic eddies, generated near the eastern
18 boundary, carry oxygen anomalies westward, eventually dissipate and may modify the
19 oxygen distribution in the OMZs. In the ETNA region such eddies are observed mainly on the
20 poleward side of the OMZs (Chelton et al., 2011). In the Eastern Tropical South Pacific
21 (ETSP) eddies are strong, particularly affecting the flow and property fields of the upper
22 several hundred meters of the ocean (e.g. Stramma et al., 2013; Czeschel et al., 2015). Three
23 types of eddies have been identified in both regions: cyclonic, anticyclonic and mode water
24 eddies (e.g. McGillicuddy Jr. et al., 2007). Cyclonic eddies have an uplift of isopycnals,
25 anticyclonic eddies a downward shift of isopycnals while mode water eddies derive their
26 name from a thick lens of water that deepens the main pycnocline while shoaling the seasonal
27 pycnocline. Mode water eddies show an uplift of the isopycnals near the surface, a downward
28 shift of isopycnals below the surface layer and the direction of rotation is the same as of
29 anticyclonic eddies.

30 To investigate time integrated diapycnal fluxes in the upper boundary of the ETNA OMZ a
31 tracer release experiment (GUTRE, Guinea Upwelling Tracer Release Experiment) was
32 performed by releasing 92 kg of CF₃SF₅ (Ho et al., 2008) between 24 and 28 April 2008 at

1 ~8°N, 23°W on the density surface $\sigma_\theta = 26.88 \text{ kg m}^{-3}$ at a depth of about 330 m (Banyte et al.,
2 2012). Three profiling floats with oxygen sensors were deployed during the tracer release.
3 Subsequently three main cruises were carried out 7, 20 and 30 months after the tracer release
4 to investigate the spreading of the tracer (Banyte et al., 2012, 2013).

5 We use the three floats deployed at the tracer release site and hydrographic measurements
6 from a cruise seven months after the tracer release, two cruise legs 20 months after the tracer
7 release in November and December 2009 and one cruise leg about 30 months after the tracer
8 release to investigate the flow field of the OMZ and eddy signatures in the ETNA. We
9 compare the float trajectories with the large scale circulation and the spreading of the tracer,
10 compare the observations with a data assimilation model and use recent measurements to
11 extend an oxygen change time series. While previous studies focused on the 23°W section or
12 on hydrographic and current meter measurements from single cruises, here we combine
13 different measurements particularly including float and tracer measurements to investigate the
14 flow field and oxygen distribution of the OMZ in the ETNA varying on intraseasonal and
15 seasonal time scales.

16

17 **2 Data sets**

18 **2.1 Shipboard measurements**

19 Cruises M80/1 and M80/2 (Fig. 1) on the German research vessel R/V *Meteor* took place in
20 November and December 2009 to investigate factors that control the intensity and areal extent
21 of the OMZ in the eastern tropical North Atlantic Ocean. The first leg (M80/1; 26 October to
22 23 November 2009; Mindelo-Mindelo (Cape Verde Islands); referred to as November 2009 in
23 the following) reoccupied a section along 23°W that had been measured before and
24 afterwards several times while the second leg (M80/2; 26 November to 22 December 2009;
25 Mindelo-Dakar, referred to as December 2009 in the following) was focussed on the GUTRE
26 spatial survey of the OMZ with tracer measurements from samples from a conventional CTD
27 rosette.

28 Two shipboard ADCP systems were used to record ocean velocity in November/December
29 2009: an RDI OceanSurveyor 75 kHz ADCP provided the velocity distribution to about 600
30 m depth, while a 38 kHz ADCP provided velocity profiles down to about 1000 m depth. In
31 November 2009 a 75 kHz ADCP was used on the 23°W section on the southward section and

1 a 38 kHz ADCP was used on the northward return leg. On M80/2 in December 2009 only the
2 75 kHz ADCP was used for current measurements.

3 A Seabird CTD system with a GO rosette with 24 10 L-water bottles was used for water
4 profiling and discrete water sampling on both cruises. The CTD system was used with double
5 sensors for temperature, conductivity (salinity) and oxygen. The CTD oxygen sensors were
6 calibrated with oxygen measurements obtained from discrete samples from the rosette
7 applying the classical Winkler titration method, using a non-electronic titration stand
8 (Winkler, 1888; Hansen, 1999). The precision of the oxygen titration determined during
9 cruise M80/1 in November 2009 was $\pm 0.34 \mu\text{mol kg}^{-1}$. The uncertainty of the CTD oxygen
10 sensor calibration was determined with an RMS of $\pm 1.28 \mu\text{mol kg}^{-1}$ in November 2009 and
11 $\pm 0.93 \mu\text{mol kg}^{-1}$ in December 2009.

12 While the two cruise legs in late 2009 are used to describe the general hydrographic and
13 velocity distribution in the OMZ region, two other tracer survey cruises (Fig. 1) are included
14 to investigate changes of the tracer distribution. A research cruise on the R/V *Merian*
15 (MSM10/1) was conducted 7 months after the tracer deployment leaving Ponta Delgada
16 (Azores) on 31 October 2008, with similar equipment as on the cruise in December 2009.
17 After a survey around the region of the tracer deployment location the cruise ended in
18 Mindelo on 6 December 2008.

19 An R/V *Meteor* cruise (M83/1) was carried out between 14 October 2010 (Las Palmas, Spain)
20 and 13 November 2010 (Mindelo, Cape Verde Islands) again with similar equipment as on
21 the cruise in December 2009. Due to some technical problems no ADCP data are available for
22 a short period at the southwestern sections of the cruise track. Four zonal sections were made
23 toward the African continent at about 15, 12.5, 10 and 8°N and the ADCP sections are used to
24 investigate the meridional current field. Furthermore, the ADCP and tracer measurements are
25 used for a comparison with the distribution in December 2009.

26 In addition to oxygen measurements along 23°W on MSM10/1 in November/December 2008,
27 M80/1 in November 2009 and M83/1 in October/November 2010 several other recent cruises
28 are used to extend the historical oxygen time series. The cruises used to extend the time series
29 from 1960 to 2007 (Stramma et al., 2008a) to the year 2014 in the region 10-14°N, 20-30°W
30 are a L'Atalante cruise (GEOMAR-4) in March 2008, a Merian cruise (MSM18/3) in June
31 2011, a Meteor cruise (M97) in June 2013 and a Meteor cruise (M106) in April 2014.

1

2 **2.2 Satellite and float data**

3 Aviso satellite derived altimeter sea surface height anomaly data (SSHA) were used to define
4 the general background distribution of the surface circulation and to identify possible eddy
5 signatures. The SSHA data used in this study are delayed time products and combine
6 available data from all satellites. The data are resampled on a regular $0.25^{\circ} \times 0.25^{\circ}$ grid and are
7 calculated with respect to a seven-year mean (<http://www.aviso.oceanobs.com>).

8 The oxygen climatological fields are taken from the CSIRO Atlas of Regional Seas (CARS)
9 2009 digital climatology (Ridgway et al., 2002) with a 0.5° geographical resolution.

10 Three profiling Argo floats with Aanderaa oxygen optode sensors were deployed on 24 and
11 26 April 2008 at about $\sim 8^{\circ}\text{N}$, 23°W at the tracer release site with parking depths at 330, 350
12 and 400 m depth, named here f330, f350 and f400 (Table 1). The cycling interval is 10 days
13 for the floats at 330 and 350 m depth and 7 days for the float at 400 m depth. The floats at 350
14 and 400 m depth were deployed at the same location and at the same time, however the first
15 data recording started 1 to 46 days after deployment (Table 1). The life-time of the floats was
16 1.6, 3.7 and 4 years. The floats drifting at 330 to 400 m were targeted to analyze the tracer
17 spreading behavior near the core of the OMZ of the eastern North Atlantic Ocean. The
18 shallow parking depth of our floats is different compared to most floats of the Argo project,
19 which have a parking depth typically between 1000 and 1500 m depth and can be used to
20 describe the flow field in the deep depths layers (e.g. Cravatte et al., 2012). In December 2009
21 CTD profiles were taken in the area of the three floats and a comparison of oxygen profiles
22 shows good agreement in the surface layer and below the OMZ while in the OMZ some
23 differences exist most likely due to time-space differences. The 100 to 800 m mean oxygen
24 difference at 1 m steps between float f330 and a CTD-oxygen profile 3 days and 68 km apart
25 result in a difference of $-1.1 \mu\text{mol kg}^{-1}$ (a negative value for higher CTD oxygen), for float
26 f350 0 days and 36 km apart of $-0.2 \mu\text{mol kg}^{-1}$ (Fig. S1 in the Supplement) while for float
27 f400 oxygen profiles are missing for the period of the ship cruise. In summary, the float
28 oxygen measurements seem to be accurate after ~ 20 months operating time and possibly
29 throughout the lifetime of the floats.

30

31 **2.3 The SODA assimilation model**

1 SODA (Simple Ocean Data Assimilation) combines Los Alamos's POP (Parallel Ocean
2 Program) model with a sequential estimation data assimilation method (Carton et al., 2000;
3 Carton and Giese, 2008). The SODA version used for this paper (SODA 2.2.4) is similar to
4 earlier versions but with some important differences as described by Giese and Ray (2011).
5 For the remainder of the paper, we use SODA to refer to the SODA 2.2.4 product mapped
6 onto a uniform $0.5^\circ \times 0.5^\circ \times 40$ level grid (22 vertical levels in the upper 800 m depth).
7 Version 2.2.4 is forced by the 20CRv2 (20th Century Reanalysis version 2) (Compo et al.,
8 2011) wind stresses from 1871 to 2011 and uses atmospheric variables from 20CRv2 for the
9 calculation of surface heat and freshwater fluxes using bulk formulae. As in earlier versions,
10 SODA 2.2.4 assimilates all available data from hydrographic stations, expandable
11 bathythermographs, and floats, but does not use satellite altimetry. In this version,
12 hydrographic observations come from WOD09 (Boyer et al., 2009) using their standard level
13 temperature and salinity data. Thus these data have been corrected for the drop-rate error as
14 described by Levitus et al. (2009). Experiments with SODA show that applying the Levitus et
15 al. (2009) drop-rate correction reduces much of the decadal variability observed in both the
16 hydrographic observations and ocean reanalysis (Giese et al., 2011). The model results from
17 this set-up will be referred to as SODA model in the following.

18 The SODA model mean flow field for the period 2001 to 2010 for the layers 50 to 200 m and
19 200 to 400 m (Fig. 2) will be used as background information of the mean circulation in the
20 tropical eastern North Atlantic and for a comparison with the observed circulation features.
21 The model results for October/November 2010 (Fig. S2) can be used for a detailed
22 comparison to the October/November 2010 Meteor M83/1 measurements and the 2001 to
23 2010 mean velocity section at 23°W (Fig. 3) to investigate the depth distribution of the zonal
24 currents. The SODA model velocity difference between the year 2009 and the mean 2001 to
25 2010 velocity at 23°W and 10°N (Fig. S3) is used to check the 2009 anomaly in the Guinea
26 Dome.

27

28 **3 Results**

29 **3.1 Circulation in the low oxygen zone of the eastern tropical North Atlantic** 30 **Ocean**

31 The SSHA along 23°W on 18 November 2009 reflects the large-scale upper ocean velocity in
32 November 2009 (Fig. 4) with westward flow south of $\sim 5^\circ\text{N}$, eastward flow between 5 and

1 11°N and both west- and eastward flow components north of 11°N. The zonal velocity and
2 oxygen distribution in November 2009 along 23°W (Fig. 4) shows enhanced oxygen content
3 in the upper 250 to 300 m in the region of the two eastward NECC bands centered at about
4 4.5 and 8°N. The westward flow component at 6-7°N carries oxygen-poor water in the 100 to
5 200 m depth range. Westward flow at 12°N north of the center of the Guinea Dome (flow
6 reversal at 23°W at about 9°N in November 2009) recirculates oxygen-rich nNECC water
7 westwards at a depth of about 200 m, but carries oxygen-poor water westward in the OMZ
8 depth range (Fig. 4). The strongest oxygen minimum is located slightly above the 27.1 kg m⁻³
9 isopycnal layer, hence in the lower part of the Central Water (σ_{θ} = 25.8 and 27.1 kg m⁻³).

10 As seen in the November 2009 velocity distribution at 23°W, most current bands from the
11 near surface layer reach down to the OMZ core, especially the nNECC and the current bands
12 north of 10°N. The NECC at about 4.5°N is sometimes weaker and can even reverse to
13 westward flow below 300 m in the mean of several ADCP sections along 23°W (Brandt et al.,
14 2015; their Fig. 6). The eastward flow component south of the Cape Verde Islands called
15 Cape Verde Current by Peña-Izquierdo et al. (2015) centered at about 13°N at 23°W is weak
16 in November 2009 compared to the mean 23°W section (Brandt et al., 2015), nevertheless
17 reaches to at least 500 m depth. A similar flow direction in the upper ocean, as well as in the
18 OMZ layer, can be seen in the mean velocity distribution at 23°W and at 18°N east of 26°W
19 (Brandt et al., 2015; their Figs. 6 and 5), and in single velocity sections at 38°W (Urbano et
20 al., 2008; their Fig. 5), at 28°W (Stramma et al., 2008b; their Fig. 3) and at 11°N east of 22°W
21 (Stramma et al. 2005; their Fig. 11). Also the SODA 10-year mean velocity section at 23°W
22 (Fig. 3) shows the current bands except for the NECC at 4-5°N to reach to depth of 800 m.
23 The eastward flow south of the Cape Verde Islands is found in two cores located at about 12
24 and 14°N, similar as in the mean 23°W section (Brandt et al., 2015).

25 Except for the sections at 18 and 11°N there is not much information about the meridional
26 circulation at the OMZ layer and schematics often show only zonal flow components (e.g.
27 Brandt et al., 2015). Mittelstaedt (1983) showed that the cyclonic circulation cell in the
28 surface layer between the Cape Verde Islands and the African continent varies in size related
29 to the wind. The circulation has a weaker southward extent of the center of the cyclonic cell
30 in winter, reaching to about 15°N, in contrast to summer when it reaches to about 12°N.
31 Several sections near the African shelf were made in October 2010 (Fig. 5). At 15°N the
32 northward currents are strongest in the upper ocean, but most of the currents reach into the

1 OMZ layer. The SODA results from the 50 to 200 m layer and 200 to 400 m layer for the 10-
2 year mean circulation (Fig. 2) as well as for October/November 2010 (Fig. S2) show that this
3 is the southernmost part of the cyclonic circulation cell, with a flow contribution from the
4 west. At 12.5°N a cyclonic circulation cell with a core of southward flow at 19.5 to 20°W
5 and northward flow at 18 to 18.5°W (Fig. 5) was observed. The SODA velocities for the same
6 time (Fig. S2) indicate a second cyclonic cell located south of the cyclonic cell east of the
7 Cape Verde Islands in October/November 2010. The northward flow at 10°N at 18.5 to 19°W
8 and the southward flow west of 19.5°W seem to be the southern component of the cyclonic
9 cell observed at 12.5°N. The southward flow at about 21°W at 8°N (Fig. 5) could be a
10 recirculation branch of the nNECC as in the SODA flow field during October/November
11 2010 (Fig. S2, bottom) while the meridional flow components in the 8°N section east of 20°W
12 are quite weak. The second cyclonic cell located at about 12.5°N in October 2010 is an
13 exceptional situation, in the long-term mean the SODA velocities show a westward excursion
14 of the 200 to 400 m layer in this region. The existing zonal and meridional ADCP velocity
15 sections and the SODA velocity distribution show that flow in the low oxygen layer generally
16 mirrors the circulation of the near-surface layer.

17 The observed velocity distribution and measured oxygen values at 350 m depth in November
18 and December 2009 (Fig. 6) are quite variable especially on the equatorial side of the OMZ.
19 This is in agreement with the SODA flow field for October/November 2010 (Fig. S2) where
20 the flow is also highly variable. The oxygen distribution measured in November and
21 December 2009 at 350 m near 30°W between 8 and 10°N is generally higher compared to the
22 CARS climatology, however it is lower for most measurements east of 30°W. In the OMZ
23 north of 10°N measured oxygen levels are mostly lower at the boundaries of the OMZ than
24 climatological ones. Near the climatological OMZ core at about 12.5°N, 20°W, measured
25 oxygen values are higher just south of 12.5°N, and are either in agreement with or lower than
26 climatology to the north of 12.5°N. The lowest oxygen concentrations in the climatology at
27 350 m depth are located in the region of the cyclonic circulation cells between the Cape Verde
28 Islands and Africa indicative of reduced water renewal in these cyclonic circulation cells. The
29 float at 350 m depth (f350) moved westward north of the OMZ core in December 2009 (Fig.
30 6) and measured oxygen values from this float agree well with climatology. The northward
31 shift of the float at about 12.5°N, 21°W seems to be connected to an increased SSHA gradient
32 connected to a strengthening of the high and low SSHA at this location (Supplement Movie
33 M1 for 1 December 2009). Because the NECC is weaker in boreal winter the variable flow

1 components south of 10°N might be related to a weakened NECC, although the oxygen
2 distribution indicates an oxygen supply to the OMZ, especially west of 20°W. In boreal
3 winter the core of the Guinea Dome should be located at 10.5°N, 22°W, however according to
4 the velocity distribution at 23°W the Guinea Dome center seems to be still south of 10.5°N.
5 While the nNECC south of the Guinea Dome is weak and variable, the westward flow with
6 low oxygen content north and east of the Guinea Dome center is more obvious (Fig. 6).

7 In general lower oxygen levels in late 2009 as compared to climatology dominate in the OMZ
8 region (Fig. 6), and is probably related to the observed trend of deoxygenation in the eastern
9 tropical North Atlantic (region 10-14°N, 20-30°W) from 1960 to 2007 (Stramma et al.,
10 2008a). The oxygen trend in the region 10-14°N, 20-30°W computed as by Stramma et al.
11 (2008a) was extended with additional shipboard measurements along 23°W from cruises in
12 November 2008, November 2009 and October 2010 as well as from March 2008, June 2011,
13 June 2013 and April 2014. Several measurements in the area within a calendar year were
14 averaged to derive a single oxygen value per year. Some improvements in the computation
15 method were made, e.g. no interpolation was made for years without measurements. The
16 computed trend confirms the deoxygenation trend in the upper OMZ layer at 100 to 300 m
17 depth up to the year 2014 with a linear trend of $-0.49 \pm 0.16 \mu\text{mol kg}^{-1} \text{ year}^{-1}$ (Fig. 7a). A
18 similar oxygen decrease for recent years was described for the 150 to 300 m layer (Brandt et
19 al., 2015). For the deeper layer (300 to 700 m) the linear trend of $-0.34 \pm 0.13 \mu\text{mol kg}^{-1}$
20 year^{-1} up to the year 2007 (Stramma et al., 2008a) slightly increased to $-0.38 \pm 0.09 \mu\text{mol kg}^{-1}$
21 year^{-1} when extended to 2014 (Fig. 7b). The 300 to 700 m layer shows an increase in oxygen
22 in recent years (Fig. 7b) as described by Brandt et al. (2015), however, the measured lower
23 oxygen content in recent years than predicted from the 1960 to 2007 trend nevertheless leads
24 to the slightly increased trend.

25

26

27 **3.2 Float measurements**

28

29 The three floats deployed with a parking depth at 330 m, 350 m and 400 m at the same
30 location (Table 1) show quite different flow paths (Fig. 8). Especially the floats drifting at 350
31 and 400 m depth that were deployed at the same location within the eastward flowing nNECC

1 (Fig. 4) and same time show quite different flow trajectories (Fig. 8). According to the mean
2 velocity distribution at 23°W (e.g. Brandt et al., 2010, 2015) the deployment site at about
3 8°10'N is in a region of strong eastward flow in the upper 100-200 m of the nNECC, weak
4 eastward flow below 200 m depth and weak westward flow at 100 to 1000 m depth to the
5 south of about 7.5-8°N.

6 All floats stayed first in the nNECC or the region south of the nNECC. The float at 330 m
7 depth (f330) first moved south-eastward through 22 December 2008 and then moved south-
8 westward through 31 May 2009 before it started to move north-eastward crossing the nNECC
9 region to 10°N where the float stopped working in late December 2009. The float at 350 m
10 depth moved north-eastward through 25 October 2008 then reversed direction to westward
11 and northward and it crossed 10°N on 10 August 2009. The float at 400 m depth stayed its
12 entire lifetime of 3.75 years in the region between 6 and 9°N between the NEUC and the
13 nNECC. Annual reversals were strongest in the western part of the f400 track when it moved
14 south-eastward through 9 December 2008 and then westward up to 9 June 2009, before it
15 mainly moved eastward. The mean eastward velocity between the deployment location and
16 the location of the final data transmission was 0.6 cm s^{-1} .

17 In the region 6 to 9°N and 25 to 20°W the floats moved eastward between about May and
18 November and westward in late winter and spring. In an early investigation of the NECC a
19 strong NECC with two eastward cores in July to September was observed, while generally
20 near surface westward flow is found during March to May (Richardson and Reverdin 1987).
21 When separating the surface drift of the three floats for the time at the surface to transmit data
22 and for the subsurface drift, the surface drift was eastward from mid-May to December and
23 westward from January to mid-May with an annual mean eastward velocity of 9 cm s^{-1} , while
24 the subsurface flow was weak year-round with a mean eastward component of 0.02 cm s^{-1} .
25 Hence the eastward shift of the floats, especially of float f400, is mainly due to the seasonal
26 surface signal of the NECC and not caused by an eastward drift at the parking depth near the
27 OMZ core. This is in agreement with ADCP velocity observations that show the flow below
28 the NECC is weak, or even westward, as well as with the velocity field of the 10-year mean
29 SODA data (Fig. 2).

30 The floats were used to determine the velocity at the surface and at the parking depth. The
31 float at 400 m depth, with a mean eastward flow of 0.6 cm s^{-1} , stayed at the surface for about
32 seven hours of the seven day diving cycle which leads to an eastward flow component of 14.4

1 cm s⁻¹ if the float had stayed at the surface for the entire time. This value is higher than the 9
2 cm s⁻¹ eastward flow computed for the three floats at the surface when they were located
3 south of 9°N but weaker compared to the mean geostrophic eastward component of 20 cm s⁻¹
4 at 6°N between 15 and 40°W for the period 1993 to 2009 computed for the core position of
5 the NECC (Hormann et al., 2012).

6 The complete record of spreading and vertical oxygen distribution of float f350 (Fig. 9) shows
7 northward movement through the low oxygen layer of the OMZ into the oxygenated North
8 Atlantic subtropical gyre. This float stayed south of 10°N for more than a year, but in July
9 2009 it suddenly moved rapidly to the north. At the same time a strong SSHA appeared in the
10 region between the two NECC branches (Supplement Movie M1), which may have triggered
11 the northward displacement of the float. The float drifted northward from 10°N on 10 August
12 2009 to 14.9°N on 28 March 2010 and crossed the OMZ with a mean speed of about 2.75 cm
13 s⁻¹. While crossing the OMZ the drift of this float had a weak south-westward component,
14 hence the entire northward drift across the OMZ of the float took place at the subsurface layer
15 near the parking depth of 350 m. Unfortunately the float f330 stopped operating at the end of
16 2009 when it moved northward into the OMZ, however the f330 northward movement
17 occurred in the subsurface while the surface drift was toward the east.

18 The float at 350 m moved in a cyclonic track around the core of the OMZ. This cyclonic
19 northward drift in summer 2009 could be caused by an anomalously strong Guinea Dome
20 event. The Guinea Dome strength is described as connected to the Atlantic Meridional Mode
21 (AMM; Doi et al., 2010). The AMM is a climate mode associated with the cross-equatorial
22 meridional gradient of the sea surface temperature (SST) anomaly in the tropical Atlantic
23 (Doi et al., 2010) with a negative AMM defined by a negative/positive SST anomaly in the
24 northern/southern hemisphere (Foltz et al., 2012). During the preconditioning phase of an
25 AMM in late fall of the previous year the dome is weaker with enhanced mixed layer depth
26 exists in the dome region. As a result a positive SST anomaly appears in early winter. In the
27 following spring the wind-evaporation-SST positive feedback is associated with anomalous
28 northward migration of the inter-tropical convergence zone (ITCZ) leading to stronger Ekman
29 upwelling and colder subsurface temperatures in the dome region in summer (Doi et al., 2010)
30 and hence to a stronger circulation around the Guinea Dome. Using monthly mean SST
31 values of the AMM
32 (<http://www.esrl.noaa.gov/psd/data/timeseries/monthly/AMM/ammsst.data>) the temperature

1 was about 2°C higher during the preconditioning time from September to December 2008 and
2 cooler, with an anomaly of up to -5.3°C, in the period from February to June 2009. The
3 SODA velocity anomalies for 2009 at 23°W and 10°N (Fig. S3) confirm a stronger cyclonic
4 circulation around the Guinea Dome located at about 10°N, 23°W below the mixed layer. A
5 strong AMM event in 2009 was also described using measurements at two moorings located
6 at 4°N, 23°W and 12°N, 23°W by Foltz et al. (2012).

7 The anticyclonic movement of the float f350 in the OMZ at about 22°W 12 to 14°N (Fig. S4)
8 between 29 November 2009 and February 2010 shows a weak upward shift of the isopycnals
9 in the upper 50 m and weak downward shift of the isopycnals down to 350 m in December
10 2009. This is a weak signature of an anticyclonic mode water eddy. Most striking during this
11 time is the shallow oxygen minimum at 100 m depth between 29 November 2009 and 28
12 January 2010. Animation of the float location relative to the SSHa (Supplement Movie M1)
13 indicates some anticyclonic structure in the SSHa field, although positive SSHa move
14 rapidly to the west. North of the Cape Verde Islands low oxygen layers just below the mixed
15 layer seem to be created by cyclonic and anticyclonic mode water eddies (Karstensen et al.,
16 2015). The low oxygen concentration at 50 to 100 m depth in November 2009 might be
17 created by a mode water eddy although the second oxygen minimum in January does not
18 show a mode water density structure. A comparison of low oxygen periods near 100 m depth
19 or the higher oxygen content in the upper 200 m (Sup. Fig. S4) with the SSHa (Supplement
20 Movie M1) and the density distribution did not provide convincing evidence for eddy
21 signatures. Mode water eddies might have a weak surface SSHa signature and might be
22 missed in altimeter data, however as only few mode water type signatures were observed in
23 the measurements a large influence on the OMZ by mode water eddies is not expected.

24 The high oxygen layer that extends down to 200 m present before December 2009
25 gradually reduces along the northward float track in the following months and oxygen of less
26 than 70 $\mu\text{mol kg}^{-1}$ dominates the region near the Cape Verde Islands through July 2011 (Fig.
27 9) although oxygen of more than 100 $\mu\text{mol kg}^{-1}$ down to 200 m was measured in August to
28 October 2010 and April to early June 2011. The SSHa movie shows a fast passage of a
29 SSHa maximum indicating an anticyclonic feature (Supplement Movie M1). The core of the
30 oxygen minimum was continuously near the isopycnal 27.0 kg m^{-3} through December 2011
31 and shifts deeper to 600 m depth and the isopycnal 27.15 kg m^{-3} at about 18°N (Fig. 9) when
32 drifting north-westward into the NEC of the subtropical gyre. The north-westward drift of the

1 float north of the Cape Verde Islands (Fig. 9) from mid-2011 to May 2012 is a combination of
2 a westward flow component at the surface and a north-westward flow component in the
3 subsurface.

4

5 **3.3 Tracer spreading**

6 The tracer distribution and float paths represent an integral effect of the velocity field since
7 deployment, while ADCP surveys yield velocity snapshots influenced by eddies, tides and
8 internal waves. Nevertheless it is interesting to investigate how different results obtained by
9 the various methods might be. Spreading of the tracer (CF_3SF_5) deployed in April 2008 at
10 $\sim 8^\circ\text{N}$, 23°W on the density surface $\sigma_\theta = 26.88 \text{ kg m}^{-3}$ at about 330 m depth through December
11 2009 was investigated using rosette water samples collected at CTD stations of Meteor cruise
12 M80/2 (Banyte et al., 2013). Close to the tracer deployment target density tracer samples were
13 taken with 10 m spacing. The spatially maximum tracer value distribution was close to the
14 target density in December 2009 in relation to the ADCP velocity at the isopycnal $\sigma_\theta = 26.88$
15 kg m^{-3} and shows the tracer spreading in the ETNA (Fig. 10). The mean depth of the
16 maximum tracer concentration in December 2009 was 314 m. The largest tracer
17 concentrations were measured east of the deployment location at locations where the ADCP
18 velocity in December 2009 is toward the east in agreement with the mean eastward flow
19 between $\sim 7.5^\circ\text{N}$ and 9.5°N in the mean 23°W section (Brandt et al., 2015; their Fig. 6c). This
20 eastward spreading differs in comparison to our floats, which showed only weak eastward
21 velocity components at the subsurface layer. Investigating a larger float data set it is found
22 that the NECC eastward component is strongest at the surface. However, at 200 and 1000 m
23 depth levels there is also an eastward component (Rosell-Fieschi et al., 2015; their Fig. 7).
24 Our three floats may underestimate the mean eastward spreading of the tracer during periods
25 of westward flow component. One reason could be that the floats drift at a greater depth than
26 the mean tracer maximum depth at 314 m. The mean zonal velocity components from about
27 25 ADCP velocity ship sections at about 23°W (Brandt et al., 2015) show that below 200 m
28 depth there is a westward flow between 5.5 and 7.5°N leading to an enhanced westward
29 component for the floats when located south of 7.5°N .

30 To the south of the deployment location large tracer signals are found at about 6°N where the
31 ADCP velocities are directed westward and indicate recirculation of the lower part of the

1 NECC or the nNECC to the west or the influence of waves in November/December 2009. The
2 higher tracer concentrations north of 9°N are connected to ADCP velocities directed
3 westward and indicate a cyclonic flow around the Guinea Dome in agreement with the
4 westward flow between 9.5 and 12°N in the mean 23°W section (Brandt et al., 2015). Also
5 near 12°N , 21°W , where the float showed a westward flow at the time of the ship passage, the
6 ADCP velocity is directed westward with higher tracer signals.

7 The three floats deployed together with the tracer in April 2008 at the same location ($\sim 8^{\circ}\text{N}$,
8 23°W) and similar depth levels show paths in the region where the largest tracer signals were
9 observed (Figure 10). At 6 to 7°N the floats have a westward drift and the tracer signal is
10 large. The ADCP flow field at 6 to 7°N in December 2009 is directed southwestward, hence
11 the flow field in December 2009 agrees with the long-term signal of the tracer spreading.
12 North of 8°N the floats have mainly an eastward drift and the strongest tracer signals were
13 measured east of the deployment site, although further east than the floats moved during the
14 time period up to December 2009. However, the float at 400 m depth had a stronger eastward
15 component after December 2009 and reached east to almost 15°W , similar to where the large
16 tracer signal was measured. The float at 350 m that experienced the cyclonic circulation
17 around the Guinea Dome was close to the ship section near 13°N , 21°W in December 2009
18 with a westward component in agreement with the westward ADCP velocity component and
19 the enhanced tracer signal in the ship measurements.

20 The tracer distribution in October/November 2010 (Fig. 11), nearly one year later, shows
21 reduced maximum tracer concentrations due to further spreading of the tracer by lateral
22 diffusivity or mesoscale eddy diffusion (Banyte et al., 2013; Gnanadesikan et al., 2013; Hahn
23 et al., 2014) and diapycnal mixing (Banyte et al., 2012). Nevertheless, the maximum tracer
24 signal confirms the circulation features described for December 2009. The largest tracer
25 signals are found in the NECC current bands near the tracer deployment location. Float f400
26 (Fig. 11; light grey) circled in this region starting at the time of deployment and is located at
27 7.49°N , 19.2°W on 9 November 2010, close to a CTD station, with enhanced tracer load
28 transported eastward. South of the NECC (south of 4°N) the tracer signal is almost zero,
29 hence there appeared to be no exchange between the NECC and the equatorial region. Tracer
30 spreading around the Guinea Dome and along the float track of float f350 is visible. Float
31 f350 was located just northeast of the Cape Verde Islands in November 2010.

1 The tracer signal between the Cape Verde Islands and Africa along 15°N is near zero except
2 for a weak signal at 21 to 22°W, indicating a weak northward flow component just east of the
3 Cape Verde Islands. Such a northward flow component is consistent with the northward drift
4 of float f350 and the flow field in October/November 2010. In an earlier description of the
5 subsurface flow in the eastern tropical North Atlantic (Stramma et al., 2008b) it was
6 mentioned that oxygen-poor water east of the Cape Verde Islands is trapped in a cyclonic
7 circulation cell, which agrees with the lowest oxygen concentrations at 350 m in the
8 climatology (Fig. 6) and the observation that the tracer signal at 15°N east of the Cape Verde
9 Islands is very low. Instead of the mean cyclonic circulation reaching from the surface to the
10 OMZ layer as observed in the mean ADCP section at 18°N (Brandt et al., 2015),
11 measurements in November 2008 between Cape Verde and the Canary Islands show cyclonic
12 flow in the upper ocean but an anticyclonic circulation cell of 0.8 Sv for the density layer $\sigma_\theta =$
13 $26.85\text{-}27.1 \text{ kg m}^{-3}$ in $\sim 300\text{-}500$ m depth (Peña-Izquierdo et al., 2012). According to the
14 SODA model water from the Guinea Dome region is transported northeastward just east of
15 the Cape Verde Islands which could explain the measurements of weak tracer transport west
16 of the African shelf and possibly slightly enhanced tracer transport east of the Cape Verde
17 Islands.

18 In November/December 2008, just seven months after the deployment the tracer was located
19 closer to the deployment region and the ship survey was carried out in a smaller region near
20 the deployment site (Fig. S5). The maximum tracer concentration seven months after the
21 deployment in November 2008 are up to 230 fm kg^{-1} , much larger than the maximum tracer
22 values of 6.5 fm kg^{-1} in November/December 2009 or 3.9 fm kg^{-1} in October/November 2010.
23 The strongest tracer values were observed northeast of the deployment site with the highest
24 values at about 9°N, 20°W and the float f350 also drifted to this region. The two other floats
25 drifted toward the southeast and in this region a westward flow component with enhanced
26 tracer values is present. Some of the tracer drifted around the Guinea Dome and spread
27 westward at 11°N.

28

29 **4 Discussion and conclusion**

30 In this study the eastern tropical North Atlantic was investigated with a focus on the upper
31 OMZ using SODA model results, ship surveys in November/December 2008,
32 November/December 2009 and October/November 2010, and 3 floats with oxygen sensors

1 which was compared to the spreading of a tracer released in April 2008 at $\sim 8^{\circ}\text{N}$, 23°W .
2 Oxygen sensors of two floats stayed well calibrated for ~ 20 months within $1 \mu\text{mol kg}^{-1}$ for the
3 100 to 800 m depth layer in comparison to near CTD-oxygen profiles.

4 The shipboard oxygen observations in 2008, 2009 and 2010 augmented by 4 other oxygen
5 cruise measurements are used to determine the deoxygenation trend near the Guinea Dome in
6 the upper OMZ through the year 2014. The linear trend for oxygen since the 1960's in a layer
7 from 100 to 300 m continues in recent years with a trend of $-0.49 \pm 0.16 \mu\text{mol kg}^{-1} \text{ year}^{-1}$.
8 For a deeper layer from 300 to 700 m a weaker, but also continuous, linear trend of $-0.38 \pm$
9 $0.09 \mu\text{mol kg}^{-1} \text{ year}^{-1}$ exists despite an oxygen increase in this layer in recent years. This trend
10 is similar to the trend of $-0.34 \pm 0.13 \mu\text{mol kg}^{-1} \text{ year}^{-1}$ for this layer calculated up to the year
11 2006 (Stramma et al., 2008a).

12 Strong eddies as observed in the eastern tropical Pacific off Peru could not be detected in the
13 data set used here south of the Cape Verde Islands. Observations of mesoscale eddies in the
14 eastern tropical Atlantic show less eddy activity than in the eastern tropical Pacific (Chelton et
15 al., 2011). Some indication of eddy activity is seen in the float time series and eddies seem to
16 affect and blur the ADCP flow fields, however the eddy signal in the ETNA is not as strong
17 and not as deep-reaching as in the eastern Pacific off Peru (Stramma et al., 2013). The
18 cyclonic and anticyclonic features in the SSHA field move westward rapidly and the floats
19 stayed in these features for only a few days, different from observations of the eastern tropical
20 South Pacific, where the floats followed the SSHA anomalies for weeks (Czeschel et al.,
21 2015). Eddy activity might be stronger north of the Cape Verde Islands where enhanced eddy
22 activity is visible in satellite data (Chelton et al., 2011) and observations of very low oxygen
23 values were observed in long-lived cyclonic and anticyclonic mode water eddies (Karstensen
24 et al., 2015).

25 Splitting of the drift components of float tracks at the surface and at subsurface depth shows
26 that the depth of largest float drift strongly depends on the geographic location. Eastward
27 spreading of the floats in the NECC region south of 9°N was governed by a drift at the
28 surface, while at the parking depth at 330 m to 400 m only weak zonal flow influenced the
29 float path. The zonal flow component showed the influence of the seasonal signal of the
30 NECC which, modulated by interannual variations, is related to long Rossby waves (Hormann
31 et al., 2012). The measured eastward surface flow component of the float with a parking depth
32 at 400 m of 14.4 cm s^{-1} was a little less than the estimated mean NECC core velocity for the

1 region 15 to 40°W of 20 cm s⁻¹ (Hormann et al., 2012). This velocity difference might be
2 caused by higher NECC velocities in the western Atlantic and the location of the float not
3 propagating in the core of the NECC. In the OMZ region between the Cape Verde Islands and
4 ~9°N the northward drift takes place near the parking depth at 350 m. The northwestward
5 drift of the float into the subtropical gyre north of the Cape Verde Islands was due to a
6 combination of westward surface drift and subsurface northwestward drift. The measured
7 oxygen and velocity distribution at 350 m depth slightly above the core of the OMZ shows the
8 signature of a cyclonic flow around about 10°N, 23°W, hence the near-surface circulation of
9 the Guinea Dome reaches down to this subsurface layer. The fast northward progression of
10 the float at 350 m depth in 2009 seems to be connected to a period of strong Guinea Dome
11 caused by an Atlantic Meridional Mode event (e.g. Doi et al., 2010). In addition to the
12 meridional mode event, the combination with a zonal mode event is responsible for a
13 northward drift of the NECC core and a current strengthening (Hormann et al., 2012) which
14 seems to be responsible for the float drift from the NECC region to the Guinea Dome region
15 in boreal summer 2009. Climatology for November/December at 350 m (Fig. 6) shows the
16 oxygen minimum at about 12.5°N, 20°W, northeast of the cyclonic flow of the Guinea Dome
17 region. This oxygen minimum is located in the anticyclonic circulation south of the Cape
18 Verde Islands indicating weak water renewal in this circulation cell at this depth.

19 In December 2009 the distribution of the tracer deployed at ~8°N, 23°W in April 2008 at the
20 isopycnal $\sigma_\theta = 26.88 \text{ kg m}^{-3}$ shows the integral effect of over 20 months of the flow
21 components of the upper OMZ. The largest tracer signal was observed east of the deployment
22 region, which is in agreement with the nearly barotropic eastward flow between 7.5 and 9.5°N
23 found in the mean 23°W section by Brandt et al. (2015). Large tracer signals southeast and
24 southwest of the deployment region show the recirculation paths south of the nNECC. Finally
25 large tracer signals north of the deployment region show cyclonic flow around the Guinea
26 Dome. The three floats were deployed together with the tracer in April 2008 at the same
27 location and the spreading paths of the floats are in good agreement with the spreading of the
28 tracer signal.

29 The different observations as well as the SODA model results show a weak mean flow field
30 (Fig. 12) by averaging the velocity field including seasonal and short-term variability. The
31 eastward flowing NECC is strong in the upper 250 m while the NEUC below is either weak
32 or flowing westward in agreement with the NECC flow fields derived by Peña-Izquierdo et al.

1 (2015). The nNECC reaches from the surface down to the OMZ layer and westward flow to
2 the south of the nNECC at subsurface layers is clearly visible in zonal velocity sections in
3 November 2009 (Fig. 5) and in the mean 23°W velocity sections (Brandt et al., 2010). The
4 velocity section at 23°W north of 10°N as well as velocity sections reported in literature
5 confirm that the near surface flow often reaches down to the low oxygen layer. The near-
6 surface cyclonic circulation cell east of the Cape Verde Islands described by Mittelstaedt
7 (1983) reaches down into the OMZ layer. The eastward flow south of the Cape Verde Islands,
8 named Cape Verde Current by Peña-Izquierdo et al. (2015), seems to be a permanent
9 eastward recirculation of the westward flow component of the Guinea Dome. The higher
10 NACW contribution compared to SACW in the lower Central Water layers south of the Cape
11 Verde Islands (e.g. Tomczak, 1984; Peña-Izquierdo et al., 2015) appears to be related to the
12 weak SACW inflow in the weak NEUC flow at about 5°N compared to strong SACW inflow
13 in the NECC above in the upper Central Water and the enhanced contribution of NACW
14 nNECC (Lumpkin and Garzoli, 2005).

15 In vector plots of snapshots of the horizontal flow field combined with oxygen and tracer
16 measurements (Figs. 6, 10 and 11) the mean large-scale circulation signal is obscured by the
17 meridional variability in the current bands as observed in the ship surveys and in the SODA
18 velocity field. It is further overlain by circulation changes caused by climate modes such as
19 the AMM and mesoscale variability. Despite large variability in snapshots of the ADCP-
20 derived flow field the different measurements used and combined here demonstrate that
21 circulation of the upper OMZ widely mirrors the near-surface circulation (Fig. 12).
22 Exceptional cases are the weak 200 to 400 m flow below the NECC and an enhanced
23 westward excursion of the 200 to 400 m flow north of the Guinea Dome at about 12°N.
24 Based on additional data sets a more detailed description of the OMZ flow field than that
25 presented by Stramma et al. (2008b) could be derived here. While the floats and tracer
26 spreading results enabled a better understanding of the circulation on a density or depth
27 surface, the mean three-dimensional flow field can only be derived by averaging repeated
28 hydrographic sections and thus removing the variability present in snapshots of the flow field
29 from single ship surveys.

30

31

1 **The Supplement related to this article is available online at DOI:10.5194/osd-12-1-2015-**
2 **supplement.**

3

4

5

6 **Acknowledgements.**

7 The Deutsche Forschungsgemeinschaft (DFG) provided support as part of the
8 “Sonderforschungsbereich 754: Climate-Biogeochemistry Interactions in the Tropical Ocean
9 and the Deutsche Bundesministerium für Bildung und Forschung (BMBF) as part of AWA
10 (01DG12073E). The altimeter data were produced by Ssalto/Duacs and distributed by Aviso
11 with support from Cnes. The Argo float profiles are available from
12 ftp://ftp.ifremer.fr/ifremer/argo/dac/coriolis. Partial funding for BSG was provided by NSF
13 from award ACI 1049075.

14

15

16

17

18 **References**

19 Banyte, D., Tanhua, T., Visbeck, M., Wallace, D. W. R., Karstensen, J., Krahnmann, G.,
20 Schneider, A., Stramma, L., and Dengler, M.: Diapycnal diffusivity at the upper boundary of
21 the tropical North Atlantic oxygen minimum zone, *J. Geophys. Res.*, 117, C09016,
22 doi:10.1029/2011JC007762, 2012.

23 Banyte, D., Visbeck, M., Tanhua, T., Fischer, T., Krahnmann, G., Karstensen, J.: Lateral
24 diffusivities from tracer release experiments in the tropical North Atlantic thermocline, *J.*
25 *Geophys. Res.*, 118, 1-15, doi10.1002/jgrc.20211, 2013.

26 Boyer, T. P., Antonov, J. I., Baranova, O. K., Garcia, H. E., Johnson, D. R., Locarnini, R. A.,
27 Mishonov, A. V. Seidov, D., Amolyar, I. V., and Zweng, M. M.: World Ocean Database 2009
28 [DVDs], Noaa Atlas Nesdis, vol. 66, edited by: S. Levitus, 216 pp, U.S. Govt. Print. Off.,
29 Washington, D.C. USA, 2009.

1 Brandt, P., Hormann, V., Körtzinger, A., Visbeck, M., Krahnemann, G., Stramma, L., Lumpkin,
2 R., and Schmid, C.: Changes in the ventilation of the oxygen minimum zone of the tropical
3 North Atlantic, *J. Phys. Oceanogr.*, 40, 1784-1801, doi:10.1175/2010JPO4301.1, 2010.

4 Brandt, P., Bange, H. W., Banyte, D., Dengler, M., Didwischus, S.-H., Fischer, T.,
5 Greatbatch, R. J., Hahn, J., Kanzow, T., Karstensen, J., Körtzinger, A., Krahnemann, G.,
6 Schmidtke, S., Stramma, L., Tanhua, T., and Visbeck, M.: On the role of circulation and
7 mixing in the ventilation of oxygen minimum zones with a focus on the eastern tropical North
8 Atlantic, *Biogeosciences*, 12, 489-512, doi:10.5194/bg-12-489-2015, 2015.

9 Carton, J. A. and Giese, B. S.: A reanalysis of ocean climate using SODA, *Mon. Weather*
10 *Rev.*, 136, 2999-3017, 2008.

11 Carton, J. A., Cherupin, G., Cao, X., and Giese, B. S.: A simple ocean data assimilation
12 analysis of the global upper ocean 1950-95. Part I: Methodology, *J. Phys. Oceanogr.*, 30, 294-
13 309, 2000.

14 Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear
15 mesoscale eddies, *Prog. Oceanogr.*, 91, 167-216, doi:10.1016/j.pocean.2011.01.002, 2011.

16 Chu, P. C., Ivanov, L. M., Melnichenko, O. V., Wells, N. C.: On long baroclinic Rossby
17 waves in the tropical North Atlantic observed from profiling floats, *J. Geophys. Res.*, 112,
18 C05032, doi:10.1029/2006JC003698, 2007.

19 Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason,
20 B. E., Vose, R. S., Rutledge, G., Bessemoulin, P., Brönnimann, S., Brunet, M., Crouthamel,
21 R. I., Grant, A. N., Groisman, P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J.,
22 Maugeri, M., Mok, H. Y., Nordli, Ø., Ross, T. F., Trigo, R. M., Wang, X. L., Woodruff, S. D.,
23 and Worley, S. J.: The Twentieth Century Reanalysis Project. *Q.J.R. Meteorol. Soc.*, 137: 1–
24 28. doi: 10.1002/qj.776, 2011.

25 Cravatte, S., Kessler, W. S., and Marin, F.: Intermediate zonal jets in the tropical Pacific
26 Ocean observed by Argo floats, *J. Phys. Oceanogr.*, 42, 1475-1485, doi:10.1175/JPO-D-11-
27 0206.1, 2012.

28 Czeschel, R., Stramma, L., Weller, R.A., Fischer, T.: Circulation, eddies, oxygen and nutrient
29 changes in the eastern tropical South Pacific, *Ocean Sci.*, 11, 455-470, doi:10.5194/os-11-455,
30 2015.

- 1 Deans, S.R.: The Radon Transform and Some of its Applications, John Wiley, Hoboken, NJ,
2 USA, 1983.
- 3 Doi, T., Tozuka, T., and Yamagata, T.: The Atlantic Meridional Mode and its coupled
4 variability with the Guinea Dome, *J. Climate*, 23, 455-475, doi:10.1175/2009JCLI3198.1,
5 2010.
- 6 Duteil, O., Schwarzkopf, F. U., Böning, C. W., and Oschlies, A.: Major role of the equatorial
7 current system in setting oxygen levels in the eastern tropical Atlantic Ocean: A high-
8 resolution model study, *Geophys. Res. Lett.*, 41, doi:10.1002/2013GL058888, 2014.
- 9 Fischer, T., Banyte, D., Brandt, P., Dengler, M., Krahlmann, G., Tanhua, T., and Visbeck, M.:
10 Diapycnal oxygen supply to the tropical North Atlantic oxygen minimum zone,
11 *Biogeosciences*, 10, 5079-5093, doi:10.5194/bg-10-5079-2013, 2013.
- 12 Foltz, G. R., McPhaden, M. J., and Lumpkin, R.: A strong Atlantic Meridional Mode event in
13 2009: The role of mixed layer dynamics, *J. Clim.*, 25, 363-380, doi:10.1175/JCLI-D-11-
14 00150.1, 2012.
- 15 Giese, B. S. and Ray, S.: El Niño variability in simple ocean data assimilation (SODA),
16 1871–2008, *J. Geophys. Res.*, 116, C02024, doi:10.1029/2010JC006695, 2011.
- 17 Giese, B. S., Chepurin, G. A., Carton, J. A., Boyer, T. P., and Seidel, H. F.: Impact of
18 bathythermograph temperature bias models on an ocean reanalysis, *J. Clim.*, 24, 84-93,
19 doi:10.1175/2010JCLI3534.1, 2011.
- 20 Gnanadesikan, A., Bianchi, D., Pradal, M.-A.: Critical role for mesoscale eddy diffusion in
21 supplying oxygen to hypoxic ocean waters, *Geophys. Res. Lett.*, 40, 5194-5198,
22 doi:10.1002/grl.50998, 2013.
- 23 Hahn, J., Brandt, P., Greatbatch, R. J., Krahlmann, G., and Körtzinger, A.: Oxygen variance
24 and meridional oxygen supply in the tropical North East Atlantic oxygen minimum zone, 43,
25 2999-3024, *Climate Dynamics*, doi:10.1007/s00382-014-2065-0, 2014.
- 26 Hansen, H. P.: Determination of oxygen. In: *Methods of Seawater analysis*, edited by:
27 Grasshoff, K. K., Ehrhardt, M., Wiley-VCH, Weinheim, Germany, 75-89, 1999.
- 28 Ho, D. T., Ledwell, J. R., Smethie Jr., W. M.: Use of SF₅CF₃ for ocean tracer release
29 experiments, *Geophys. Res. Lett.*, 35, L04602, doi:10.1029/2007GL032799, 2008.

- 1 Hormann, V., Lumpkin, R., and Foltz, G. R.: Interannual North Equatorial Countercurrent
2 variability and its relation to tropical Atlantic climate modes, *J. Geophys. Res.*, 117, C04035,
3 doi:10.1029/2011JC007697, 2012.
- 4 Kalvelage, T., Lavik, G., Lam, P., Contreras, S., Arteaga, L., Löscher, C. R., Oschlies, A.,
5 Paulmier, A., Stramma, L., and Kuypers, M. M. M.: Nitrogen cycling driven by organic
6 matter export in the South Pacific oxygen minimum zone, *Nature Geoscience*, 6, 228-234,
7 doi:10.1038/NGEO1739, 2013.
- 8 Karstensen, J., Stramma, L., and Visbeck, M.: Oxygen minimum zones in the eastern tropical
9 Atlantic and Pacific oceans, *Prog. Oceanogr.*, 77, 331-350, 2008.
- 10 Karstensen, J., Fiedler, B., Schütte, F., Brandt, P., Körtzinger, A., Fischer, G., Zantopp, R.,
11 Hahn, J., Visbeck, M., and Wallace, D.: Open ocean dead zones in the tropical North Atlantic
12 Ocean, *Biogeosciences*, 12, 2597-2605, doi:10.5194/bg-12-2597-2015, 2015.
- 13 Levitus, S., Antonov, J. I., Boyer, T. P., Locarnini, R. A., Garcia, H. E., and Mishonov, A. V.:
14 Global ocean heat content 1955-2007 in light of recently revealed instrumentation problems,
15 *Geophys. Res. Lett.*, 36, L07608, doi:10.1029/2008GL037155, 2009.
- 16 Lumpkin, R. and Garzoli, S. L.: Near-surface circulation in the tropical Atlantic Ocean, *Deep-*
17 *Sea Res., Pt. I*, 52, 495-518, 2005.
- 18 Machin, F. and Pelegrí J. L.: Northward penetration of Antarctic Intermediate Water off
19 Northwest Africa, *J. Phys. Oceanogr.*, 39, 512-535, 2009.
- 20 McGillicuddy Jr., D. J., Anderson, L. A., Bates, N. R., Bibby, T., Buesseler, K. O., Carlson,
21 C. A., Davis, C. S., Ewart, C., Falkowski, P. G., Goldthwait, S. A., Hansell, D. A., Jenkins,
22 W. J., Johnson, R., Kosnyrev, V. K., Ledwell, J. R., Li, Q. P., Siegel, D. A., and Steinberg, D.
23 K.: Eddy/wind interactions stimulate extraordinary mid-ocean plankton blooms, *Science*, 316,
24 1021-1026, 2007.
- 25 Mittelstaedt, E.: The upwelling area off Northwest Africa – a description of phenomena
26 related to coastal upwelling, *Prog. Oceanogr.*, 12, 307-331, 1983.
- 27 Peña-Izquierdo, J., Pelegrí, J. L., Pastor, M. V., Castellanos, P., Emilianov, M., Gasser, M.,
28 Salvador, J., and Vasquez-Dominguez, E.: The continental slope current system between
29 Cape Verde and the Canary Islands, *Sci. Mar.*, 76, 65-78, 2012.

1 Peña-Izquierdo, J., van Sebille, E., Pelegrí, J. L., Sprintall, J., Mason, E., Llanillo, P. J., and
2 Machín, F.: Water mass pathways to the North Atlantic oxygen minimum zone, *J. Geophys.*
3 *Res. Oceans*, 120, 3350-3372, doi:10.1002/2014JC010557, 2015.

4 Richardson, P. L. and Reverdin G.: Seasonal cycle of velocity in the Atlantic North Equatorial
5 Countercurrent as measured by surface drifters, current meters, and ship drift, *J. Geophys.*
6 *Res.*, 92(C4), 3691-3708, 1987.

7 Ridgway, K. R., Dunn, J. R., and Wilkin, J. L.: Ocean interpolation by four-dimensional least
8 squares-Application to the waters around Australia, *J. Atmos. Ocean. Technol.*, 19, 1357-
9 1375, 2002.

10 Rosell-Fieschi, M., Pelegrí, J. L., and Gourrion, J.: Zonal jets in the equatorial Atlantic Ocean,
11 *Prog. Oceanogr.*, 130, 1-18, doi:10.1016/j.pocean.2014.08.008, 2015.

12 Schott, F., Fischer, J., and Stramma, L.: Transports and pathways of upper-layer circulation in
13 the western tropical Atlantic, *J. Phys. Oceanogr.*, 28, 1904-1928, 1998.

14 Siedler, G., Zangenberg, N., Onken, R., and Morliere, A.: Seasonal changes in the tropical
15 Atlantic circulation: Observations and simulations of the Guinea Dome, *J. Geophys. Res.*, 97,
16 703-715, 1992.

17 Stramma, L. and Schott, F.: The mean flow field of the tropical Atlantic Ocean, *Deep-Sea*
18 *Res. Pt. II*, 46, 279-303, 1999.

19 Stramma, L., Hüttl, S., and Schafstall, J.: Water masses and currents in the upper tropical
20 northeast Atlantic off northwest Africa, *J. Geophys. Res.*, 110, C12006,
21 doi:10.1029/2005JC002939, 2005.

22 Stramma, L., Johnson, G. C., Sprintall, J., and Mohrholz, V.: Expanding oxygen-minimum
23 zones in the tropical oceans, *Science*, 320, 655-658, doi:10.1126/science.1153847, 2008a.

24 Stramma, L., Brandt, P., Schafstall, J., Schott, F., Fischer, J., and Körtzinger, A.: Oxygen
25 minimum zone in the North Atlantic south and east of the Cape Verde Islands, *J. Geophys.*
26 *Res.*, 113, C04014, doi:10.1029/2007JC004369, 2008b.

27 Stramma, L., Visbeck, M., Brandt, P., Tanhua, T., and Wallace, D.: Deoxygenation in the
28 oxygen minimum zone of the eastern tropical North Atlantic, *Geophys. Res. Lett.*, 36,
29 L20607, doi:10.1029/2009GL039593, 2009.

1 Stramma, L., Prince, E. D., Schmidtko, S., Luo, J., Hoolihan, J. P. Visbeck, M., Wallace, D.
2 W., Brandt, P., and Körtzinger, A.: Expansion of oxygen minimum zones may reduce
3 available habitat for tropical pelagic fishes, *Nature Climate Change*, 2, 33-37,
4 doi:10.1038/NCLIMATE1304, 2012.

5 Stramma, L., Bange, H. W., Czeschel, R., Lorenzo, A., and Frank M.: On the role of
6 mesoscale eddies for the biological productivity and biogeochemistry in the eastern tropical
7 Pacific off Peru, *Biogeosciences*, 10, 7293-7306, doi:10.5194/bg-10-7293-2013, 2013.

8 Tomczak, M.: Ausbreitung und Vermischung der Zentralwassermassen in den
9 Tropengebieten der Ozeane. 1: Atlantischer Ozean, *Oceanol. Acta*, 7, 145-158, 1984.

10 Urbano, D. F., Jochum, M., and da Silveira, I. C. A.: Rediscovering the second core of the
11 Atlantic NECC, *Ocean Model.*, 12, 1-15, 2006.

12 Urbano, D. F., De Almeida, R. A. F., and Nobre, P.: Equatorial Undercurrent and North
13 Equatorial Countercurrent at 38°W: A new perspective from direct velocity data, *J. Geophys.*
14 *Res.*, 113, C04041, doi:10.1029/2007JC004215, 2008.

15 Winkler, L.W.: Bestimmung des im Wasser gelösten Sauerstoffs, *Ber. Dtsch. Chem. Ges.*, 21,
16 2843-2855, 1888.

17
18
19
20
21


22 **Table 1.** Argo floats with oxygen sensors used here. The APEX floats had cycle intervals of
23 10 days, the PROVOR float of 7 days. The APEX floats stayed at the surface for about 13
24 hours while the PROVOR float stayed at the surface for about 7 hours. According to the
25 parking depths the floats are named here f330, f350 and f400.

26

27 Float type	serial #	deployment position; date	comment
28 Named here	Argo ID	time period data delivery	
29 WRC APEX	6013/3936	08°04'N, 22°59'W; 24 April 2008	parking depth: 330 m

1	f330	6900524	4 June 2008 -16 December2009	max. depth:1000 m
2	WRC APEX	6014/3937	08°11'N, 22°51'W; 26 April 2008	parking depth: 350 m
3	f350	6900525	27 May 2008 – 6 May 2012	max. depth: 1000 m
4	Martec PROVOR		08°11'N, 22°51'W; 26 April 2008	parking depth: 400 m
5	f400	6900629	27 April 2008 – 24 January 2012	max. depth: 2000 m

6
7




8
9
10

11 **Figure 1.** Aviso sea level height anomaly (in cm) for 25 November 2009, cyclonic features
12 are shown in bright color, anticyclonic ones in dark grey. The CTD stations of R/V *Merian*


1 cruise MSM10/1 in November/December 2008 (yellow dots), of R/V *Meteor* cruise M80/1 in
2 November 2009 (red dots), of cruise M80/2 in December 2009 (cyan dots) and of R/V *Meteor*
3 cruise M83/1 in October/November 2010 (blue dots) are included. The white line off Africa
4 marks the 200 m depth contour. The white cross with a black frame at $\sim 8^{\circ}\text{N}$, 23°W marks the
5 location of the tracer release. Some upper ocean current bands based on earlier schematics
6 (e.g. Mittelstaedt, 1983; Stramma and Schott, 1999; Stramma et al., 2008b; Peña-Izquierdo et
7 al., 2012; Brandt et al., 2015) are shown as solid black lines. For current names please refer to
8 the text. Cyclonic as well as anticyclonic eddies are indicated on a location with
9 corresponding sea level height anomaly.

10



1
2
3
4
5

Figure 2. SODA model mean circulation for the period 2001 to 2010 for the layer 50 to 200 m (a) and 200 to 400 m (b).



1


2 **Figure 3.** SODA model mean velocity section in cm s^{-1} for the period 2001 to 2010 along
 3 23°W (positive eastward).

4

5

6


1




2

3 **Figure 4.** Sea surface height anomaly on 18 November 2009 in cm (top), zonal velocity
4 component 15 to 21 November 2009 in cm s^{-1} (middle) and oxygen content in $\mu\text{mol kg}^{-1}$
5 (bottom) at 23°W . Selected isopycnals $\sigma_\theta = 25.8$ and 27.1 kg m^{-3} for Central Water
6 boundaries and $\sigma_\theta = 26.88 \text{ kg m}^{-3}$ for the tracer release density are included as black (middle
7 panel) and white (lower panel) lines.

8




1
 2 **Figure 5.** Meridional velocity component in cm s^{-1} (positive northward) at about 15°N east of
 3 23°W , at 12.5°N east of 21°W at 10°N east of 23°W and at 8°N east of 22.4°W and west of
 4 17°W all measured in October 2010 (cruise tracks can be seen in Fig. 11).



1
 2 **Figure 6.** Horizontal distribution of ADCP velocity vectors converted to 350 m depth
 3 recorded in November and December 2009 with current vectors colored with oxygen (in μmol
 4 kg^{-1}) of the accompanying CTD oxygen measurements at this depth. The oxygen distribution
 5 of the background field is from CARS 2009 climatology (Ridgway et al., 2002) for the mean
 6 of the November and December distribution at 350 m depth. The float track (white line) and
 7 surfacing location (white squares) drifting at 350 m depth and the oxygen measured by the
 8 float at 350 m depth (color in the square) is shown for the months November and December
 9 2009.

10




1

2 **Figure 7.** Mean dissolved oxygen concentration time series ($\mu\text{mol kg}^{-1}$) for the area $10\text{-}14^\circ\text{N}$,
 3 $20\text{-}30^\circ\text{W}$ with fitted linear trend and 95% confidence interval for **(a)** $100\text{--}300\text{ m}$ ($-0.49 \pm$
 4 $0.16 \mu\text{mol kg}^{-1} \text{ year}^{-1}$) and **(b)** $300\text{--}700\text{ m}$ ($-0.35 \pm 0.16 \mu\text{mol kg}^{-1} \text{ year}^{-1}$).

5


6

7



1
 2 **Figure 8.** Annual mean climatological oxygen distribution (grey shaded contours, $\mu\text{mol kg}^{-1}$)
 3 at 350 m depth from CARS 2009 climatology (Ridgway et al., 2002) with trajectories (in
 4 color) of three floats deployed in April 2008 (red 330 m (f330), green 350 m (f350), blue 400
 5 m (f400)). First measurement cycle is shown as solid dot and the last cycle as open circle
 6 (Table 1).

7




1
 2 **Figure 9.** Measurements of float f350 in the tropical eastern North Atlantic between April
 3 2008 and May 2012 with float path (color coded) in time (top) and oxygen distribution
 4 (bottom) in $\mu\text{mol kg}^{-1}$ (color) vs. time in the upper 900 m, density contours are marked in
 5 white, the mixed layer depth defined for the depth where the density is 0.125 kg m^{-3} larger

1 than at the surface as thick white line. Some dates bounding anomalous oxygen distributions
2 are marked on top of the figure.

3

4




5

6

7 **Figure 10.** Horizontal distribution of ADCP velocity vectors converted to the isopycnal $\sigma_0 =$
8 26.88 kg m^{-3} recorded in November and December 2009 with current vectors interpolated on
9 0.25° intervals colored with the maximum tracer (CF_3SF_5) concentration (in fmol kg^{-1}) of the
10 nearest CTD bottle measurements near this isopycnal. Open arrows only show velocity

1 information because no tracer was sampled. The black x with a white frame at $\sim 8^\circ\text{N}$, 23°W
2 shows the deployment location where the tracer and the three floats were deployed in April
3 2008. The three float tracks for the period April 2008 to December 2009 are included as red,
4 green and blue curves with a circle for the December 2009 location. The main current field is
5 indicated by dashed black arrows.




8 **Figure 11.** Horizontal distribution of ADCP velocity vectors converted to the isopycnal $\sigma_\theta =$
9 26.88 kg m^{-3} recorded in October/November 2010 with current vectors interpolated on 0.25°
10 intervals colored with the maximum tracer (CF_3SF_5) concentration (in fmol kg^{-1}) of the
11 nearest CTD bottle measurements near this isopycnal. Colored dots show the maximum tracer

1 at all tracer measurement locations even if no ADCP measurements are available. The black x
2 with a white frame at $\sim 8^\circ\text{N}$, 23°W shows the deployment location where the tracer and the
3 three floats were deployed in April 2008. The three float tracks for the period April 2008 to
4 November 2010 are included as red, green and blue curves with a circle for the November
5 2010 location. Float f330 is shown through the transmission end on 16 December 2009 at
6 10.04°N , 23.12°W as red curve. The main current field is indicated by dashed black arrows.
7 Please note that the tracer color scale is different to the color scale in Fig. 10.

8

9



10

11 **Figure 12.** Schematic flow field based on the results from this investigation and on earlier
12 schematics (e.g. Mittelstaedt, 1983; Stramma and Schott, 1999; Stramma et al., 2008b; Peña-

1 Izquierdo et al., 2012; Brandt et al., 2015) shown as solid black lines for the upper ocean to
2 about 200 m depth and as dashed black lines for the layer 200 to 400 m. The two dash-dotted
3 lines indicate possible different paths of the 200 to 400 m layer, thin lines indicate weak or
4 reversing currents. For current names please refer to the text. The white cross at $\sim 8^{\circ}\text{N}$, 23°W
5 marks the location of the tracer release. Annual mean climatological oxygen distribution
6 (color) at 350 m depth from CARS 2009 climatology (Ridgway et al., 2002).