

Upwelling characteristics in the Gulf of Finland (Baltic Sea) as revealed by Ferrybox measurements in 2007-2013

Villu Kikas, Urmas Lips

Marine Systems Institute at Tallinn University of Technology

Akadeemia tee 15a, 12618 Tallinn, Estonia

Tel: +3726204315, Fax: +3726204301

e-mail: villu.kikas@msi.ttu.ee

Abstract. Ferrybox measurements are carried out between Tallinn and Helsinki in the Gulf of Finland (Baltic Sea) in a regular basis since 1997. The system measures autonomously water temperature, salinity, chlorophyll *a* fluorescence and turbidity and takes water samples for further analyses at a pre-defined time interval. We aimed to show how the Ferrybox technology could be used to study the coastal upwelling events in the Gulf of Finland. Based on the introduced upwelling index and related criterion, 33 coastal upwelling events were identified in May-September 2007-2013. The number of events as well as frequency of their occurrence and intensity, expressed as a sum of daily average temperature deviations in the 20-km wide coastal area, were almost equal near the northern and southern coast. It is shown that the wind impulse needed to generate upwelling events of similar intensity differ between the two coastal areas whereas this difference is related to the average wind forcing in the area. Two types of upwelling events were identified – one characterized by a strong temperature front and the other revealing gradual decrease of temperature from the open to coastal area with maximum temperature deviation close to the shore.

Key words: Ferrybox, coastal upwelling, upwelling index, cumulative wind stress, Gulf of Finland

1. INTRODUCTION

Unattended monitoring of marine environment using ships of opportunity has been implemented in many regions of the World Ocean (e.g. Paerl et al., 2009; Hardman-Mountford et al., 2008) including the Baltic Sea and the Gulf of Finland (Rantajärvi, 2003). The measurement systems installed on board commercial ferries or other ships are called “Ferryboxes” and they consist of various sensors, devices creating water flow through the sensors and software packages controlling the system and managing the data. The commonly used Ferryboxes measure temperature, salinity and chlorophyll *a* fluorescence in the sea water pumped through the system from the surface layer along the ship track. First trials of using ships of opportunity for environmental monitoring in the Gulf of Finland were made by Estonian and Finnish scientists between Tallinn and Helsinki in 1990-1991 (Rantajärvi, 2003). Regular Ferrybox measurements along this route were started in 1997 while the longest data series of Ferrybox measurements (since 1993) is available along the ferry route Helsinki-Travemünde (Petersen, 2014).




Figure 1. Map of the Baltic Sea (a) and the study area (b) with the Ferrybox transect and Kalbadagrund meteorological station.

The Gulf of Finland (GoF) lies in the northeastern part of the Baltic Sea (Fig. 1). It is an elongated basin with a length of about 400 km and a maximum width of 135 km (Alenius et al., 1998). The long-term residual circulation in the surface layer of the gulf is characterized by a relatively low speed and by a cyclonic pattern. According to the latter, the saltier water of the northern Baltic Proper intrudes into the gulf along the Estonian (southern) coast and the seaward flow of gulf water, which is less saline due to the large freshwater inflow in the eastern end of gulf (the Neva River), occurs along the Finnish (northern) coast. The circulation is more complex

56 at time scales from days to weeks mainly due to the variable wind forcing. A variety of
57 mesoscale processes/features (fronts, eddies, upwelling/downwelling), which significantly affect
58 the biological production, retention and transport, have been observed in the Gulf of Finland (e.g.
59 Talpsepp et al., 1994; Pavelson et al., 1997; Lips et al., 2009).

60
61 The vertical stratification in the gulf is characterized by a quasi-permanent halocline at [the](#)
62 depths of 60-80 m, and a seasonal thermocline, which forms in spring-summer at the depths of
63 10-20 m (e.g. Liblik and Lips, 2011). While high concentrations of dissolved inorganic nitrogen
64 (DIN) and phosphorus (DIP) are observed in winter, the concentrations of DIN and DIP are
65 usually below the detection limit in summer in the upper mixed layer but still high just below the
66 seasonal thermocline. In general, the most prominent features in the seasonal dynamics of
67 phytoplankton in the Gulf of Finland are the spring bloom in April-May dominated by
68 dinoflagellates/diatoms and the late summer bloom in July (or late June to mid-August)
69 dominated by cyanobacteria (Kononen et al., 1996). However, the variations in bloom intensities
70 and their spatial distributions are very high between the years and within the season that is often
71 related to the physical forcing and especially to the mesoscale processes, including upwelling
72 events (Lips and Lips, 2008; Vahtera et al., 2005).

73
74 Dynamics and characteristics of upwelling events have been studied in the Gulf of Finland based
75 on in-situ measurements (e.g. Haapala, 1994), remote sensing (e.g. Uiboupin and Laanemets,
76 2009) and modelling (e.g. Myrberg and Andrejev, 2003). Most prominent upwelling events that
77 were captured by measurements are an event along the northern coast in July 1999 (Vahtera et
78 al., 2005) and an event along the southern coast in August 2006 (Lips et al., 2009). The
79 following characteristic features of upwelling events in the Gulf of Finland are suggested:

- 80
81
- 82 1) the Finnish coastal sea in the north-western GoF is one of the main upwelling areas in the
83 Baltic Sea (Myrberg and Andrejev, 2003) where upwelling frequency in May-September
84 1990-2009 has been up to 15% (Lehmann et al., 2012); almost the same upwelling
85 frequency is suggested by the latter authors for the central GoF along the Estonian
86 (southern) coast;

- 2) mean upwelling area detected on the basis of 147 maps during the period of 2000-2009 was 5642 km² (19% of the GoF surface area) along the northern coast and 3917 km² (13% of the GoF surface area) along the southern coast (Uiboupin and Laanemets, 2015), while the largest area covered by the upwelling water was identified as 12140 km² (data from 2000-2006; Uiboupin and Laanemets, 2009); the authors' estimate of the mean cross-shore extent of upwelling area was 20-30 km off the northern coast and varied between 7 and 20 km off the southern coast;
- 3) the intensity of upwelling events depends on the values of cumulative upwelling-favorable wind stress and strength of vertical stratification; Haapala (1994) suggested that at least 60 h long wind event has to exist to create an upwelling event; based on the wind data analysis from 2000-2005 and taking the threshold value for cumulative wind stress of 0.1 N m⁻² d, on average, about 2 upwelling events should appear off the southern coast and 4 events off the northern coast (Uiboupin and Laanemets, 2009);
- 4) it is suggested that the difference in topography off the southern and northern coast of the GoF results in differing upwelling dynamics along the opposite coasts – in case of similar wind stress (but in opposite directions) the transport of waters from deeper layers starts earlier and is larger along the southern coast (Väli et al., 2011).

The motivation of the present paper is to show how the Ferrybox technology can be used to study mesoscale processes, especially coastal upwelling events in the Gulf of Finland. We describe the approach, its advantages and limits, and present statistical characteristics of upwelling events on the basis of data collected in 2007-2013. The main aim is to relate the observed variability and dynamics of upwelling events to the atmospheric forcing and reveal the differences in upwelling behaviour in the two (the one opposite to the other) coastal areas.

2. THE MEASUREMENT SYSTEM AND METHODS

2.1. Ferrybox system

Temperature (T), salinity (S), chlorophyll *a* fluorescence and turbidity data and water samples for nutrients and phytoplankton chlorophyll *a* (Chl *a*), species composition and biomass analyses are collected unattended on passenger ferries, travelling between Tallinn and Helsinki (Fig. 1) since 1997. Due to the internal arrangements at the ferry company Tallink Silja and its predecessors, several ferries were used as the platforms for Ferrybox measurements, which also differ in regard of water intake features. A flow-through system from 4H-Jena, Germany with the water intake attached to the sea chest of the ferry is in use since 2006. The water enters the sea chest through a grating with a total surface area of 0.84 m² located at about 4 m depth below the waterline. The water flow from the sea chest into the system is forced by the hydrostatic pressure since the Ferrybox is located on the lower deck about 3 meters below the waterline. To restrict larger particles to get into the measurement system a mud filter (pore size 1 mm) is used just after the water intake. Prior to the sensors a debubbler is installed to avoid air bubbles to affect the measurements of conductivity, turbidity and Chl *a* fluorescence. The flow rate through the sensors is stabilized by an internal pump, which is controlled by a pressure sensor in the system. Water samples are taken by a sampling device (Hach Sigma 900 MAX) whereas the water is pumped from the debubbler into the bottles using an internal pump of the water sampler.

For temperature measurements, a PT100 temperature sensor is used that is installed close to the water intake to diminish the effect of warming of water while flowing through the tubes onboard. The sensor has a measuring range from -2 to +40 °C and accuracy of ±0.1% of the range, thus 0.04 °C. For salinity measurements a FSI Excell thermosalinograph (temperature and conductivity meter) and for Chl *a* fluorescence and turbidity measurements a SCUFA submersible fluorometer (Turner Designs) with a flow-through cap is used. The system starts the measurements and data recording when the ferry is away from the harbor more than a control distance of 0.7 nautical miles (controlled by a GPS device in the system) and stops when it is closer than this distance in order to avoid the sediments getting into the system.

The data are recorded during every crossing (twice a day) every 20 seconds that corresponds to a horizontal resolution of approximately 160 m.

2.2. Quality assurance and pre-processing of data

148 The sensors have been calibrated at the factory before the installation and if necessary sent for an
149 additional laboratory calibration. Since the system contains two temperature sensors, the
150 performance of them is routinely followed by comparison of data acquired from the sensors. The
151 quality of thermosalinograph data is guaranteed by taking a series of water samples (14-17
152 samples) and analyzing them using a high-precision salinometer AUTOSAL 2-4 times a year.
153 The analyses have shown, that a correction of 0.08 (units in Practical Salinity Scale; the value
154 has been stable over the years) must be added to the recorded salinity. While the raw salinity is
155 recorded ~~(and presented in Fig. 2)~~ in units according to the Practical salinity Scale 1978, the
156 results on salinity distribution and variability are given later in this paper in g kg^{-1} (Sections 3
157 and 4). Special care is taken to calibrate the SCUFA fluorometer; however, since we do not use
158 the fluorometer data in this study the used routine is not described here.

159
160 The data acquired by the Ferrybox system are recorded with a time step of 20 s and stored in an
161 onboard terminal. In order to synchronize the measurements performed by the sensors having
162 different sampling frequencies and GPS, the acquired data within every 19 s interval are
163 averaged and recorded as measurements at every 20th second. The data are automatically
164 delivered to the on-shore ftp-server once a day when the ferry is in the harbor using a GSM
165 connection. The performance of the system is validated by the control parameters, such as the
166 flow rate and pressure in the system, and the data are checked for unrealistic values against the
167 criteria set for every parameter on the basis of known natural variation of them in the Gulf of
168 Finland.

169
170 One of the procedures, which has to be carried out when using the Ferrybox data, is the shifting
171 of data points to the actual positions of the water intake. The problem arises since the position
172 attached to a data record is the position of the ferry at the time of measurement but the water is
173 taken in earlier at a different position. Since various systems of water intake are applied, this
174 procedure is unique for each combination of a Ferrybox and a ferry.

175 As described above, in our design the sea water enters first a relatively large sea chest and the
176 flushing through time of it is unknown. While the water flows through the sea chest and into the
177 tubes and debubbler with a flow rate of $12\text{-}15 \text{ l min}^{-1}$, the ferry moves on at an average speed of

16 knots. We solved the problem of position correction taking into account the advantage of having two crossings a day.




Figure 2. Measured salinity distribution along the ferry route Tallinn-Helsinki from the forth and backward journey on 27 July 2010. Raw data are presented in the left panel and the processed data in the right panel where the shifting of data points by 3 minutes and 20 seconds was applied; x-axis shows the distance from the Tallinn Bay (latitude 59.48 N) in km along the meridional transect.

Analysis of data from forth and backward journeys allowed us to introduce a rough position correction procedure – the best result is achieved by shifting the measured data points against the GPS time by 3-4 minutes depending on the ferry and exact intake installation. This relatively long time shift is obviously related to the water exchange in the sea chest. Due to an almost constant cruising speed of the ferry outside the harbor areas, the applied procedure gives acceptable results – as it is seen for instance in Fig. 2. In the shown example, the data points are shifted by 3 minutes and 20 seconds. This period is found by seeking for the best correlation between the data from forth and backward journeys. The comparison of data along the ferry line from Tallinn to Helsinki and back from Helsinki to Tallinn obtained on the same day is one of the applied quality assurance procedures – the profiles containing unexplained deviations are marked by a quality flag indicating a possible quality problem.

2.3. Data and calculation methods

201 Temperature and salinity data collected along the ferry line Tallinn-Helsinki from May to
202 September in 2007-2013 are used for analysis purposes. In 2008, the system on board the
203 passenger ferry “Galaxy” was in use until 13 July and the measurements started again on 13
204 August when the system was installed on board the ferry “Baltic Princess”. However, due to
205 some technical problems, the regular measurements were successful since 2 September 2008.

206
207 A failure of the system occurred late August 2012 and therefore the data are not available ~~from~~
208 ~~September 2012 (since from~~ 29 August ~~until the end of September 2012)~~. In early 2013, the next
209 ferry (“Silja Europa”) came to this line and the system was moved again causing a break in the
210 measurements until 15 July 2013. The number of crossings with the full data coverage is given in
211 Table 1. One can conclude that among the seven analyzed years (from May to September), four
212 years – 2007, 2009, 2010 and 2011 – were the years with almost full data coverage while most of
213 the data were not available in the second half of July and August 2008, in September 2012 and in
214 May, June and first half of July 2013. Thus, the data from all months from May to September
215 were analyzed at least from six years in 2007-2013.

216
217 Collected raw data were preliminarily processed, including shifting of measurements as
218 described in Section 2.2, quality checked and stored in the database. This data set was used to
219 draw the maps of temporal variations of horizontal distributions of T and S for all studied years
220 (Fig. 32). In order to assess upwelling intensities the data set was transformed from the matrix
221 with constant time step into the matrix with a constant spatial resolution. A step (cell width) of
222 0.5 km along the south-north oriented line was used. The fixed south-north orientation was
223 applied to eliminate the influence of differences in orientation of the ship track in the southern,
224 central and northern parts of the route (see route configuration in Fig. 1) and of possible
225 deviations from the ordinary route. As a result, the extent of upwelling area is presented below in
226 south-north direction and a coefficient has to be applied to convert these values into the
227 upwelling extent in cross-shore direction (as cosine of the angle between the south-north
228 direction and a perpendicular line to the shore – approximately 20 degrees).

229
230 ~~In order to characterize the upwelling intensity, temperature deviations were calculated against~~
231 ~~the daily average temperature values over the entire ferry route from harbor to harbor.~~ An

upwelling index was introduced for the 20-km wide coastal areas both off the southern coast (UI_S) and off the northern coast (UI_N). ~~The width of 20 km was defined on the basis of the analysis of all available temperature data from Tallinn-Helsinki ferry line in 2007-2013 (see Section 3.1 for details).~~ For each crossing, the average water temperature and horizontal profile of temperature deviations from the average were found. The upwelling index was calculated as a sum of negative temperature deviations in the 20-km coastal areas ~~as as a measure of the upwelling intensity in the temperature signal off the southern and northern coast was obtained, respectively:~~

$$UI_S = \sum_{\Delta T_i < 0}^{i=1 \dots 40} |\Delta T_i| \text{ and } UI_N = \sum_{\Delta T_i < 0}^{i=101 \dots 140} |\Delta T_i| \quad (1)$$

where ΔT_i is the temperature deviation at 0.5-km cell i from the average temperature of the crossing. ~~The width of 20 km was defined selected on the basis of the analysis of all available temperature data from Tallinn-Helsinki ferry line in 2007-2013 (see Section 3.1 for details). The daily indexes were obtained by averaging the two upwelling indexes from a single day (from forth and backward journey of the ferry). A calculated upwelling index value equal to zero or below a certain threshold value that can be defined means that there was no upwelling observed in one or the other coastal area.~~ The cumulative upwelling index (CUI) can be calculated by summing up upwelling index values for certain periods. ~~When summing up the upwelling index~~ The obtained CUI values were divided ~~them~~ by 40, which is the number of data cells in the 20-km wide coastal area, to keep the meaning of CUI as the sum of average negative temperature deviations, having a unit of [$^{\circ}\text{C day}$]:

$$CUI_S(n1 \dots n2) = \sum_{j=n1}^{j=n2} \left(\frac{1}{40} UI_{Sj} \right) \text{ and } CUI_N(n1 \dots n2) = \sum_{j=n1}^{j=n2} \left(\frac{1}{40} UI_{Nj} \right) \quad (2)$$

where $n1$ and $n2$ are the start and the end day number of the selected period, for which the cumulative upwelling index is calculated, and UI_{Sj} and UI_{Nj} are the upwelling indexes at day j off the southern and northern coast, respectively. ~~The daily indexes were obtained by averaging the two upwelling indexes from a single day (from forth and backward journey of the ferry).~~ This approach of the CUI calculation is similar to those used previously in the studies of upwelling events and their influence to the phytoplankton dynamics in the Gulf of Finland (see e.g. Lips and Lips, 2008; Myrberg et al., 2008).

Thus, an upwelling event can be characterized by the cumulative upwelling index whereas the first and the last day of the event can be defined as the start and end of the period when the upwelling index (UI_N or UI_S) exceeded a certain threshold value. We have defined this threshold value as 40 °C, which corresponds e.g. to a 20-km wide upwelling with an average negative temperature deviation of 1 °C or a 10-km wide upwelling with an average negative temperature deviation of 2 °C. This choice is explained in more details ~~when presenting the results~~ in Section 3.2. Although the precision of the temperature sensor is obviously better than its accuracy, we estimated the uncertainty of the calculated index values based on the absolute accuracy of PT100. The accuracy of the temperature measurements of 0.04 °C gives a maximum uncertainty of 1.6 °C in the upwelling index estimates (it is 25 times less than the selected threshold) and a maximum uncertainty of 0.4 °C day in the cumulative upwelling index estimates (considering a 10-day upwelling event).

Wind data were obtained from the HIRLAM (High Resolution Limited Area Model) version of the Estonian Meteorological and Hydrological Institute with the spatial resolution of 11 km and time interval of 3 h (Väli, 2011; Männik and Merilain, 2007). Model data point close to Kalbådagrund, where also a meteorological weather station is located (Finnish Meteorological Institute), was chosen to represent the wind conditions in the study area. The data from Kalbådagrund weather station or from the closest HIRLAM model point have been also used in the earlier studies describing wind conditions in the Gulf of Finland (Lips et al., 2008; Uiboupin and Laanemets, 2009). According to Keevallik and Soomere (2010) the HIRLAM model matches well with the measured wind speed as well wind directions, whereas to obtain the wind direction at 10 m height the measured wind direction at Kalbådagrund (measured at 32 m) is advised to turn by 20° counter-clockwise.

Wind stress (in $N\ m^{-2}$) is calculated for the wind component along the axis of the Gulf of Finland, which corresponds to the direction turned by 70 degrees clockwise from the north direction, as:

$$\tau_{70} = C_D \rho_a |U| U_{70} \quad (3)$$


where U is the wind speed (in $m\ s^{-1}$) and U_{70} is its component in the along-gulf direction, C_D is the drag coefficient (a value of $1.2 \cdot 10^{-3}$ was chosen in the present study) and ρ_a is the air density

(1.2 kg m⁻³). Accordingly, positive values of the wind stress should initiate ~~across gulf~~
~~southward~~ Ekman transport in the surface layer ~~from north to south~~ and vice versa. The
cumulative wind stress (in N m⁻² day) was calculated based on daily averages of wind stress. If
the cumulative wind stress is large enough, upwelling events occur along the northern coast in
case of the positive wind stress and along the southern coast in case of the negative wind stress.

3. RESULTS

3.1 General variability and distribution patterns

The ~~seasonal~~ course of the surface layer temperature in the Gulf of Finland in the warm season is
characterized with temperature about 5 °C in the beginning of May, a maximum above 20 °C in
late July – early August and a drop below 15 °C in late September. Within the analyzed years
2007-2013, the surface layer temperature was the highest in summer 2010 (Fig. ~~32~~). The period
when surface layer temperature exceeded 20 °C was the longest in 2010 and 2011, while in the
other years the periods with water temperature above 20 °C were very short – only a few days.
On the background of seasonal course and simultaneous shorter-term increases or decreases of
temperature over the whole study transect, the periods with clearly lower temperature off the
northern or southern shore are observed. Such situations are related to the coastal upwelling
events – their characteristic time scale was several days to 1-2 weeks and they extended towards
the open sea by 15-20 km (Fig. ~~32~~).







Figure 32. Temporal changes of temperature (in °C) and salinity (in g kg^{-1}) distributions between Tallinn and Helsinki from 1 May to 30 September in 2007 (a, b), 2008 (c, d), 2009 (e, f), 2010 (g, h), 2011 (i, j), 2012 (k, l) and 2013 (m, n); y-axis shows the distance from the Tallinn Bay (latitude 59.48 N) in km along the meridional transect.

Inter-annual variations of the surface layer salinity in 2007–2013 were high with the highest salinity in 2011 and the lowest in 2009. The surface layer salinity exceeded 6.5 g kg^{-1} for a longer period only in 2011 in the southern half of the study transect (Fig. 32j) and for shorter periods of several days in case of coastal upwelling events off the southern shore (e.g. Figs. 3b, 2b and 3d). Note that in the case of coastal upwelling events seen in the temperature distributions off the northern coast, simultaneous increase in salinity was not well visible. As a rule, the surface layer salinity was higher near the southern coast than that near the northern coast. However, often the lowest salinity was measured in the middle of the transect – it means in the open sea areas of the Gulf of Finland (e.g. Figs. 3f, 2f and 3h). Seasonal course in of salinity variation differed between the studied years remarkably. While usually, the lowest surface layer salinity was observed in June–July, in 2008, the salinity was the lowest in May and in 2010 and 2011, it was the lowest in August.






Figure 229. Distributions of temperature (in °C) and salinity (in g kg⁻¹) deviations from the daily transect mean value along the ferry route Tallinn-Helsinki for all measurements in May-September 2007-2013 (a, b), in 2009 (c, d) and in 2010 (e, f). Mean values on each 0.5-km cell (solid curves) and plus/minus RMSE (dashed curves) are shown; x-axis shows the distance from the Tallinn Bay (latitude 59.48 N) in km along the meridional transect.

To characterize the spatial distribution and variability of the surface layer temperature and salinity along the study transect, the average temperature and salinity deviations from the daily mean value of temperature and salinity as well as their root mean square errors (RMSE) were calculated for each 0.5 km cell. On average, the temperature deviations from the mean value along the transect Tallinn-Helsinki were close to zero along the entire study transect (Fig. 4a3a) – the absolute values of average deviation were six times less than estimated RMSE of temperature for corresponding cells. However, nevertheless, the surface layer temperature has been slightly warmer in the open Gulf of Finland than that in approximately 20-km wide coastal areas (Fig. 34a). This could be related to the coastal upwelling events. For instance, in

2009, when coastal upwelling events were observed off the both coasts (~~somewhat more intensive events off the northern coast~~), the average temperature deviations ~~are~~ were negative ~~off~~ near the both coasts (Fig. 34c). In 2010, when upwelling events occurred mostly off the southern coast, the negative values of average temperature deviations were detected only in the southern part of the transect (Fig. 34e).

It is remarkable that, on average, the variability of temperature deviations (~~expressed as RMSE of temperature deviations from the mean value of the certain crossing~~) was much higher near the coasts than in the central part of the study transect (Fig. 34a). In the case of upwelling events off the southern coast and their absence off the northern coast (in 2010), this high variability of temperature was concentrated only in the 20-km wide coastal area off the southern shore (Fig. 34e). Since the area of high temperature variability, which mostly could be related to the upwelling activity, extended about 20 km from the shores, ~~we-it was~~ suggested to ~~take temperature values from such areas into account when~~ estimating the intensity of upwelling events based on data from these 20-km wide coastal zones.

The average distribution of the surface layer salinity along the transect was characterized by higher salinity values in the southern gulf and lower values in the northern gulf (Fig. 4b3b). The salinity deviations ~~from the transect mean value was~~ were positive in the 28-km wide area off the southern coast (with clearly higher salinity in the first 10 km) ~~while it was~~ and negative along the rest of the study transect. However, the minimum of the surface layer salinity was observed at about 20 km from the northern shore (or at a distance of 50 km from the starting point of the study transect in the Tallinn Bay) almost in every year (Fig. 34b, d and f). The only exception was the year 2007 when the lowest salinity was observed on average in the cell closest to the northern shore. The low salinity water at the distance of 50 km indicates that, in summer, the outflow of the less saline Gulf of Finland surface waters occurs mostly in the northern part of the open gulf. The spatial differences in variability of the surface layer salinity ~~are~~ were not so distinct than in variability of the surface layer temperature. One can recognize slightly higher variability (RMSE) of the surface layer salinity in the coastal areas and in the southern part of the open gulf at ~~a the~~ distance of 20-30 km ~~from the southern end of the transect in the southern part of the open gulf~~.

3.2 Upwelling characteristics

Upwelling events along both coasts of the Gulf of Finland can be identified by temperature deviation maps (Fig. 4). ~~The occurrence and intensity of upwelling events along the both coasts of the Gulf of Finland can be qualitatively identified from the temperature deviation maps presenting temperature deviations from the daily mean value along the transect between Tallinn and Helsinki (Fig. 5).~~ As it ~~is seen from the maps~~, the years 2007 and 2009 had a similar pattern – the upwelling events occurred off the southern coast in the first half of the season and off the northern coast in the second half. In 2008, upwelling events were observed near the southern coast in May and September while in June the upwelling events appeared near the northern coast. The year 2010 was an exceptional year when the upwelling events occurred mostly along the southern coast. It was exceptional also because the sea surface temperature outside the upwelling waters was the highest among the studied summers. A sequence of consecutive upwelling events near the northern and southern coast was observed in 2011. Upwelling events occurred mostly off the northern coast in 2012 and 2013.




Figure 54. Temporal changes of spatial distributions of temperature deviations (in °C) from the daily transect mean value between Tallinn and Helsinki from 1 May to 30 September in 2007 (a), 2008 (b), 2009 (c), 2010 (d), 2011 (e), 2012 (f) and 2013 (g); y-axis shows the distance from the Tallinn Bay (latitude 59.48 N) in km along the meridional transect.

We selected a criterion to detect whether an upwelling event occurs or not as the value of upwelling index (UI) exceeding $40\text{ }^{\circ}\text{C}$ (in absolute values, while UI is by definition a negative number). ~~This value corresponds to the situation when an average negative temperature deviation of at least $-1\text{ }^{\circ}\text{C}$ in a 20-km wide coastal area is measured.~~ The upwelling events found using the selected criterion were also the occasions when the maximum negative temperature deviation from the transect mean value was at least $-2\text{ }^{\circ}\text{C}$ (except one event on 10-17 September 2007 when the maximum deviation was $-1.97\text{ }^{\circ}\text{C}$). Furthermore, no other cases with negative temperature deviations exceeding $-2\text{ }^{\circ}\text{C}$ were detected. Thus, the criterion $UI < -40\text{ }^{\circ}\text{C}$ gives quite similar results as would yield if using the criterion based on the maximum negative temperature deviation of $-2\text{ }^{\circ}\text{C}$.

We identified in May-September 2007-2013 altogether 33 upwelling events, approximately half of them (17) near the northern coast and half (16) near the southern coast (Table 2).

424 The events lasted from 3 days to 3 weeks ~~and (the longest event occurred~~ was observed on 11-31
425 August 2013). On average five events a year were registered and the maximum number of events
426 (eight) was observed in 2011. Based on available data, the number of days with the upwelling
427 near the northern coast was 150 and near the southern coast 140. As the total number of days
428 with measurements ~~in the present study~~ was 838, the upwelling occurred on 18 % and 17 % of
429 days off the northern and southern coast, respectively. The maximum negative temperature
430 deviation from the transect mean value was detected ~~in-in August~~ in August 2010 near the southern coast
431 when it reached -7.78 °C ~~during the event from 17 to 23 August 2010~~. The largest temperature
432 deviation in the case of upwelling events near the northern coast of -6.15 °C was detected during
433 the event from ~~in~~ 17 July ~~to 1 August~~ 2013. The average of maximum temperature deviations was
434 larger for the upwelling events near the southern coast than near the northern coast – -4.64 °C
435 and -3.60 °C, respectively.

436
437 While the maximum temperature deviation characterizes the peak of the upwelling, the
438 introduced cumulative upwelling index takes also into account the extent of the upwelling in
439 space and time. In regard of *CUI* the largest upwelling events were observed in 2013 – on 15-30
440 September 2013 off the southern coast (*CUI* = -40.2 °C day) and on 11-31 August 2013 off the
441 northern coast (*CUI* = -39.7 °C day). The upwelling events with the largest temperature deviation
442 in July-August 2010 were relatively short events lasting 7 days and gave respective *CUI* value as
443 -15.7 °C day and -20.8 °C day. The average *CUI* value ~~for of 47 all~~ for all upwelling events off the
444 northern coast was -14.5 °C day and off the southern coast -16.2 °C day. The Summing up of
445 *CUI* values ~~for of~~ for all detected upwelling events ~~gave the total CUI for the events~~ off the northern
446 coast ~~as was~~ was -247.0 °C day and off the southern coast -258.4 °C day. ~~We also found the total sum~~
447 ~~of upwelling indexes (divided by 40) off the both coasts regardless whether the set upwelling~~
448 ~~criterion was met or not.~~




Figure 45. Temporal changes of upwelling index off the northern coast (upper columns; °C) and off the southern coast (lower columns, °C) and along-gulf wind stress (black curve in the middle; N m^{-2}) in May-September 2007 (a), 2008 (b), 2009 (c), 2010 (d), 2011 (e), 2012 (f) and 2013 (g).

~~We also found that the total sum of upwelling indexes (divided by 40) off the both coasts regardless whether the set upwelling criterion was met or not. The resulting~~ total *CUI* for all measurement days in 2007-2013 was -405.3 °C day for the northern coastal area and -356.6 °C day for the southern coastal area. Thus, the negative temperature deviations from the transect mean ~~are~~ were more common for the northern coastal sea area while the upwelling events were more intense in the southern coastal sea area. This feature is also well seen in Fig. 6-5 where e.g. ~~for in~~ 2007 relatively low values of UI_N were found in most of the days near the northern coast but only three upwelling events were revealed according to the criterion set in the present study.

Seasonal variation of the frequency of occurrence and intensity of upwelling events was revealed. The highest number of events was observed in July – 10 events, 5 off the northern coast and 5 off the southern coast, and the lowest in May – 4 events. The sum of *CUI* values of all events in July and August were -185.3 °C day and -187.9 °C day, respectively, while it was only -28.6 °C day in May. In June and September, the *CUI* of all events had intermediate magnitude – -107.5 °C day and -137.0 °C day, respectively. Obviously, the revealed seasonal course was partly related to the temperature difference between the surface layer and the cold layer beneath the seasonal thermocline, which has its maximum in the Gulf of Finland in July-August (Liblik and Lips, 2011).

3.3 Upwelling characteristics in relation to wind forcing

The occurrence of coastal upwelling events in the Gulf of Finland can be related quite well to the variations of the along-gulf wind stress (Fig. 65). The upwelling events appeared ed after a certain favorable wind pulses with long enough duration and magnitude. In the case of upwelling events off the northern coast the positive along-gulf wind stress was usually observed a few days before the event and in the case of upwelling events off the southern coast the wind stress was negative for a few days (Fig. 65).

490 The estimated cumulative wind stress for the detected upwelling events varied between 0.31 and
491 1.37 N m⁻² day for westerly winds and between -0.09 and -1.08 N m⁻² day for easterly winds
492 (Table 2). The cumulative wind stress associated with each upwelling event was calculated based
493 on daily average wind stress values by summing them up from the first day with favorable wind
494 stress (within a period of 1 week before the event) to the last day with favorable wind stress
495 before the end of the event. If within the favorable wind stress series only one day with opposite
496 wind stress appeared in a sequence then the calculation period was not broken. The average
497 value of the cumulative wind stress for an upwelling event off the northern coast was 0.71 N m⁻²
498 day and off the southern coast -0.44 N m⁻² day. It suggests that in order to produce a coastal
499 upwelling event of an equal magnitude the required favorable along-gulf wind stress has to be
500 larger for the upwelling events off the northern coast than for the events off the southern coast.
501 This conclusion is drawn by taking into account the above result that the average upwelling
502 intensity (estimated as *CUI*) was similar for the both coastal areas with slightly higher values of
503 *CUI* for the upwelling events off the southern coast. [This suggestion is supported also by](#)
504 [comparison of relationships between the *CUI* and cumulative wind stress \(*CWS*\) related to the](#)
505 [upwelling events near the opposite coasts \(Fig. 6\). Although the results are quite scattered, the](#)
506 [upwelling events off the southern coast occurred in conditions of lower *CWS* values and the](#)
507 [maximum *CUI* near the southern coast was also found at lower *CSW* than near the northern](#)
508 [coast.](#)
509




Figure 6. Relationship between the cumulative upwelling index (CUI) and cumulative along-gulf wind stress (CWS) based on 33 detected upwelling events in May-September 2007-2013. Red symbols indicate the events off the southern coast and blue symbols the events off the northern coast; circles correspond to the events with pronounced upwelling front (N_UF and S_UF) and triangles the events with gradual decrease of temperature towards the coast (N_GD and S_GD).

The average along-gulf wind stress for the entire study period from May to September in 2007-2013 was 0.016 N m^{-2} (based on wind data with the time step of 3 hours). The seasonal averages had positive values in all studied years indicating that the westerly-south-westerly winds prevailed in the region. The average values of wind stress varied between 0.001 N m^{-2} in 2010 and 0.029 N m^{-2} — the value, which was obtained for three years — in 2007, 2009 and 2012. The year 2010 was the year when in May-September 2010, when five upwelling events occurred off the southern coast and only one event off the northern coast. However, the average along-gulf wind stress was close to zero indicating that the cumulative wind forcing was almost equal from both directions — had almost similar occurrence. It is interesting to mention that Furthermore, the wind stress averaged over the all observed upwelling events in 2007-2013 was 0.015 N m^{-2} , which is very close to the average wind stress over the entire study period. This latter estimate was obtained based on the mean length of upwelling events of 8.8 days and mean cumulative wind stress values of 0.71 and $-0.44 \text{ N m}^{-2} \text{ day}$ off the northern and southern coasts, respectively.

529 ~~Thus, the average values of wind stress during the entire study period and during the occurred~~
530 ~~upwelling events were very close to each other.~~ It can be concluded that the difference between
531 the wind impulses needed for the generation of ~~coastal~~ upwelling events with similar intensity
532 near the opposite coasts ~~with a comparable intensity in regard of the introduced cumulative~~
533 ~~upwelling index is related comparable~~ to the average wind stress value in the region.

534
535
536 Usually the upwelling events occurred one or a few days after the start of the favorable wind
537 pulse and the maximum of upwelling intensity (upwelling index) was reached one or a few days
538 after the maximum wind stress (Fig. ~~65~~). Daily measurements are too scarce yet to describe the
539 temporal evolution of upwelling events in detail since the time required to initiate Ekman
540 transport is shorter than or close to the inertial period (e.g. Lehmann and Myrberg, 2008) ~~that in~~
541 ~~the Gulf of Finland~~ is approximately 14 hours in the Gulf of Finland (about half a day). Instead,
542 we made an attempt to reveal characteristic spatial temperature and salinity distributions in the
543 surface layer from coast to coast at times of the maximum intensity of upwelling events.
544 Surprisingly, the results did not differ significantly between the northern and southern coast –
545 two characteristic shapes of upwelling events in the temperature distribution were identified for
546 both coastal areas.

547




Figure 57. Characteristic distributions of temperature and salinity along the ferry route Tallinn-Helsinki with coastal upwelling events off the northern coast (a, b) and off the southern coast (c, d); x-axis shows the distance from the Tallinn Bay (latitude 59.48 N) in km along the meridional transect.

The most intense—Mostly the upwelling events were characterized by a sharp and very intense temperature front between the upwelling waters and the rest of the transect (see Fig. 7 the yellow and orange curves). Typical for such events were an almost uniform temperature outside the upwelling area and the temperature minimum (maximum temperature deviation) close to the upwelling front. The other distribution pattern (blue curves in Fig. 7) exposed a gradual decrease of temperature towards the upwelling waters. Typical for the latter events were the irregularities in temperature distribution with characteristic scale of a few kilometers and the temperature minimum (maximum temperature deviation) in the cell closest to the shore. In some cases, e.g. the event near the northern coast with maximum intensity on 18 August 2013 (see light blue curve in Fig. 7 upper left panel), the observed temperature deviations were as large as during the upwelling events with pronounced temperature front. There was also a third type of temperature distribution when the upwelling waters were not attached to the shore (see red curve in Fig. 7 upper left panel) at least according to the measurements along the ferry route. All these types of upwelling events are well recognized on the maps of temporal changes of temperature and temperature deviation along the ferry route Tallinn-Helsinki (Figs. 3-2 and 54).

The spatial distribution of salinity in the surface layer from coast to coast drastically differed between the upwelling events near the northern coast and the events near the southern coast (Fig. 7 right panels). In the latter case, both the salinity difference across the gulf and the spatial variability at scales of a few to ten kilometers were much larger than in the former case. It is also interesting that in the case of southern upwelling events, the salinity minimum along the transect can be situated [either](#) very close to the upwelling front (e.g. on 28 July 2010) or near the northern coast (e.g. 8 July 2011). Although such diverse patterns are partly related to the history of water movements in the gulf, the salinity minimum (at least local minimum) close to the upwelling front obviously is caused by the westward current jet along the front [as revealed also by model experiments \(Laanemets et al., 2011\)](#). The salinity distribution across the gulf associated with the northern upwelling events is very uniform with some variability at scales of a few to ten kilometers, which have the amplitude several times less than ~~similar variability in~~ spatial salinity ~~distribution variations~~ associated with the southern upwelling events.

4. DISCUSSION AND CONCLUSIONS

Several studies have shown how the Ferrybox measurements are successfully used for different applications, such as for monitoring of coastal waters in combination with remote sensing (Petersen et al., 2008), estimating carbon fluxes and primary productivity (Schneider et al., 2014) and detecting cyanobacterial blooms (Seppälä et al., 2007). However, not enough attention is paid to the systems itself especially to the question how the results are affected by different technical solutions used in the systems (like water intake depth and construction, piping etc). Furthermore, the particularities of geographical location as well as the ferry route and schedule often determine the most suitable applications and requirements for the data treatment. A good example of taking advantage on the geographical location and ferry route (schedule) is demonstrated by Buijsman and Ridderinkhof (2007) who estimated the water and suspended matter exchange between the Wadden Sea and the North Sea using data collected along the ferry route Den Helder – Texel.

The ferry route between Tallinn and Helsinki across the elongated Gulf of Finland and the schedule consisting of two cruises a day and a short 1.5-hour stay in Helsinki makes it possible

600 to introduce a rough procedure for correction of coordinates of measurement points and an
601 additional quality check routine for the collected data. The correlation between the data from the
602 two crossings on the same day should be high enough; otherwise, the data can be marked as
603 suspicious. We found that the highest correlation between the two data sets is achieved when the
604 data points are shifted by 3-4 minutes depending on the intake installation and the ferry. We
605 suggest that in all Ferrybox systems such coordinate correction procedure should be used. This
606 analysis also demonstrates the confidence of the applied Ferrybox system even though the water
607 is taken in through a relatively large sea chest. Furthermore, the ferry route across the relatively
608 narrow gulf from coast to coast is very convenient to collect data on the offshore extension and
609 intensity of coastal upwelling events.

610
611 Various methods have been applied to reveal characteristic features of coastal upwelling events
612 in the Baltic Sea ~~(including the Gulf of Finland)~~ based on data mainly from remote sensing and
613 numerical models. Certain temperature isoline as the border of upwelling area was used by
614 Uiboupin and Laanemets (2009) and a temperature deviation (2 °C) from the mean temperature
615 along zonal transects was used by Lehmann et al. (2012). The latter method is similar to the
616 approach applied in the present study, but we argue that the analysis of temperature deviations
617 along meridional transects is more appropriate in the Gulf of Finland. This conclusion is justified
618 by the fact that, on average, the north-south temperature gradient is negligible in the gulf (see
619 Fig. 4a3a) ~~and when using meridional transects the results are not biased by the different~~
620 ~~temporal dynamics of surface layer temperature in the open Baltic Proper and in the relatively~~
621 ~~narrow Gulf of Finland. For instance while the west-east temperature gradient could exist~~
622 ~~between the shallower and narrower Gulf of Finland and the deeper and wider Northern Baltic~~
623 ~~Proper due to differential warming and cooling, the seasonal warming and cooling are faster in~~
624 ~~the gulf than in the open Baltic.~~

625
626 Nevertheless, it is interesting that our results on upwelling frequencies of about 17-18 % near the
627 northern and southern coast are very close to the results of Lehmann et al. (2012) if their results
628 based on remote sensing data were considered. They concluded that upwelling events are present
629 more than 15 % of time near the northern coast and about 15 % of time near the southern coast.
630 At the same time, the estimates of corresponding upwelling frequencies based on numerical

631 experiments differ from those values and the results of the present study. Based on model
632 results. ~~Usually~~ the northern coastal area ~~is~~ has been suggested as the main upwelling area in the
633 Gulf of Finland with the upwelling occurrence up to 30 % of time (Lehmann et al., 2012;
634 Myrberg and Andrejev, 2003) while near the southern coast downwelling should prevail (e.g.
635 Myrberg and Andrejev, 2003). It shows that the models ~~, in regard with~~ their current resolution
636 and parameterization of sub-grid processes should be improved.

637
638 Analysis of wind data has also suggested that the coastal upwelling events should occur more
639 often off the northern coast of the Gulf of Finland than off the southern coast (Lehmann et al.,
640 2012; Uiboupin and Laanemets, 2009). The data set consisting of 838 days of measurements
641 from coast to coast used in the present analysis has revealed that, on average, the frequency of
642 upwelling events and their ~~intensity~~ is-are similar near the northern and southern coast of the
643 gulf although the wind data from the same period suggest prevalence of upwelling events off the
644 northern coast. Partly, this outcome can be explained by the higher position of the thermocline,
645 steeper slope and greater depths in the southern part of the gulf as suggested by some earlier
646 studies (e.g. Väli et al., 2011; Laanemets et al. 2009). However, one could suggest that the
647 thermohaline structure of the Gulf of Finland is adapted to the general prevalence of westerly-
648 south-westerly winds. Thus, the wind impulse needed for the generation of a coastal upwelling
649 event of similar intensity near the southern coast can have smaller magnitude. We suggest that
650 rather the deviation from the average wind forcing than the absolute value of it should be
651 considered. This suggestion is supported by the comparison of average upwelling intensities
652 expressed as cumulative upwelling index values and cumulative wind stress values for the all
653 upwelling events recorded in 2007-2013 near the opposite costs of the gulf. Similar conclusion
654 suggestion was made by Liblik and Lips (2016) on the basis of data analysis from 35 cross-gulf
655 CTD surveys conducted in 2006-2013.

656
657 ~~As a side product we also described~~ The average cross-gulf distributions of temperature and
658 salinity deviations from the transect mean value was described based on the 7-year data set of
659 horizontal profiles. On average, the surface layer temperature ~~does~~ did not have any horizontal
660 gradient while the surface layer salinity was higher in the southern part than in the northern part
661 of the gulf. The result that the surface water with the lowest salinity ~~is~~ was on average at about

Formatted: Highlight

20 km from the northern coast supports the suggested general circulation scheme in the Gulf of Finland (e.g. Andrejev et al., 2004). At the same time, if the wind forcing favorable for upwelling events near the southern coast prevailed (as it was observed in summer 2010) the low salinity water appeared in the southern part of the open gulf, close to the upwelling front. This phenomenon was also observed during an intense upwelling event in August 2006 when a very intense upwelling event developed near the Estonian coast (Lips et al., 2009), it was modelled by Laanemets et al. (2011) and has been noted by Liblik and Lips (2016) when analyzing based on an analysis data of CTD data from surveys across the gulf from in 2006-2013.




Figure 68. Wind roses Polar histogram of wind stress vectors ($N m^{-2}$) based on the wind data from a weekly period before the peak of upwelling events off the Estonian coast on 19 August 2010 and 8 July 2011.

676 The most intense upwelling events in regard of temperature deviations were observed near the
677 southern coast as it was also found by Uiboupin and Laanemets (2009, 2015). However, we did
678 not identify clear differences in the temperature distribution patterns between the upwelling
679 events off the two coasts. Instead, near the both coasts the classical distribution with a sharp
680 temperature front as well as the distribution characterized with a gradual decrease of temperature
681 towards the coast have been observed. We analyzed the wind data to find out whether the forcing
682 would be the reason of such different outcomes. Since often the wind conditions were quite
683 variable before the upwelling events, it was not possible to suggest any quantitative criterion for
684 wind forcing generating one or the other type of temperature distribution.

685
686 However, the wind speed was usually higher before the upwelling events with the sharp
687 temperature front. For instance, the wind roses for a weekly period before the upwelling event
688 shown in Fig. 8 are very similar except the wind speed distribution – the period before the
689 upwelling event with sharp temperature front observed on 19 August 2010 has a large share of
690 wind speed in the range of 10 to 15 m s⁻¹. It allows us to suggest that the observed variability in
691 spatial temperature distribution at the scales of a few kilometers could be related to sub-
692 mesoscale motions, which are made visible if due to the slightly lower forcing the mesoscale
693 dynamics do not fully dominate. These suggestions have to be studied further in the future by
694 combining Ferrybox data and data on vertical structure of the water column and its temporal
695 evolution since the upwelling development is very much dependent on the vertical structure of
696 the water column before the event.

697
698 In conclusion, we showed that Ferrybox data from the Tallinn-Helsinki ferry route could be
699 successfully employed to describe the characteristics of coastal upwelling events in the Gulf of
700 Finland. ~~We took~~An advantage of the geographical location of the ferry route across the
701 relatively narrow gulf and the schedule consisting of two crossings a day allowed to control the
702 quality of the data and introduce the ~~daily~~ upwelling index based on the data from a single
703 crossing and the cumulative upwelling index. In total, 33 coastal upwelling events were
704 identified in May-September 2007-2013. It is shown that the upwelling occurrences ~~on of~~ 18 %
705 ~~of days near the northern coast~~ and 17 % of days ~~near the southern coast~~ as well as intensities of
706 upwelling events are similar near the northern and southern coast. The most intense events occur

Formatted: Highlight

707 in July-August, most probably because of the warmest surface layer (strongest thermocline)
708 during those months. It is shown that the wind impulse needed to generate upwelling events of
709 similar intensity differ between the two coastal areas. We suggest that the thermohaline structure
710 of the Gulf of Finland is adapted to the prevailing forcing and rather the deviation from the
711 average wind forcing than the absolute value of it should be considered when comparing the
712 wind impulses related to the upwelling generation. Two types of upwelling events were
713 identified – one characterized by a strong temperature (upwelling) front and the other revealing
714 gradual decrease of temperature from open to coastal area with maximum temperature deviation
715 very close to the shore. We suggest that the latter case is a result of slightly lower wind speeds,
716 which are not able to generate strong enough Ekman drift in the entire surface layer down to the
717 seasonal thermocline. This conclusion is supported by the presence of spatial variations in
718 temperature with scales of a few kilometers, which could be signs of sub-mesoscale motions
719 associated with the development of such upwelling events.

Formatted: Highlight

727 Acknowledgements

728
729 We thank our colleagues, especially Inga Lips and Fred Buschmann, for their help in maintaining
730 the Ferrybox system, and Taavi Liblik for his suggestions in regard of data processing. This
731 work was supported by institutional research funding IUT19-6 of the Estonian Ministry of
732 Education and Research and by EU Regional Development Foundation, Environmental
733 Conservation and Environmental Technology R&D Programme project VeeOBS (3.2.0802.11-
734 0043).

736 References

737

738 Alenius, P., Myrberg, K., Nekrasov, A. 1998. The physical oceanography of the Gulf of Finland:
 739 a review. *Boreal Environ. Res.*, 3, 97-125.

740 ~~Alenius, P., Nekrasov, A., Myrberg, K. 2003. Variability of the baroclinic Rossby radius in the~~
 741 ~~Gulf of Finland. *Cont. Shelf Res.*, 23, 563-573.~~

742 Andrejev, O., Myrberg, K., Alenius, P., Lundberg, P.A., 2004. Mean circulation and water
 743 exchange in the Gulf of Finland - a study based on three-dimensional modeling. *Boreal*
 744 *Environ. Res.*, 9(1), 1-16.

745 Buijsman, M.C., Ridderinkhof, H., 2007. Long-term ferry-ADCP observations of tidal currents
 746 in the Marsdiep inlet. *J. Sea Res.*, 57, 237-256.

747 Haapala, J. 1994. Upwelling and its influence on nutrient concentration in the coastal area of the
 748 Hanko Peninsula, entrance of the Gulf of Finland. ~~*Estuarine, Coastal and Shelf*~~
 749 ~~*Science*~~, 38(5), 507-521.

750 Hardman-Mountford, N. J., Moore, G., Bakker, D. C. E., Watson, A. J., Schuster, U., Barciela,
 751 R., Hines, A., Moncoiffe', G., Brown, J., Dye, S., Blackford, J., Somerfield, P. J., Holt,
 752 J., Hydes, D. J., and Aiken, J. 2008. An operational monitoring system to provide
 753 indicators of CO₂- related variables in the ocean. – *ICES Journal of Marine Science*, 65,
 754 1498-1503.

755 Keevalik, S., ~~and~~ Soomere, T., 2010. Towards quantifying variations in wind parameters across
 756 the Gulf of Finland. *Estonian Journal of Earth Sciences*, 59(4), 288 - 297.

757 ~~Kononen, K. 1992. Dynamics of the cyanobacterial blooms in the Baltic Sea. PhD thesis. *Finn.*~~
 758 ~~*Mar. Res.* 261, 3-36.~~

759 ~~Kononen, K., Kuparinen, J., Mäkela, K., Laanemets, J., Pavelson, J., Nömmann, S., 1996.~~
 760 ~~*Initiation of cyanobacterial blooms in a frontal region at the entrance to the Gulf of*~~
 761 ~~*Finland, Baltic Sea. *Limnol. Oceanogr.*, 41, 98-112.*~~

762 ~~Laanemets, J., Väli, G., Zhurbas, V., Elken, J., Lips, I., Lips, U., 2011. Simulation of mesoscale~~
 763 ~~structures and nutrient transport during summer upwelling events in the Gulf of Finland~~
 764 ~~in 2006 *Boreal Environ. Res.*, 16A, 15-26.~~

765 Laanemets, J., Zhurbas, V., Elken, J., Vahtera, E., 2009. Dependence of upwelling-mediated
 766 nutrient transport on wind forcing, bottom topography and stratification in the Gulf of
 767 Finland: model experiments. *Boreal Environ. Res.*, 14, 213-225.

768 [Lehmann, A., Myrberg, K., Höfllich, K., 2012. A statistical approach to coastal upwelling based](#)
 769 [on the analysis of satellite data for 1990-2009. *Oceanologia*, 54, 369-393.](#)
 770 [Lehmann, A., Myrberg, K., 2008. Upwelling in the Baltic Sea – A review. *J. Marine Syst.*, 74,](#)
 771 [S3-S12.](#)
 772 Liblik, T., Lips, U., 2016⁵. Variability of pycnoclines in a three-layer, large estuary: the Gulf of
 773 Finland. *Est. Coast. Shelf Sci. Boreal Environ. Res.* (in press).
 774 **Liblik, T., Lips, U., 2012.** Variability of synoptic-scale quasi-stationary thermohaline
 775 stratification patterns in the Gulf of Finland in summer 2009. *Ocean Sci.*, 8, 603 - 614.
 776 Liblik, T., Lips, U., 2011. Characteristics and variability of the vertical thermohaline structure in
 777 the Gulf of Finland in summer. *Boreal Environ. Res.*, 16A, 73-83.
 778 Lips, I., Lips, U. 2008. Abiotic factors influencing cyanobacterial bloom development in the
 779 Gulf of Finland (Baltic Sea). *Hydrobiologia*, 614, 133–140.
 780 Lips, I., Lips, U., Liblik, T. 2009. Consequences of coastal upwelling events on physical and
 781 chemical patterns in the central Gulf of Finland (Baltic Sea). *Cont. Shelf Res.* 29, 1836-
 782 1847.
 783 Lips, U., Lips, I., Kikas, V., Kuvaldina, N., 2008. Ferrybox measurements: a tool to study meso-
 784 scale processes in the Gulf of Finland (Baltic Sea). *US/EU-Baltic Symposium, Tallinn,*
 785 *27-29 May, 2008. IEEE, (IEEE Conference Proceedings)*, 1 - 6.
 786 Männik, A., Merilain, M., 2007. Verification of different precipitation forecasts during extended
 787 winter-season in Estonia. *HIRLAM Newsletter*, No. 52, 65–70.
 788 [Myrberg, K., Lehmann, A., Raudsepp, U., Szymelfenig, M., Lips, I., Lips, U., Matciak, M.,](#)
 789 [Kowalewski, M., Krezel, A., Burska, D., Szymanek, L., Ameryk, A., Bielecka, L.,](#)
 790 [Bradtke, K., Galkowska, A., Gromisz, S., Jedrasik, J., Kaluzny, M., Kozłowski, L.,](#)
 791 [Krajewska-Soltys, A., Oldakowski, B., Ostrowski, M., Zalewski, M., Andrejev, O.,](#)
 792 [Suomi, I., Zhurbas, V., Kauppinen, O.-K., Soosaar, E., Laanemets, J., Uiboupin, R.,](#)
 793 [Talpsepp, L., Golenko, M., Golenko, N., Vahtera, E., 2008. Upwelling events, coastal](#)
 794 [offshore exchange, links to biogeochemical processes – Highlights from the Baltic Sea](#)
 795 [Science Congress at Rostock University, Germany, 19-22 March 2007. *Oceanologia*, 50,](#)
 796 [95-113.](#)
 797 Myrberg, K., Andrejev, O. 2003. Main upwelling regions in the Baltic Sea – a statistical analysis
 798 based on three-dimensional modeling. *Boreal Environ. Res.*, 8(2), 97-112.

Formatted: Highlight

799 [Paerl, H.W., Rossignol, K.L., Guajardo, R., Hall, N.S., Joyner, A., Peierls, B.L., Ramus, J.S.](#)
800 [Paerl, H., et al.](#) 2009. FerryMon: Ferry-Based Monitoring and Assessment of Human and
801 Climatically Driven Environmental Change in the Albemarle-Pamlico Sound System.
802 *Environ. Sci. Technol.*, 43, 7609–7613

803 Pavelson, J., Laanemets, J., Kononen, K., S. Nõmman, 1997. Quasi-permanent density front at
804 the entrance to the Gulf of Finland: Response to wind forcing. *Cont. Shelf Res.*, 17, 253-
805 265.

806 Petersen, W., 2014. FerryBox systems: State-of-the-art in Europe and future development.
807 *J.ournal of Marine Syst.*, 140, 4-12.

808 [Petersen W., Wehde, H., Krasemann, H., Colijn, F., Schroeder, F., 2008. FerryBox and MERIS –](#)
809 [Assessment of coastal and shelf sea ecosystems by combining in situ and remotely sensed](#)
810 [data. *Est. Coast. Shelf Sci.*, 77, 296-307.](#)

811

812 Rantajärvi, E. (Ed.) 2003. Alg@line in 2003: 10 years of innovative plankton monitoring and
813 research and operational information service in the Baltic Sea. *MERI – Report Series of*
814 *the Finnish Institute of Marine Research*, No. 48, 1-36.

815 Schneider, B., Gülzow, W., Sadkowiak, B., Rehder, G., 2014. Detecting sinks and sources of
816 CO₂ and CH₄ by ferrybox-based measurements in the Baltic Sea: Three case studies. *J.*
817 *Marine Syst.*, 140, 13-25.

818 Seppälä, J., Ylöstalo, P., Kaitala, S., Hällfors, S., Raateoja, P., Maunula, P., 2007. Ship-of-
819 opportunity based phycocyanin fluorescence monitoring of the filamentous cyanobacteria
820 bloom dynamics in the Baltic Sea. *Est. Coast. Shelf Sci.*, 73, 489-500.

821 Talpsepp, L., Nõges, T., Raid, T., Kõuts, T. 1994. Hydrophysical and hydrobiological processes
822 in the Gulf of Finland in summer 1987 – characterization and relationship. *Cont. Shelf*
823 *Res.*, 14, 749-763.

824 Uiboupin, R., Laanemets, J., 2009. Upwelling characteristics derived from satellite sea surface
825 temperature data in the Gulf of Finland, Baltic Sea, *Boreal Environ. Res.*, 14 (2), 297-
826 304.

827 Uiboupin, R., Laanemets, J., 2015. Upwelling parameters from bias-corrected composite satellite
828 SST maps in the Gulf of Finland (Baltic Sea). *IEEE Geoscience and Remote Sensing*
829 *Letters*, 12, 592-596.

830 Vahtera, E., Laanemets, J., Pavelson, J., Huttunen, M., Kononen, K., 2005. Effect of upwelling
831 on the pelagic environment and bloom-forming cyanobacteria in the Western Gulf of
832 Finland, Baltic Sea. *J. Marine Syst.*, 58, 67-82.
833 Väli, G., 2011. Numerical experiments on matter transport in the Baltic Sea. PhD thesis, Tallinn
834 Technical University Press.
835 Väli, G., Zhurbas, V., Laanemets, J., Elken, J., 2011. Simulation of nutrient transport from
836 different depths during an upwelling event in the Gulf of Finland. *Oceanologia*, 53, 431-
837 448.

838
839

840
841
842

843 **Table 1.** Periods of measurements along the ferry route Tallinn-Helsinki in 2007-2013, number
844 of days with measurements and number of days with upwelling events off the northern coast (N)
845 and off the southern coast (S).

Year	Ferry	Period	Number of days with data	Number of days with upwelling	
				N	S
2007	Galaxy	1 May – 30 September	141	26	21
2008	Galaxy	1 May – 13 July	90	8	11
	Baltic Princess	13 August – 30 September			
2009	Baltic Princess	1 May – 30 September	145	33	30
2010	Baltic Princess	1 May – 30 September	140	5	32
2011	Baltic Princess	1 May – 30 September	135	19	30
2012	Baltic Princess	1 May – 28 August	113	22	0
2013	Silja Europa	15 July – 30 September	74	37	16

846
847

848 **Table 2.** Characteristics of detected upwelling events; dates, coastal area (N – off northern coast;
849 S – off southern coast), [type \(UF – with strong upwelling front, GD - - with gradual decrease of](#)
850 [temperature\)](#), maximum temperature deviation from the transect mean value, cumulative

851 upwelling index calculated for each event and cumulative along-gulf wind stress calculated for
852 upwelling favourable winds before and during the upwelling event.

No	Dates	Coast	Maximum temperature deviation (°C)	Cumulative upwelling intensity (°C day)	Cumulative wind stress (N m ⁻² day)	
1.	3-14 June 2007	S	-4.12	-19.8	-0.49	
2.	8-16 July 2007	S	-3.02	-12.6	-0.34	
3.	21-27 July 2007	N	-4.02	-13.9	0.93	
4.	29 July—8 August 2007	N	-3.64	-16.5	0.38	
5.	10-17 September 2007 ⁽¹⁾	N	-1.97	-7.5	0.75	
6.	26-28 May 2008 ⁽²⁾	S	-2.52	-3.9	-0.20	
7.	11-15 June 2008	N	-2.73	-7.2	0.62	
8.	27-29 June 2008	N	-2.27	-6.2	0.53	
9.	10-17 September 2008	S	-5.42	-23.0	-1.08	
10.	9-16 June 2009	S	-4.77	-14.8	-0.27	
11.	24 June—14 July 2009	S	-5.78	-36.1	-0.42	
12.	16-22 August 2009	N	-3.20	-10.7	0.54	
13.	28 August—9 September 2009	N	-2.74	-14.1	0.56	
14.	17-30 September 2009 ⁽³⁾	N	-3.09	-19.3	1.28	
15.	20-24 May 2010	S	-2.21	-5.1	-0.56	
16.	12-13 June 2010 ⁽⁴⁾	S	-2.60	-2.3	-0.19	
17.	20-24 July 2010	N	-4.70	-9.3	0.31	
18.	26 July—1 August 2010	S	-6.19	-15.7	-0.34	
19.	17-23 August 2010	S	-7.78	-20.8	-0.66	
20.	2-12 September 2010	S	-5.27	-16.0	-0.25	
21.	4-12 May 2011 ⁽⁵⁾	S	-2.22	-9.3	-0.09	
22.	31 May—8 June 2011	N	-2.32	-10.3	0.60	
23.	11-15 June 2011	S	-3.12	-6.0	-0.38	
24.	24-27 June 2011	N	-2.40	-4.8	0.41	
25.	5-10 July 2011	S	-5.05	-10.6	-0.38	
26.	29 July—7 August 2011	S	-4.69	-22.2	-0.62	
27.	14 September 2011 ⁽⁶⁾	N	-4.90	-3.1	0.47	
28.	26-30 September 2011 ⁽⁷⁾	N	-3.27	-13.8	1.26	
29.	18-27 July 2012 ⁽⁸⁾	N	-4.55	-22.4	1.37	
30.	2-13 August 2012	N	-4.17	-22.2	0.58	
31.	17 July—1 August 2013 ⁽⁹⁾	N	-6.15	-26.0	0.63	
32.	11-31 August 2013	N	-5.03	-39.7	0.92	
33.	15-30 September 2013	S	-7.34	-40.2	-0.71	
No	Dates	Coast	Type	Maximum temperature deviation (°C)	Cumulative upwelling intensity (°C day)	Cumulative wind stress (N m ⁻² day)

1.	3-14 June 2007	S	UF	-4.12	-19.8	-0.49
2.	8-16 July 2007	S	GD	-3.02	-12.6	-0.34
3.	21-27 July 2007	N	UF	-4.02	-13.9	0.93
4.	29 July – 8 August 2007	N	GD	-3.64	-16.5	0.38
5.	10-17 September 2007⁽¹⁾	N	GD	-1.97	-7.5	0.75
6.	26-28 May 2008⁽²⁾	S	UF	-2.52	-3.9	-0.20
7.	11-15 June 2008	N	UF	-2.73	-7.2	0.62
8.	27-29 June 2008	N	UF	-2.27	-6.2	0.53
9.	10-17 September 2008	S	UF	-5.42	-23.0	-1.08
10.	9-16 June 2009	S	UF	-4.77	-14.8	-0.27
11.	24 June – 14 July 2009	S	GD	-5.78	-36.1	-0.42
12.	16-22 August 2009	N	UF	-3.20	-10.7	0.54
13.	28 August – 9 September 2009	N	UF	-2.74	-14.1	0.56
14.	17-30 September 2009⁽³⁾	N	UF	-3.09	-19.3	1.28
15.	20-24 May 2010	S	GD	-2.21	-5.1	-0.56
16.	12-13 June 2010⁽⁴⁾	S	UF	-2.60	-2.3	-0.19
17.	20-24 July 2010	N	UF	-4.70	-9.3	0.31
18.	26 July – 1 August 2010	S	UF	-6.19	-15.7	-0.34
19.	17-23 August 2010	S	UF	-7.78	-20.8	-0.66
20.	2-12 September 2010	S	GD	-5.27	-16.0	-0.25
21.	4-12 May 2011⁽⁵⁾	S	GD	-2.22	-9.3	-0.09
22.	31 May – 8 June 2011	N	UF	-2.32	-10.3	0.60
23.	11-15 June 2011	S	UF	-3.12	-6.0	-0.38
24.	24-27 June 2011	N	UF	-2.40	-4.8	0.41
25.	5-10 July 2011	S	GD	-5.05	-10.6	-0.38
26.	29 July – 7 August 2011	S	GD	-4.69	-22.2	-0.62
27.	14 September 2011⁽⁶⁾	N	UF	-4.90	-3.1	0.47
28.	26-30 September 2011⁽⁷⁾	N	UF	-3.27	-13.8	1.26
29.	18-27 July 2012⁽⁸⁾	N	GD	-4.55	-22.4	1.37
30.	2-13 August 2012	N	UF	-4.17	-22.2	0.58
31.	17 July – 1 August 2013⁽⁹⁾	N	UF	-6.15	-26.0	0.63
32.	11-31 August 2013	N	GD	-5.03	-39.7	0.92
33.	15-30 September 2013	S	UF	-7.34	-40.2	-0.71

853

854 ⁽¹⁾ temperature deviation was less than -2 °C during the event on 10-17 September 2007

855 ⁽²⁾ data absent before 26 May 2008 for more than 1 day

856 ⁽³⁾ data analysed until 30 September 2009 (upwelling event did further)

857 ⁽⁴⁾ data absent before 12 June 2010 for more than 1 day

858 ⁽⁵⁾ early spring with possible contribution of difference in surface water warming

859 ⁽⁶⁾ no data available after 14 September 2011

860 ⁽⁷⁾ no data available before 26 September 2011, wind data missing on 24-26 September 2011

861 ⁽⁸⁾ wind data on 14-15 July 2012 not available
862 ⁽⁹⁾ ferrybox data on 20-21 July 2013 not available
863
864
865
866
867
868
869
870
871
872

873 **Figure captions**
874

875 **Figure 1.** Map of the Baltic Sea (a) and the study area with the Ferrybox transect and
876 Kalbadagrund meteorological station.
877

878 ~~**Figure 2.** Measured salinity distribution along the ferry route Tallinn-Helsinki from the forth and~~
879 ~~backward journey on 27 July 2010. Raw data are presented in the left panel and the processed~~
880 ~~data in the right panel where the shifting of data points by 3 minutes and 20 seconds was applied;~~
881 ~~x-axis shows the distance from the Tallinn Bay (latitude 59.48 N) in km along the meridional~~
882 ~~transect.~~
883

884 **Figure 23.** Temporal changes of temperature (in °C) and salinity (in g kg⁻¹) distributions between
885 Tallinn and Helsinki from 1 May to 30 September in 2007 (a, b), 2008 (c, d), 2009, (e, f), 2010
886 (g, h), 2011 (i, j), 2012 (k, l) and 2013 (m, n); y-axis shows the distance from the Tallinn Bay
887 (latitude 59.48 N) in km along the meridional transect.
888

889 **Figure 43.** Distributions of temperature (in °C) and salinity (in g kg⁻¹) deviations from the daily
890 transect mean value along the ferry route Tallinn-Helsinki for all measurements in May-
891 September 2007-2013 (a, b), in 2009 (c, d) and in 2010 (e, f). Mean values on each 0.5-km cell
892 (solid curves) and plus/minus RMSE (dashed curves) are shown; x-axis shows the distance from
893 the Tallinn Bay (latitude 59.48 N) in km along the meridional transect.

894
895 | **Figure 54.** Temporal changes of spatial distributions of temperature deviations (in °C) from the
896 daily transect mean value between Tallinn and Helsinki from 1 May to 30 September in 2007 (a),
897 2008 (b), 2009, (c), 2010 e), 2011 (f), 2012 (g) and 2013 (h); y-axis shows the distance from the
898 Tallinn Bay (latitude 59.48 N) in km along the meridional transect.

899
900 | **Figure 65.** Temporal changes of upwelling index off the northern coast (upper columns; °C) and
901 off the southern coast (lower columns, °C) and along-gulf wind stress (black curve in the middle;
902 N m^{-2}) in May-September 2007 (a), 2008 (b), 2009 (c), 2010 (d), 2011 (e), 2012 (f) and 2013 (g).

903
904 | [Figure 6. Relationship between the cumulative upwelling index \(CUI\) and cumulative along-](#)
905 [gulf wind stress \(CWS\) based on 33 detected upwelling events in May-September 2007-2013.](#)
906 [Red symbols indicate the events off the southern coast and blue symbols the events off the](#)
907 [northern coast; circles correspond to the events with pronounced upwelling front \(N_UF and](#)
908 [C_UF\) and triangles the events with gradual decrease of temperature towards the coast \(N_GD](#)
909 [and S_GD\).](#)

910
911 | **Figure 7.** Characteristic distributions of temperature and salinity along the ferry route Tallinn-
912 Helsinki with coastal upwelling events off the northern coast (a, b) and off the southern coast (c,
913 d); x-axis shows the distance from the Tallinn Bay (latitude 59.48 N) in km along the meridional
914 transect.

915
916 | **Figure 8.** ~~Wind roses~~ [Polar histogram of wind stress vectors \(\$\text{N m}^{-2}\$ \)](#) based on the wind data from
917 a weekly period before the peak of upwelling events off the Estonian coast on 19 August 2010
918 and 8 July 2011.

919