

Interactive comment on “Acoustic Doppler Current Profiler observations in the southern Caspian Sea: shelf currents and flow field off Freidoonkenar Bay, Iran” by P. Ghaffari and V. Chegini

Y. Liu

yliu18@gmail.com

Received and published: 12 March 2010

Depth-averaged long-term mean flow tends to orient along-isobath on a continental shelf, and the vertical profiles of the mean across-shore currents usually show a two-layer structure (e.g., offshore flow in the near-surface layer and onshore flow in the near-bottom layer, and vice versa) on the inner shelf (e.g., Lentz 2008; Weisberg et al. 2009). However, as shown in Figure 7, the mean across-shore current component has positive values (northward) throughout the water column, which indicates that the depth-averaged mean across-shore current does not vanish. Actually, the depth-

C1137

averaged mean across-shore current is about 1 cm/s, not that smaller than the depth-averaged mean along-shore current (3~4 cm/s) in Figure 7. This is counter-intuitive, given the fact that the time series is not short (57 days) and the water depth is only 14 m at this site.

The along-shore direction is defined as “the coast direction”, 12 degrees from the east. Note that the across-shore component of the current on a continental shelf is sensitive to the choice of the coordinate system. It would be indicative to see the difference between this choice (12 deg) and those determined from conventional methods: (1) principal axis of subtidal depth-averaged currents (e.g., Liu and Weisberg 2005) and (2) direction of mean depth-averaged currents (e.g., Lentz 2008; Weisberg et al. 2009).

Also in Figure 7, the mean across-shore current shows a much smaller value at the top-most level (2 m). For a bottom-mounted ADCP with 1-m bin, the data at this 2-m level (below the surface) may not be reliable at all, because it may be affected by the surface reflections of the acoustic echoes. The data at the top 2-m levels are usually discarded for scientific analysis (e.g., Liu and Weisberg 2005, 2007; Weisberg et al. 2009).

In Figure 8, the units of the power spectra should be $(\text{m/s})^2/\text{cph}$ and $(\text{cm/s})^2/\text{cph}$ for the winds and currents, respectively.

References:

Lentz, S. J., 2008. Observations and a model of the mean circulation over the Middle Atlantic Bight continental shelf. *Journal of Physical Oceanography* 38, 1203–1221, doi:10.1175/2007JPO3768.1.

Liu, Y., Weisberg, R.H., 2005. Momentum balance diagnoses for the West Florida Shelf. *Continental Shelf Research*, 25, 2054-2074.

Liu, Y., Weisberg, R.H., 2007. Ocean currents and sea surface heights estimated across the West Florida Shelf. *Journal of Physical Oceanography*, 37(6), 1697-1713.

C1138

Weisberg, R.H., Liu, Y., Mayer, D.A., 2009. West Florida Shelf mean circulation observed with long-term moorings. *Geophysical Research Letters*, 36, L19610, doi:10.1029/2009GL040028.

Interactive comment on *Ocean Sci. Discuss.*, 6, 3019, 2009.

C1139