

This discussion paper is/has been under review for the journal Ocean Science (OS).  
Please refer to the corresponding final paper in OS if available.

# A Study of the hydrographic conditions in the Adriatic Sea from numerical modelling and direct observations (2000–2008)

**P. Oddo and A. Guarnieri**

Istituto Nazionale di Geofisica e Vulcanologia, Viale Aldo Moro 44, Bologna, Italy

Received: 11 February 2011 – Accepted: 15 March 2011 – Published: 22 March 2011

Correspondence to: P. Oddo (odd@bo.ingv.it)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**OSD**

8, 565–611, 2011

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The inter-annual variability of Adriatic Sea hydrographic characteristics is investigated by means of numerical simulation and direct observation. The period investigated runs from the beginning of 2000 to the end of 2008. The model used to carry out the simulation is derived from the primitive equation component of the Adriatic Forecasting System (AFS). The model is based on the Princeton Ocean Model (POM) adapted in order to reproduce the features of the Adriatic. Both numerical findings and observations agree in depicting a strong inter-annual variability in the entire Adriatic Sea and its sub-basins. The dense water formation process has been found to be intermittent. In addition to inter-annual variability, a long-scale signal has been observed in the salinity content of the basin as a consequence of a prolonged period of reduced Po river runoff and high evaporation rates. As a result, the temperature and salinity of the northern Adriatic dense water vary considerably between the beginning and the end of the period investigated.

1 Introduction

The hydrographic characteristics of the Adriatic Sea (Fig. 1) are known to be subject to great spatial and temporal (both seasonal and inter-annual) variability. The nature of this variability is related to the characteristics of its geometry, its geographical position, the geography of the surrounding areas and, clearly, to the surface and lateral forcing functions acting on the basin. The Adriatic Sea extends northwest from 40° to 45°5' North, with an extreme length of about 800 km and an average width of about 160 km. The southern limit is delimited by the Strait of Otranto, which connects it to the south with the Ionian Sea.

The northern part is characterised by a shallow mean depth (about 30 m) with a very weak bathymetric gradient (Russo and Artegiani, 1996). The central part of the basin, spanning approximately from the 100 m contour to the Palagruža sill (about

Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



170 m depth), is characterised by two depressions, the Pomo (or Jabuka) Pits, having a maximum depth of about 270 m. The southern part of the Adriatic Sea extends from the Palagruža sill to the Otranto sill; both western and eastern coasts have a narrow continental shelf and a steep continental slope reaching a maximum depth of 1270 m (often referred to as the south Adriatic Pit) (Russo and Artegiani, 1996). The river runoff is a significant component of the basin hydrological cycle and is responsible for its net freshwater gain (Raicich, 1994, 1996), implying an average estuarine thermohaline circulation. The freshwater discharge is concentrated particularly in the northern sub-basin, where the river Po constitutes the main freshwater source. However, in the southern basin, Albanian and Croatian rivers provide a significant freshwater input (Raicich, 1994; Oddo et al., 2005).

Prevailing wind regimes are the Bora (northeast), the Sirocco (southern) and Maestral (western). The Sirocco wind, characterized by a high humidity content, blowing from the southeast drives the sea level of the north Adriatic Sea. The Bora wind, which can be dry, cold, and strong, blows over the Adriatic Sea from the north and north-east. In summertime, besides local breezes, the dominant wind – the Maestral – comes from the northwest.

The climatological annual heat budget of the Adriatic basin is negative. Several values have been proposed in the literature; these vary as a consequence of different methodologies used and different periods investigated. Mantziafou and Lascaratos (2008) found values ranging between  $-6$  to  $-34 \text{ W m}^{-2}$ ; similar values have also been suggested by Cardin and Gacic (2003); Maggiore et al. (1998) suggest values in the range between  $-5$  and  $-10 \text{ W m}^{-2}$ ; Chiggiato et al. (2005) by comparing different data source and different computational methods found values in the range of  $-18$  and  $-54 \text{ W m}^{-2}$ . All the authors agree in suggesting that the total heat flux exhibits a strong inter-annual variability, particularly marked during the periods of negative heat budget (autumn–winter).

The resulting general circulation of the Adriatic Sea is cyclonic and highly variable according to the season (Artegiani et al., 1997a, b; Malanotte-Rizzoli and

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bergamasco, 1983; Zavatarelli et al., 2002; Zavatarelli and Pinardi, 2003). The climatological circulation pattern is composed of well-known currents and gyre structures (Artegiani et al., 1997b; Poulain, 2001; Zavatarelli et al., 2002; Zavatarelli and Pinardi, 2003). The three main gyres (Southern, Middle and Northern Adriatic Gyre, 5 Artégiani et al., 1997b) are interconnected (with seasonally varying characteristics) by two coastal currents, one flowing southward along the Italian coast from the Po delta to the Strait of Otranto (Western Adriatic Coastal Current or WACC), the other flowing northward from the Strait of Otranto along the eastern coast and reaching the central Adriatic (Eastern Southern Adriatic Current or ESAC) (Oddo et al., 2005).

10 The Adriatic Sea is usually divided into four sub-basins (Artegiani et al., 1997a; Artégiani et al., 1999): the WACC region, between the Po delta and Strait of Otranto along the Italian coast; the Northern region, delimited by 100m bathymetry north of the Jakuba Pit; the Middle Adriatic region, with the southern boundary across the basin starting from the Gargano Peninsula; and the Southern Adriatic region. The limits of 15 these sub-regions are depicted in Fig. 1.

The Adriatic Sea has been the main source of dense water masses for the Eastern Mediterranean basin for many decades, until the late nineteen-eighties, when the Aegean Sea became the main supplier of Eastern Mediterranean Dense Water (EMDW) (Roether et al., 1996). Nevertheless, it has been demonstrated that although the Adriatic contribution to EMDW might have been temporarily reduced, it has never ceased (Klein et al., 2000; Nardelli and Salusti, 2000; Sellschop and Alvarez, 20 2003).

Dense waters are formed locally in the Northern, Middle and Southern Adriatic. However, while the formation of the dense water mass in the Southern and Northern sub-basins is reasonably well detected in the hydrographic observations, the local formation process in the Middle sub-basin is not well documented from in-situ data (Bergamasco et al., 1999). The Northern Adriatic dense waters (hereafter NAdw) are formed in winter during outbreaks of the bora winds that provoke strong surface cooling and evaporation processes (Bergamasco et al., 1999), and are also at the same time under 25

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a significant inter-annual variability (Gacic et al., 2002; Mantziafou and Lascaratos, 2008).

Formation rates of the Adriatic dense water have been investigated in several studies (Artegiani et al., 1989; Artegiani and Salusti, 1987; Vested et al., 1998; Mantziafou and Lascaratos, 2004, 2008). Numerically- and observationally-based estimates of NAdw formation rates suggest values of about 0.03–0.07 Sv; similar derivations for the MAdw are about 0.06–0.07 Sv, whereas in the Southern Adriatic, dense water formation rate ranges from 0.18 to 0.64 Sv. In all three sub-basins, large inter-annual variability in the dense water formation rate has been observed (Mantziafou and Lascaratos, 2008; Vilibic and Supic, 2005).

This work explores the inter-annual variability of the Adriatic Sea hydrographic conditions connected to atmospheric forcing through the analysis of numerical simulation results and direct and remote observations. We concentrate on the period 2000–2008, during which several dedicated oceanographic cruises extensively monitored the Adriatic Sea, and satellite-derived Sea Surface Temperature data were also available. The Adriatic Sea hydrodynamic is simulated by a three-dimensional model already used by Oddo et al. (2005) to investigate the response of the Adriatic Sea circulation to variable atmospheric forcing; this later implemented in the operational chain of the Adriatic Forecasting System (AFS, Oddo et al., 2006) and continuously developed within the framework of international projects (Guarnieri et al., 2008). The version of the model used in this work has been developed within the ECOOP (European COastal sea Operational observing and forecasting system) project as modelling component of the new AFS. Major differences with previous model versions regard the increased horizontal and vertical resolution and a new parameterization of the vertical and lateral open boundary conditions. The current model version has been used to produce on daily bases operational products (simulation and forecast) for the Adriatic basin.

Section 2 gives a general description of the model's implementation. In Sect. 3 the model results and available observed data are discussed. Summary and conclusions are offered in the last section.

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2 Model description

Numerical integration was carried out using a pre-operational version of the Primitive Equation module of the Adriatic Forecasting System (AFS, <http://gnoo.bo.ingv.it/afs>, Oddo et al., 2006). This model is based on the Princeton Ocean Model (POM, Blum-  
berg and Mellor, 1987), and was implemented with a regular horizontal grid having  
approximately 2.2 km resolution ( $1/45^\circ$  of latitude and longitude) and 27 sigma layers  
in the vertical. The eddy viscosity coefficients are provided by the Smagorinsky (1993)  
parameterization implemented into POM according to Mellor and Blumberg (1985) and  
corresponding diffusivity terms are obtained through the Prandtl number (which is set  
equal to 0.001). The vertical mixing coefficients for momentum and tracers are calcu-  
lated using the Mellor and Yamada (1982) turbulence closure scheme.

The model lateral open boundary is south of the Strait of Otranto along the  $39^\circ$   
parallel. The bathymetry was obtained from US Navy data (horizontal resolution  $1/60^\circ$ );  
the minimum depth is 10 m and between 0 and 10 m depth the topography slope was  
flattened (that is, the modelled Adriatic has a realistic coastline).

The surface fluxes are interactively computed using model predicted sea surface  
temperature and realistic atmospheric data. The model uses the same bulk formulae  
to compute surface fluxes as described in Oddo et al. (2005). Solar radiation is com-  
puted by means of an astronomical formula (Reed, 1975, 1977), the net long-wave  
radiation flux is computed according May (1986), sensible and latent heat fluxes are  
computed by classical bulk formulae parameterized according to Kondo (1975). The  
wind stress computation uses a drag coefficient computed according to Hellerman and  
Rosenstein (1983). Details on the bulk formulae used can be found in Maggiore et  
al. (1998) and Castellari et al. (1998).

The atmospheric data (air temperature, relative humidity, cloud cover, mean sea level  
pressure and both wind components) are operational analyses with  $0.5^\circ$  horizontal res-  
olution and 6 h frequency provided by the European Centre for Medium-Range Weather  
Forecasts (ECMWF). Despite, the relatively coarse resolution of the atmospheric fields,

OSD

8, 565–611, 2011

### Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

it has been already proved that are sufficient to investigate the inter-annual variability of the Adriatic Sea with good accuracy (Oddo et al. 2005).

Real freshwater input was used in the surface boundary condition for the vertical velocity, which reads:

$$w|_{z=h} - \left( \frac{\partial h}{\partial t} + \bar{v} \cdot \nabla h \right) \bigg|_{z=h} = (E - P - R)$$

where  $w$  is the vertical velocity,  $h$  is the surface elevation,  $E$  is the evaporation,  $P$  is the precipitation and  $R$  indicates the river runoff. The two critical components of the water flux are the precipitation and the runoff. For precipitation we use the climatological monthly mean values from Legates and Wilmott (1990) according to the operational procedures. For river runoff, daily observed values are used for the Po while only monthly climatological values are considered for the other forty-eight sources (rivers and springs) (Raich, 1994). Following the hypothesis made in Oddo et al. (2005) concerning a possible overestimation of these climatological values in the past ten years, we reduced the prescribed data for the Croatian part by a factor of 0.3 (this factor derives from sensitivity studies).

The Po runoff is specified daily taking the values at the closing point of the drainage basin (Pontelagoscu) and partitioned over several grid points approximately representing the proportion of the fresh water discharge through the different branches of the delta (Provini et al., 1992).

Thus, inter-annual variability of the surface water (and consequently salinity) flux is driven only by variation in the evaporation flux and in Po river regimes (which, however, is more than 1/3 of the total runoff of the Adriatic Sea).

A further difference with the previous POM simulations in the Adriatic Sea (Oddo et al., 2005, 2006; Zavatarelli et al., 2002; Zavatarelli and Pinardi, 2003) is related to the advection scheme. A flux limiting advection scheme (MUSCL, Monotonic Upwind Scheme for Conservation Laws, Van Leer, 1979) was implemented, allowing a better reproduction of the horizontal and vertical gradients.

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In order to reduce the pressure gradient error associated with sigma coordinates, a horizontally averaged density before computing the baroclinic integrals is subtracted (Mellor et al., 1998).

For initial and lateral boundary conditions (temperature, salinity and velocity fields), data are taken from a 1/16° horizontal resolution operational analysis of the Mediterranean Forecasting System (MFS, Tonani et al., 2008). The lateral boundary data are provided daily. The definition of the nested open boundary conditions is based on Oddo et al. (2005) and Oddo and Pinardi (2008).

The simulation spans the period from January 1999 to December 2008; results for the year 1999 are not shown as this year is considered to represent the model spin up.

### 3 Results and discussion

Based on the simulated and observed oceanographic characteristics of the Adriatic Sea (Artegiani et al., 1997a, b), 4 regions were defined and analysed separately: the WACC region, between the Po delta and Strait of Otranto along the Italian coast; the Northern region, delimited approximately by 100 m isobaths north of the Jabuka Pit; the Middle region, with the southern boundary across the basin starting from the Gargano Peninsula; and the Southern Adriatic region. The limits of these sub-regions are depicted in Fig. 1. It has to be stressed that the WACC region is defined here as a relatively wide area along the Italian Peninsula potentially also including the western limbs of the cyclonic gyres. This definition is similar to that adopted by Poulain (2001) but differs from that used in Artegiani et al. (1999) and Hopkins et al. (1999).

In addition to satellite-derived Sea Surface Temperature (Marullo et al., 2007), a total of 1617 CTD stations, collected between January 2001 and August 2006, were used to support the model results. However, as hydrographic observations are sparse in space and time (Table 1 gives the total number of CTD casts per region and per year, while in Fig. 1 the sampling positions are shown), direct calculation of inter-annual variability of basin and sub-basin averages will be affected by a large sampling error. When they

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were compared, model results were sampled in the grid point nearest to the sampling position and then both observations and numerical results were averaged over the intervals listed in Table 1.

#### 4 Heat and water fluxes

The analysis of the estimated surface forcings for the period under examination shows an important inter-annual variability in both heat (in Fig. 2 the time series of the total surface heat flux is shown, while in the upper panel of Fig. 3 the advective heat flux at the Strait of Otranto is drawn) and freshwater fluxes (Fig. 3, bottom panel, presents the Po river runoff and Fig. 4 the time series of the evaporation rate).

In the figures, ordinates indicate the monthly values, while colours indicate the amplitude of the anomalies calculated using the monthly climatology obtained from the same simulation. In agreement with previous findings (Oddo et al., 2005; Mantziafou and Lascaratos, 2008) the total surface heat flux variability is larger during cooling events (from November to February) than in periods of net heat gain. In general, the spring-summer heat gain does not change significantly from year to year. On the contrary, the autumn and winter cooling exhibits a strong inter-annual variability mainly due to the latent heat flux component of the surface heat balance. Thus, the inter-annual variability in the maximum volume averaged temperature (Fig. 6), usually occurring in August, is principally a consequence of the heat gains or losses during the preceding months in addition to the differences in the spring-summer heat fluxes. In terms of basin average, only summer 2003 is characterized by surface heat flux significantly larger than the climatology.

The mean annual heat budget of the basin oscillates between negative and positive values, varying from  $-9\text{ W m}^{-2}$  in 2005 to  $19\text{ W m}^{-2}$  in 2003, and thus the advective contribution at the Strait of Otranto (Fig. 3, upper panel) is a fundamental component of the overall heat budget of the basin. Unlike previous studies (Gacic et al., 2002a, b; Manca et al., 2002; Maggiore et al., 1998; Chiggiato et al., 2005; Cardin and Gacic,

### Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2003; Mantziafou and Lascaratos 2008) the surface heat flux averaged over the entire simulated period (January 2000 to December 2008) is slightly positive. There are several possible sources for this difference: primarily, it is worth mentioning that all the previous studies analysed different periods (mostly ranging between 1990 and 1999, with the sole exception of Mantziafou and Lascaratos 2008 which also studied the nineteen-eighties); in addition there are differences in the atmospheric data sets, in the bulk formulae used, in the Sea Surface Temperature (SST) data, as well as possible deficiencies in the model (and its set-up) used in the present study. Unfortunately, the lack of observational field does not allow a direct validation of the model heat budget estimates.

However, the year-to-year pattern is similar in all sub-basins, with the WACC region (Fig. 2b) characterized by the smallest amplitude of both seasonal and inter-annual variability.

In all the basins, the largest anomaly is detected in Autumn 2001 (November and December) with a seasonal heat loss of about  $-295 \text{ W m}^{-2}$ , ranging from  $-315 \text{ W m}^{-2}$  in the Northern region to  $-234 \text{ W m}^{-2}$  in the WACC region. Climatological estimates for the same season are smaller (i.e., Maggiore et al., 1998 suggest values of about  $-225 \text{ W m}^{-2}$ ). During the same period advective heat flux at the Strait of Otranto (Fig. 3, upper panel) is negative ( $-17 \text{ W m}^{-2}$ ), thus acting in the same direction as the surface heat fluxes and amplifying the overall cooling of the basin.

Autumns 2005 and 2007 are also characterized by large heat losses but with different dynamics at the Strait of Otranto. During November and December 2005 the advective Otranto contribution to the heat budget is positive ( $+12 \text{ W m}^{-2}$ ), while during the same months in 2007 a net heat loss of  $20 \text{ W m}^{-2}$  characterizes the model results. The advective contribution can thus strongly influence the thermodynamic of the Adriatic system, amplifying or mitigating the effect of the surface heat exchanges.

Winters between 2003 and 2006 are characterized by large heat losses that are particularly marked during January and February. With the sole exception of the 2003, a contemporaneous net positive advective flux at the Strait of Otranto is observed,

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

partially compensating the local heat losses.

As in previous studies (Oddo et al., 2005), in the period between August 2000 and March 2001 surface heat fluxes are continuously above the climatological values, with a stronger warming during summer and a weaker cooling in autumn and winter. During this period large heat anomalies are not as large as in other periods, however the extended length of this episode has largely affected the thermohaline characteristics of the basin.

Similarly, the period between autumn 2006 and winter 2007 is also characterized by weaker heat losses at the surface; in the Northern region this event is particularly strong in autumn, while in the Middle and Southern sub-basins it is more relevant in winter (2007) and persists until summer.

Starting from 2005 the seasonal amplitude of the heat flux through the Strait of Otranto grows, reaching values almost twice as high as in the previous period; however, the annual mean does not differ significantly between the two periods. The values of the Otranto mass fluxes (not shown) are slightly higher than previous numerical (Mantzi-afou and Lascaratos, 2008) and observational (Gacic et al., 1996) estimates, ranging between 1.3 Sv in 2001 and 1.7 Sv in 2008.

The large inter-annual variability depicted in the total heat flux is mainly due to the variations in the latent heat flux component and consequently to the evaporation rate (Fig. 4). The seasonal cycle of the evaporation flux (Fig. 4) is stable over the years simulated, showing the largest inter-annual variability during the autumn-winter period (and thus strongly contributing to the inter-annual variability of the total heat flux). Starting from autumn–winter 2001–2002, a large positive anomaly is diagnosed by the model during all the cool seasons. This anomaly is larger in the Southern basin than in the coastal areas (WACC and Northern regions). However, in terms of water balance, 2001 and 2002 are characterized by a very large Po river runoff (Fig. 3 bottom panel), compensating the large mass loss occurring at the surface. In the period between 2003 and 2007, on the other hand, the Po regime (Fig. 3 bottom panel) is always below the climatological mean.

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Water mass exchanges at the Strait of Otranto also play a crucial role in the salinity budget of the Adriatic Sea. The relatively fresh waters formed locally, spilling as bottom density currents through the Otranto sill, are compensated by an inflow at intermediate depths of waters of Mediterranean origin. Among others, the Modified Levantine Intermediate Water is fundamental in conditioning the thermohaline characteristics of the Adriatic Sea. The Hovmoeller diagram of the salinity profile zonally averaged along the Strait of Otranto is shown in Fig. 5. The diagram shows low salinity at surface and bottom layers, indicating the outflow of fresh and dense waters, and high salinity at intermediate depths. The black solid line identifies the MLIW and thus a source of salty waters for the Adriatic Sea. The presence of the salty MLIW is irregular. During summer 2000 a small quantity of MLIW is found between 200 and 400 m depths; during summer 2001 a larger amount of MLIW is present at deeper layers. Starting from winter 2002 and until summer 2005 MLIW are absent. Summers 2005 and 2006 are characterized by a noticeable presence of very salty MLIW reaching 800 m depth. Later in 2006 the signal weakens, having a less pronounced salinity maximum, and it tends to disappear in late winter 2008. The described inter-annual variability can be ascribed to the dense waters productions in the Aegean Sea and their effect on the pathway of the LIW (Malanotte-Rizzoli et al., 1999). It is worth mentioning that lateral boundary data are provided by the MFS operational system (Tonani et al., 2008), and thus the number and position of assimilated data, which strongly influence the quality of the operational products, can vary considerably from year to year. However, as will be discussed in the following section, there is observational evidence of MLIW intrusion during late summer 2002.

Again, it is worth to point out that the inter-annual variability in the water budget includes only Po river runoff, evaporation rate and Otranto mass exchanges, since all the other components (precipitation and all the other rivers) are climatological.

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5 Sub-basin temperature and salinity

The time series of volume- averaged temperature for each sub-region and for the whole basin are shown in Fig. 6. As with Figs. 2–3 and 4, the abscissa indicates monthly values, while colours indicate the monthly anomalies computed against the monthly climatology obtained from the simulation.

Starting from late summer 2000 and until winter 2003, the basin-averaged volume temperature is constantly higher than the climatological values. This permanent anomalous condition strongly depends on the high warming rate simulated during spring 2000 and the missing cooling in the autumn–winter 2000–2001 seasons (Oddo et al., 2005). The basin volume temperature rise between the two annual minima in 2000 and 2001 ( $0.8^{\circ}\text{C}$ ) is large enough to mitigate the effect of the strong cooling depicted during autumn 2001 and the volume-averaged temperature persists above the climatological values until spring 2003. Year 2003 is a transition period between 2/3 consecutive years of positive anomalies and 3 years of negative anomalies. However, the first positive period is characterized by relatively short intervals with strong anomalies in the surface fluxes that will then affect a longer period. The second period (2004–2006), characterized by negative anomalies, is stable, on the other hand, with minor year-to-year differences. Like 2000–2001, the period spanning from autumn 2006 to summer 2007 is characterized by weak cooling and strong warming, and thus the volume-averaged temperature minimum in 2007 (occurring on February) and maximum (occurring in late August) are similar to those occurring in 2002.

The described inter-annual variations are strongly modulated in the different Adriatic sub-basins. The sub-basin time-series obtained from the model simulation and the comparison between available observations and model results are shown in Fig. 6 (panels b-c-d and e). Model results were first sampled in the observational positions and then both simulated and observed values were grouped by region and averaged in time according to the time intervals listed in Table 1.

OSD

8, 565–611, 2011

### Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The WACC region (Fig. 6 panel b), extending from North to South and characterized by relatively shallow depths, reacts faster than the deeper regions to the surface fluxes. Differently from the entire Adriatic basin, this region is characterized by an alternation of positive and negative anomalies. The largest inter-annual variability is found in correspondence with the minima of the annual cycle; this is particularly pronounced during two episodes also detected in the whole Adriatic average: the warm autumn-winters 2000–2001 and 2006–2007. As also discussed for the whole basin, the successive summers are also warmer than the climatology. A general good agreement characterizes the comparison with the available observations, particularly during the autumn-winter seasons. During summer a warm bias characterizes the model results. Analysing the vertical structure of the water column (Fig. 7), it is evident that the summer bias (magenta for September–October 2002 and cyan for August 2006) is due to an excessive vertical mixing. The model reproduces the surface values well but between 20 and 50m depth the simulated thermocline is weaker than observed. During winters the surface waters are generally cooler than the bottom layers, thus requiring a compensation effect from salinity in order to maintain the water column stable.

The northern region, as a consequence of the very shallow bathymetry, is characterized by the largest amplitude of the volume-averaged temperature seasonal cycle (Fig. 6 panel c) with minimum values below 10 °C and summer maxima exceeding 21 °C. The anomaly pattern is similar to that depicted for the whole basin, with prolonged periods of positive and negative signs. Moreover, the amplitude of the anomalies is the highest recorded in all the sub-basins. The cooling event occurring in autumn 2001 strongly influences the mean temperature of this region as a consequence of its shallow depth. In fact, even though the previous summer (2001) is characterized by temperature values higher than the climatology, autumn 2001 and early winter 2002 sub-basin-averaged temperatures are approximately 10 °C (similar to other years, excluding 2001 and 2007). Summers 2001 and 2007, both preceded by anomalous warm autumns and winters, are characterized by values significantly higher than 20 °C. The vertical structure of the water column (Fig. 7) is similar to that observed in the WACC

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



region, but vertical gradients are weaker. In this region too the positive summer bias is due to an overestimation of the vertical mixing during the stratified season and below 20 m depth the model produces temperature higher than the observation. During winter the model slightly underestimates temperature but is capable of reproducing the vertical shape of the observed profiles and the inter-annual variability.

The Middle Adriatic volume-averaged temperature (Fig. 6 panel d) ranges between 12 and 17 °C. The large heat flux anomaly occurring between summer 2000 and spring 2001 determines temperature minima during February 2001 to be almost 2 °C higher than the climatological values. The strong cooling event occurring in autumn 2001 produces an abrupt decrease in temperature but not enough to bring the successive winter values close to the climatological estimates. This sub-basin requires a longer time to recover from the anomalous condition occurring in 2000. Starting from summer 2001, there is a progressive decrease in volume-averaged temperature that persists until summer 2006. Unlike the WACC and the Northern regions, temperature vertical profiles (Fig. 7) are always characterized by a negative gradient from top to bottom (even though it becomes very small during winters as the water column is very well mixed).

The large inter-annual variability predicted by the model simulation is also confirmed by the observations. The model predicted difference between winter 2001 (the warmest) and winter 2006 (the coolest) matches the observational evidence. Both model and observation depict a well-mixed water column with a nearly constant temperature that during 2001 is of 13.5 °C and in 2006 is about 14.7 °C. During late summer 2002 (magenta line in Fig. 7) the model fails to reproduce the observed mixed layer, which extends from the surface to 20 m depth; whereas below the mixed layer modelled and observed temperature agree.

The Southern Adriatic volume is approximately 24 500 km<sup>3</sup> (77% of the entire basin). As a consequence of this large volume, the sub-basin responses to surface and lateral fluxes are slower than in the rest of the basin and the amplitude of the seasonal and inter-annual variability signal is smaller than in the other sub-basins. In particular, in contrast with the other sub-basins, the 2007 event is less pronounced. The 2000–

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2001 event increases the sub-basin volume-averaged temperature of about 0.5 °C, and this region persists in its positive anomalous condition until the beginning of 2004. The highest summer temperature is recorded in 2001. The successive strong cooling occurring during autumn is partially compensated by the advective flux through the Strait of Otranto (Fig. 3, upper panel), and thus temperature minima in winter 2002 are similar to the previous year. Only in 2003 do the combined effects of surface and lateral fluxes start bringing the volume-averaged temperature back to normal values, which are reached at the beginning of 2004.

Observed values (Fig. 7) show that most of the seasonal cycle signal is confined to the upper 50–60 m of the water column, while the model produces a weaker thermocline and tends to overestimate vertical mixing, propagating down to 100 m the seasonal warming cycle. Below 200–250 m depth model results and observations agree, having values of about  $13.5 \pm 0.1$  °C during all the periods sampled. At these depths, observed and predicted temperatures depend more on the inter-annual variability than on seasonal oscillation. Observed and modelled temperature differences, in the upper layers, in winter 2001 and 2006 are almost 1 °C.

Analysed and modelled sea surface temperatures (Fig. 8) are both characterized by a large inter-annual variability that differs from the volume-averaged temperature analysed series. In general, the model is able to reproduce the observed inter-annual variability. The agreement between the two datasets is very good, having a Nash-Sutcliffe model efficiency coefficient (Nash and Sutcliffe, 1970) of about 0.95 considering the entire signal, and about 0.8 after removing the monthly climatology. The model results are characterized by a negative bias, which is more pronounced during summer seasons. The largest anomaly is observed during summer 2003 by both modelled and observed datasets. As already noted by Grbec et al. (2007), the spring/summer 2003 period was extremely warm and the response of the sea was remarkable. It is interesting to note this anomalous warm summer has a very weak signal in the volume-averaged temperature. The anomalous warm winter 2001 and 2007 signal is also present in the SST time-series, with temperatures 1 °C higher than the corresponding climatological

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

monthly mean.

During the period under investigation, the inter-annual variability of the volume-averaged salinity (Fig. 9) is larger than the seasonal oscillations. From January 2000 until summer 2003 the values oscillate seasonally around 38.5 psu. From summer 2003 to winter 2006 a constant increase in salinity content of the basin is observed; in late winter 2006 a new plateau is reached and basin-averaged values are constantly slightly higher than 38.6 psu. Large sea surface salinity values during summer 2003 have been detected in previous studies analysing observed data collected in the Middle Adriatic (Grbec et al., 2007). This progressive salinification of the basin is strongly driven by the Po runoff inter-annual variability (Fig. 3, bottom panel). In fact, from January 2003 to spring 2008 the Po regime is always below its climatological values. The effect of the missing freshwater input from the Po is also amplified by the concomitant high evaporation rates simulated by the model during winters 2003–2004–2005 and 2006 (Fig. 4). The surface fluxes compensate the missing intrusion of MLIW at the Strait of Otranto that becomes significant only in summer 2005 (Fig. 5).

In the WACC and Northern regions, the effects of the large Po runoff event (Fig. 3 bottom panel) occurring during autumn 2000 and continuing until spring 2001 are evident, and a freshening of about 0.5 psu is observed in both the sub-basins during this period (Fig. 9, panels b and c). Within the WACC region observational and modelled data are consistent. As already discussed for temperature data, model results are affected by an excessive vertical mixing (Fig. 10) in the upper layers of the water column. However, the inter-annual variability signal detected in the observations is reproduced well by the model simulation. The winter 2001 data are those with lower salinity values at the surface, as a consequence of the large Po runoff, with values lower than 36.0 psu in both the datasets.

Observed and modelled stratification progressively decrease as a consequence of the reduced input from the Po and simultaneous large evaporation rates. These conditions, persisting for several years, strongly modify the salt content of the WACC region especially in the upper layers, reducing stratification (Fig. 10).

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the Northern region the evaporation rates (Fig. 4c) strengthen the effects of the reduced Po river regime (Fig. 3, bottom panel) and a strong inter-annual variability signal characterizes this sub-basin. Differences between minimum and maximum values modelled during the simulated period are larger than 1 psu. Agreement between observed and modelled values is weaker than in the WACC region; however, the long-period trend seems to be reproduced well by the model. The high salinity values diagnosed by the model, starting from summer 2003, are partially confirmed by observational evidence in 2006. Similarly to the WACC region, the salinification resulting from the reduced river input and large evaporation rate produces a weaker stratification (Fig. 10) with the surface layers significantly saltier during 2006 than 2001–2002 and 2003. The difference between modelled and observed values seems to be a consequence of an excessive vertical, and probably horizontal, mixing. Unlike the WACC region, the lowest surface salinity values were observed during the 2003 cruise, while the 2000 Po event signal is confined to a very thin surface layer (Fig. 10). The general positive model bias could be a consequence of the only partially resolved water budget deriving from the usage of climatological river input (except for the Po) and precipitation rate. This missing surface freshwater input could reinforce vertical stratification preventing excessive mixing.

Similarly to the volume averaged temperature, the amplitude of the inter-annual variability signal decreases increasing the volume of the analysed sub-basin. In the Middle Adriatic differences between maximum and minimum salinity values are smaller than 0.3 psu. With the sole exception of August 2006, in this region model results are characterized by a negative bias. We can thus argue that the model overestimates the exchanging processes (both advective and diffusive) between the Northern and Middle sub/basins, the former being characterized by a positive bias. However, long-term variation seems to be reproduced by the model simulations.

The low salinity values characterizing the model simulation in 2001 are in good agreement with observations from surface to about 100 m depth; in the deeper layers the model underestimates salinity content by about 0.1 psu. The largest difference

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

between modelled and observed data is probably that recorded in 2002. During this year there is a clear observational evidence of MLIW intrusion in the intermediate layers of the Middle Adriatic. In addition to the small amount of MLIW present at the Strait of Otranto (Fig. 5) during this period, the missing reproduction of this intrusion can be amplified as a consequence of the excessive mixing described above. During this period the model results are characterized by a strong negative bias with salinity values that are systematically lower than the observations. However, modelled and observed differences between the years 2002 and 2003 are similar. During 2003 MLIW intrusion is absent and a general freshening of the sub-basin is observed. Thus the missing reproduction of the 2002 intrusion event does not seem to affect the quality of the simulation in the following years.

The increase in salinity content of the sub-basin is also confirmed by the observations, with the sole exception of the August 2006 cruise. Data obtained from this cruise are characterized by a marked surface minimum that the model fails to reproduce. However, the sampling positions (Fig. 1) suggest that the major model deficiency is in the representation of the river runoff along the East coast and probably during this period the climatological values used are not representative of the real freshwater input.

Inter-annual variability in the Southern sub-basin is very small; volume-averaged salinity values oscillate around 38.6 and difference between maximum and minimum recorded values is about 0.1 psu. Similarly to the Middle Adriatic, the model results are characterized by a negative bias, which is constant below 200 m depth (Fig. 10). The differences between modelled and observed values in the upper layers seem to be related to the sampling positions (Fig. 1) and can be ascribed to the only partially resolved inter-annual variability in the surface water fluxes.

At intermediate depths, in addition to the MLIW intrusions occurring in late summer 2002 and found also in the Middle sub-basin, a subsurface salinity maximum not detected in the Middle Adriatic is observed in spring 2003 characterized by an unusual uplifted horizon that may be ascribed to MLIW intrusion. The model fails to reproduce both the events, since this peculiar subsurface salinity maximum cannot be detected in

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the corresponding model results.

In the bottom layers, the increased salinity content is also confirmed by observations. During 2006 cruises bottom salinity is 0.1 psu larger than the previous years and model results confirm this finding. Unlike the other sub-basins, the Southern Adriatic mean salinity does not show a marked increase between 2003 and 2006. Instead, the salinity content grows in late summer 2005 as a consequence of the intrusion of the MLIW (Fig. 5). This salinification, both observed and modelled, may probably have been strengthened as a consequence of the missing freshwater input in the Northern sub-basins and the consequent formation of dense waters saltier than usual filling the deep part of the Southern Adriatic sub-basin later.

## 6 Dense waters

Dense Water (hereafter DW) volume ( $\text{km}^3$ ) and formation/dissipation rate (Sv) are investigated here for each sub-basin with the sole exception of the WACC region. The WACC region is omitted from the calculation since preliminary analysis has shown that in this region dense water is only poorly present and is not formed locally. The analysis was performed on a daily basis setting the threshold sigma-t limit to 29.2 in all the sub-basins and results are shown in Fig. 11. In addition, mean characteristics (temperature and salinity) of the newly-formed water masses are presented in Fig. 12. Positive values of formation rate indicate new DW locally formed or advected into the sub-basin; negative values, on the other hand, indicate the occurrence of local processes acting in the direction of decreasing density (positive buoyancy fluxes) or advective processes removing dense water from the sub-basin under investigation. The formation rate is computed as the local tendency of the volume of dense water having sigma-t larger than the threshold on a daily basis. This definition of formation rate differs from others presented in the literature in which the amount of volume of dense water is divided by the seconds elapsed in the entire year; we will refer to the values computed using this method as “seasonal”.

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DWs are formed in the Northern sub-basin (Northern Adriatic dense water, NAdw) in all years with the sole exception of 2000–2001; however, daily values of formation rate (Fig. 11) and mean properties (Fig. 12) differ from year to year as a consequence of the surface and river buoyancy fluxes discussed. Seasonal formation rates (not shown) are in good agreement with previous numerical findings. This “seasonal” formation rate is on average about 0.04/0.05 Sv, as already found by Mantziafou and Lascaratos (2008). At seasonal scales, inter-annual variability of the formation rate is small. In other words, when NAdw are formed, this formation process has similar velocities in all the years investigated. However, the duration of the formation processes differs from year to year as well as the advective processes and the positive buoyancy fluxes, and thus the final volume of NAdw formed is different in different years. Maximum formation rate occurs between the end of December and the beginning of January with values varying from year to year.

The occurrence and the intensity of the NAdw formation process are directly related to the amplitude and timing of the negative surface heat flux modulated by the Po regime. The anomalous large Po river runoff together with the missing cooling during both autumn 2000 and winter 2001 have prevented DW formation in winter 2000–2001. The two strong cooling events occurring in autumn 2001 and 2007 (Fig. 2, panel c) produce large formation rates, both higher than 2 Sv. However, even though cooling was stronger in the 2001–2002 formation event, the high Po runoff input compensated the overall buoyancy flux mitigating the maximum NAdw formation velocity, which is, during this period, significantly lower than in 2007–2008. Thus temperature and salinity characterizing these newly formed NAdw strongly differ between the two events. The NAdw formed in 2000–2001 event are characterized by very low salinity values, ranging between 37.6 psu at the beginning of the event to 38.3 psu at the end of the event. Those values are significantly lower than corresponding values reported in literature (Zore-Armanda, 1963; Artegiani et al., 1997b; Vilibic and Orlic 2002), which range between 38.3 and 38.5 psu (Vilibic and Orlic, 2002). In 2007–2008 the Northern sub-basin is strongly saltier than the previous period and the NAdw formed also has a

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



salinity content higher than 38.6. As a consequence waters with sigma-t larger than 29.2 start to be produced even when associated temperatures are higher than 12 °C. Moreover, the 2001–2002 NAdw formation event is followed by a period characterized by negative values of NAdw formation rate as a consequence of a strongly positive surface buoyancy flux.

Despite the fact that autumn 2006 and the successive winter are characterized by a weak cooling at the sea surface similar to the 2000–2001 anomaly, NAdw formation process occurs during this period. This is clearly related to the high salinity values that characterize the Northern sub-basin in this year. In fact, the NAdw temperature is about 12 °C, significantly higher than the 10.6 °C indicated by Vilibic and Orlic (2002), or 11.3 °C as suggested by Artegiani et al. (1997a), and associated salinity is higher than 38.5 psu. All the other NAdw formation events are characterized by similar behaviour, with salinity slightly increasing from year to year prolonging the NAdw formation period and consequently increasing the associated maximum temperature values (Fig. 12). It is interesting to note that a reversal of this tendency is diagnosed by the model simulation at the end of 2008, associated with the Po regime, which is back to climatological values.

Inter-annual variability of DW formation in the Middle Adriatic (Middle Adriatic Dense Water, MAdw) is very large. In addition to the absence of formation in 2001, 2002–2003 and 2007 are also anomalous, and “seasonal” formation rates are smaller than other simulated years. The “seasonal” formation rate varies from 0.02 Sv in the above years to 0.1 Sv in 2000–2004–2005–2006 and 2008. The values and the inter-annual variability computed at seasonal scales are in agreement with previous findings in Mantziafou and Lascaratos (2004, 2008).

In addition to the surface flux activity, the intrusion of the MLIW in the Middle Adriatic is a key factor in determining the occurrence of MAdw formation events (Bergamasco et al., 1999). The three consecutive (2001–2002–2003) years of missing MAdw formation are due to the concomitant effect of surface buoyancy fluxes and absence of MLIW in the sub-basin. The total volume of MAdw during this period is significantly small, and

OSD

8, 565–611, 2011

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the temperature and salinity associated with these waters (Fig. 12) suggest that the DW is advected into the sub-basin from the North. In fact, MADw temperature (Fig. 12) is significantly lower than 12 °C and the corresponding salinity (Fig. 12) is below 38.5 psu. It is important to mention that the data collected in late summer 2002 indicate the presence of MLIW and highlight a model deficiency in reproducing this event. The model therefore probably underestimates formation processes in this period.

In agreement with previous findings (Zore-Armanda, 1963; Mantziafou and Lascaratos, 2004, 2008) MADw formation occurs later in the Middle Adriatic than the same processes do in the Northern sub-basin. Large temperature and reduced input of MLIW during winter 2007 inhibit MADw formation. MADw volume diagnosed by the model in 2004–2005–2006 and 2008 clearly indicates that DW formation occurred locally. The corresponding temperature and salinity (Fig. 12) are also in good agreement with previous estimates (Zore-Armanda, 1963; Artegiani et al., 1997a; Vilibic and Orlic 2002). However, similarly to the Northern region, modelled and observed increase in salt content of the basin allows dense waters to be formed with progressively higher temperature. In fact, temperature and salinity characterizing dense water formed during 2000 are the closest to climatological and observed values (Artegiani et al., 1997a).

The dimensions of the Southern sub-basin affect the memory of the systems, and as a consequence the missing cooling occurring in 2000 and the absence of MLIW strongly impact DW formation rates until 2005 (Fig. 11). The small formation rates recorded between 2001 and 2004 are clearly due to DW advected from the northern sub-basins. This is particularly evident analysing the mean properties of DW. As a matter of fact, during this period DW found in the Southern sub-basin are cool and fresh (Fig. 12).

In winter 2004 the temperature of the basin reverts to standard or climatological values; however, salty waters of Levantine origin are still absent in the basin and the model starts the production of DW again only in winter 2005. This is confirmed both by the amount of DW present in the sub-basin and the associated temperature and salinity. During the periods of local DW formation (2005, 2006 and 2008), DW characteristics

are similar to previous finding with temperatures of about 13°C and salinity slightly higher than 38.65 psu (Zore-Armanda, 1963; Artegiani et al., 1997a; Vilibic and Orlic 2002). Similarly to what happens in the Middle sub-basin, the local formation of DW stops in 2007 as a consequence of the surface buoyancy fluxes and of the reduced inflow of MLIW.

## 7 Summary and conclusions

The inter-annual variability of the Adriatic Sea hydrographic characteristics was investigated for the period 2000–2008 using both numerical simulation results and direct observations. The model used is derived from the Adriatic Forecasting System (<http://gnoo.bo.ingv.it/afs/>) and was also developed within the ECOOP project. The primitive equations model was implemented on a regular horizontal grid having approximately 2.2 km of resolution and 27 vertical sigma layers. In order to reproduce the inter-annual variability the model was forced at the air-sea interface using interactively computed fluxes (both momentum and buoyancy) by means of the model predicted sea surface temperature, also making use of the atmospheric variables needed from the operational products of the ECMWF, with a 6 h frequency and 0.5° horizontal resolution. Observational data used in support of numerical findings include CTD casts and satellite-derived sea surface temperature. As per a well-known Adriatic characteristics, the basin was divided in four sub-basins (WACC, Northern, Middle and Southern Adriatic) and the data were analysed separately.

Total heat flux during the period under investigation is characterized by a marked inter-annual variability (Fig. 2), which is larger during the cooling periods and mostly due to the variability in the latent heat flux component (Fig. 4).

A remarkable positive anomaly was detected, in all the sub-basins, during autumn–winter 2000–2001. During the same period the Po river runoff was significantly larger than the climatological values (Fig. 3, bottom panel) and the MLIW absent at the Strait of Otranto (Fig. 5). The basin (and sub-basin) reaction was noticeable in terms of

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



temperature (both volume-averaged and surface, Figs. 6 and 8 respectively), salinity (Fig. 9) and consequent DW production (Fig. 11). Temperature minima, during early winter 2001, are significantly higher than those occurring in previous and following years. The whole basin volume-averaged temperature rise between early winter 2000 and early winter 2001 is almost  $0.8^{\circ}\text{C}$ ; the amplitude of the variation is larger in the Northern and Middle Adriatic sub-basins, reaching almost  $2^{\circ}\text{C}$ , and smaller in the Southern basin ( $0.5^{\circ}\text{C}$ ). During the same period the SST anomaly is almost  $1^{\circ}\text{C}$  (Fig. 8). The response of the salinity field to the anomalous Po river runoff and to the missing evaporation is particularly evident in the WACC and Northern basins but, as a consequence of the induced vertical stability, it is mostly confined in a very thin surface layer of the water column (Figs. 9 and 10). Consequently, no DW formation occurred in the entire Adriatic Sea. The river Po continues to supply a large amount of fresh waters to the basin until the end of 2002 (Fig. 3 bottom panel). The successive autumn of 2001 was very cold – the coldest of the entire period under investigation – and was characterized by a strongly negative heat flux (Fig. 2) as a response to the large evaporation rate (Fig. 4). Temperature in the WACC and Northern sub-basin quickly reverted to climatological values in early autumn and was below the climatological values by the end of the year. The relatively large volumes of the Middle and Southern sub-basins mitigate the cooling effect and the corresponding averaged temperature response, which, in the Southern basin, persists above the climatological values (Fig. 6). Volume-averaged salinity in the Northern and Middle Adriatic are now very low as a consequence of the prolonged period characterized by high Po runoff rates. The low temperature reached in the Northern basin allows DW formation process to restart in early 2002 (Fig. 11); the newly formed DW are characterized by very cold and fresh properties (Fig. 12).

The year 2002 was relatively mild, with a winter-autumn total heat flux above the climatological values, and spring-summer ones slightly below (Figs. 2 and 3). This caused the volume temperature to be warmer than climatology again during the entire year and over the entire basin (Fig. 6). The signal is very weak in the sea surface

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

temperature (Fig. 8). The volume-averaged salinity is now stable in all the sub-basins with the sole exception of the northernmost region, which is now slightly saltier than the previous year (Fig. 9). This is due to the fact that the Po runoff, even though above the climatological mean, is significantly reduced compared to autumn 2000 (Fig. 3). During summer there is observational evidence of MLIW intrusion (Fig. 10) in the Middle and Southern sub-basins not reproduced by the model, and thus the salinity content is probably underestimated by the model simulation. The year 2003 starts with a very cold winter characterized by high evaporation rates (Fig. 4); however, the Po river now enters a long prolonged period of very low runoff (Fig. 3), and consequently evaporation effects are remarkable in the salinity content of the basin. The cold winter restores the volume-averaged temperature to climatological values in all the basins with the sole exception of the Southern region, which has longer response times as a consequence of its larger volume. However, as a consequence of the mild 2002 autumn, DW formation occurs only in the Northern sub-basin (Fig. 11). Despite the cold winter, the SST during the summer is the highest recorded in the entire period investigated (Fig. 8). High evaporation and almost absent Po runoff produce a net salinity growth in all the sub-basins except for the Southern where MLIW contribution is still missing (Fig. 5). The concomitant effect of the strong summer stratification amplifies the increased salinity signal at the sea surface (Fig. 10).

The years 2004 and 2005 are similar to each other. Both are characterized by negative total heat flux anomalies all year round, with the exception of a few months during the late-summer/early-autumn periods (Fig. 2), reduced Po runoff and high evaporation rates. Temperature is generally below the climatological values except in the shallower regions during the warmer period mentioned above (Fig. 6). At the beginning of 2004 the Southern basin temperature is also restored to climatological values and encounters a relatively long period with values constantly below the climatology (Fig. 6). During this two consecutive years the volume average salinity continuously grows (Fig. 9). Southern growth only starts in summer 2005 with the arrival of the MLIW at the Strait of Otranto (Fig. 5). DW formation process is now also active in the Middle sub-basin

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and a small production also starts in the Southern Adriatic in winter 2005 (Fig. 11). The increased salinity content allows NAdw to reach the 29.2 threshold even with warmer temperatures (Fig. 12).

The cold period persists until August 2006. Temperatures are now below, and salinity above, the climatological values over the entire Adriatic Sea. The characteristic freshwater input from the Po river is still absent and a considerable signature of MLIW is observed at the Strait of Otranto. DW formation is very pronounced in all the basins and also in the South in winter 2006 (Fig. 11); DW volumes formed in this period are the largest of the entire simulation. Similarly to 2000, 2006 ends with a significantly warm autumn, which induces temperature to be higher than climatology in all the basins except for the Southern region (Fig. 6). The warm period started in autumn 2006 continues in winter 2007 (Fig. 2), which is characterized by very mild atmospheric conditions (similar to the period 2000–2001). This, in addition to the reduced amount of MLIW intruding into the basin (Fig. 6), interrupts DW production in the Middle and Southern sub-basins. A weak NAdw formation is observed as a consequence of the reduced Po runoff and of the high salinity content characterizing the sub-basin during these years (Figs. 11 and 12). Volume and surface temperatures remain significantly higher than the climatology until spring/early-summer (Figs. 6 and 8). Similarly to 2000–2001, the following autumn is particularly cold as a consequence of the high evaporation rates and an abrupt drop in temperatures is observed (Figs. 6 and 8). As a consequence, even though under the influence of a relatively mild winter, DW are formed in all the sub-basins at the beginning of 2008 (Fig. 11). In Spring 2008 the Po river runoff reverts to considerable regimes (Fig. 3) and salinity tends to decrease in the WACC, in the Northern, and in the Middle, sub-basins (Fig. 9). The starting of a DW formation process is observed in the Northern basin at the end of 2008 with salinity characterizing the newly-formed water masses that are significantly lower than the preceding years (Fig. 12).

The period investigated shows significant inter-annual variability in all the considered quantities and DW formation processes. However, a long-scale signal has been

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

observed in the basin as a consequence of the reduced river runoff and high evaporation rate. The basin significantly increased the salinity content and the temperature and salinity characterizing DW, especially in the Northern sub-basin, are modified consequently.

In terms of inter-annual variability, the proposed modelling framework enables qualitative studies of the main characteristics of the basin. The nesting procedure, with the operational MFS model, allows for a correct reproduction of the inter-annual variability related to the large-scale flow and the remote influences of the Eastern Mediterranean dynamics.

Even though characterized by a relatively coarse resolution, the atmospheric forcing seems to be adequate for the inter-annual variability investigations.

The weaker modelling components are related to the freshwater balance. The absence of an observational dataset capable of capturing the runoff inter-annual variability of the major Adriatic rivers (with the exception of the Po) seems to affect the quality of the simulation, especially along the east coast. However, the present model configuration allows for an accurate reproduction of the Northern Adriatic sub-basin dynamics, while the inaccuracy of the climatological river data affects mostly the east coasts of the Middle and Southern Adriatic sub-basins.

There is a need for further modelling efforts in order to fill the runoff observational data gap.

The study confirmed that the Adriatic variability depends upon many factors: river input, atmospheric forcing and mass exchanges through the Strait of Otranto.

The Po river influence seems to be confined in the Northern and WACC sub-regions, while the Middle and Southern sub-basins seem to be marginally affected by Po river variability. The Middle and Southern sub-basin salinity is strongly dependent on the river regime along the east coast and on the intrusion of the MLIW.

In the Northern Adriatic sub-basin, the freshwater input from the Po determines the characteristics of the locally formed DW, while the occurrence of the DW production processes is totally driven by the surface heat fluxes. In the Middle and Southern

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sub-basins, however, the key factor in determining the DW production processes is the presence of sufficiently salty water of Mediterranean origin. In fact, even under strong atmospheric conditions the DW formation processes do not start if MLIW are missing.

*Acknowledgements.* The authors wish to thank M. Vichi for constructive discussion. The SST data were obtained from the GOS (Gruppo di Oceanografia Sperimentale) of the National Research Council. The CTD data used have been kindly provided by NURC (NATO Undersea Research Centre). This work was partially funded by ECOOP (European Coastal sea Operational observing and Forecasting system). ECOOP is funded by the European Commission's Sixth Framework Programme under the priority Sustainable Development, Global Changes and Ecosystem Contract No. 36355. We also thank S. Gualdi for his continuing support and suggestions.

## References

- Artegiani, A. and Salusti, E.: Field observation of the flow of dense water on the bottom of the Adriatic Sea during the winter of 1981, *Oceanol. Acta.*, 10, 387–391, 1987.
- Artegiani, A., Azzolini, R., and Salusti, E.: On the dense water in the Adriatic Sea. *Oceanol. Acta.*, 12, 151–160, 1989.
- Artegiani, A., Bregant, D., Paschini, E., Pinardi, N., Raicich, F., and Russo, A.: The Adriatic Sea general circulation, part I: Air-sea interactions and water mass structure, *J. Phys. Oceanogr.*, 27, 1492–1514, 1997a.
- Artegiani, A., Bregant, D., Paschini, E., Pinardi, N., Raicich, F., and Russo, A.: The Adriatic Sea general circulation. part II: Baroclinic circulation structure, *J. Phys. Oceanogr.*, 27, 1515–1532, 1997b.
- Artegiani, A., Pariente, R., Paschini, E., Russo, A., and Totti, C.: Seasonal Evolution of the physical water properties observed from Senigallia transect, in: *The Adriatic Sea, Ecosystem Report*, edited by: Hopkins, T. S., 32, European Commission, Brussels, 59–70, 1999.
- Bergamasco, A., Oguz, T., and Malanotte-Rizzoli, P.: Modeling dense water mass formation and winter circulation in the northern and central Adriatic Sea, *J. Mar. Syst.*, 20, 279–300, 1999.
- Bignami, F., Salusti, E., and Schiarini, S.: Observation on a bottom vein of dense water in the Southern Adriatic and Ionian Seas, *J. Geophys. Res.*, 95(C5), 7249–7259, 1990.

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

## Abstract

## Introduction

## Conclusions

## References

## Tables

## Figures



[Back](#)

Close

Full Screen / Esc

[Printer-friendly Version](#)

## Interactive Discussion



- Blumberg, A. F. and Mellor, G. L.: A description of a three-dimensional coastal ocean circulation model, in: Three-dimensional coastal ocean models, edited by: Heaps, N. S., Washington, DC, American Geophysical Union, 208 pp., 1987.
- Cardin, V. and Gacic, M.: Long term heat flux variability and winter convection in the Adriatic Sea, *J. Geophys. Res.*, 108(C9), n. 8103, 2003.
- Castellari, S., Pinardi, N., and Leaman, K.: A model study of air-sea interactions in the Mediterranean Sea, *J. Mar. Syst.*, 18, 89–114, 1998.
- Chiggiato, J., Zavatarelli, M., Castellari, S., and Deserti, M.: Interannual variability of surface heat fluxes in the Adriatic Sea in the period 1998–2001 and comparison with observations, *Sci. Total Environ.*, 353, 89–102, 2005.
- Gacic, M., Kovacevic, V., Manca, B., Papageorgiou, E., Poulain, P.M., Scarazzato, P., and Vetrano, A.: Thermohaline properties and circulation in the Otranto Strait, *Bulletin Institut Oceanographie Monaco*, No. Spec. 17, 117–145, 1996.
- Gacic, M., Lascaratos, A., Manca, B., and Mantziafou, A.: Adriatic deep water and interaction with the Eastern Mediterranean Sea, in: Physical Oceanography of the Adriatic Sea, in: Past, Present and Future, Cushman-Roisin, B., Gacic, M., Poulain, P.-M., Artegiani, A., Kluwer Academic Publishers, 111–142 (Chapter 4), 2002.
- Gacic, M., Civitarese, G., Miserocchi, S., Cardin, V., Crise, A., and Mauri, E.: The open-ocean convection in the Southern Adriatic: a controlling mechanism of the spring phytoplankton bloom, *Cont. Shelf. Res.*, 22, 1897–1908, 2002.
- Grbec, B., Vilibic, I., Bajic, A., Morovic, M., Bec Paklar, G., Matici, F., and Dadic, V.: Response of the Adriatic Sea to the atmospheric anomaly in 2003, *Ann. Geophys.*, 25, 835–846, 2007, <http://www.ann-geophys.net/25/835/2007/>.
- Guarnieri, A., Oddo, P., Pastore, M., and Pinardi, N.: The Adriatic Basin Forecasting System: new model and System development. Coastal to Global Operational Oceanography: Achievements and Challenges, edited by: Dahlin, H., Fleming, N. C., and Petersson, S. E., Proceeding of 5th EuroGOOS Conference, Exeter, 20–22, 2008.
- Hellerman, S. and Rosenstein, M.: Normal monthly wind stress over the world ocean with error estimates, *J. Phys. Oceanogr.*, 13, 1093–104, 1983.
- Hendershott, M. C. and Rizzoli, P.: The winter circulation of the Adriatic Sea, *Deep-Sea Res.*, 23, 353–370, 1976.
- Hopkins, T. S., Artegiani, A., Kinder, C., and Pariente, R.: A discussion of the northern Adriatic circulation and flushing as determined from ELNA hydrography, in: The Adriatic Sea.

# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

## Abstract

## Introduction

## Conclusions

## References

## Tables

## Figures



[Back](#)

Close

Full Screen / Esc

Printer-friendly Version

## Interactive Discussion



- Ecosystem Report, edited by: Hopkins, T. S. et al., 32, European Commission, Brussels, 85–106, 1999.
- Klein, B., Roether, W., Civitarese, G., Gacic, M., Manca, B.B., and Ribera d'Alcala, M.: Is the Adriatic returning to dominate the production of Eastern Mediterranean Deep Water?, *Geophys. Res. Lett.*, 27, 3377–3380, 2000.
- Kondo, J.: Air-sea bulk transfer coefficients in diabatic conditions, *Bound.-Lay. Meteorol.*, 9, 91–112, 1975.
- Legates, D. R. and Wilmott, C. J.: Mean seasonal and spatial variability in a gauge corrected global precipitation, *Int. J. Climatol.*, 10, 121–7, 1990.
- Maggiore, A., Zavatarelli, M., Angelucci, M. G., and Pinardi, N.: Surface heat and water fluxes in the Adriatic Sea: seasonal and interannual variability, *Phys. Chem. Earth*, 23, 561–7, 1998.
- Malanotte-Rizzoli, P. and Bergamasco, A.: The dynamics of the coastal region of the northern Adriatic Sea, *J. Phys. Oceanogr.*, 13, 1105–1130, 1983.
- Malanotte-Rizzoli, P., Manca, B., Ribera d'Alcala, M., Theocharis, A., Brenner, S., Budillon, G., and Ozsoy, E.: The eastern Mediterranean in the 80s and the 90s: the big transition in the intermediate and deep circulations, *Dynam. Atmos. Oceans*, 29, 365–395, 1999.
- Manca, B.B., Kovacevic, V., Gacic, M., and Viezzoli, D.: Dense water formation in the Southern Adriatic Sea and spreading into the Ionian Sea in the period 1997–1999, *J. Marine Syst.* 33–34, 133–154, 2002.
- Mantziadou, A. and Lascaratos, A.: An eddy resolving numerical study of the general circulation and deep-water formation in the Adriatic Sea, *Deep-Sea Res.* I 51, 921–952, 2004.
- Mantziadou, A. and Lascaratos, A.: Deep-water formation in the Adriatic Sea: Interannual simulation for years 1979–1999, *Deep-Sea Res. Pt. I*, 55, 11, 1403–1426, 2008.
- Marullo, S., Buongiorno Nardelli, B., Guarracino, M., and Santoleri, R.: Observing the Mediterranean Sea from space: 21 years of Pathfinder-AVHRR sea surface temperatures (1985 to 2005): re-analysis and validation, *Ocean Sci.*, 3, 299–310, doi:10.5194/os-3-299-2007, 2007.
- May, P.W.: A brief explanation of Mediterranean heat and momentum flux calculations NORDA Code 322, Stennis Space Center, Miss. 7 Nav Oceanogr Atmos Res Lab, 5 pp., 1986.
- Mellor, G. L. and Blumberg, A. F.: Modeling vertical and horizontal diffusivities with the sigma coordinate ocean models, *Mon. Weather Rev.*, 113, 1379–1383, 1985.
- Mellor, G. L. and Yamada, T.: Development of a turbulence closure submodel for geophysical

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

fluid problems, *Rev. Geophys. Space Phys.*, 20, 851–75, 1982.

Mellor, G. L., Oey, L. Y., and Ezer, T.: Sigma Coordinate Pressure Gradient Errors and the Seamount Problem, *J. Atmos. Oceanic Technol.*, 15, 1122–1131, 1998.

Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I – A discussion of principles, *J. Hydrol.*, 10(3), 282–290, 1970.

Nardelli, B. B. and Salusti, E.: On dense water formation criteria and their application to the Mediterranean Sea, *Deep-Sea Res. Pt. I*, 47(2), 193–221, 2000.

Oddo, P. and Pinardi, N.: Lateral Open Boundary Conditions for Nested Limited Area Models: Precess selective approach, *Ocean Modeling*, 20(2), 134–156, 2008.

Oddo, P., Pinardi, N., and Zavatarelli, M.: A numerical study of the interannual variability of the Adriatic Sea (1999–2002), *Sci. Total Environ.*, 353, 39–56, 2005.

Oddo, P., Pinardi, N., Zavatarelli, M., and Colucelli, A.: The Adriatic Basin Forecasting System, *Acta Adriatica*, 47 (suppl), 169–184, 2006.

Ovchinnikov, I. M., Zats, V. I., Krivosheya, V. G., and Udodov, A. I.: Formation of deep eastern Mediterranean waters in the Adriatic Sea, *Oceanology*, 25, 704–707, 1985.

Poulain, P. M.: Adriatic Sea surface circulation as derived from drifter data between 1990 and 1999, *J. Marine Syst.*, 29, 3–32, 2001.

Provini, A., Crosa, G., and Marchetti, R.: Nutrient export from the Po and Adige river basin over the last 20 years, *Sci. Total Environ.*, 291–313(suppl.), 1992.

Raichich, F.: On the fresh water balance of the Adriatic Sea, *J. Marine Syst.*, 9, 305–319, 1996.

Raichich, F.: Note on flow rates of the Adriatic rivers, Istituto Talassografico Sperimentale Trieste, Technical Report, RF02:94, 8, 1994.

Reed, R. K.: An evaluation of formulas for estimating clear sky insulation over the ocean, NOAA-ERL 352-PMEL Technical Report, 26, 1–25, 1975.

Reed, R. K.: On estimating insolation over the ocean, *J. Phys. Oceanogr.*, 1, 482–485, 1977.

Roether, W. and Schlitzer, R.: Eastern Mediterranean Deep Water renewal on the basis of chlorofluoromethane and tritium data, *Dyn. Atmos. Oceans*, 15, 333–354, 1991.

Roether, W., Manca, B. B., Klein, B., Bregant, D., Georgopoulos, D., Beitzel, V., Kovacevic, V., and Luchetta, A.: Recent changes in eastern Mediterranean deep waters, *Science*, 271, 333–335, 1996.

Russo, A. and Artegiani, A.: Adriatic Sea hydrography, *Sci. Mar.*, 60, 33–43, 1996.

Sellschopp, J. and Alvarez, A.: Dense low-salinity outflow from the Adriatic Sea under mild (2001) and strong (1999) winter conditions, *J. Geophys. Res.*, 108(C9), 8104,

doi:10.1029/2002JC001562, 2003.

Smagorinsky, J.: Some historical remarks on the use of nonlinear viscosities, in: Large eddy simulations of complex engineering and geophysical flows, edited by: Galperin, B., Orszag, S. A., New York, USA, Cambridge Univ. Press, 3–36, 1993.

5 Steinfeldt, R.: Age and spectra of Eastern Mediterranean Deep Water, *J. Mar. Syst.*, 48, 67–81, 2004.

Supic, N. and Vilibic, I.: Dense water characteristic in the northern Adriatic in the 1967–2000 interval with respect to surface fluxes and Po river discharge rates, *Estuar. Coast Shelf. S.*, 66, 3–4, 580–593, 2006.

10 Tonani, M., Pinardi, N., Dobricic, S., Pujol, I., and Fratianni, C.: A high-resolution free-surface model of the Mediterranean Sea, *Ocean Sci.*, 4, 1–14, doi:10.5194/os-4-1-2008, 2008.

Van Leer, B.: Towards the Ultimate Conservative Difference Scheme, V., A Second Order Sequel to Godunov's Method, *J. Comput. Phys.*, 32, 101–136, 1979.

Vested, H. J., Perg, P., and Uhrenholdt, T.: Dense water formation in the Northern Adriatic, *JMS*, 18, 1–3, 135–160, 1998.

Vilibic, I. and Orlic, M.: Adriatic water masses, their rates of formation and transport through the Otranto Strait, *Deep-Sea Res. I*, 49, 1321–1340, 2002.

Vilibic, I. and Supic, N.: Dense water generation on a shelf: the case of the Adriatic Sea, *Ocean Dyn.*, 55(5–6), 403–415, 2005.

20 Zavatarelli, M. and Pinardi, N.: The Adriatic Sea modelling system: a nested approach, *Ann. Geophys.*, 21, 345–364, 2003, <http://www.ann-geophys.net/21/345/2003/>.

Zavatarelli, M., Pinardi, N., Kourafalou, V. H., and Maggiore, A.: Diagnostic and prognostic model studies of the Adriatic Sea circulation Seasonal variability, *J. Geophys. Res.*, 107(C1) (art 3004), 2002.

25 Zore-Armanda, M.: Les masses d'eau de la mer Adriatique, *Acta Adriat.*, 10, 5–88, 1963.

**OSD**

8, 565–611, 2011

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri


**Table 1.** Number and time intervals of CTD sampling for each year.

	2000	2001	2002	2003	2004	2005	2006	2007	2008
WACC	/	28 Jan–10 Feb (52)	16 Sep–6 Oct (111)	28 Apr–2 May (37)	/	/	15–27 Jan (59) 3–21 Mar (81) 14–27 Aug (75)	/	/
North	/	28 Jan–10 Feb (162)	16 Sep–9 Oct (489)	27 Apr–4 May (159)	/	/	14–20 Jan (44) 14–18 Aug (45)	/	/
Middle	/	4 Feb–7 Feb (24)	20 Sep–10 Oct (25)	27 Apr (17)	/	/	19–22 Jan 5–22/Mar (44) 17–29 Aug (39)		
South	/	5 Feb–6 Feb (13)	19 Sep–11 Oct (24)	26–27 Apr (12)	/	/	25–27 Jan (22) 2–22 Mar (33) 16–26 Aug (19)	/	/

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarneri



**Fig. 1.** Adriatic coastline, bathymetry and sub-region boundaries are shown (WR = WACC region; NR = Northern Adriatic Region; MR = Middle Adriatic Region, SR = Southern Adriatic Region). Locations of CTD sampling are also shown. Green dots indicate CTD positions for the 2001 cruise; magenta dots the 2002 cruise; blue dots 2003; red, black and cyan dots indicate first, second and third 2006 cruises respectively. Exact dates are listed in Table 1. T-S diagrams from CTDs data are also shown for the different regions (NR, WR, MR and SR from left to right, respectively).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarneri




**Fig. 2.** Time series of total Heat Flux, units are  $\text{W m}^{-2}$ . Colour indicates monthly anomaly values. A: whole Adriatic Sea; B: WACC region; C: Northern Adriatic Region; D: Middle Adriatic Region; E: Southern Adriatic Region.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri




**Fig. 3.** Top Panel: time series of advective heat flux through Strait of Otranto, units are  $\text{W m}^{-2}$  and Colour indicates monthly anomaly values. Bottom panel: time series of daily Po river runoff, units are  $\text{m}^3 \text{s}^{-1}$ , colour indicates monthly anomaly values.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarneri



**Fig. 4.** Time series of Evaporation rate, units are mm/day. Colour indicates monthly anomaly values. A: whole Adriatic Sea; B: WACC region; C: Northern Adriatic Region; D: middle Adriatic Region; E: Southern Adriatic Region.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 5.** Hovmoeller diagram of mean salinity profile at Strait of Otranto. Y-axis indicates depth in metres; X-axis indicates time and colour indicates salinity in psu. The solid black line indicates water having salinity content larger than 38.68 psu.

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarneri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 6.** Time series of volume-averaged temperature, units are °C. Colour indicates monthly anomaly values. A: whole Adriatic Sea; B: WACC region; C: Northern Adriatic Region; D: Middle Adriatic Region; E: Southern Adriatic Region. In panels B-C-D and E dots indicate the vertically- and spatially-averaged values of observations (black dots) and model results (white dots). Dates of the sampling are listed in Table 1; positions of the sampling are displayed in Fig. 1. The model results were sampled in the same position and at the same time as the observation before averaging.

## Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion


# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri




**Fig. 7.** Temperature Vertical profiles obtained averaging both observations (solid lines) and corresponding model results (dashed lines) spatially and temporally. Colours indicate different cruises listed in Table 1 following Fig. 1.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




**Fig. 8.** Top panel: time series of satellite (red line) and modelled (blue line) Sea Surface Temperature. Bottom panel: time series of satellite (red line) and modelled (blue line) Sea Surface Temperature anomalies.



**Fig. 9.** Time series of volume averaged salinity, units are  $\text{PSU}$ . Colour indicates monthly anomaly values. A: whole Adriatic Sea; B: WACC region; C: northern Adriatic Region; D: middle Adriatic Region; E: southern Adriatic Region. In panels B-C-D and E dots indicate the vertically- and spatially-averaged values of observations (black dots) and model results (white dots). Dates of the sampling are listed in Table 1; positions of the sampling are displayed in Fig. 1. The model results were sampled in the same position and at the same time as the observation before averaging.

# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarnieri




**Fig. 10.** Salinity Vertical profiles obtained averaging both observations (solid lines) and corresponding model results (dashed lines) spatially and temporally. Colours indicate different cruises listed in Table 1 following Fig. 1.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarneri




**Fig. 11.** Time-series of total volume of water ( $\text{km}^3$ ) with  $\sigma_{\text{t}29.2}$  (blue line and left Y-axis) in Northern, Middle and Southern sub-basins, together with the relative formation/dissipation rate expressed in Sv (black line and right Y-axis).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

# Study of the hydrographic conditions in the Adriatic Sea

P. Oddo and A. Guarneri



**Fig. 12.** Time-series of mean Temperature (blue lines, left Y-axis) and Salinity (black lines, right Y-axis) of waters masses having  $\sigma_t > 29.2$  in the Northern, Middle and Southern sub-basins.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)