

This discussion paper is/has been under review for the journal Ocean Science (OS).  
Please refer to the corresponding final paper in OS if available.

# On the influence of wind and waves on the underwater light field

M. Hieronymi<sup>1,2,3</sup> and A. Macke<sup>1</sup>

<sup>1</sup>Leibniz Institute for Tropospheric Research (IFT), Leipzig, Germany

<sup>2</sup>Helmholtz Centre for Ocean Research (GEOMAR), Kiel, Germany

<sup>3</sup>Institute of Coastal Research, Helmholtz Centre Geesthacht (HZG), Germany

Received: 1 March 2012 – Accepted: 7 March 2012 – Published: 20 March 2012

Correspondence to: M. Hieronymi (mhieronymi@hzg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

OSD

9, 1231–1272, 2012

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

We present a detailed study of the influence of various wind and wave conditions on the distribution of downwelling irradiance within the upper ocean mixed layer down to 100 m water depth. The work is based on a two-dimensional Monte Carlo radiative transfer model with high spatial resolution. We treat conditions that are favorable for the development of extreme light fluctuations, e.g. light at 490 nm and very clear oceanic water. Local wind determines the steepness of capillary-gravity waves which in turn dominate the irradiance variability near the surface. Maximum irradiance peaks that exceed the mean irradiance by a factor of more than 10 can be observed at low wind speeds up to  $5 \text{ m s}^{-1}$ . Sea states influence the light field much deeper; gravity waves can cause considerable irradiance variability even at 100 m depth. The simulation results show that under realistic conditions 50% radiative enhancements compared to the mean can still occur at 30 m depth.

## 1 Introduction

The solar radiative transfer into the ocean is strongly influenced by air-sea interactions. Wind generates waves at the free water surface that can range in size from small ripples, so called capillary waves, to huge waves over 25 m height (e.g. Sterl and Caires, 2005). The wind strength, the areal and temporal wind impact, and the water depth influence the formation of a wind sea. Sunlight is refracted at the irregularly shaped water surface. Individual waves that are superposed at the surface can act as lenses that focus the light at various water depths. For this reason, the underwater light field, in this work characterized by the downwelling irradiance, is subject to considerable variance that characteristically depends on the surface waves and thus on the prevailing wind condition and the sea state.

Besides the mentioned wave conditions, underwater light fluctuations also depend on the spectral range of sunlight, the sun altitude, diffuse sky radiation, clouds, and

OSD

9, 1231–1272, 2012

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on the inherent optical properties (IOPs) of the water (e.g. Stramski, 1986; Walker, 1994; Stramska and Dickey, 1998; Gernez and Antoine, 2009; Gege and Pinnel, 2011). Compared to other natural light regimes (e.g. in a forest), the variability of irradiance in water can be very large, e.g. near the surface irradiance peaks can exceed the time-averaged irradiance by a factor of 15 (Darecki et al., 2011).

The generation mechanisms of irradiance fluctuations and their decreasing intensity with increasing water depth are known (e.g. Schenck, 1954; Snyder and Dera, 1970; Nikolayev and Khulapov, 1976; Stramski and Dera, 1988; Gernez et al., 2011). Whereas, less is known about the wind-dependency of light fluctuations and very little is known about its sea state-dependency. There are several papers on the relationship between wind conditions and underwater light field fluctuations (e.g. Nikolayev et al., 1972; Dera and Stramski, 1986; Gernez and Antoine, 2009; Weber, 2010; Hieronimi and Macke, 2010; Darecki et al., 2011). The general view is that the most favorable conditions for light focusing by waves prevail at light to moderate winds between 2 and  $7 \text{ m s}^{-1}$ . With this work we will show that other wind and wave conditions also can lead to extreme irradiance fluctuations, in particular if we look at different water depths. We provide the first detailed analysis on the influence of local wind on the light variability in the near-surface layer and in addition on the impact of fully developed sea states on the light regime within the upper ocean mixed layer down to 100 m depth.

The work is based on a two-dimensional Monte Carlo model (Hieronimi et al., 2012) whose capability has been verified by in-situ measurements of the underwater light field and by inter-comparison with the widely-used radiative transfer code *HydroLight* (Mobley, 1994). The Monte Carlo model is optimized for investigations of spatially high resolved light fields underneath any desired wave profiles.

**Underwater light field fluctuations**M. Hieronimi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2 Applied methods

### 2.1 Radiative transfer models

The radiative transfer in water is simulated by means of two model approaches that are explained in detail by Hieronimi et al. (2012). At small depths, the underwater light regime is governed by high-frequency and small-scale intensity changes (e.g. Snyder and Dera, 1970; Hieronimi and Macke, 2010; Darecki et al., 2011). For this reason, we choose a high spatial resolution of  $dx = 2.5$  mm for our near-surface model; some current irradiance collectors have sensor head diameters of 2.5 mm (e.g. Darecki et al., 2011). In very clear seawater the fraction of total downwelling irradiance due to scattering in the first metres of the water body is small compared to the direct light beam and, furthermore, most of this scattered light is located very close to the initial light path because of the predominance of the forward scattering (Hieronimi, 2011). Thus, we neglect the diffuse light in our near-surface model (which is valid down to approximately 5 m) and just superpose narrow single rays according to the wave geometry. This approach has been used several times for example by Schenck (1957), Nikolayev et al. (1972), Stramski and Dera (1988), or Zaneveld et al. (2001). In our (near-surface) model, the intensity of each single ray is continuously attenuated by absorption of the medium

$$I = I_0 \exp(-aI_z) \quad (1)$$

where  $I_0$  is the initial intensity just after entering the water,  $a$  is the total absorption coefficient (sum of the absorption coefficients of seawater, particles and dissolved colored matter), and  $I_z$  is the covered distance with respect to the depth level  $z$ .

The second model approach is designated for larger water depths down to 100 m, where light scattering is not negligible. In this model, the spatial expansion and attenuation of single light beams is simulated on the basis of a Monte Carlo (MC) method (Hieronimi, 2011). The model domain is 100 m deep and 100 m wide, and has a spatial

## Underwater light field fluctuations

M. Hieronimi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 resolution of 10 cm horizontally and vertically. Individual photon transport calculations have been conducted for air-to-water transmission angles between 0° and 70° (relative to the downward vertical) with an angular spacing of 0.05° to 1°, each with  $2 \times 10^5$  photons. The outputs of the MC simulations are 2-D grids of downwelling irradiance for single light beams with varying transmission angles, which are stored in a database. Equivalent light fields are also calculated for isotropic diffuse skylight. In order to investigate the focusing effect of waves, individual (independent) single-beam patterns of downwelling irradiance are superposed according to the ratio of diffuse (skylight) to direct solar irradiation and the exact structure of the sea surface (including the vertical wave deflection).

10 The specific feature of our model is that any desired water surface structure can be implemented into the model. The accuracy of the results of the used radiative transfer model has been verified by irradiance measurements at sea and through intercomparison with the widely-used *HydroLight* software (Hieronymi et al., 2012).

### 15 2.1.1 Model input parameters

The model input parameters are listed in Table 1. They are selected in such a manner that maximum light fluctuations can be achieved (Dera and Stramski, 1986; Walker 1994; Gernez and Antoine, 2009). The radiative transfer is calculated for a single wavelength of 490 nm, which can be considered representative for the blue-green spectral band. Light of this spectral range can penetrate especially deep into oligotrophic waters. Different measurements have shown a wavelength-dependency of light fluctuations in shallow water (e.g. Gernez and Antoine, 2009; Darecki et al., 2011; Gege and Pinnel, 2011). However, orange-red light (580–700 nm) is strongly attenuated by absorption in water (e.g. Zielinski et al., 2002). Below 10 m water depth, the blue-green spectral components yield the vast majority of the photosynthetically active radiation (PAR) (400–700 nm). For this reason, we assume that the fluctuation characteristics at 25 490 nm can be considered representative for the variability of the entire PAR value.

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The presented simulations are carried out for perpendicular solar irradiation and for an extreme clear atmosphere, where the diffuse skylight due to atmospheric *Rayleigh* and *Mie* scattering accounts for 10 % only. The water body under consideration is very clear and oligotrophic; the (wavelength-depending) inherent optical properties of the well-mixed seawater are derived from a chlorophyll *a* concentration *Chl* of  $0.1 \text{ mg m}^{-3}$  (Morel et al., 2007; Morel, 2009). The resulting diffuse attenuation coefficient for the downward irradiance  $K_d$  (490 nm) is approximately  $0.038 \text{ m}^{-1}$ , relating to a flat sea surface.

**2.2 Description of the sea surface**

For generating two-dimensional irregular wave profiles, we apply the superposition principle of solitary sine waves, whose amplitudes are determined by the omnidirectional elevation spectrum. First, consideration is given to the effects of locally limited wind. The wind friction velocity basically defines the steepness of short waves of less than about a half metre length. In particular capillary-gravity and capillary waves in the wavelength range of 0.7–3 cm are most affected by the wind friction (Jähne and Riemer, 1990). In order to distinguish the pure wind effects on the light field, we use the short-wave part of the omnidirectional elevation spectrum as it is proposed by Elfouhaily et al. (1997). Figure 1 shows the utilized elevation spectra for wind speeds  $U_{10}$  (10 m above the sea level) from 3 to  $15 \text{ m s}^{-1}$  (solid lines). The corresponding curvature spectra exhibit a gravity-capillary peak at 1.7 cm wavelength for all wind speeds. With increasing wind, the spectra move together. This implies that the wave profiles that are generated out of the spectra (with random phase position) have almost the same statistical characteristics at strong wind, and thus the corresponding light fields must be very similar.

In reality, the short-wave spectra in Fig. 1 are modified by the long-wave regime of the present sea state. In particular at lower wind speeds, the unified long and short wave spectra adapt to the high-wind spectra with increasing wavelength (dashed lines). In case of weak wind over an unlimited fetch, ultra-gravity waves ( $> 2 \text{ cm}$ ) have actually higher amplitudes compared to a restricted fetch, which has to do with the wave growth

and the so called *wave age*. We bear this long-wave interaction in mind, but we primarily use the short-wave part of the spectra (solid lines) and apply this to waves of 6 mm to 50 cm length. We suggest that this spectral range is directly associated with the term “local wind”, although wind in general can also originate much longer waves, as ordinary gravity waves or swell waves.

In a second step, the significance of distinctive sea states is considered. Sea states are characterized by the significant wave height  $H_S$  and by a mean wave period  $T$ . Figure 2 (left) shows combinations of wave height and period together with the prevailing averaged wind speed  $U_{10}$  as it appears at the global ocean. The data are derived from the *KNMI/ERA-40 wave atlas* (Sterl and Caires, 2005). The corresponding wavelength  $L$  is additionally marked at the top of the chart. It is deduced from the dispersion relation for deep-water waves (*Airy wave theory*):

$$L = \frac{g}{2\pi} T^2 \quad (2)$$

in which  $g$  is the acceleration of gravity. The ratio of wave height to length  $H/L$  is the wave steepness, which is theoretically 0.14 at maximum; steeper waves break. The white areas at the bottom left in both diagrams of Fig. 2 stand for waves too steep to occur in nature (i.e. they are not in the wave atlas); these wave combinations are not considered in this study. Sea states with wave periods longer than 11 s (about 190 m wave length) are not considered either, since they are rather insignificant in terms of underwater light field variability.

Figure 2 right shows the frequency of occurrence (probability density function) of sea states in the global ocean (Sterl and Caires, 2005). Considerable differences arise in regional and monthly climatologies. In general, extremely steep seas occur rarely and wave systems with periods between 5 and 9 s and wave heights between 1 and 3 m occur most frequently (60 % of all cases). This range is framed white in the right diagram of Fig. 2.

With regard to the model assumption of homogeneous water properties, it should be mentioned that surface waves cause a mixing of the upper ocean layer. Simplistically,

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



it can be assumed that the sea water within this layer has homogeneous physical and bio-optical properties. Beside the fact that marine phytoplankton needs sufficient light (supplied from above) as well as nutrients (supplied from below), the mixed layer depth (MLD) is an important factor associated with the accumulation of phytoplankton biomass and the development of deep chlorophyll *a* maxima. In oligotrophic waters, where the surface mixed layer is poor in nutrients, chlorophyll *a* maxima are often found between 20 and 150 m depth with maximum concentrations of generally 3 to 10 times of those in surface waters (e.g. Anderson, 1969; Cullen, 1982; Furuya, 1990; Zielinski et al., 2002; Huisman et al., 2006). The depth of mixing  $Z_{\text{MLD}}$  can be predicted based on the knowledge of the wave climate. According to Babanin (2006), the wave-induced MLD can be approximated by:

$$Z_{\text{MLD}} = \frac{g}{2\omega^2} \ln \left( \frac{a_0^2 \omega}{Re_{\text{cr}} \nu} \right), \quad (3)$$

where  $g$  is the acceleration of gravity,  $\omega$  is the wave angular frequency,  $a_0$  is the wave amplitude,  $Re_{\text{cr}}$  is the critical Reynolds number (chosen as  $Re_{\text{cr}} = 3000$ ), and  $\nu$  the kinematic viscosity of ocean water (chosen as  $\nu = 1.35 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$ ). Figure 3 shows the mixed layer depth (Eq. 3) for the introduced wave classes. For a given wave period  $Z_{\text{MLD}}$  increases with growing wave height. The wave-induced upper ocean mixed layer can be more than 100 m deep. The figure provides an interesting hint to the depth of a deep chlorophyll *a* maximum, which might be affected by deeply penetrating light fluctuations. In addition, an abrupt rise of biomass concentration is associated with a rapid change of the IOPs of the water body, i.e. the light beam attenuation (absorption and scattering) increases.

The irregular sea state profiles used in this work are generated by means of unified spectra for long and short waves. The wind-dependent high-frequency part is based on the work of Elfouhaily et al. (1997) (and their references). The color-coding in Fig. 2 (left) shows the underlying “sea state typical” wind velocities  $U_{10}$ . We employ the two-parameter *Pierson-Moskowitz* spectrum for the respective long-wave regime as it is



proposed by the *International Towing Tank Conference* (Pierson and Moskowitz, 1964; ITTC, 2002):

$$S(f) = \frac{5H_S^2}{16T_P^4 f^5} \exp\left(-\frac{5}{4T_P^4 f^4}\right) \quad (4)$$

where  $f$  is the wave frequency,  $H_S$  is the significant wave height, and  $T_P$  is the spectral peak period. The two input parameters wave height and period (for ease without indices) are taken from Fig. 2 (left), where  $H$  is 0.5 to 7 m and  $T$  is 3 to 11 s, respectively. As seen in Fig. 2 (left), different sea states can arise from equal prevailing wind velocities, which is due to different stages of wave growth, wave energy dissipation, or due to superposition of different wave systems. In some cases, especially in low-wind situations, the transition between short and long wave spectra can be subject to a discontinuity. This particularly applies to the range of small gravity waves, which are not unimportant in the context of subsurface light variability. The issue is addressed by insertion of an intermediate function into the spectrum that approximately fits to observations (e.g. Leykin and Rozenberg, 1984; Donelan et al., 1985; Jähne and Riemer, 1990).

All wave profiles under consideration are 600 m long with a horizontal resolution  $dx$  of 0.1 mm. The corresponding modeled wave slopes are always normally distributed due to the superposition of stochastic independent elementary waves. Under realistic conditions, the steepness of capillary or gravity-capillary waves is modulated by longer waves in such way, that the short waves are steeper, on average, when riding on the forward faces of the longer waves (Longuet-Higgins, 1982). This observation explains the actual wind-dependency of the skewness of the wave slope distribution (Cox and Munk, 1954), which is not regarded in this work.

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.3 Fluctuation parameters

The variability of the downwelling irradiance  $E_d$  in water is based on spatial datasets with horizontal resolutions  $dx$  of 2.5 mm and 10 cm, respectively. The water depth  $z$  is positive downwards, vertical deflections of the sea surface (around the mean waterline at  $z = 0$  m) are treated exactly. The vertical length of a narrow water column is defined as reference or true water depth  $z_t$ , the following light field analysis refers to this reference depth.

Light fluctuations are commonly described by the coefficient of variation

$$CV = 100 \frac{\sigma_E}{\bar{E}_d} \quad (5)$$

given as the percentage ratio of the standard deviation  $\sigma_E$  and the averaged downwelling irradiance  $\bar{E}_d$  at the reference depth.  $E_d$  time series are typically normalized with the mean level irradiance in order to evaluate extreme values. The normalized downwelling irradiance (related to spatial  $E_d$  variability) is denoted as

$$\chi(x) = \frac{E_d(x)}{\bar{E}_d} \quad (6)$$

Irradiance pulses that exceed the mean irradiance by a factor  $\chi$  of more than 1.5 are termed underwater light flashes (Dera and Stramski, 1986). We define extreme values by means of the maximum normalized downwelling irradiance  $\chi_{\max}$  and by the “significant irradiance enhancement”  $\chi_{1/10}$ . The latter describes the mean of the 10 % highest irradiance values; the labeling is motivated by the definition of the significant wave height  $H_s$  (or  $H_{1/3}$ ), which is the average height of the one-third highest waves. On the one hand the significant irradiance enhancement provides a statistically smoothed function of extreme intensity peaks over the water depth; on the other hand  $\chi_{1/10}$  can be seen as a rough estimate for measured  $E_d$  maximum values, taking into account

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that the sampling rates of some radiometers may be insufficient for high-frequency irradiance measurements and that sensor integration times can elongate with increasing water depth (we did  $E_d$  measurements with a *RAMSES-ACC-VIS* radiometer (*TriOS*, Germany) with a spectral range of 320 to 950 nm and effective sampling rates of 2 to 8 s (Hieronymi et al., 2012)).

The modeled spatial light fluctuations are subject to a wavenumber analysis (analogous to frequency analysis) in order to characterize the statistical dynamics of the underwater light field and thereby draw conclusions on the influence of associated wave regimes at the sea surface. The power spectral density of  $\chi$  fluctuations (variance spectrum) is computed by means of a fast *Fourier* transformation. The spectral peak length  $L_p$  marks the wavelength that contributes most to the variance. Nevertheless, an accurate determination of  $L_p$  is difficult for example in cases with a broad spectral maximum or if individual peaks are located in close vicinity. We therefore additionally use the mean (wave-) length of fluctuations  $L_m$  which comprises more of the relevant spectral band. It is determined by using the  $n$ -th spectral moments

$$m_n = \int_0^{\infty} k^n S_{\chi}(k) dk \quad (7)$$

where  $k$  is the (angular) wavenumber and  $S_{\chi}(k)$  the spectral density of the  $\chi$ -profile. The mean fluctuation length is the area below the spectrum  $m$  divided by the spectral center of gravity  $m_1$

$$L_m = \frac{m_0}{m_1} \quad (8)$$

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3 Results and discussion

#### 3.1 Direct influence of local wind conditions

The influence of spatially very limited, so called “local”, wind on the underwater light field is accessed by using the near-surface model with  $dx = 2.5$  mm horizontal resolution. The evaluated model domain is 5 m deep and 20 m wide. The implemented irregular wave profiles with random phase are generated from the wave spectra that are shown in Fig. 1 (solid lines). Figure 4 shows two 1 m wide sections of resulting spatial  $E_d$  distributions for a “light breeze” and a “near gale” wind situations ( $3$  and  $15$   $\text{m s}^{-1}$ ). The red colors indicate a radiative enhancement of more than 100% of the surface insolation which is caused by the wave-focusing; bluish colors stand for a reduction due to defocusing. Waves of the capillary and capillary-gravity range of 0.7–3 cm length play a key role in near-surface light focusing. Their spectral densities strongly increase with the wind friction velocity. In contrast, small gravity waves are less dependent on the local wind speed (Jähne and Riemer, 1990). Very close to the surface, the light field variance is dominated by the de-/focusing of waves in this order of magnitude, visible as clear stripes of  $E_d$  enhancements. Local wind determines the steepness of these waves, affecting the depth of the (first) focal point, i.e. the more wind, the closer the focal point comes to the surface. The red single rays are further deflected by overlaying small gravity waves, which again are steeper at stronger winds. This leads to intensified light beam grouping at different depths and together with the occurrence of secondary and further focal points that are caused by neighboring capillary-gravity waves, those larger waves are responsible for very intense irradiance fluctuations within the top 5 m layer. With increasing depth the narrow stripes of radiative concentration are geometrically scattered, light focusing is reduced, and in addition the intensity of each single ray is attenuated.

Some statistical characteristics of the two light fields (Fig. 4) are compared in Fig. 5. The occurrence frequencies of downwelling irradiance values are shown in Fig. 5a, b. The mean values  $\bar{E}_d$  are almost identical, mainly because the surface albedo (i.e. irra-

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



diance reflectance) is wind-independent at high sun (Preisendorfer and Mobley, 1986);  $\bar{E}_d$  decreases exponentially with increasing water depth. However, one can clearly identify differences in the depth-dependence of the probability density functions *PDF*. Because of wave-focusing both probability distributions are highly skewed to the right and heavily tailed. The wind-dependent surface geometry affects the occurrence depth and intensity of extreme irradiance values. The *PDF* skewness  $\gamma_1$  is around 1.5 at depths between 0.5 and 2 m in the low wind situation, but only little more than 1 between 0.1 and 1.5 m depth in the case of strong wind. In both cases the excess kurtosis  $\gamma_2$  can take maximum values of 3, but  $\gamma_2$  decays faster at strong wind, i.e. the probability distribution faster approaches a *Gaussian* shape and less variance results from extreme intensity peaks.

Figure 5c, d show the frequency of intensity peaks  $N$  that exceed an irradiance level. In general, we see stronger irradiance peaks at low wind speed exceeding  $7\bar{E}_d$  at the maximum. According to the model, light flashes of  $6\bar{E}_d$  appear down to 5 m water depth at moderate wind speeds, whereas the flash intensity is evidently reduced at the same depth at strong wind.

The corresponding wavenumber analysis is illustrated in Fig. 5e, f. Red colors stand for spectral ranges that contribute large amounts to the variability of the normalized irradiance  $\chi$ . White and bluish colors denote negligible and small amplitudes of the power spectral density  $S_\chi$ . We see stronger and distinctly deeper reaching variance at  $3 \text{ m s}^{-1}$  wind speed. In both cases irradiance fluctuations near the surface are governed by small waves of less than 5 cm length (the bounds of the wavelength range with most wind friction dependency (0.007 and 0.03 m) are additionally marked). With increasing depth the capillary wave ( $L < 1.7 \text{ cm}$ ) influence fades away and larger waves dominate the fluctuations, i.e.  $L_p$  and  $L_m$  grow.

Statistical characteristics of near-surface light field variability in dependence of local wind are summarized in Fig. 6. The maximum coefficient of variation  $CV_{\max}$  is associated with the geometrical depth of the (first) focal point of capillary-gravity and small gravity waves. At a wind speed of  $3 \text{ m s}^{-1}$ ,  $CV_{\max} = 83 \%$  at a water depth of 1.25 m;

$CV_{\max} = 81\%$  at 25 cm depth and  $15 \text{ m s}^{-1}$  wind. Below 80 cm water depth, irradiance variability decreases with growing wind. The same wind dependency shows up in the significant irradiance enhancement  $\chi_{1/10}$  (Fig. 6b). All maximum values reach the same order of magnitude (approximately 2.8). At 5 m water depth,  $\chi_{1/10}$  is between 1.7 and 2.4. The characteristic values  $L_m$  and  $L_p$  of the spectral analysis (Fig. 6c, d) confirm the principal trend of increasing influence of longer waves with increasing water depth. In 5 m depth,  $L_p = 9 \text{ cm}$  and  $34 \text{ cm}$  for wind speeds of  $3 \text{ m s}^{-1}$  and  $15 \text{ m s}^{-1}$ , respectively. These averaged distances between two clustered irradiance maxima are clearly visible in Fig. 4.

The wave spectra in Fig. 1 move closer together with increasing wind speed. Thus, the wave and the underwater light field characteristics approach each other. In terms of the pure geometrical influence of surface waves, differences in the statistics of the underwater irradiance fluctuations become insignificant with wind speeds of over  $12 \text{ m s}^{-1}$ . Apart from that, we have to acknowledge that the description of the sea surface using linear superposition of spectrally weighted harmonics (*Gaussian* surface) becomes increasingly inaccurate in higher wind speeds. Generally, wave crests are higher and sharper and the troughs are shallower and flatter. Nonlinear wave interactions, including the generation of “parasitic capillaries” on the downwind faces of gravity waves alter the hydrodynamic properties and the shape of the free water surface (e.g. Longuet-Higgins, 1982; Zhang, 1995).

The model results may be biased because they lack wind-stress-dependent bubble injection into the water column. Occasional wave breaking, which generally provides the dominant source of foam at the surface and bubbles in the water, can be already observed at wind speeds about  $3 \text{ m s}^{-1}$ . If the wind speed exceeds  $7 \text{ m s}^{-1}$ , horizontally uniform bubble layers can evolve in the first metres and persist over hours (Thorpe, 1992). At the wind speeds under consideration ( $3\text{--}15 \text{ m s}^{-1}$ ), the contribution of foam to the (broadband) surface albedo, and thus irradiance transmission, is very limited despite its high reflectance (Zhang et al., 2006). More relevant are air bubbles in the upper water layer as they cause enhanced reflectance (in clear water the reflectance

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



has a strong dependence on the light spectrum). But in contrast to the upwelling light, bubbles induce only very small enhancement in downwelling irradiance within the top several tens of centimetres just beneath the surface and below that layer,  $E_d$  is reduced compared to bubble-free water (Stramski and Tegowski, 2001). For this reason we must consider that air bubbles in water impair the effectiveness of wave lensing, and thus damp the intensity of the described fluctuations, in particular at increasing wind speed. Nevertheless, for the sake of model simplicity and a better inter-comparison of the model results, whitecaps and bubbles are neglected within all presented radiative transfer simulations.

### 3.1.1 The influence of growing waves

It is obvious that no underwater light fluctuations can evolve from a perfectly flat water surface, i.e. at  $0 \text{ m s}^{-1}$  wind speed we have the absolute fluctuation minimum in terms of  $CV$ , et cetera. A threshold wind speed or friction velocity is required to actually produce waves. The reason is clearly that energy input from the wind does not exceed viscous dissipation below this threshold (Donelan and Plant, 2009). The further growth of small-scale waves depends on the energy input by the turbulent wind field (i.e. the wave field develops with increasing fetch or duration), on the energy transfer between waves of different length by nonlinear wave-wave interaction, and on the energy dissipation by wave breaking, viscous dissipation, and turbulent diffusion (Phillips, 1985). The simulation results that are shown in the previous Sect. 3.1 are based on the short-wave spectra as used by Elfouhaily et al. (1997); they are plotted in Fig. 1 (solid lines). Over the open ocean, we must consider all the just mentioned aspects of wave field interactions and assume a long-wave modification of the short-wave spectra (dashed lines in Fig. 1). This in particular concerns the wave fields at low and moderate wind conditions, where ultra-gravity waves (wavelength range approximately 1.7 cm to 1.6 m) have been stronger developed.

Figure 7 shows the corresponding influence on the underwater light field in terms of the significant irradiance enhancement. In the case of strong wind, the fluctuation

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 statistics remain unchanged, because the wave spectra have almost the same shape. Whereas at low wind speed of  $3 \text{ m s}^{-1}$ , we observe clearly intensified irradiance peaks due to the lensing effect of steeper ultra-gravity waves. Within the depth range of 0.5 to 3 m, maximum light flashes exceed  $10\bar{E}_d$  ( $\chi_{\max} = 12$ ), the skewness of the  $E_d$  probability distribution is above 2 (instead of 1.5 in the reference case), the excess kurtosis exhibits values of 7 (instead of 3), and  $CV$  is around 100%. Consequently, the spectral characteristics of light fluctuations change too;  $L_m$  and  $L_p$  grow faster with increasing depth, e.g.  $L_p(z_t = 5 \text{ m}) = 28 \text{ cm}$  (instead of 9 cm). The principal wind-dependency of light field statistics (Fig. 6) is retained, but with amplified values at low and moderate wind speeds.

### 3.2 Influence of the sea state on the underwater light field

15 Now we focus on light field changes due to variations of the long-wave part of the sea spectrum which is referred to as sea state. The directly wind-dependent short-wave part is also included in the applied spectra. The evaluable model domain covers 500 m horizontally and up to 100 m vertically (water depth); the model resolution is 10 cm in both directions. The model takes into account all fractions of direct and scattered radiation.

20 We would like to point out that the use of the two irradiance “detector sizes” of 2.5 mm for the near-surface model and 10 cm for the deep-water model may lead to differing results in the statistical analysis. The rather coarse resolution in the deep-water model makes sense to cover the large study area (of  $50\,000 \text{ m}^2$ ), and it takes into consideration the fact that the saturation time of radiometers typically increases with increasing water depth (e.g. the integration time of the spectrally measuring *RAMSES-ACC-VIS* can be up to 8 s). The broader detector averages out substantial parts of the high-frequency fluctuations and diminishes the amplitudes of light flashes near the surface. 25 If we carefully look at Fig. 4 and visually average the spatial  $E_d$  distribution over 10 cm horizontally, we will find actually higher variability at 5 m depth in the case of strong wind which contradicts the statements in Fig. 6a. Darecki et al. (2011) compare the

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





effects of different collector diameters (ranging from 2.5 to 23 mm) on temporal  $E_d$  measurements; their data from 2.7 m depth do not reveal a clear effect of the collector size on the  $E_d(t)$  signal (different IOPs and irradiation conditions as in our case). On the one hand this is due to the scattering processes in water and the associated increasing diffuseness of light, and on the other hand it shows the vanishing influence of the high-frequency contribution to the light field variance which depends on the water surface geometry (as shown in Fig. 6c, d). Below approximately 10 m water depth, the used model resolution of  $dx = 10$  cm is perfectly sufficient to characterize the light field fluctuations.

Figure 8 shows the computed irradiance distribution beneath the most prevalent sea state with  $H_s = 2$  m,  $T_p = 8$  s (both input parameters for the spectrum Eq. 4), and  $U_{10} = 4.8$  m s<sup>-1</sup>. The 10 and 5%  $E_d$  contour lines are each highlighted to illustrate the depth-effect of the irregularly deflected sea surface. In the example, the wave-induced mixed layer depth, where under certain circumstances enhanced biomass concentration (a deep chlorophyll *a* maximum) can be expected, is approximately at 35.5 m (Eq. 3). Here, the averaged downwelling irradiance is 27.7% (of the surface value), and individual  $E_d$  values vary between 22.4 and 39.2% ( $\chi_{\max} = 1.44$ ,  $CV = 8.15\%$ ). According to the model, light flashes ( $\chi = 1.5$ ) can appear down to 30 m water depth. Wave-induced light variability can be observed even at the lower boundary of the model domain, i.e. at 90 m depth  $\bar{E}_d = 3.2\%$  (the  $E_d$  minimum and maximum is at 2.9 and 3.4%, respectively),  $CV = 2.84\%$ , and the mean (peak) fluctuation period is around 6 s.

The statistical characteristics of the simulated irradiance variability with respect to different sea states are summarized in Fig. 9. The data refer to four water depths with  $\bar{E}_d = 50, 25, 10,$  and  $5\%$  of the surface irradiance value, which approximately correspond to 20, 40, 60, and 80 m, respectively. Figure 10 provides additional information on vertical changes of selected fluctuation parameters, where both the sea state related minimum and maximum values are identified. The red lines in Fig. 10a–c mark

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the parameter values that are averaged over the range of highest occurrence probability shown by the white frame in Fig. 2.

With regards to the  $CV$  (Figs. 9a–d and 10a), the strongest light fluctuations appear at particular steep sea states with high  $H/L$  values but low probability of occurrence (see Fig. 2). Indirectly the statement is also true for high wind speeds, since wind speed and sea state are correlated. It is obvious that fully developed seas shape larger lens surfaces with enhanced depth effect. Figure 9d shows the  $CV$  at 80 m water depth, ranging from 1.1 % ( $H_S = 0.5$  m,  $T = 9$  s,  $H/L = 0.004$ ) to 14.3 % ( $H_S = 3.5$  m,  $T = 5$  s,  $H/L = 0.09$ ), but a  $CV$  around 4.7 % being most likely. A special feature noticeable at the 50 %  $E_d$  level with  $H_S = 0.5$  m (Fig. 9a) is the slight increase of fluctuations in the case of a wave period of 11 s compared to the case with 5 s, where  $CV$  is 16.7 and 13.4 %, respectively. The wave steepness is smaller, thus the enhanced  $CV$  must be due to the slightly enhanced local wind speed  $U_{10}$  that is characteristic for the 11 s sea state (see Fig. 2). This observation is an indication of the still present depth effectiveness of a local wind (and thus of capillary-gravity waves) at 20 m depth.

With regards to  $\chi_{1/10}$  and  $\chi_{\max}$  (Figs. 9e–l and 10b), we see the same depth changes as in  $CV$ . As mentioned before,  $\chi_{1/10}$  is a good benchmark of the maximum measurable irradiance enhancement (with e.g. a *RAMSES-ACC-VIS*). At 20 m water depth, our comparative measurements mostly showed  $E_d$  maxima being in the order of  $1.3$ – $1.4\bar{E}_d$  which absolutely fits to the simulated  $\chi_{1/10}$  of the corresponding sea states (Fig. 9e). In rare events irradiance values of more than  $1.5\bar{E}_d$  were measured at this depth level under moderate sea conditions (Veal et al., 2010; Hieronymi et al., 2012). According to Figs. 9i and 10b, the mean value of the simulated irradiance peaks  $\chi_{\max}$  within the range of high occurrence probability is around 2, i.e. still at 20 m depth,  $E_d$  can achieve maximum values of 100 % of the surface irradiation. Theoretically,  $\chi_{\max}$  can be up to 2.5 (at  $H_S = 2.5$  m,  $T = 4$  s,  $H/L = 0.1$ ). In case of perfect single waves,  $\chi_{\max}$  can be 10 at 20 m and 1.5 at 80 m depth, respectively (Hieronymi, 2011). In reality, the surface roughness causes considerable noise in the light regime that dampens the lensing effectiveness of longer waves. However, deep-reaching light flashes originate from the

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



superposition with fully developed gravity waves. They can reach 30 m depth under moderate and prevailing sea conditions but down to 75 m in extreme cases (Fig. 10b).

The wavenumber analysis of underwater irradiance fluctuations confirms the increasing importance of growing surface waves (Figs. 9m–t and 10c).  $L_m$  provides a clearer picture of the recurrence of radiative peaks as a function of different sea states compared to the rather noisy  $L_p$ . But  $L_p$  gives a direct hint on the mean distance between two intense irradiance peaks, and thus about the magnitude of surface wavelengths that dominate the irradiance variability. As seen in Fig. 9r, the light variability of flatter sea states is (even at 40 m depth) still dominated by ultra-gravity waves, i.e. it is associated with local wind. The differences in  $L_m$  increase with water depth. At 80 m depth,  $L_m$  is between 5 and 45 m, the mean value is around 30 m. The dependency of the mean fluctuation length on the wave steepness becomes clear, if we recall the focusing effect of single waves again (Hieronymi et al., 2012). The steeper a wave, the closer is its focal point to the sea surface, i.e. if we look at a certain depth level, than the focal point of a steeper wave must be associated with a longer wavelength. This mechanism is mirrored for example in Fig. 9p, where steeper sea states are associated with larger distances between relatively strong irradiance enhancements.

Figure 11 points out the influence of surface waves on the irradiance fluctuations at certain depths, or more precisely it marks the wavelengths that contribute at least 0.1 % to the total spectral variance of the signal. Both used model resolutions and all investigated cases are considered in this figure. The lower boundary of the spectral wavelength range is partly dashed which indicates that shorter wavelengths occur, but due to resolution limitations no statements can be made about this range. Local wind primarily affects the light fluctuations down to roughly 10 m depth, if we consider the most wind-dependent capillary and capillary-gravity waves of 0.7 to 3 cm length only. The depth-impact of ultra-gravity waves (up to 1 s wave period or 1.5 m length), which are also closely related to the prevailing wind regime, is restricted to approximately 50 m depth. Waves at the surface must be at least 4 m long to being able to affect the irradiance variability at a water depth of 80 m. On the other hand, waves longer than

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



20 cm do not contribute considerably to the irradiance variability at 1 m depth. The light field influence due to ordinary gravity waves ( $> 1$  s period) starts at 13 m water depth. And even swell waves potentially influence the light field at greater depths, e.g. a 10 s swell-dominated sea state affects the light field below 40 m depth.

### 5 3.3 Discussion

We now want to put the simulation results into context with previous publications and measurements. Remember that our model assumes optimal conditions for wave-induced underwater light fluctuations, for example an extreme clear atmosphere that causes 10 % diffuse surface irradiation only and the high standing sun ( $\theta_S = 0^\circ$ ).

#### 10 3.3.1 Discussion of the applied methods

Our Monte Carlo model is based on a completely novel approach (Hieronymi et al., 2012). It is optimized for investigations of the light field below arbitrary sea surfaces. The model yields reliable results on the spatial  $E_d$  distribution, from which all statistical properties of the light field including the periodicity of extreme values can be deduced. In comparison to other models (e.g. Deckert and Michael, 2006; D'Alimonte et al., 2010; You et al., 2010; Shen et al., 2011) our model provides a high information density with high spatial resolution down to much greater depths (100 m). And yet our model is relatively fast (light fields below more than 100 different wave profiles were investigated for this study).

The description of the sea surface and its implementation into the radiative transfer model remains a critical factor that still leaves room for improvements. Wave spectra, as the *Pierson-Moskowitz* (PM) spectrum, have been utilized with this regard (e.g. by Nikolayev et al., 1972; Weber, 2010; You et al., 2010). However, the PM spectrum applies to gravity waves in a fully developed sea, which means that the wave growth has reached a point of equilibrium with the wind input. The PM spectrum does actually not apply to capillary and ultra-gravity waves that are most important in terms of light

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



field variability (at least near the surface); the validity of such a long-wave spectrum is limited to wavenumbers up to approximately 10 times its spectral peak  $k_p$  (e.g. Leykin and Rozenberg, 1984; Elfouhaily et al., 2009). Over the open ocean, sea states are very often influenced or even dominated by swells (e.g. as seen in Fig. 2, wave classes with large periods appear despite the fact that there is not enough prevailing wind to actually generate these long waves). In the presence of swells, the wave spectrum exhibits a narrow peak, or in combination with a wind sea it shows a bimodal shape (e.g. ITTC, 2002); both spectral shapes differ from the PM shape. Nevertheless, the spectra that we use include all relevant classes of waves ranging from capillary to swell waves; the long-wave part is always described by the *Pierson-Moskowitz* spectrum (Eq. 4) and the wind-specific short-wave part is additionally accounted for.

In some radiative transfer models (e.g. *HydroLight*), the wind-dependent roughness of the sea surface is implemented by the statistical slope distribution of water facets according to Cox and Munk (1954). Our wave profiles, which are generated from the unified spectra for long and short waves, have a slope distribution that can differ from the *Cox-Munk* distribution. There is a good match at moderate wind speeds around  $5 \text{ m s}^{-1}$ , but at less wind we observe a narrower distribution and at higher wind speeds we have more frequently steeper slopes compared to *Cox-Munk*. One explanation for the deviations is that we partly consider extreme sea events with significant wave heights up to 7 m. However, there are clear differences at high wind speeds ( $> 10 \text{ m s}^{-1}$ ) that raise the question on the validity of our approach with linear wave theory and the disregard of nonlinearities. Just recently, there have been two studies on polarized underwater light fields by You et al. (2011) and Xu et al. (2011), where three-dimensional wave elevations were derived from high-resolution wave slope measurements and from a numerical “high-order spectral method”, respectively. Both realizations of the sea surface sound promising and could be applied in future studies. With this regard, we would expect that the light field variance rather weakens at strong wind and increases at light wind in comparison with our model.

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Our model considers the exact shape of wave profiles in the  $x$  and  $z$  directions (with  $dx = 0.1$  mm). Viewed statistically, the sea states investigated can have maximum wave amplitudes of more than 5 m around the mean waterline (e.g. with  $H_s = 7$  m). As far as we know, there is only one comparable model by D'Alimonte et al. (2010) that includes vertical wave elevations as well. Regarding the precise wave geometry gives a more realistic picture of the underwater irradiance distribution, i.e. otherwise, in case of a  $z$ -invariant light input, the lensing pattern would be distorted. However, the differences in the light field statistics are generally minor.

### 3.3.2 Discussion of the simulation results

With regards to the magnitude of irradiance enhancements, temporally high resolution measurements of the downwelling irradiance show that near the surface instantaneous light pulses can exceed  $10\bar{E}_d$ ; the amplitude of the strongest flash was in excess of  $15\bar{E}_d$  (measured with a 2.5 m<sup>2</sup> collector at 0.86 m water depth) (Darecki et al., 2011; Gernez et al., 2011). Our near-surface model is capable of reproducing such irradiance enhancements for light to moderate wind conditions and under the presence of steeper ultra-gravity waves. In theory, irradiance pulses can exceed the mean irradiance by a factor of 40 at a water depth of 1 m under the assumption of an ideal steep wave of 80 cm length (Hieronymi et al., 2012).

According to our model, light flashes can occur much deeper than observed so far. Under realistic conditions, in terms of the sea state, irradiance peaks with  $1.5\bar{E}_d$  should be still possible at 30 m water depth (Fig. 10b), but their occurrence frequency is low. Up to now, light flashes were recorded down to a depth of 21 m only (Veal et al., 2010; Hieronymi et al., 2012). Irradiance variability ( $CV$ ) of around 10% has been detected at the depth range of 30 to 35 m (Stramska and Dickey, 1998; Veal et al., 2010) which absolutely fits to our simulation results for moderate sea states (Fig. 10a). However, the simulated deep-water maximum values that are presented in this article should be verified with more precisely and temporally high resolution in-situ measurements.

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







riod, rather than wind speed, and that between 3 and 25 m water depth waves with  $H_s$  of 1.5 to 2.5 m provoke the strongest intensity fluctuations (Hieronymi and Macke, 2010). In this sense a quite similar conclusion can be drawn from data of Nikolayev and Yakubenko (1978). With regard to a water depth of 20 m, seas with small wave heights ( $< 0.4$  m) cause considerably less light fluctuation compared to higher waves (of e.g. 1.5 m height). The simulation results of this study are in line with these observations if we consider the most likely sea states during the measuring campaigns (see Fig. 9a). But the findings also suggest that it is rather the wave steepness  $H/L$  of the characteristic wave of a sea state that influences the light variability at greater depths, namely the steeper a wave system, the stronger are the underwater light field fluctuations. This is exactly the same relation for single waves (Hieronymi et al., 2012) and of course it applies for each constant wavelength  $L$ .

Steep wave systems are mostly associated with strong wind (Fig. 2). In this case it has to be assumed that in reality the lensing efficiency is considerably reduced because of nonlinear hydrodynamic interactions at the surface and air bubbles that are induced by breaking waves (e.g. Stramski and Tegowski, 2001). Thus, it has to be assumed that at high wind speed our model overestimates the strength of irradiance variability down the water column. However, the model assumption of vertical solar insolation ( $\theta_S = 0^\circ$ ) can only occur within the tropics where the yearly mean of the wind speed does not exceed  $9 \text{ m s}^{-1}$  (Sterl and Caires, 2005). Our model generally shows very good agreement with observations under such light to moderate wind conditions.

Another question is whether different sea states at constant wind speed actually have an impact on the mean downwelling irradiance in the water column. We have slightly different slope distributions due to the differing long-wave part of the spectrum, i.e. the presence of a more developed sea is comparable with slightly higher wind speeds (related to the *Cox-Munk* distribution). Within our simulations we have no considerable changes of  $\bar{E}_d$  due to wind or the sea state. This is mainly because of the perpendicular light input where the irradiance reflectance is very low and almost wind-independent

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





(Preisendorfer and Mobley, 1986). But we would conclude that there is an effect at low sun positions, because then more light is reflected at the stronger roughened surface.

## 4 Summary

This paper presents the first systematic analysis of the influence of various wind and wave regimes on the downwelling irradiance variability within the upper ocean mixed layer down to 100 m depth. The study is based on a two-dimensional Monte Carlo radiative transfer model the principle suitability of which is proven by comparisons with several corresponding in-situ measurements and with the radiative transfer model *HydroLight* (Hieronymi et al., 2012). We assume model conditions that favor the development of extreme light fluctuations within the entire lit water column, i.e. monochromatic light at 490 nm, high standing sun ( $0^\circ$ ), a very low portion of diffuse sky radiation (10%), and very clear well-mixed sea water with a low chlorophyll *a* content of  $0.1 \text{ mg m}^{-3}$ . Any arbitrary wave profile can be implemented into the model in order to simulate the resulting underwater light field. Two model resolutions are chosen, the near-surface model is resolved with 2.5 mm horizontally and the deep-water model with 10 cm, respectively. The basic difference is that the latter considers all direct and diffuse radiation in the water, while the near-surface model considers the direct light beam only.

In a first step we look at the influence of wind on the light field near the surface. As a locally occurring event, wind primarily affects the steepness of small waves in the transition region from capillary to ultra-gravity waves. The near-surface light field is dominated by the focusing effect of these small waves. In case of high wind speeds, we have steep capillary-gravity waves that build up focal points closer to the surface than flatter waves that develop at low wind speeds. For this reason we have a vertical shift of the maximum light fluctuations towards the surface at growing wind. However, maximum irradiance peaks can be observed at relatively low wind speeds of  $3 \text{ to } 5 \text{ m s}^{-1}$  (where  $3 \text{ m s}^{-1}$  is the lowest wind speed under consideration); within the depth range of 0.5 to 3 m, maximum light flashes can exceed  $7\bar{E}_d$ . At low wind, the strength of ir-

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



radiance fluctuations can be even amplified under the influence of further developed ultra-gravity waves, thereby  $\chi_{\max}$  exceeds 12.

In the second step we investigate fully developed sea states that can occur in the open sea (some of the cases occur very rarely and are of rather theoretical interest).

The applied wave spectra cover all magnitudes of ocean waves, ranging from directly wind-dependent capillaries to swell waves which are independent of local wind. Gravity waves influence the light field to much greater depths; they are the reason for (low-amplitude) irradiance variability at 80 m and deeper. According to the model it should be possible to observe light flashes (with  $1.5\bar{E}_d$ ) still at 30 m water depth under realistic sea conditions; theoretically light flashes can reach 75 m. Our simulations show that the light variability at greater depths more clearly depends on the wave steepness  $H/L$  of the characteristic wave of a sea state than on the wave height; the steeper a wave system, the stronger are the underwater light field fluctuations.

*Acknowledgements.* This work was supported by the German Research Foundation *DFG* (contract MA 2225/11-1).

## References

- Anderson, G. C.: Subsurface chlorophyll maximum in the Northeast Pacific Ocean, *Limnol. Oceanogr.*, 14, 386–391, 1969.
- Babanin, A. V.: On a wave-induced turbulence and a wave-mixed upper ocean layer, *Geophys. Res. Lett.*, 33, L20605, doi:10.1029/2006GL027308, 2006.
- Cox, C. and Munk, W.: Measurements of the roughness of the sea surface from photographs of the sun's glitter, *J. Opt. Soc. Am.*, 44, 838–850, 1954.
- Cullen, J. J.: The deep chlorophyll maximum: comparing vertical profiles of chlorophyll *a*, *Can. J. Fish. Aquat. Sci.*, 39, 791–803, doi:10.1139/f82-108, 1982.
- D'Alimonte, D., Zibordi, G., Kajiyama, T., and Cunha, J. C.: Monte Carlo code for high spatial resolution ocean color estimation, *Appl. Optics*, 49, 4936–4950, doi:10.1364/AO.49.004936, 2010.

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Darecki, M., Stramski, D., and Sokolski, M.: Measurements of high-frequency light fluctuations induced by ocean surface waves with an underwater porcupine radiometer system, *J. Geophys. Res.*, 116, C00H09, doi:10.1029/2011JC007338, 2011.
- Deckert, R. and Michael, K. J.: Lensing effect on underwater levels of UV radiation, *J. Geophys. Res.*, 111, C05014, doi:10.1029/2005JC003332, 2006.
- Dera, J. and Stramski, D.: Maximum effects of sunlight focusing under a wind-disturbed sea surface, *Oceanologia*, 23, 15–42, 1986.
- Donelan, M. A. and Plant, W. J.: A threshold for wind-wave growth, *J. Geophys. Res.*, 114, C07012, doi:10.1029/2008JC005238, 2009.
- Donelan, M. A., Hamilton, J., and Hui, W. H.: Directional spectra of wind-generated waves, *Philos. T. Roy. Soc. A*, 315, 509–562, doi:10.1098/rsta.1985.0054, 1985.
- Elfouhaily, T., Chapron, B., Katsaros, K., and Vandemark, D.: A unified directional spectrum for long and short wind-driven waves, *J. Geophys. Res.*, 102, 15781–15796, doi:10.1029/97JC00467, 1997.
- Furuya, K.: Subsurface chlorophyll maximum in the tropical and subtropical Western Pacific Ocean: vertical profiles of phytoplankton biomass and its relationship with chlorophyll *a* and particulate organic carbon, *Mar. Biol.*, 107, 529–539, doi:10.1007/BF01313438, 1990.
- Gege, P. and Pinnel, N.: Sources of variance of downwelling irradiance in water, *Appl. Optics*, 50, 2192–2203, doi:10.1364/AO.50.002192, 2011.
- Gernez, P. and Antoine, D.: Field characterization of wave-induced underwater light field fluctuations, *J. Geophys. Res.*, 114, C06025, doi:10.1029/2008JC005059, 2009.
- Gernez, P., Stramski, D., and Darecki, M.: Vertical changes in the probability distribution of downwelling irradiance within the near-surface ocean under clear sky conditions, *J. Geophys. Res.*, 116, C00H07, doi:10.1029/2011JC007156, 2011.
- Hieronymi, M.: Solar radiative transfer into the ocean: A study on underwater light fluctuations due to surface waves, Ph.D. Thesis, Christian-Albrechts-Universität zu Kiel, Germany, 103 pp., 2011.
- Hieronymi, M. and Macke, A.: Spatiotemporal underwater light field fluctuations in the open ocean, *J. Eur. Opt. Soc.-Rapid*, 5, 10019S, doi:10.2971/JEOS.2010.10019S, 2010.
- Hieronymi, M., Macke, A., and Zielinski, O.: Modeling of wave-induced irradiance variability in the upper ocean mixed layer, *Ocean Sci.*, 8, 103–120, doi:10.5194/os-8-103-2012, 2012.

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Huisman, J., Thi, N. N. P., Karl, D. M., and Sommeijer, B.: Reduced mixing generates oscillations and chaos in the oceanic deep chlorophyll maximum, *Nature*, 439, 322–325, doi:10.1038/nature04245, 2006.

ITTC: The specialist committee on waves – Final report and recommendations to the 23rd ITTC, in: *Proceedings of the 23rd ITTC*, Vol. 2, 505–551, Venice, Italy, 8–14 September 2002, 2002.

Jähne, B. and Riemer, K. S.: Two-dimensional wave number spectra of small-scale water surface waves, *J. Geophys. Res.*, 95, 11531–11546, doi:10.1029/JC095iC07p11531, 1990.

Leykin, I. A. and Rozenberg, A. D.: Sea-tower measurements of wind-wave spectra in the Caspian Sea, *J. Phys. Oceanogr.*, 14, 168–176, doi:10.1175/1520-0485(1984)014<0168:STMOWW>2.0.CO;2, 1984.

Longuet-Higgins, M. S.: On the skewness of sea-surface slopes, *J. Phys. Oceanogr.*, 12, 1283–1291, doi:10.1175/1520-0485(1982)012, 1982.

Mobley, C. D.: *Light and Water: Radiative Transfer in Natural Waters*, Academic Press, San Diego, 592 pp., 1994.

Morel, A.: Are the empirical relationships describing the bio-optical properties of case 1 waters consistent and internally compatible?, *J. Geophys. Res.*, 114, C01016, doi:10.1029/2008JC004803, 2009.

Morel, A., Gentili, B., Claustre, H., Babin, M., Bricaud, A., Ras, J., and Tieche, F.: Optical properties of the “clearest” natural waters, *Limnol. Oceanogr.*, 52, 217–229, doi:10.4319/LO.2007.52.1.0217, 2007.

Nikolayev, V. P. and Khulapov, M. S.: Use of a nonstatistical model to explain the mechanism of underwater illumination fluctuations, *Izv. Atmos. Ocean. Phys. New York.*, 12, 993–997, 1976.

Nikolayev, V. P. and Yakubenko, V. G.: On the relationship between the statistical characteristics of the underwater light field and the wave state characteristics, *Izv. Atmos. Ocean. Phys.*, 14, 88–92, 1978.

Nikolayev, V. P., Prokopov, O. I., Rozenberg, G. V., and Shevernev, V. I.: Statistical properties of the underwater illumination, *Izv. Atmos. Ocean. Phys.*, 8, 936–944, 1972.

Phillips, O. M.: Spectral and statistical properties of the equilibrium range in wind-generated gravity waves, *J. Fluid Mech.*, 156, 505–531, doi:10.1017/S0022112085002221, 1985.

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pierson Jr., W. J. and Moskowitz, L.: A proposed spectral form for fully developed wind seas based on the similarity theory of S. A. Kitaigorodskii, *J. Geophys. Res.*, 69, 5181–5190, doi:10.1029/JZ069i024p05181, 1964.
- Preisendorfer, R. W. and Mobley, C. D.: Albedos and glitter patterns of a wind-roughened sea surface, *J. Phys. Oceanogr.*, 16, 1293–1316, doi:10.1175/1520-0485(1986)016, 1986.
- Schenck Jr., H.: On the focusing of sunlight by ocean waves, *J. Opt. Soc. Am.*, 47, 653–657, doi:10.1364/JOSA.47.000653, 1957.
- Shen, M., Xu, Z., and Yue, D. K. P.: A model for the probability density function of downwelling irradiance under ocean waves, *Opt. Express*, 19, 17528–17538, doi:10.1364/OE.19.017528, 2011.
- Snyder, R. L. and Dera, J.: Wave-induced light field fluctuations in the Sea, *J. Opt. Soc. Am.*, 60, 1072–1079, doi:10.1364/JOSA.60.001072, 1970.
- Sterl, A. and Caires, S.: Climatology, variability and extrema of ocean waves: the Web-based NKMI/ERA-40 wave atlas, *Int. J. Climatol.*, 25, 963–977, doi:10.1002/JOC.1175, 2005.
- Stramska, M. and Dickey, T. D.: Short-term variability of the underwater light field in the oligotrophic ocean in response to surface waves and clouds, *Deep-Sea Res. Pt. I*, 45, 1393–1410, doi:10.1016/S0967-0637(98)00020-X, 1998.
- Stramski, D.: The effect of daylight diffuseness on the focusing of sunlight by sea surface waves, *Oceanologia*, 24, 11–27, 1986.
- Stramski, D. and Dera, J.: On the mechanism for producing flashing light under a wind-disturbed water surface, *Oceanologia*, 25, 5–21, 1988.
- Stramski, D. and Tegowski, J.: Effects of intermittent entrainment of air bubbles by breaking wind waves on ocean reflectance and underwater light field, *J. Geophys. Res.*, 106, 31345–31360, doi:10.1029/2000JC000461, 2001.
- Thorpe, S. A.: Bubble clouds and the dynamics of the upper ocean, *Q. J. Roy. Meteor. Soc.*, 118, 1–22, doi:10.1002/qj.49711850302, 1992.
- Veal, C. J., Carmi, M., Dishon, G., Sharon, Y., Michael, K., Tchernov, D., Hoegh-Guldberg, O., and Fine, M.: Shallow-water wave lensing in coral reefs: a physical and biological case study, *J. Exp. Biol.*, 213, 4304–4312, doi:10.1242/jeb.044941, 2010.
- Walker, R. E.: *Marine Light Field Statistics*, Wiley Series in Pure and Applied Optics, 675 pp., ISBN 0471310468, New York, USA, 1994.
- Weber, V. L.: Coefficient of variation of underwater irradiance fluctuations, *Radiophys. Quantum El.*, 53, 13–27, 2010.

- Xu, Z., Yue, D. K. P., Shen, L., and Voss, K.: Patterns and statistics of in-water polarization under conditions of linear and nonlinear ocean surface waves, *J. Geophys. Res.*, 116, C00H12, doi:10.1029/2011JC007350, 2011.
- 5 You, Y., Stramski, D., Darecki, M., and Kattawar, G. W.: Modeling of wave-induced irradiance fluctuations at the near-surface depths in the ocean: a comparison with measurements, *Appl. Optics*, 49, 1041–1053, doi:10.1364/AO.49.001041, 2010.
- 10 You, Y., Kattawar, G. W., Voss, K. J., Bhandari, P., Wei, J., Lewis, M., Zappa, C. J., and Schultz, H.: Polarized light field under dynamic ocean surfaces: numerical modeling compared with measurements, *J. Geophys. Res.*, 116, C00H05, doi:10.1029/2011JC007278, 2011.
- Zaneveld, J. R. V., Boss, E., and Barnard, A.: Influence of surface waves on measured and modeled irradiance profiles, *Appl. Optics*, 40, 1442–1449, doi:10.1364/AO.40.001442, 2001.
- Zhang, X.: Capillary-gravity and capillary waves generated in a wind tank: observations and theories, *J. Fluid Mech.*, 289, 51–82, doi:10.1017/S0022112095001236, 1995.
- 15 Zhang, X., He, M. X., Yang, Q., and Zeng, K.: Effects of wind on ocean color, IGARSS 2006, Denver, USA, 31 July–4 August, 2006, 4056–4059, doi:10.1109/IGARSS.2006.1040, 2006.
- Zielinski, O., Llinás, O., Oschlies, A., and Reuter, R.: Underwater light field and its effect on one-dimensional ecosystem model at station ESTOC, north of the Canary Islands, *Deep-Sea Res. Pt. II*, 49, 3529–3542, doi:10.1016/S0967-0645(02)00096-6, 2002.
- 20

**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Underwater light field fluctuations**M. Hieronymi and  
A. Macke**Table 1.** Underlying data for radiative transfer simulations.

Wavelength of light	$\lambda$	[nm]	490
Sun zenith angle	$\theta_s$	[°]	0
Refractive index of seawater	$n$	[–]	1.34
Fraction of diffuse sky	Dif	[%]	10
Chlorophyll <i>a</i> concentration	<i>Chl</i>	[mg m <sup>-3</sup> ]	0.1

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 1.** Wave spectra for different local wind conditions that are used for our near-surface underwater light field simulations (short-wave part only); the dashed lines show the modification due to the long-wave part of the unified spectra.

**Underwater light field fluctuations**

M. Hieronymi and  
A. Macke

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion





## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

**Fig. 2.** Sea states under consideration for underwater light field simulations with  $dx = 10$  cm horizontal resolution; left: input parameters (wave height  $H$ , wave period  $T$ , and wind speed  $U_{10}$ ) for the generation of irregular wave profiles; right: probability of the occurrence of wave classes averaged over all seasons and regions (Sterl and Caires, 2005).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







Fig. 3. Wave-induced mixed layer depth  $z_{MLD}$  for the wave classes under consideration (Eq. 2).

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

**Fig. 4.** Spatial distribution of the downwelling irradiance  $E_d$  due to local wind speeds of 3 and  $15 \text{ m s}^{-1}$  that are simulated with the near-surface model with  $dx = 2.5 \text{ mm}$ ,  $E_d$  above the surface is 100 % (logarithmic color scale).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion




## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

**Fig. 5.** Statistics of the underwater light fields from Fig. 4 with wind velocities of  $3 \text{ m s}^{-1}$  (left) and  $15 \text{ m s}^{-1}$  (right); top: probability density function  $PDF$  of downwelling irradiance  $E_d$ ; middle row: number of flashes  $N$  above a certain irradiance threshold  $x_{th}$ ; lower panels: power spectral density  $S_x$  with respect to the normalized irradiance.

## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

**Fig. 6.** The influence of local wind on the underwater light field near the water surface; **(a)** coefficient of variation  $CV$  of  $E_d$  fluctuations; **(b)** significant irradiance enhancement  $\chi_{1/10}$ ; **(c)** mean fluctuation length  $L_m$ ; **(d)** spectral peak length  $L_p$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 7.** Significant irradiance enhancement  $\chi_{1/10}$  for two wind velocities with (dashed) and without (solid lines) long-wave modification of the applied wave spectra (Fig. 1).

**Underwater light field fluctuations**

M. Hieronymi and  
A. Macke

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Underwater light field fluctuations**M. Hieronymi and  
A. Macke

**Fig. 8.** Spatial distribution of downwelling irradiance  $E_d$  for the sea state of highest occurrence probability with a resolution of  $dx = 10$  cm, the 10 and 5% irradiance levels are additionally marked.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

**Fig. 9.** Statistical characteristics of the influence of the sea states (Fig. 2) on the underwater irradiance variability at the 50, 25, 10, and 5 %  $E_d$  depth level ( $z_t = 19.7, 38.5, 62.0,$  and  $79.2$  m, respectively) based on the model with  $dx = 10$  cm resolution; **(a–d)**: coefficient of variation  $CV$ ; **(e–h)**: significant irradiance enhancement  $\chi_{1/10}$ ; **(i–l)**: maximum normalized irradiance  $\chi_{\max}$ ; **(m–p)**: mean fluctuation length  $L_m$ ; **(q–t)**: spectral peak length  $L_p$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion





## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

**Fig. 10.** Summary of light field statistics of all sea states under consideration with the  $dx = 10\text{ cm}$  model; the gray lines show the respective minimum and maximum values, the red lines present the average of the values due to the sea states with highest occurrence probability (in the white framed in Fig. 2); **(a)**: coefficient of variation  $CV$ ; **(b)**: maximum normalized downwelling irradiance  $\chi_{\max}$ ; **(c)**: mean fluctuation length  $L_m$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Underwater light field fluctuations

M. Hieronymi and  
A. Macke

**Fig. 11.** Range of waves that influence the light field at depth; the red lines represent the minimum and maximum wavelengths that contribute at least 0.1 % (one per-mille) to the first spectral moment  $m_1$  of  $S_\chi$  (integral of the wavenumber-weighted variance spectrum of  $\chi$  fluctuations, Eq. 6); top: near-surface model with  $dx = 2.5$  mm; below:  $dx = 10$  cm model resolution.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

