

This discussion paper is/has been under review for the journal Ocean Science (OS).
Please refer to the corresponding final paper in OS if available.

Optimal adjustment of the atmospheric forcing parameters of ocean models using sea surface temperature data assimilation

M. Meinvielle, J.-M. Brankart, P. Brasseur, B. Barnier, R. Dussin, and J. Verron

Laboratoire des Ecoulements Géophysiques et Industriels, CNRS/Université de Grenoble, France

Received: 4 June 2012 – Accepted: 3 July 2012 – Published: 20 July 2012

Correspondence to: M. Meinvielle (marion.meinvielle@legi.grenoble-inp.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

OSD

9, 2493–2533, 2012

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In ocean general circulation models, near surface atmospheric variables used to specify the atmospheric remain one of the main sources of error. The objective of this research is to constrain the surface forcing function of an ocean model by Sea Surface Temperature (SST) data assimilation. For that purpose, a set of corrections for ERAinterim (hereafter ERAi) reanalysis data is estimated for the period from 1989 to 2007 using a sequential assimilation method, with ensemble experiments to evaluate the impact of uncertain atmospheric forcing on the ocean state. The control vector of the assimilation method is extended to atmospheric variables to obtain monthly mean parameter corrections by assimilating monthly SST and Sea Surface Salinity (SSS) climatological data in a low resolution global configuration of the NEMO model. In this context, the careful determination of the prior probability distribution of the parameters is an important matter. This paper demonstrates the importance of isolating the impact of forcing errors in the model to perform relevant ensemble experiments.

The results obtained for every month of the period between 1989 and 2007 show that the estimated parameters produce the same kind of impact on the SST as the analysis itself. The objective is then to evaluate the long term time-series of the forcing parameters focusing on trends and mean error corrections of air-sea fluxes. Our corrections tend to equilibrate the net heat flux balance at the global scale (highly positive in ERAi database), and to remove the potentially unrealistic negative trend (leading to ocean cooling) in the ERAi net heat flux over the whole time period. More specifically in the intertropical band, we reduce the warm bias of ERAi data by mostly modifying the latent heat flux by wind velocity intensification. Consistently, when used to force the model, the corrected parameters lead to a better agreement between the mean SST produced by the model and mean SST observations over 1989–2007 in the intertropical band.

OSD

9, 2493–2533, 2012

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Near surface variables from atmospheric reanalyses (air temperature and humidity, wind velocity, downward radiation and precipitation) are commonly used to specify surface boundary condition in Ocean General Circulation Models (hereafter OGCMs) required for operational forecasts, ocean reanalyses, or hindcast simulations of the recent ocean variability (the last 50 years). However, these atmospheric variables are characterized by significant uncertainties at global scale as shown in different studies like Milliff et al. (1999), Wang and McPhaden (2001), Smith et al. (2001) or Sun et al. (2003). For example, the use of two different databases (e.g. from numerical weather prediction centers like ECMWF or NCEP) to compute the mean ocean-atmosphere net heat flux can lead to discrepancies of the order of at least 10 W m^{-2} , while the signal of a global warming of the world ocean corresponds to a value of 0.5 W m^{-2} Josey (2011). It is therefore important to reduce the atmospheric variables uncertainties in order to obtain better agreement between the models and the real ocean. Sea Surface Temperature (SST) is more accurately observed from space than most near surface atmospheric variables or air-sea fluxes assimilated in atmospheric models to construct the reanalyses. However, this observed variable, intrinsically linked to air-sea exchanges, is not involved in the boundary condition of OGCMs except when explicitly assimilated. In brief, models do not benefit in their forcing from one of the best observed ocean surface variables.

One of the approaches to incorporate SST information into ocean simulations consists in assimilating observed SST products to correct the model state. This method can lead however to some inconsistency between the “assimilated” solution of the model and the “forced” one, and to the rejection of the information contained in the SST. Since atmospheric forcing and particularly atmospheric variables are an important source of surface errors, an alternative is to use SST data assimilation to constrain the surface atmospheric input variables of the atmospheric reanalyses. The aim is to correct the source of errors and not the consequence. An improvement of fluxes estimation is

OSD

9, 2493–2533, 2012

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



crucial to perform realistic ocean simulations and can also help improving the atmospheric reanalyses since their atmospheric state estimation could benefit from the link with the ocean dynamics via the ocean model. Moreover, this approach proposes an alternative to other corrections of atmospheric forcing realized following ad-hoc considerations (e.g. Large and Yeager, 2004; Brodeau et al., 2010).

This idea has been explored in previous studies using two different types of data assimilation schemes. On the one hand, Stammer et al. (2004) used a four dimensional variational data assimilation scheme by including air-sea fluxes in the control vector. Over a ten-year period, they assimilated a large variety of oceanic observations, computed air-sea flux corrections, and carried out an important validation effort to compare the corrected fluxes to other independent estimates. This pioneering work has been the first demonstration that it is possible to make estimates of the atmospheric forcing fields by ocean data assimilation in a realistic case, even if the authors also show that the forcing correction may sometimes compensate for other errors in the ocean model. On the other hand, fluxes corrections can also be computed using sequential assimilation methods. These methods are widely used in ocean data assimilation systems, and are easier to implement than their variational counterpart. Skachko et al. (2009) and Skandrani et al. (2009) explored the capability of such methods to estimate forcing parameters corrections by using ocean observations data. On the basis of a reduced order Kalman filter, they carried out idealized experiments that differ from the parameters included in the control vector, the construction of the synthetic observations to be assimilated, and the construction of initial condition. They showed that in an idealized case where errors are assumed to be entirely due to forcing parameters, it is possible to implement a sequential data assimilation method to estimate objective corrections of these parameters. While Skachko et al. (2009) used temperature and salinity profiles that simulate the horizontal and temporal distribution of Argo floats in twin experiments, Skandrani et al. (2009) assimilated Sea Surface Temperature and Sea Surface Salinity extracted from a Mercator-Ocean reanalysis (Ferry et al., 2010). The first study by Skachko et al. (2009) showed that this procedure leads to accurate

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



estimations of parameters included in the control vector. And following these first results, Skandrani et al. (2009) in a more realistic context, estimated forcing parameters corrections leading to reduced differences between the free simulation and the reanalysis. However, these two studies also suggest that the other sources of model errors (due to the coarse resolution or to the initial condition for example) can lead to unrealistic forcing parameters corrections and must be considered very carefully. In particular, they point out the importance of a proper determination of the prior probability distribution of forcing parameters and of their associated error covariance matrix to obtain good parameters estimation. These results are thus an appropriate starting point for further developments of this approach, investigating its feasibility in a more realistic context. In this case the sources of errors are more diverse and require a particular attention.

The purpose of this research is thus to constrain (within observation-based air-sea flux uncertainties) the surface forcing function of an ocean model (i.e. surface atmospheric input variables from atmospheric reanalyses) by using a methodology based on advanced statistical assimilation methods (Ensemble Kalman Filter) to take into account SST satellite observations. In other words, the objective of this work is to take advantage of an ocean model to correct near surface atmospheric variables, and to ensure their consistency with ocean surface dynamics. With respect to the studies briefly described before, this work presents the originality to be carried out for longer timescale, and with real SST observations assimilated in realistic global ocean model simulations.

In the present paper, we make one step further than the past idealized studies to estimate a set of corrections for the atmospheric input data from the ERAinterim (ERAi hereafter) reanalysis for the period from 1989 to 2007. We use a sequential method based on the SEEK filter. Over experiments of one month duration, observed monthly SST product (Hurrell et al., 2008) and SSS seasonal climatology data (Levitus et al., 1998) are assimilated to obtain monthly corrections of the atmospheric forcing parameters (air temperature and humidity, zonal and meridional wind speed, and downward

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



radiation). The expected outcome of our experiment is thus to obtain a comprehensive set of estimated parameters for every month of the period between 1989 and 2007, so that they can be used later in a free model simulation.

We will first describe in Sect. 2 the forcing function of the model and the Kalman filter methodology used as a basis for this work. Section 3 concerns the principal adjustments necessary to fit our realistic context and the long term focus of this study. Finally, a selection of results that illustrate the strengths and weaknesses of the method will be presented in Sect. 4.

2 Estimation method

2.1 Ocean model and forcing by the atmosphere

The OGCM is the first key ingredient to estimate relevant parameters corrections, and to evaluate their validity and their impact in free (i.e. without data assimilation) model runs. Since the estimation problem requires a large number of model simulations (to perform the ensemble experiments), and since the goal is to explore the problem at a global scale, we chose to use a coarse resolution OGCM: the 2° resolution configuration of NEMO (Madec, 2008), with 46 vertical levels. In previous studies, Skachko et al. (2009) and Skandrani et al. (2009) chose this coarse resolution approach to set up their idealized experiments before considering applying it to higher resolution configurations. Implementing this kind of methodology with high resolution models would indeed be numerically too expensive to be considered as a first step.

Different methods exist to force an ocean-only model. Here the forcing function is computed in the model using global aerodynamic formulation (bulk) as described by Large et al. (1997). All fluxes are calculated for each model grid point from near-surface atmospheric variables (air temperature and humidity, zonal and meridional wind speed and precipitation), downward fluxes (short wave and long wave downward radiation)

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and the ocean surface variables calculated by the model (SST and surface currents velocity).

The components of the forcing function are the freshwater flux (F_w), the momentum flux (τ), and the net heat flux (Q_{net}) which is compiled as the sum of the short wave solar radiation flux (Q_{sw}), the long wave radiation flux (Q_{lw}), the sensible heat flux (Q_{sens}) and the latent heat flux (Q_{lat}).

Air sea fluxes estimation in the model requires the SST of the model (T_s), the atmospheric state, the downward radiation, and the precipitation. Atmospheric state and downward radiation are involved in the fluxes computation as follows:

- The turbulent fluxes involve the knowledge of $\mathbf{X_A} = (\theta_a, q_a, U_{10})$ the atmospheric state, with θ_a being the potential temperature, q_a the specific humidity and U_{10} the relative wind speed:

$$\begin{aligned} Q_{\text{sens}} &= \rho_a C_H(T_s, \mathbf{X_A}) C_p |\Delta U_{10}| [\theta_a - T_s] \\ Q_{\text{lat}} &= \rho_a C_E(T_s, \mathbf{X_A}) L_v |\Delta U_{10}| [q_a - q_{\text{sat}}(T_s)] \\ \tau &= \rho_a C_D(T_a, \mathbf{X_A}) |\Delta U_{10}| \Delta U_{10} \\ E &= -Q_{\text{lat}}/L_{\text{vap}} \end{aligned} \quad (1)$$

where ρ_a is the air density considered as a constant of 1.29 kg m^{-3} , C_p the air specific heat ($\sim 1000 \text{ J kg}^{-1} \text{ K}^{-1}$), L_v the latent heat ($2.26 \times 10^6 \text{ J kg}^{-1}$), and q_{sat} the saturation specific humidity.

- The radiative fluxes involve the knowledge of $\mathbf{X_R} = (\text{rad}_{\text{sw}}, \text{rad}_{\text{lw}})$ the downward radiation, with rad_{sw} being the downward shortwave radiation, and rad_{lw} the downward longwave radiation:

$$\begin{aligned} Q_{\text{sw}} &= (1 - \alpha) \text{rad}_{\text{sw}} \\ Q_{\text{lw}} &= \text{rad}_{\text{lw}} - \epsilon \sigma T_s^4 \end{aligned} \quad (2)$$

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where α is the ocean surface albedo (~ 0.066), $-\epsilon\sigma T_s^4$ is the infrared radiation flux from the ocean surface with σ the Stefan–Boltzmann constant (5.67×10^{-8}), and ϵ the seawater emissivity (0.98).

- The freshwater flux involves the knowledge of precipitation (precip):

$$F_w = -E + \text{precip} + R \quad (3)$$

with R the continental contribution to the freshwater budget.

Near surface variables (air temperature and humidity, wind velocity, downward radiation and precipitation) from atmospheric reanalysis (here ERAi) used to specify surface boundary condition of the model are characterized by large uncertainties at global scale. Temperature and humidity are usually available 2 m above the ocean surface and will be mentioned as t_2 and q_2 , respectively thereafter. Zonal and meridional wind speed, available 10 m above the surface will be noted u_{10} and v_{10} . We thus propose here to estimate corrections of t_2 , q_2 , u_{10} , v_{10} , rad_{sw} , rad_{lw} and precip.

C_H , C_E and C_D are the bulk transfer coefficients for sensible heat, humidity and momentum, respectively. Several parameterizations exist to compute these coefficients. All of them present some uncertainties but the one used in our ocean model NEMO (hereafter LY04) is the one described by Large and Yeager (2004). Even if they are impacted by forcing modifications, we chose to focus here on the atmospheric variables corrections, instead of following the approach of Skachko et al. (2009) and Skandrani et al. (2009) used in previous work, because as shown in Fig. 1, the contribution of atmospheric variable error on the net heat flux uncertainty is more important than the contribution of bulk coefficient uncertainty Brodeau (2007).

2.2 Kalman filtering for parameter estimation

In the standard current state of the art of ocean data assimilation systems, sequential data assimilation methods are most often used to correct the model state (e.g.

Drévilion et al., 2008). However, these traditional implementations do not actually correct the source of errors but the consequence and can lead to possible inconsistencies between the assimilated solution and the forcing. The method developed here proposes an alternative to this classical approach by estimating corrections of a source of model errors: the forcing parameters. We use here the same methodology (based on a Kalman filter analysis) as developed by Skandrani et al. (2009) to estimate atmospheric forcing parameter corrections. With respect to the classical formulation of the Kalman filter (Kalman, 1960), the control vector containing initially the model state is extended to the forcing parameters that we want to estimate (Eq. 4). The control vector becomes:

$$\hat{\mathbf{x}} = \begin{bmatrix} [\mathbf{x}] \\ [\mathbf{p}] \end{bmatrix} \text{ avec } [\mathbf{p}] = \begin{bmatrix} t_2 \\ q_2 \\ u_{10} \\ v_{10} \\ \text{rad}_{\text{sw}} \\ \text{rad}_{\text{lw}} \\ \text{precip} \end{bmatrix} \quad (4)$$

with $\mathbf{p}$ the parameters vector that we aim to correct, where t_2 is the 2 m air temperature, q_2 the 2 m air humidity, u_{10} the 10 m zonal wind velocity, v_{10} the 10 m meridional wind velocity, rad_{sw} the downward shortwave radiation, rad_{lw} the downward longwave radiation, and precip the precipitation.

The background control vector $\hat{\mathbf{x}}^f$ is the model output from a simulation forced by the first guess atmospheric parameters over the assimilation window extended to these atmospheric parameters. We can then obtain the best estimate $\hat{\mathbf{x}}^a$ of the control vector by taking into account available observations vector $\mathbf{y}$:

$$\hat{\mathbf{x}}^a = \hat{\mathbf{x}}^f + \hat{\mathbf{K}}(\mathbf{y} - \hat{\mathbf{H}}\hat{\mathbf{x}}^f) \quad (5)$$

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with $\hat{\mathbf{x}}^f$ the background simulation, $\hat{\mathbf{H}}$ the observation operator, and $\hat{\mathbf{K}}$ the Kalman gain given by:

$$\hat{\mathbf{K}} = \hat{\mathbf{P}}^f \hat{\mathbf{H}}^T (\hat{\mathbf{H}} \hat{\mathbf{P}}^f \hat{\mathbf{H}}^T + \mathbf{R})^{-1} \quad (6)$$

with $\mathbf{R}$ the observation error, and $\hat{\mathbf{P}}^f$ the forecast error covariance matrix.

The formulation of the Kalman filter applied to an extended state vector needs to know the forecast error covariance matrix $\hat{\mathbf{P}}^f$ in the augmented space, given by:

$$\hat{\mathbf{P}}^f = \hat{\mathbf{M}} \hat{\mathbf{P}}^a \hat{\mathbf{M}}^T + \hat{\mathbf{Q}} \quad (7)$$

with $\hat{\mathbf{P}}^a$ the background error covariance matrix, or initialisation error, $\hat{\mathbf{M}}$ the model operator, and $\hat{\mathbf{Q}}$ the model error.

The knowledge of the $\hat{\mathbf{P}}^f$ matrix (and thus $\hat{\mathbf{P}}^a$ and $\hat{\mathbf{Q}}$) is crucial in statistical estimation approaches. As already pointed out by Skachko et al. (2009) the impact of the background error on the model results is even more important when the model state correction is not applied. In their more realistic study, Skandrani et al. (2009) also highlight the fact that making appropriate assumptions about the forecast error statistics is increasingly important as the problem becomes more realistic.

However, the objective is here different from these two previous studies, as it focuses on a mean long-term correction of the forcing parameters. The aim is not to correct the model state and improve the short-term forecast of the model, but to estimate long term atmospheric parameter corrections to be used in another independent free simulation. For this reason, we decided to use a different strategy and to carry out an independent ensemble experiment for each assimilation window instead of running a classical chain of analysis cycles. This process simplifies the estimation of the forcing error covariance matrix (described in Sect. 3). Moreover, under the assumption that all the forecast error is due to forcing error and that the forcing correction is ideal, the background error covariance matrix for the next assimilation window is assumed very

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



small. In practice, if we consider that the initialization state of each experiment is close enough to assimilated observations, we can then neglect the initial error covariance error and set $\hat{\mathbf{P}}^a$ to zero in the expression of $\hat{\mathbf{P}}^f$ (Eq. 7).

The expression of the $\hat{\mathbf{P}}^f$ matrix will become:

$$\hat{\mathbf{P}}^f = \hat{\mathbf{Q}} \quad (8)$$

In this approximation, the estimation of $\hat{\mathbf{P}}^f$ involves only the knowledge of the model error $\hat{\mathbf{Q}}$ generated by forcing errors during assimilation window. As this matrix must be representative of forcing parameters errors only, we have to reduce all the other error sources in the ensemble experiments to avoid compensation effects in the parameters estimation. In this context, the main difficulty is to preserve the robustness of the methodology previously developed. Contrary to Skachko et al. (2009) and Skandrani et al. (2009), the observation errors need to be taken into account. We do not know the true value of forcing parameters, and neither the perfect initial condition that will ensure the efficiency of assimilation experiments. We also have to make the distinction between forcing errors and other potential errors in the system to obtain the most versatile correction possible. Indeed, the solution will depend on the model used and it could be difficult to apply the same parameter corrections to other configurations where the nature of the model error is different. All these matters will be addressed to evaluate the relevance of the method and results in Sect. 4.1.

Finally, two additional refinements of the forecast error statistics will be used in the experiments. First, to avoid spurious long range influence of the observations, the ensemble covariance matrix will be localized as described in Brankart et al. (2011), using a horizontal cutting length scale of three grid points (i.e. about 600 km in longitude along the equator). Second, to avoid excessive and non-physical parameters corrections, the forecast probability distribution will be assumed to be a truncated Gaussian probability distribution, as developed by Lauvernet et al. (2009), and as already used in Skandrani et al. (2009). In practice, all parameter corrections above 1.5 standard deviations will be truncated.

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3 Challenge of a realistic application

The first challenge of a realistic application of this method is that available observations of the real ocean are sparse in space and time. We chose to assimilate monthly SST data, that is one of the most accurate observations of the ocean, and SSS climatological data to constrain the freshwater budget of the system, and thus to limit unrealistic impact of the correction on the SSS. We use the Hurrell database (Hurrell et al., 2008) giving SST monthly mean between 1989 and 2007, and a climatology of monthly SSS for the same period from (Levitus et al., 1998). Since the available information is now more limited than in idealized cases, this application involves some crucial adjustments of the methodology (see Sect. 3).

The time resolution of available data implies to consider assimilation windows of one month, and thus monthly parameters corrections. However we do not have access to the corresponding subsurface mean state to construct the ideal initial condition in order to ensure the validity of the assumption of $\hat{\mathbf{P}}^a$ equal to zero. For our application, this will be the first adaptation to make to the initial methodology.

With these observations, we will compute corrections to the atmospheric parameters from the ERAi reanalysis available between 1989 and 2007. ERAi is the most recent reanalysis produced by the European Center for Meteorological and Weather Forecasting (ECMWF). The atmospheric variables are available every 6 h for $\mathbf{X}_A$ and every day for $\mathbf{X}_R$ and precipitation at the same spatial resolution as the model, but we do not have access to the associated uncertainties. The estimation of the prior probability distribution of the forcing parameters is the second difficulty of a realistic experiment. The monthly uncertainties associated to the parameters have to be characterized. Finally, to run Monte-Carlo experiments to estimate the matrix $\hat{\mathbf{Q}}$, the forcing error in the model has to be isolated to avoid potential compensation effects.

In practice, the whole procedure is repeated independently for each month of the period between 1989 and 2007, which represents an important amount of simulations and could be a critical difficulty with a higher resolution model.

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Error statistics

The forecast error covariance matrix $\hat{\mathbf{P}}^f$ reflects the impact of the forcing parameters errors on the model state. One way to estimate this matrix is to run Monte-Carlo experiments, or ensemble experiments, as performed by Skachko et al. (2009) and Skandrani et al. (2009). For each month, we run 200 experiments from the same initial condition. These experiments only differ by the atmospheric parameters used to force the model. $\hat{\mathbf{P}}^f$ is then deduced from a statistical analysis of the 200 forecast states ensemble. However, to be consistent with prior assumptions needed to apply the filter methodology, the ensemble experiments have to be representative of the impact of the forcing parameters error only. We need thus to take care of three constraints that represent the most important methodological developments necessary for a realistic application, and that will be described in the following sections:

- Minimization of the model errors that are not due to the forcing function by a robust diagnostic approach;
- Minimization of the initial condition errors;
- Computation of parameter perturbations representative of the forcing uncertainties.

3.1 Robust diagnostic ensemble experiments

The 200 model states ensemble have to be representative of the impact of forcing errors on monthly time scale ocean dynamics, and particularly on the surface where observations are assimilated. To ensure the consistency of this assumption, other possible errors have to be minimized while performing ensemble experiments. Indeed, the coarse resolution model used here involves parametrization and approximations to compensate unresolved processes. It thus contains errors that are not attributable to forcing, which lead in particular to a bad positioning of the water masses in strong

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



advection regions such as western boundary currents. If this type of error is present in the ensemble runs, it will lead to an inconsistent estimation of $\hat{\mathbf{P}}^f$, and finally to unphysical parameters compensating the non-forcing errors. As a consequence, ensemble runs have to be as close as possible to the real ocean, and be representative of the forcing errors only.

We chose to constrain the model by using a robust diagnostic approach (Sarmiento and Bryan, 1982) consisting in a three dimensional relaxation of the solution to temperature and salinity climatologies (Levitus et al., 1998). Relaxation terms are added to heat, mass and salt conservation equations and create artificial sources and sinks to bring back the simulation to the corresponding climatologies. The relaxation strength is adapted through its time constant. A large value of this time constant corresponds to a light relaxation, and a little one to a strong relaxation. In our case, we adjust the relaxation time constant to allow the model to react to atmospheric forcing without getting to far from the climatology. Two domains are thus specified where the relaxation characteristics are different:

- The first 400 m from the surface are likely to be impacted by a forcing modification at monthly time scale. The time constant of the relaxation is one year in this domain to constrain the large scale feature while allowing possible impact of the atmospheric fluxes on the dynamics.
- Below 400 m depth, the relaxation applied is strong with a one-month relaxation constant. The large scale feature is thus strongly constrained at depth to ensure a good consistency with the observed ocean and limit non-forcing error sources.

The robust diagnostic approach is a tool used here to produce a density field consistent with an incomplete set of observations. The purpose is to help the identification of a particular set of atmospheric parameters and not to produce a relevant description of ocean circulation. In the constrained simulations, the large scale water masses positioning in the model solution (e.g. the structure of gyres) is more consistent with observed ocean (figure not shown). The first use of the method was to evaluate the

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ocean response to a given atmospheric CO₂ field (Sarmiento and Bryan, 1982). This first approach presents some similarities with our objective to identify the impact on the ocean surface of a given atmosphere.

A robust diagnostic simulation introduces artificial sources and sinks of heat and freshwater in the conservation equations and can lead to an unrealistic representation of some processes. However, since the relaxation is lighter in the region directly affected by atmospheric forcing, the relevance of our approach is still valid in this special zone of interest. This three dimensional relaxation ensures a limited effect of non-forcing errors in the ensemble experiments, so that the $\hat{\mathbf{P}}^f$ matrix is only representative of the errors in the atmospheric parameters. The validity of this hypothesis is crucial to estimate consistent parameter corrections and to reduce possible compensation effects.

3.2 Initialization procedure

The second assumption of this study is that the initial error for every month (the $\hat{\mathbf{P}}^a$ matrix in Eq. 7) can be neglected. If $\hat{\mathbf{P}}^a$ is non-negligible, then the final equation of $\hat{\mathbf{P}}^f$ (Eq. 8) cannot be considered valid. In absolute terms, to ensure that $\hat{\mathbf{P}}^a$ is equal to zero we should initialize ensemble experiments with the true ocean state. However this true three dimensional ocean state is not available. To obtain a reasonable approximation of an initial condition with the correct SST, a simulation over the 1989–2007 period strongly constrained to fit surface observation is constructed. From this simulation we then extract the initial condition that will be used for ensemble experiments. To carry out this simulation, a strong relaxation is applied to the surface in addition to the robust diagnostic approach described in the previous section. The surface relaxation is prescribed with a time constant of two hours to deprive the model of surface evolution freedom. However the objective is to construct initial condition for the ensemble experiments. This particular initial condition construction is the best compromise that we have found to initialize the ensemble experiments, and it is an important additional

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



procedure allowing to carry out a realistic application of the methodology developed by Skandrani et al. (2009).

For each month the corresponding initial condition is extracted from the strongly constrained simulation. It is then used to initialize ensemble experiments. To build consistent parameters corrections, the background simulation of one month that gives the background control vector $\hat{x}^f$ (Eq. 5) has to be initialized with the same ideal conditions.

3.3 Parameter perturbations computation

The last important assessment to make concerning the ensemble experiments set up is to ensure that the perturbations applied to atmospheric parameters are actually representative of the real forcing uncertainties. It is assumed that parameters uncertainties are comparable to their intra-seasonal and interannual variability in the ERAi reanalysis. The parameters prior probability distribution is defined as Gaussian (strictly speaking, a truncated Gaussian to avoid extreme parameter corrections, see Sect. 2.2) and its covariance matrix is derived from the ERAi reanalysis intra-seasonal and interannual variability between 1989 and 2007. Perturbations are specific to a given month and constant over each monthly assimilation window. For example, to estimate a set of perturbations for April, we consider the reanalysis signal corresponding to all three month windows March-April-May of the reanalysis over the 1989–2007 period (60 states). This assumption is made to characterize the covariance of atmospheric parameters. The choice of intra-seasonal and interannual variability rather than the total variability is made to avoid in the perturbations of April for example, a variability that is characteristic of winter time. A random sample of 200 perturbations is then constructed for each month.

Figure 2 illustrates the standard deviation of the parameter perturbations for a given month ensemble (here January). To ensure relevant $\hat{P}^f$ estimation, this amplitude has to be representative of the uncertainty of each atmospheric variable. One can see that the standard deviation associated to each parameter corresponds to monthly

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



parameter uncertainties at the global scale, with values in the range of 0 to 1.5°C for the temperature, 10 to 25 Wm⁻² for the downward shortwave radiation, and 0.5 to 3 ms⁻¹ for the zonal and meridional wind velocities. However, at high latitudes, the ensemble dispersion of temperature and shortwave radiation is larger (up to 6°C for the temperature and 70 Wm⁻² for the shortwave radiation). These values are clearly excessive to represent the real uncertainty, which illustrates the limits of considering the intra-seasonal and interannual variability of parameters to quantify the uncertainties. The assumption made to generate the parameter perturbations in the ensemble experiments could thus be irrelevant in high latitudes regions because changes from one month to another in the three month period can be much larger than the uncertainties. As a consequence, the results in these regions must be interpreted cautiously.

4 Results

The objective of this section is to analyze the strength and weaknesses of the methodology that has been adapted to correct ERAi variables. The relevance of specific assumptions described above is evaluated by selecting some diagnostic at global and regional scales. As a first step, we will focus on the capability of the method to isolate the forcing impact in the model and compute realistic parameter corrections. Second, we will look at the results in terms of global scale heat budget resulting from our corrections. And third, we will discuss the impact of the computed corrections in the intertropical band where the method presents the maximum reliability.

4.1 Assessment of the method

The first important step of the method validation is to look at the results over a specific month (January 2004) by assimilation window. We compare (Fig. 3, top panel) the SST resulting from the analysis step of the Kalman filter (i.e. the output of Eq. 5), and (Fig. 3, bottom panel) the SST computed by a free model run forced by the newly estimated

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



parameters (hereafter ERAcor). This diagnostic will determine if the use of corrected parameters in a free simulation has the same impact as the correction computed by the analysis scheme. In other words, the objective here is to evaluate if we have properly determined the $\hat{\mathbf{P}}^f$ matrix which describes the impact of forcing errors in the model.

Figure 3 compares the increment of the analysis step to the increment produced in a free run model forced by ERAcor atmospheric variables. In both cases, the resulting SST shows a better agreement with its observed counterpart (not shown). Furthermore, one can see that both increments are very similar in structure and intensity. This result means that the error covariance matrix $\hat{\mathbf{P}}^f$ constructed here is consistent with the actual impact of forcing errors in the model.

The second step to evaluate the reliability of our method to improve the forcing for free model simulations is to look at the consistence of the amplitude of parameter corrections estimated by the method with the uncertainties on atmospheric variables. Since the focus is here to correct potential long-term trends and mean error, it is of primary importance to assess the regional relevance of the corrections temporal mean between 1989 and 2007 (Fig. 4).

In Fig. 4, we notice that for the major part of the world ocean the corrections are consistent with the assumed forcing uncertainties (e.g. -1 to 1°C for the temperature, 1.5 to 1.5 ms^{-1} for wind velocities, or -20 to 20 W m^{-2} for shortwave downward radiation). Furthermore, temperature and humidity corrections have a similar behavior, consistently with the strong correlation of these two variables. The same comment can be made regarding shortwave and longwave radiation corrections. On the other hand, the large scale structure of the corrections is mostly zonal. For example, we obtain a mean shortwave radiation (rad_{sw}) correction of approximately $+15\text{ W m}^{-2}$ in the intertropical band, and -15 W m^{-2} for mid-latitudes. Longwave radiation corrections have an opposite behavior with negative corrections in the intertropical band (-5 W m^{-2}) and positive corrections at mid-latitudes (5 W m^{-2}). The corrections obtained with our method are thus physically reasonable in terms of large scale intensity and structure.

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



However, a critical analysis of Fig. 4 allows identifying some persisting problems. These correction fields present some small scale features (especially at high latitudes), which are not appropriate in forcing corrections. These structures are particularly obvious in the Southern Ocean in the temperature, shortwave radiation and wind speed corrections, and are likely to be the result of the local implementation of the analysis. These small scale structures make difficult to interpret the corrections obtained in the Southern Ocean, and highlight one limit of the use of local data assimilation. The same problem has already been evidenced by Stammer et al. (2004) in a variational approach. Small scale structures in the Southern Ocean lead to important gradients in the correction and can result in instability propagation when applied to a free model run. However critical gradients of 20 to 40 W m^{-2} for the shortwave radiation in this region are not only explained by the localized analysis. Indeed, we noticed in Sect. 3.3 that the assumption made to compute parameters perturbations (and thus parameter prior probability distribution) was not relevant at high latitudes, which can lead to spurious corrections. These spurious corrections have obviously not been totally eliminated by the truncation of the prior probability distribution mentioned in Sect. 2. The combination of this problem with the covariance localization hampers the relevance of the correction in the high latitudes.

Figure 4 also allows looking more precisely at the results in strong advection regions such as the Gulf Stream, the Kuroshio region, or the Agulhas Current region but also the Antarctic Circumpolar Current region. As mentioned in Sect. 3.1, in these regions, using a coarse resolution model often leads to a shifted positioning of the water masses and fronts. Despite of the use of a robust diagnostic approach to compute ensemble experiments, these non-forcing errors are still present in the model, with the consequence that estimated corrections in these regions are often excessive (with for example orders of magnitude of 4°C for the temperature, 3 ms^{-1} for the zonal wind speed, and up to 60 W m^{-2} for the shortwave radiation). These values are typical of compensation effects. We have here an over-estimation of parameter modifications compensating for other model errors. Considering that larger observation errors are not sufficient to

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



bypass this problem, we cannot be very confident in the realism of estimates made in strong advection regions.

Finally, we observe that the corrections computed for the precipitation field are negligible (figure not shown). This is consistent with the fact that the SSS assimilated aims to avoid unbalanced corrections, and not to actually generate significant precipitation corrections.

In summary, we have identified in this section the limits of the method implemented to compute upgraded atmospheric variables. And the conclusion is that even if the results look rather consistent with parameter uncertainties in the major part of the world ocean, we have to be cautious with regional corrections computed in the high latitudes and strong advection regions. However these problems remain quite local and we can still look at the global impact of the correction on the ocean heat budget.

4.2 Global results

To study this global impact we have to compute air-sea fluxes corresponding to this new forcing parameters dataset using the bulk formulas. The objective here is to evaluate this new dataset reliability to force a free model run. The tool used to compute air-sea fluxes offline, using the same formulae as the ocean model and a given atmospheric dataset, was developed by Brodeau (2007). Instead of using the model SST, this diagnostic involves a prescribed SST (here the Hurrell SST). Inter-comparison between different atmospheric dataset to quantify the parameter modifications impact is thus easier than in the model where the SST feedback on the flux computation is taken into account. All fluxes presented in the following are computed using this offline approach. Global results concerning the net heat flux will be presented first, before focusing on the intertropical band to quantify the distribution of this integrated modification over the different net heat flux components.

The net heat flux integrates all modifications applied to the latent, sensible and radiative fluxes. The diagnostic of the parameter corrections impact on this flux thus reflects the global effect of forcing modifications. The first criterion used to evaluate a forcing

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dataset is to look at the global heat balance over the considered period (here 1989 to 2007). A balanced budget is a sign of quality for the given forcing. Figure 5 shows that the corrections modify the net heat flux budget from a positive value of almost 15 W m^{-2} (left) to approximately 2 W m^{-2} (right). This result means that forcing the model with ERAi atmospheric variables would lead to an excessive heating of the ocean, while our corrections reduce drastically this heat excess and maintain the heat budget close to equilibrium.

Furthermore, a clear negative trend is present in the ERAi time-series of the net heat flux. This trend is inconsistent with the observed global warming for the last twenty years (Bindoff et al., 2007). Moreover, this negative trend is not present any more in the time-series of the net heat flux resulting from the corrected parameters. This diagnostic leads to two crucial results since the methodology does not involve any explicit constraint to obtain a balanced heat budget or to remove potential unrealistic trends.

4.3 Intertropical band

Results commented in Sect. 4.1 allowed to identify regions where reliability of the method is questionable. To evaluate locally the impact of computed corrections on each net heat flux component, we chose now to focus on the region between 20N and 20S, called intertropical band. This region has the advantage not to be affected by spurious corrections or biases in the model circulation mentioned in Sect. 4.1. Furthermore the model used here has 0.5° meridional resolution along the equator and thus captures the critical scales of the equatorial current system (Madec and Imbard, 1996). As a consequence we expect that the results there reach the maximum reliability. It is thus interesting to go further from the net heat flux diagnostic and to look at the impact of parameter corrections on the sensible, latent, longwave radiation and shortwave radiation fluxes.

To summarize the main parameter modifications in this region, the results presented in Sect. 4.1 (Fig. 4) evidenced the following mean corrections:

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Reduced temperature and humidity of 0 to 1 °C and 0 to 1 gkg⁻¹, respectively,
- Increased resultant wind speed of 0.5 to 1.5 ms⁻¹ (Fig. 7),
- Increased downward shortwave radiation of 5 to 10 Wm⁻²,
- Reduced downward longwave radiation of 5 to 10 Wm⁻².

Figure 6 illustrates the impact of these modifications on the mean of the different net heat flux components over the period 1989–2007. As expected, given the reduced global heat budget, we observe a reduction of the net heat flux of about 20 Wm⁻². This tends to reduce the warm bias in the intertropical band that can be evidenced by comparing the model SST to the observations (see Fig. 9 described later). Furthermore, since we compute corrections separately for every atmospheric variable, we can access to valuable information regarding the relative contribution of each component of the net heat flux to this reduction of the net heat flux. From this diagnostic, it appears that to reduce heat gain in the intertropical band, the radiative flux ($Q_{sw} + Q_{lw}$) remains almost the same (augmentation about 5 Wm⁻²), while the heat loss through turbulent fluxes is clearly increased. As a consequence, the net heat flux decrease is mostly due to a modification of the turbulent fluxes. All variable corrections in the intertropical band contribute together to an increased heat loss by turbulent fluxes (decreased temperature and humidity, and increased wind speed).

Among these contributions, wind speed is identified by our method to be the essential mechanism involved in the increase of the turbulent heat loss (Fig. 7).

4.4 Long term free model simulation

The last diagnostic used here to assess the method is the analysis of a free model run forced by the corrected parameters. We showed in the previous sections that our corrections present some unreliable characteristics that could propagate in the model. Parameter corrections are thus post-processed before being used in the model, in

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



order to smooth unrealistic small scale features, and to avoid spurious corrections to be applied in the model.

To smooth small scale features, an anisotropic boxcar filter is applied to each monthly correction. The box size is 10° in latitude and 20° in longitude in order to keep the zonal signal structure while limiting local gradients. The large scale correction characteristics thus persist but the parasite structures observed for example in the Southern Ocean are smoothed out as shown in Fig. 8. Since spurious corrections are still present at high latitudes, the corrections are set to 0 for latitudes larger than 60° in both hemispheres because the error is considered to be mostly due to other processes than forcing (essentially the fact that the method does not take into account the ice cover in these regions). Resulting 1989–2007 mean corrections are shown in Fig. 8.

Even if this procedure does not allow to test the corrections actually computed by the data assimilation method, we can evaluate the corrections in terms of impact in the model focusing on the results in the intertropical band where this post-processing has not a drastic impact on the solution since it is the region of correction maximum validity.

In Sect. 4.3 we had shown that the integrated impact of our parameter corrections is to reduce the net heat flux in the intertropical band. A reduced SST in the free model run is thus expected when replacing ERAi forcing by corrected parameters. In Fig. 9, we first observe that the simulation forced by ERAi variables presents a long term mean warm bias ranging between 0.5 and 2°C in the intertropical band (top). Reducing this warm bias to reach values below 0.5°C for most of the region (consistently with a reduced net heat flux), our correction thus leads to a better agreement with observed SST from a long-term point of view (bottom). Moreover, the standard deviation of the monthly differences between observed and simulated SST is also reduced by up to 0.2°C on average in the intertropical band (figure not shown). The strategy developed here to compute independent monthly corrections is thus consistent with our initial objective to improve the long term mean forcing.

Besides the consistence between observed and simulated SST in the intertropical band, the interannual variability of the model response is not modified by the forcing

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

corrections as shown in Fig. 10. This figure shows the interannual variability of the global means of the SST, the Sea Surface Height (Fig. 10, top panel), and the Meridional Overturning Cell intensity at 30° N (Fig. 10, bottom panel). It is clear that despite the discrepancies between the mean values, the year to year variations are in a very good agreement. This result ensure that the corrections applied do not hampers the interannual variability of the initial ERAi forcing.

5 Conclusions

In this study the objective has been to explore the feasibility of a sequential data assimilation methodology to estimate monthly corrections of ERAi forcing parameters using real SST observations. Independent monthly data assimilation experiments have been performed over the period 1989–2007 to compute a corrected forcing dataset ERAcor (without correcting the model state) to be used in a free model run. The prior probability distribution of the parameters has been characterized by the intra-seasonal and interannual variability of ERAi reanalysis, and ensemble experiments of 200 members were performed to estimate the forecast error covariance matrix. Implementing such a methodology in a realistic case is challenging. Assumptions and methodology developments made to reach these corrections have to be treated with special care. We needed to keep a close attention to the initialization procedure of the experiments to avoid the propagation of potential initial condition errors unrelated to the forcing. This has been done by extracting initial condition to make it as consistent as possible with the surface observations from a strongly constrained simulation. Moreover, the impact of the forcing on the model state had to be isolated from other model errors in the ensemble experiments with a robust diagnostic approach. Finally, the residual excessive corrections potentially linked to some compensation effects have been limited by a truncation of the prior probability distribution of the parameters as described in Skandrani et al. (2009).

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The results have shown that it is possible to identify properly the monthly forcing impact on the model state. First, for every given month, the computed corrections applied to a free model run have been demonstrated to have the same impact as the analysis step on the model SST. Second, the net heat flux budget resulting from ERAcor (highly positive in ERAi database) is balanced at the global scale, and the potentially unrealistic negative trend observable in the time-series of this flux in the ERAi database (leading to ocean cooling) over the whole time period has been removed. Finally, parameters corrections computed improve the long term mean state of a free (without data assimilation) simulation with respect to ocean surface observations.

More generally, this paper illustrates the benefit that can be obtained by using an objective method to estimate forcing corrections. Indeed, as far as the multivariate analysis used to compute $\hat{\mathbf{P}}^f$ is consistent enough with real uncertainties, it gives the necessary information about the impact of forcing errors on the different components of the net heat flux. Furthermore, these results can guide typical ad hoc forcing corrections mostly used to adjust forcing parameters (e.g. Brodeau et al., 2010). It is thus consistent to use SST observations to constrain the forcing function of a model using a statistical updating scheme. As a further step, one could envisage applying a similar method to an operational system for short term forecast in present day operational systems. Indeed, the ocean state correction can present some inconsistency with the forcing parameters used, whereas our approach proposes an alternative to keep the interactive ocean-atmosphere link while correcting efficiently the ocean surface state.

The diagnostic of the computed correction itself has highlighted some limits of the method. Small scale features probably due to the local application of the Kalman filter analysis appear in the correction fields. We also computed some spurious corrections in strong advection regions and at high latitudes despite the effort made to reduce non-forcing uncertainties in ensemble experiments and to define an appropriate parameter prior probability distribution. Improving the identification of forcing error in the model is essential to expand the validity domain of the corrections from the intertropical band to the whole ocean (sea ice covered regions would probably need to be

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



treated separately). In our study, the problem of non-forcing errors limitation has been addressed by using a robust diagnostic approach. To go further one could consider to constrain the water masses by the same type of relaxation but using an ocean reanalysis as reference instead of temperature and salinity climatology. As there are now many atmospheric reanalyses available (ERAi, JRA25, NASA MERRA, NCEP/DOE...) the definition of parameter prior probability distribution could also involve cross differences between different atmospheric reanalyses instead of intra-seasonal and interannual variability (as in Lucas et al., 2008). This could represent a substantial improvement of the method since it would provide a more realistic description of the parameter uncertainties and would reduce spurious corrections at high latitudes.

Furthermore, the model convergence to assimilated observation strongly depends on the presence of the initialization procedure in the experiments. Parameters computed via this method are thus not necessarily appropriate to reduce the instantaneous discrepancies between model state and observations when applied to a free model long-term simulation. However, our results show that these corrections have a positive impact on the long-term mean values of fluxes and ocean surface state. The heat budget resulting from corrected parameters is closer to zero than the one computed with original ERAi variables, with a heat excess reduced by more than 10 W m^{-2} . In addition, our corrections remove the unrealistic negative trend observed in the time-series of the global net heat flux computed with ERAi forcing parameters and Hurrell SST. In the intertropical band where we have a maximum validity of the corrections, the warm bias is reduced which leads to a better agreement with surface observations. The diagnostics of the impact on each net heat flux component allowed identifying the turbulent heat flux as the main actor of this reduced warming, via wind speed increase in the intertropics essentially. These results are not explicitly prescribed by the method and only result from the ensemble description of the correlations between the forcing variables and the SST.

Several perspectives can now be proposed to go further from this study. On the one hand the method can be improved to maximize the impact of observations. In terms

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of methodology, the first difficulty is that the problem is under-determined: we estimate seven forcing parameters using only two observed variables. This under-determination could be partly reduced by assimilating real SSS observations like SMOS or AQUARIUS when available instead of SSS climatology, as well as other ocean profile observations databases like ARGO or TAO. All available observations of the surface, the mixed layer or the deep convection regions that are directly influenced by air-sea fluxes would be beneficial to constrain our system. Recent progresses in intensive ocean observation give a strong potential to our methodology in addressing the problem of the objective computation of forcing correction.

On the other hand further work could be developed in order to take advantage of the results already available from our study. Our method is numerically expensive since it involves numerous simulation experiments, and could not reasonably be used for higher resolution models. Furthermore, even if the results are inhomogeneous in quality depending on the region considered, the method is certainly able to give valuable insights to guide typical ad hoc forcing adjustments. Results should be first subjected to further evaluation by comparison with available atmospheric or fluxes observations and reconstructions especially in the intertropical band like OAflux (Yu and Weller, 2007), TROPFlux (Praveen Kumar et al., 2011) or NOCS (Berry and Kent, 2009) databases. This work would make possible to identify more precisely the strengths and weaknesses of the method, to have a more critical look on the results, and to identify the benefit of the corrections in every particular application.

A wider perspective concerns the implementation of ocean reanalyses. In the construction of an ocean reanalysis, it is certainly useful to extend the ocean state control vector to the forcing parameters to avoid the propagation of forcing errors in the system. For example, the work of Cerovecki et al. (2011) addresses this type of problematic with a variational approach. Conversely, our methodology could also be valuable to provide improved ocean boundary information for the implementation of atmospheric reanalyses, since our approach is not only dedicated to the simulation of the ocean state but can also be viewed as a way of objectively controlling the air-sea interactions.

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. This research has been conducted with the support of the CNES (French Space Agency) and the CNRS (French National Research center), and received partial support from the EU-funded MyOcean Project (Grant FP7-SPACE-2007-1-CT-218812-MYOCEAN). The calculations were performed using HPC resources from GENCI-IDRIS (Grant 2010-011279).

References

- Berry, D. I. and Kent, E. C.: A new air-sea interaction gridded dataset from ICOADS with uncertainty estimates, *B. Am. Meteorol. Soc.*, 90, 645–656, doi:10.1175/2008BAMS2639.1, 2009. 2519
- Bindoff, N. L., Willebrand, J., Artale, V., Cazenave, A., Gregory, J. M., Gulev, S., Hanawa, K., Le Quéré, C., Levitus, S., Nojiri, Y., Shum, C. K., Talley, L. D., and Unnikrishnan, A. S.: Observations: Oceanic Climate Change and Sea Level, *Climate Change 2007: The Physical Science Basis. Contribution of Working Group 1 to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, edited by: Solomon, S., Qin, D., Manning, M., Marquis, M., Averyt, K. B., Tignor, M., Miller, H. L., and Chen, Z.: Cambridge, UK and NY, USA, 385–432, ISBN 978-0-521-70596-7, 2007. 2513
- Brankart, J.-M., Cosme, E., Testut, C.-E., Brasseur, P., and Verron, J.: Efficient local error parameterizations for square root or ensemble Kalman filters: application to a basin-scale ocean turbulent flow, *Mon. Weather Rev.*, 139, 474–493, doi:10.1175/2010MWR3310.1, 2011. 2503
- Brodeau, L.: Contribution à l'Amélioration de la Fonction de Forçage des Modèles de Circulation Générale Océanique, Ph.D. thesis, Université Joseph Fourier – Grenoble 1, 2007. 2500, 2512, 2524
- Brodeau, L., Barnier, B., Treguier, A.-M., Penduff, T., and Gulev, S.: An ERA40-based atmospheric forcing for global ocean circulation models, *Ocean Model.*, 31, 88–104, doi:10.1016/j.ocemod.2009.10.005, 2010. 2496, 2517
- Cerovecki, I., Talley, L., and Mazloff, M.: A comparison of Southern Ocean air-sea buoyancy flux from an ocean state estimate with five other products, *J. Climate*, 24, 6283–6306, doi:10.1175/2011JCLI3858.1, 2011. 2519

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dré villon, M., Bourdallé-Badie, R., Derval, C., Drillet, Y., Lellouche, J.-M., Rémy, E., Tranchant, B., Benkiran, M., Greiner, E., Guinehut, S., Verbrugge, N., Garric, G., Testut, C.-E., Laborie, M., Nouel, L. Bahurel, P., Bricaud, C., Crosnier, L., Dombrowsky, E., Durand, E., Ferry, N., Hernandez, F., Le Galloudec, O., Messal, F., and Parent, L.: The GODAE/Mercator-Ocean global ocean forecasting system: results, applications and prospects, *J. Oper. Oceanogr.*, 1, 51–57, 2008. 2501
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B.: Bulk parameterization of air-sea fluxes: updates and verification for the COARE algorithm, *J. Climate*, 16, 571–591, doi:10.1175/1520-0442(2003)016<0571:BPOASF>2.0.CO;2, 2003. 2524
- Ferry, N., Parent, L., Garric, G., Barnier, B., Jourdain, N. C., and the Mercator Ocean team: Mercator global eddy permitting ocean reanalysis GLORYS1V1: description and results, *Mercator Ocean Q. Newsl.*, 36, 15–27, 2010. 2496
- Hurrell, J. W., Hack, J. J., Shea, D., Caron, J. M., and Rosinski, J.: A new sea surface temperature and sea ice boundary dataset for the Community Atmosphere Model, *J. Climate*, 21, 5145–5153, doi:10.1175/2008JCLI2292.1, 2008. 2497, 2504
- Josey, S.: Air-Sea Fluxes of Heat, Freshwater and Momentum, *Operational Oceanography in the 21st Century*, p. 155, 2011. 2495
- Kalman, R. E.: A new approach to linear filter and prediction problems, *J. Basic Eng.*, 82, 35–45, 1960. 2501
- Large, W. and Yeager, S.: Diurnal to decadal global forcing for ocean and sea-ice models: the data sets and flux climatologies, NCAR Tech., Note TN-460+ STR, 2004. 2496, 2500
- Large, W., Danabasoglu, G., Doney, S., and McWilliams, J.: Sensitivity to surface forcing and boundary layer mixing in a global ocean model: annual-mean climatology, *J. Phys. Oceanogr.*, 27, 2418–2447, 1997. 2498
- Lauvernet, C., Brankart, J.-M., Castruccio, F., Broquet, G., Brasseur, P., and Verron, J.: A truncated Gaussian filter for data assimilation with inequality constraints: application to the hydrostatic stability condition in ocean models, *Ocean Model.*, 27, 1–17, doi:10.1016/j.ocemod.2008.10.007, 2009. 2503
- Levitus, S., Boyer, T., Conkright, M., O'Brien, T., Antonov, J., Stephens, C., Stathoplos, L., Johnson, D., and Gelfeld, R.: *World Ocean Database 1998*, NOAA Atlas NESDID 18, US Government Printing Office, Washington, D.C., 1998. 2497, 2504, 2506

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lucas, M. A., Ayoub, N., Barnier, B., Penduff, T., and de Mey, P.: Stochastic study of the temperature response of the upper ocean to uncertainties in the atmospheric forcing in an Atlantic OGCM, *Ocean Model.*, 20, 90–113, doi:10.1016/j.ocemod.2007.07.006, 2008. 2518
- Madec, G.: NEMO reference manual, ocean dynamics component: NEMO OPA, Preliminary version, Notes techniques de modélisation, Tech. Rep. 27, ISSN No 1288–1619, Institut Pierre Simon Laplace (IPSL), France, 2008. 2498
- Madec, G. and Imbard, M.: A global ocean mesh to overcome the North Pole singularity, *Clim. Dynam.*, 12, 381–388, doi:10.1007/BF00211684, 1996. 2513
- Milliff, R. F., Large, W. G., Morzel, J., Danabasoglu, G., and Chin, T. M.: Ocean general circulation model sensitivity to forcing from scatterometer winds, *J. Geophys. Res.*, 104, 11337–11358, doi:10.1029/1998JC900045, 1999. 2495
- Praveen Kumar, B., Vialard, J., Lengaigne, M., Murty, V., and McPhaden, M.: TropFlux: air-sea fluxes for the global tropical oceans-description and evaluation, *Clim. Dynam.*, 38, 1521–1543, doi:10.1007/s00382-011-1115-0, 2011. 2519
- Sarmiento, J. L. and Bryan, K.: An ocean transport model for the North Atlantic, *J. Geophys. Res.*, 87, 394–408, doi:10.1029/JC087iC01p00394, 1982. 2506, 2507
- Skachko, S., Brankart, J.-M., Castruccio, F., Brasseur, P., and Verron, J.: Improved turbulent air-sea flux bulk parameters for controlling the response of the ocean mixed layer: a sequential data assimilation approach, *J. Atmos. Ocean. Tech.*, 26, 538–555, doi:10.1175/2008JTECHO603.1, 2009. 2496, 2498, 2500, 2502, 2503, 2505
- Skandrani, C., Brankart, J.-M., Ferry, N., Verron, J., Brasseur, P., and Barnier, B.: Controlling atmospheric forcing parameters of global ocean models: sequential assimilation of sea surface Mercator-Ocean reanalysis data, *Ocean Sci.*, 5, 403–419, doi:10.5194/os-5-403-2009, 2009. 2496, 2497, 2498, 2500, 2501, 2502, 2503, 2505, 2508, 2516
- Smith, S. R., Legler, D. M., and Verzone, K. V.: Quantifying uncertainties in NCEP reanalyses using high-quality research vessel observations, *J. Climate*, 14, 4062–4072, doi:10.1175/1520-0442(2001)014<4062:QUINRU>2.0.CO;2, 2001. 2495
- Stammer, D., Ueyoshi, K., Kohl, A., Large, W. G., Josey, S. A., and Wunsch, C.: Estimating air-sea fluxes of heat, freshwater, and momentum through global ocean data assimilation, *J. Geophys. Res.*, 109, C05023, doi:10.1029/2003JC002082, 2004. 2496, 2511
- Sun, B., Yu, L., and Weller, R. A.: Comparisons of surface meteorology and turbulent heat fluxes over the Atlantic: NWP model analyses versus moored buoy observations*, *J. Climate*, 16, 679–695, doi:10.1175/1520-0442(2003)016<0679:COSMAT>2.0.CO;2, 2003. 2495

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wang, W. and McPhaden, M. J.: What is the mean seasonal cycle of surface heat flux in the equatorial Pacific?, J. Geophys. Res., 106, 837–857, doi:10.1029/1999JC000076, 2001. 2495

5 Yu, L. and Weller, R. A.: Objectively analyzed air-sea heat fluxes for the global ice-free oceans (1981–2005), B. Am. Meteorol. Soc., 88, 527–539, doi:10.1175/BAMS-88-4-527, 2007. 2519

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.




Fig. 1. Top: Mean difference between turbulent heat flux ($Q_{\text{turb}} = Q_{\text{lat}} + Q_{\text{sens}}$, in W m^{-2}) climatologies (1984–2000) estimated from different bulk algorithms (different transfer coefficients parametrizations): COA3 described by Fairall et al. (2003) and LY04. These climatologies are computed with the same atmospheric forcing field and the same prescribed SST. Bottom: mean difference between net heat flux (in W m^{-2}) climatologies (1984–2000) estimated with downward radiation from different datasets (ISCCP and ERA40). These climatologies only differ by the radiation used and are computed with the LY04 bulk algorithm (from Brodeau, 2007).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.




Fig. 2. January ensemble standard deviation in terms of air temperature (top left), shortwave radiation (top right), zonal wind velocity (bottom left), and meridional wind velocity (bottom right).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.




Fig. 3. SST differences between the free model run forced by ERAi and the result of the analysis step (top), and between the free model run forced by ERAi and the free model run forced by ERAcor (bottom) for January 2004.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Optimal adjustment
of atmospheric
forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. 1989–2007 mean computed corrections of temperature (t_2), humidity (q_2), zonal wind speed (u_{10}), meridional wind speed (v_{10}), downward shortwave radiation (rad_{sw}), and downward longwave radiation (rad_{lw}).

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.




Fig. 5. 1989–2007 time-series of global net heat fluxes monthly means (in red) computed with ERAi variables (left) and ERAcor variables (right). The black lines represent the global mean of the net heat flux over the whole 1989–2007 period.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 6. 1989–2007 means of net heat flux correction, radiative heat fluxes corrections and turbulent heat fluxes corrections in the 20° N–20° S latitude band.

Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.




Fig. 7. 1989–2007 mean wind speed correction in the 20° N–20° S latitude band.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Optimal adjustment
of atmospheric
forcing parameters

M. Meinvielle et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 8. 1989–2007 mean corrections of temperature (t_2), humidity (q_2), zonal wind speed (u_{10}), meridional wind speed (v_{10}), downward shortwave radiation (rad_{sw}), and downward longwave radiation (rad_{lw}) after anisotropic boxcar smoothing and masking high latitudes values.

**Optimal adjustment
of atmospheric
forcing parameters**

M. Meinvielle et al.




Fig. 9. 1989–2007 mean differences between Hurrell SST and free model run forced by ERAi (top), and between Hurrell SST and free model run forced by ERAcor (bottom) in the 20° N–20° S latitude band.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Optimal adjustment of atmospheric forcing parameters

M. Meinvielle et al.




Fig. 10. Comparison between the simulations forced by ERAI (black) and ERAcor (red). Top: time-series between 1995 and 2007 of the global mean of the sea surface height in m (left), and of the global mean of the temperature in °C (right). Bottom: time-series between 1995 and 2007 of the maximum intensity of the meridional overturning circulation at 30° S in Sv.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)