

Generation of large-scale intrusions at baroclinic fronts: an analytical consideration with a reference to the Arctic Ocean

Natalia Kuzmina

5 Shirshov Institute of Oceanology, 36 Nakhimovsky Ave, 117997 Moscow, Russia

Correspondence to: Natalia Kuzmina (kuzmina@ocean.ru)

Abstract. Analytical solutions are found for the problem of instability of a weak geostrophic flow with linear velocity shear accounting for vertical diffusion of buoyancy. The analysis is based on the potential vorticity equation in a long-wave approximation when the horizontal scale of disturbances is considered much larger than the local baroclinic Rossby radius. It is hypothesized that the solutions found can be applied to describe stable and unstable disturbances of the planetary scale with respect, especially, to the Arctic Ocean, where weak baroclinic fronts with typical temporal variability periods of the order of several years or more have been observed and the β -effect is negligible. Stable (decaying with time) solutions describe disturbances that, in contrast to the Rossby waves, can propagate to both the west and east, depending on the sign of the linear shear of geostrophic velocity. The unstable (growing with time) solutions are applied to explain the formation of large-scale intrusions at baroclinic fronts under the stable–stable thermohaline stratification observed in the upper layer of the Polar Deep Water in the Eurasian Basin. The suggested mechanism of formation of intrusions can be considered a possible alternative to the mechanism of interleaving at the baroclinic fronts due to the differential mixing.

1 Introduction

The study of intrusions in oceanic frontal zones is required to understand the mechanism of ventilation and mixing in the ocean interior (see, for example, Zhurbas et al., 1983, 1987; Rudels et al., 1999, 2009; Kuzmina and Zhurbas, 2000; Walsh and Ruddick, 2000; Merryfield, 2000; Radko, 2003; Richards and Edwards, 2003; Kuzmina et al., 2005, 2011; Smyth and Ruddick, 2010). Intrusive layering, as a rule, results from the instability of oceanic fronts. One of the major mechanisms responsible for the instability of both thermohaline and baroclinic fronts is related to the double diffusion (Stern, 1967; Ruddick and Turner, 1979; Toole and Georgi, 1981; McDougall, 1985a, 1985b; Niino, 1986; Yoshida et al., 1989; Richards, 1991; Kuzmina and Rodionov, 1992; May and Kelley, 1997; Kuzmina, 2000). However, in the Eurasian Basin of the Arctic Ocean there are baroclinic and thermohaline fronts within the upper layer of the Polar Deep Water (PDW) populated with intrusive layers of vertical length scale as large as 30 m and with horizontal scale reaching up to more than 100 km (Rudels et al., 1999, 2009; Kuzmina et al., 2011) observed at the stable–stable stratification (i.e., for the mean salinity increasing while the mean temperature decreasing with depth). It can be suggested that the thermohaline intrusions within the upper layer of PDW are driven by differential mixing. Merryfield (2002) was the first to show satisfactory agreement between calculations of unstable modes from a 3D interleaving model that accounted for the differential mixing at a non-baroclinic front and observations of intrusive layering at a pure thermohaline front in the PDW. Merryfield’s (2002) findings were confirmed by Kuzmina et al. (2014). However, the 2D model of interleaving driven by differential mixing at the baroclinic front failed to simultaneously fit three modelled parameters, namely the vertical scale, the growth time and the slope of the fastest growing mode with observations of intrusions in a frontal zone with a substantial baroclinicity in the upper PDW layer (Kuzmina et al., 2014). In particular, it was found that the vertical scale of the most unstable mode was about two to three times smaller than the vertical scale of intrusions observed in the baroclinic front. Furthermore, it is worth noting that the 2D models of double-diffusive

interleaving, as applied to typical baroclinic fronts in the ocean, are able to forecast intrusive layers with vertical length scale of no more than 10 m (Kuzmina and Rodionov, 1992; May and Kelley, 1997, 2001; Kuzmina and Zhurbas, 2000; Kuzmina and Lee, 2005; Kuzmina et al., 2005). Therefore, despite the proven-by-simulation hypothesis of intrusions of small vertical scale merging into larger structures (Radko, 2007), new approaches to the mathematical description of the formation of large intrusions in the area of baroclinic fronts become relevant.

We suggest that the interleaving at a baroclinic front may be considered as a result of 3D instability of weak geostrophic current due to the combined effects of vertical shear and diffusion of density (buoyancy).

The effect of vertical diffusion of buoyancy on the baroclinic instability of geostrophic zonal wind has been studied theoretically by Miles (Miles, 1965). Proceeding from the analogy between the equations describing the dynamics of large-scale atmospheric perturbations and the Orr–Sommerfeld equation (Lin, 1955; Stern, 1965), Miles (1965) analyzed the instability of the critical layer (a very thin layer in which the phase velocity of a disturbance equals the velocity of zonal flow). He resulted in an analytical asymptotic solution accounting for a very small, though finite, vertical diffusion of buoyancy. Based on the analysis, Miles (1965) concluded that the effect of vertical diffusion of buoyancy on the destabilisation of zonal wind is negligible in comparison with the baroclinic instability (the generation of cyclones and anticyclones) for typical atmospheric geostrophic winds. One could assume, however, that other conditions can occur in the deep ocean. Indeed, in the Polar zones, for example, in the Eurasian Basin of the Arctic, very weak geostrophic currents have been observed in deep layers (Aagaard, 1981). These currents can have a large horizontal (transverse) scale and large time scale of variability, the latter being estimated to exceed one year (Aagaard, 1981). Taking into account, that the influence of the β -effect on the dynamics of large-scale disturbances is negligible in the Polar Ocean, it seems reasonable to suggest, that the contribution of diffusion of buoyancy to the destabilization of weak geostrophic currents can be important. Therefore, in such circumstances one would expect the formation of intrusions, rather than vortices.

The present work is devoted to seeking analytical unstable (increasing with time) and stable (decreasing with time) solutions based on the potential vorticity equation describing the 3D dynamics of a weak baroclinic front, with the vertical diffusion of buoyancy included. Hopefully the results will provide an opportunity to obtain some new insight into the causes of the formation of large intrusions, particularly in the regions of the Arctic Ocean with the stable–stable stratification.

2 Problem formulation, derivation of basic equation, and solution search

Let us consider the problem of the 3D instability of a baroclinic front based on the linearized equations of motion in quasi-geostrophic approximation (see, for example, Pedlosky, 1992; Cushman-Roisin, 1994):

$$U = -\frac{1}{f} \frac{\partial P}{\partial y}, \quad V = 0, \quad W = 0, \quad \frac{\partial P}{\partial z} = -g\bar{\rho} \quad (1)$$

$$\frac{\partial p}{\partial z} = -g\rho, \quad u = -\frac{1}{f} \frac{\partial p}{\partial y}, \quad v = \frac{1}{f} \frac{\partial p}{\partial x} \quad (2)$$

$$\left(\frac{\partial}{\partial t} + U \frac{\partial}{\partial x} \right) \left(\frac{1}{f} \Delta p \right) + \beta v - f \frac{\partial w}{\partial z} = \frac{\tilde{K}}{f} \frac{\partial^2}{\partial z^2} \Delta p \quad (3)$$

$$\left(\frac{\partial}{\partial t} + U \frac{\partial}{\partial x}\right) \rho + v \frac{\partial \bar{\rho}}{\partial y} - \frac{N^2}{g} w = K \frac{\partial^2}{\partial z^2} \rho, \quad (4)$$

where U and V are zonal and meridional components of the geostrophic velocity, P and $\bar{\rho}$ are the mean pressure and density both divided by the reference density, N , f and g are the buoyancy frequency, Coriolis parameter and gravity acceleration, u , v and w are velocity fluctuations along the x , y and z axes, respectively, p and ρ are the pressure and density fluctuations both divided by the reference density, $\beta = \partial f / \partial y$, $\Delta = \partial^2 / \partial x^2 + \partial^2 / \partial y^2$, and the x , y and z axes are directed eastward, northward and upward, respectively. The vertical friction with a constant coefficient \tilde{K} is considered in the vorticity equation Eq. (3). The density balance equation Eq. (4) takes into account, apart from the advection terms, only the vertical diffusion with a constant coefficient K . The constant coefficients \tilde{K} and K are treated as the average values over the ocean layer under investigation.

Let us take the distribution of mean density, divided by the reference density, as follows:

$$\bar{\rho}(z, y) = fsyz / g + f\tilde{s}y / g - \frac{N_0^2}{g} z + 1, \quad (5)$$

where $N_0 = \text{const} > 0$ is a characteristic value of the buoyancy frequency in the frontal zone, and \tilde{s} and s are dimensional constants, either positive or negative, that characterize the cross-front gradient of density and the vertical shear of the basic geostrophic flow.

The first term on the right of Eq. (5) has not been taken into account in the interleaving models describing both the 2D (see, for example, Kuzmina, Rodionov, 1992; May, Kelley, 1997; Kuzmina, Zhurbas, 2000) and 3D (Eady, 1949; Miles, 1965; Smyth, 2008) instabilities of the oceanic baroclinic fronts. Meanwhile, the oceanic fronts can be characterized by not only the cross-front gradient of density, but also by the cross-front gradient of the buoyancy frequency. This is the case described by Eq. (5): the squared buoyancy frequency $N^2 = -gd\bar{\rho}/dz$, is a linear function of y . This dependence is assumed to be weak: $|s|fL \ll N_0^2$, where L is the characteristic lateral length scale (width) of the frontal zone ($0 \leq y \leq L$). However, even a weak lateral change in the buoyancy frequency indicates the existence of a quadratic dependence of geostrophic velocity on the vertical coordinate z . Indeed, if the mean density distribution is expressed by Eq. (5), the geostrophic current velocity will be

$$U = U_1 + U_2 + U_3, \quad U_1 = sz^2 / 2, \quad U_2 = \tilde{s}z, \quad U_3 = \text{const}, \quad (6)$$

where U_1 and U_2 are the constituents of geostrophic velocity with linear (U_1) and constant (U_2) vertical shear: $dU_1/dz = sz$, $dU_2/dz = \tilde{s}$ and U_3 is the barotropic (constant) velocity addition.

The equation of evolution of potential vorticity, derived from Eqs. (1)–(4) under the assumptions of $|fsy/N_0^2| \leq |fsL/N_0^2| \ll 1$ and $U/fL \ll 1$, is

$$\left(\frac{\partial}{\partial t} + U \frac{\partial}{\partial x}\right) \left(\frac{\partial^2 p}{\partial z^2} + \frac{N_0^2 \Delta p}{f^2}\right) + \frac{\beta v N_0^2}{f} - vsf = K \frac{\partial^4}{\partial z^4} p + \tilde{K} \frac{N_0^2}{f^2} \frac{\partial^2}{\partial z^2} \Delta p. \quad (7)$$

Note that the differentiation of Eq. (4) with respect to z cancels out the terms $(\partial U / \partial z) \cdot (\partial \rho / \partial x)$ and $(\partial v / \partial z) \cdot (\partial \bar{\rho} / \partial y)$, since according to Eqs. (1) and (2) they are equal in magnitude and opposite in sign.

As it can be seen from Eq. (7), the last term on the left can strengthen or weaken, depending on the sign of s , the impact of the β -effect on the dynamics of disturbances.

We will consider at $K \approx \tilde{K}$ the long-wave disturbances (i.e. perturbations of the planetary scale) of weak geostrophic current which satisfy the following relationship between the vertical and horizontal length scales (H and \tilde{L} , respectively): $\tilde{L} \gg L_R$, where $L_R = N_0 H / f$ is the baroclinic Rossby radius of deformation. If we apply Eqs. (1)–(4) to describe the motion in the Arctic Basin, the β -effect term can be ignored because $\beta \approx 0$ in the vicinity of the North Pole.

Taking into account the abovementioned conditions, we may use the method of series expansion at small parameter $\delta^2 = N_0^2 H^2 / (\tilde{L}^2 f^2) = Bu$, where Bu is the Burger number (see e.g. Cushman-Roisin, 1994). For $\delta^2 \sim 10^{-3} - 10^{-4}$ ($\tilde{L} \sim 10^4 - 10^5$ m, $20 < H < 100$ m, $N_0 \sim 10^{-3} \text{ s}^{-1}$, $f \sim 10^{-4} \text{ s}^{-1}$), it is reasonable to consider only the first term of the series. In this case, we can rewrite the potential vorticity equation in the simplified form:

$$\left(\frac{\partial}{\partial t} + U \frac{\partial}{\partial x} \right) \left(\frac{\partial^2 p}{\partial z^2} \right) - vsf = K \frac{\partial^4}{\partial z^4} p. \quad (8)$$

The introduced by the procedure relative error of the solution is expected to be of the order of δ^2 , and the smaller δ^2 the smaller the error.

According to our approximation, Eq. (8) corresponds to the density balance equation Eq. (4) for $w = w_0 = const$:

$$\left(\frac{\partial}{\partial t} + U \frac{\partial}{\partial x} \right) \rho + v \frac{\partial \bar{\rho}}{\partial y} - \frac{N_0^2}{g} w_0 = K \frac{\partial^2}{\partial z^2} \rho. \quad (9)$$

The correspondence between Eqs. (8) and (9) can be checked by differentiating Eq. (9) with respect to z and taking into account that $\partial p / \partial z = -g\rho$.

Thus, the vorticity equation Eq. (3) drops out of consideration. Indeed, given that the diffusivity of mass, K , in the oceanic interior (particularly in the deep water of the Arctic Ocean) probably does not exceed the value of $1 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ and the vertical length scale of the intrusions, H , which this theory is applied to, is approximately equal to $H \approx 20 - 100$ m, the ratio of $U / \tilde{L} \sim K / H^2$ is estimated as $U / \tilde{L} < 3 \cdot 10^{-8} \text{ s}^{-1}$. Based on the latter estimate, one can suggest that the vertical circulation caused by the frictional force and temporal change of vorticity, will not significantly affect the dynamics of large-scale disturbances. This hypothesis will be tested a posteriori by analysing the solutions obtained. For further discussion it is important to underline that the geostrophic Richardson number $Ri = N_0^2 H^2 / \tilde{U}^2$, where \tilde{U} is the characteristic scale of geostrophic velocity, is much larger than unity for very slow currents.

Given that $w_0 = const$, we can take $w_0 = 0$ in Eq. (9), and therefore rewrite it as

$$\left(\frac{\partial}{\partial t} + U \frac{\partial}{\partial x} \right) \rho + v \frac{\partial \bar{\rho}}{\partial y} = K \frac{\partial^2}{\partial z^2} \rho. \quad (10)$$

Based on the reasoning above, we can conclude that the slow extra-large-scale disturbances of weak geostrophic flow are described by the quasi-stationary system of Eqs. (2) and (10).

Let us now pay attention to an important issue. Namely, if we suppose that $U(z) = 0$ in Eq. (10) and consider salt fingering instead of diffusion of buoyancy, then, in addition to Eq. (2), it will be necessary to write the following two equations instead of Eq. (10):

$$\frac{\partial \rho}{\partial t} = K_s (1 - \gamma) \tilde{\beta} \frac{\partial^2 S}{\partial z^2}, \quad \frac{\partial S}{\partial t} + v \frac{\partial \bar{S}}{\partial y} = K_s \frac{\partial^2 S}{\partial z^2}, \quad (10')$$

5 where S and \bar{S} are the salinity disturbance and mean, K_s and $\gamma = \tilde{\alpha} F_T / \tilde{\beta} F_s < 1$ are the vertical diffusivity of salinity and the flux ratio for salt finger convection, F_T and F_s are the vertical fluxes of temperature and salinity, $\tilde{\alpha}$ and $\tilde{\beta}$ are the temperature expansion and salinity contraction coefficients, respectively.

10 Eqs. (10') along with (2) constitute the system of equations that was used by Stern (1967) to obtain the polynomial dependence between the growth rate of unstable perturbations, wave numbers and hydrological parameters [see Eq. (4) of (Stern, 1967)]. Therefore, the proposed model, which consists of Eqs. (2) and (10), can in a certain sense be regarded as an analogue of the model by Stern (1967) for investigating the interleaving on a large horizontal scale.

15 From the point of view of the author of this paper, a simple quasi-stationary (geostrophic) system of equations accurately describes the large scale movement especially in the Arctic Ocean, where the influence of the β -effect is not significant, the baroclinic fronts of large width in the ocean interior are often not intense (Kuzmina et al., 2011), and the baroclinic radius of deformation, HN/f , at $H \sim 100$ m, does not exceed 2–5 km (see also Section 3).

To analyse the instability of the geostrophic flow in the frame of Eqs. (2) and (10), let us take a layer with the vertical scale of $2H_0$, move the z -axis origin to the middle of the layer, and consider a symmetric relative to the midline geostrophic flow with quadratic z -dependence of velocity:

$$U = U_1 + U_3, \quad U_1 = sz^2/2, \quad U_3 = -\text{sign}(s) \cdot sH_0^2/2.$$

20 A parabolic z -dependence of the geostrophic flow velocity can be observed in the rotary flow of the intra-pycnocline vortices, as well as in many other ocean flows. In any case, as mentioned above, in the oceanic frontal zones it is likely to observe changes of the buoyancy frequency in the cross-front direction indicating the presence of linear shear of geostrophic velocity. The consideration of the instability of geostrophic flow with the velocity profile of $U = U_1 + U_2 + U_3$ is also possible by analytical methods, but this issue falls out of the scope of the present study.

25 Let us discuss the conditions on the boundaries of the layer in relation to the ocean. Keeping in mind the Eady problem (Eady, 1949), one has to set the vertical velocity vanishing at the layer boundaries. Our approximation meets this condition.

30 Due to our model accounting for the vertical diffusion, it appears reasonable to accept the conditions of zero buoyancy flux (for density perturbations) at the layer boundaries: $p_{zz} = 0$ at $z = \pm H_0$ (the type 1 boundary conditions). It is reasonable to consider another type of condition too, namely, the slippery boundary conditions or equivalent of density disturbances vanishing at boundaries: $dv/dz = du/dz = \rho = 0$ at $z = \pm H_0$ (the type 2 boundary conditions). Under the type 2 boundary conditions, it is necessary to set up the absence of convergence or divergence of buoyancy flux within the layer: $p_{zz}(z = H_0) = p_{zz}(z = -H_0)$. This condition is necessary, because the convergence or divergence of the buoyancy flux within the layer may increase or, conversely, decrease the stability of the layer.

Using Eq. (2), we rewrite Eq. (10) as:

$$\left(\frac{\partial}{\partial t} + U \frac{\partial}{\partial x}\right) \left(\frac{\partial p}{\partial z}\right) - \frac{\partial p}{\partial x} s z = K \frac{\partial^3}{\partial z^3} p, \quad (11)$$

where $U = U_1 + U_3$.

To analyse the instability of geostrophic flow, we will seek the solution of Eqs. (2) and (11) for the lateral boundary conditions of $v|_{y=0} = v|_{y=L} = 0$, accordingly to Eq. (7), in the form

$$p = \text{Re}\left\{F(z)e^{ik(x-ct)} \sin(\pi y/L)\right\}, \quad (12)$$

where k is the wave number along the x axis and c is the growth rate. For the positive imaginary part of c , $\text{Im}(c) > 0$, the solution will be unstable, i.e. increasing with time.

The substitution of Eq. (12) into Eq. (11) yields the following equation:

$$ik(U_1 + U_3 - c) \left(\frac{dF(z)}{dz}\right) - F(z)iks z - K \frac{d^3}{dz^3} F(z) = 0. \quad (13)$$

10 We are interested in finding an answer to the following question: is it possible to make certain judgements about the possibility of instability of geostrophic flow in a finite vertical layer, based on the analytical solutions of Eq. (13) at some values of parameter c ?

It is easy to verify that the following functions are partial solutions of Eq. (13):

$$F_1(z) = e^{-az^2/2}, F_2(z) = az^2 + D, \quad (14)$$

15 where $a^2 = ks/2K$, $D = -2a(c - U_3)/s$, $ikc = ik(c_1 + ic_2) = 5a \cdot K + ikU_3$, $c_1 = \text{Re}c$, $c_2 = \text{Im}c$, and $U_1 + U_3 - \text{Re}c \neq 0$ at an arbitrary point $z = z_0$ in the layer domain.

To test partial solutions Eq. (14), one has to substitute $F_1(z)$ and $F_2(z)$ from Eq. (14) into Eq. (13), reduce the latter to a cubic polynomial $P(z) = A_3 z^3 + A_2 z^2 + A_1 z^1 + A_0 z^0$ and evaluate coefficients A_0 , A_1 , A_2 , and A_3 . It is easy to make sure, that this polynomial is identically zero (i.e. $A_0 \equiv 0$, $A_1 \equiv 0$, $A_2 \equiv 0$, and $A_3 \equiv 0$).

20 Proceeding from the theory of ordinary differential equations (see, for example, Polyanin and Zaitsev, 2001), due to the linearly independent functions $F_1(z)$ and $F_2(z)$, we can express the general solution of Eq. (13) for $ikc = 5a \cdot K + ikU_3$ in the form:

$$\begin{aligned} F(z) &= B_1 F_1(z) + B_2 F_2(z) + B_3 F_3(z) \\ F_3(z) &= F_1(z) \cdot \int F_2(z) \varphi(z) dz - F_2(z) \cdot \int F_1(z) \varphi(z) dz, \\ \varphi(z) &= (F_1(z) \cdot dF_2(z)/dz - F_2(z) \cdot dF_1(z)/dz)^{-2} \end{aligned} \quad (15)$$

25 where B_1 , B_2 , B_3 are arbitrary constants. It is important to note two facts. Firstly, the functions $F_1(z)$ and $F_2(z)$ are even functions, while $F_3(z)$ is an odd function. Secondly, despite the singularity of integrands in Eq. (15) at $z = 0$, the function $F_3(z)$ is differentiable at this point. (The latter becomes evident from the asymptotic analysis of function $F_3(z)$ for $z \rightarrow 0$.)

Let us now consider the unstable and stable solutions Eq. (15).

2.1 Unstable solutions

According to the expression for parameter c (see Eq. (14)), $ikc = ik(c_1 + ic_2) = 5a \cdot K + ikU_3$, solution Eq. (15) can be unstable for both $s > 0$ and $s < 0$. The real and imaginary parts of kc for the unstable (i.e. growing with time) solutions are

$$kc_1 = 2.5 \cdot \sqrt{|s|kK} + kU_3, \quad kc_2 = 2.5 \cdot \sqrt{|s|kK} \quad \text{for } s < 0, \quad (16 \text{ a})$$

$$5 \quad kc_1 = -2.5\sqrt{skK} + kU_3, \quad kc_2 = 2.5 \cdot \sqrt{skK} \quad \text{for } s > 0. \quad (16 \text{ b})$$

Eqs. (16) demonstrate that the condition $U_1 + U_3 - \text{Re}c \neq 0$ is satisfied for $z \in (-\infty, +\infty)$.

According to Eqs. (16), the unstable solution is realized for $\text{Re}a < 0$, and hence, for any finite wave number k , the function $F_1(z)$ and all its derivatives increase infinitely, if $z \rightarrow \pm\infty$. On the other hand, the function $F_3(z)$ and all its derivatives vanish, if $z \rightarrow \pm\infty$. (It can be seen from asymptotic analysis of the integrals that define $F_3(z)$, if $z \rightarrow \pm\infty$.) Therefore, to prove the instability in a finite layer, it is necessary to show that $F(z)$ for $\text{Re}a < 0$ is an eigenfunction of the eigenvalue problem with the boundary conditions of type 1 or 2 introduced above. To construct physically correct solutions we will consider the following two cases. Case 1, where the vertical scale of the layer corresponds to our approximation: $2H_0 \sim H \ll fL/N$. Case 2, where the vertical scale of the layer significantly exceeds the vertical scales of the disturbances for which our approximation holds true: $H \ll 2H_0$.

To satisfy the boundary conditions of type 1 and 2 in case 1, we have to take $B_3 = 0$, because $F_3(z)$ is an odd function. The type 1 boundary conditions are reduced to $F_{zz} = 0$ for $z = \pm H_0$. Thus, the following equality should be met:

$$e^{-aH_0^2/2}(-1 + aH_0^2) + 2B_2/B_1 = 0. \quad (17)$$

Given that $2B_2/B_1$ can have different values, the instability in the framework of solution Eq. (15) does exist, because in a wide range of typical ocean values of H_0 , s , and K , there is a wave number $k_0 \ll f/(2N_0H_0)$ for which Eq. (17) is satisfied.

The type 2 boundary conditions are reduced to $F_z = 0$ at $z = \pm H_0$. Under such conditions, the requirement of the absence of the buoyancy flux convergence/divergence within the layer is met: in case of parity of $F(z)$ for $B_3 = 0$ and for the flow symmetry relative to the midline, the values of buoyancy flux at the boundaries are of the same magnitude and direction (sign). Under the type 2 boundary conditions the following equality should be met:

$$e^{-aH_0^2/2} - 2B_2/B_1 = 0. \quad (18)$$

Obviously, in this case, as in the case of Eq. (17), there is a wave number $k_0 \ll f/(2N_0H_0)$ for which Eq. (18) is satisfied.

Figure 1 presents graphic images of unstable solutions in the form of density disturbances $\text{Re} \rho = \text{Re}(dF/dz) = \tilde{\rho}$ for different boundary conditions for the case 1. When building the solutions, typical values of hydrological parameters in relation to the Arctic Basin were used (see Subsection 2.3 and Section 3).

In the case 2, we have to take $B_1 = 0$, $B_2 = 0$, and consider $F_3(z)$ as the solution of the eigenvalue problem. Indeed, $F_3(z)$ and all its derivatives sharply decrease, if $z \rightarrow \pm\infty$, and, consequently, on the boundaries of the large vertical scale layer the

function $F_3(z)$ and all its derivatives should be infinitesimally small. Therefore, the boundary conditions of type 1 and 2 are satisfied. Indeed, the characteristic vertical scale of the decrease of $F_3(z)$ at $z \rightarrow \pm\infty$ can be evaluated as $h \sim (ks/K)^{-1/4}$. If H_0 is n times as large as h , the value of $|F_3(z)|$ at $|z|=H_0$ will be e^{n^2} times (!) as small as the maximum value of $|F_3(z)|$. Note, that the maximum value of geostrophic velocity U_{\max} increases with H_0 . However, U_{\max} should satisfy the condition of $U_{\max}k/f \ll 1$, which can be rewritten as $n^2(ksK)^{1/2}/f \ll 1$. It is easy to see that this condition is met in a wide range of k , s and K for $n=5-10$.

A plot of the disturbances of density $\tilde{\rho} = \text{Re}(dF_3/dz)$ corresponding to unstable solutions is presented in Fig. 2. It is worth noting that the function $\rho = dF_3/dz$ is differentiable at $z=0$ likewise the function $F_3(z)$.

Thus, in accordance with Eq. (11) that was obtained from Eqs. (1)-(10) for $\text{Bu} \ll 1$, $\text{Ri} \gg 1$ and $\text{Pr} = 1$, the large-scale disturbances can be unstable. Such instability has to be distinguished from the diffusive instability (McIntyre, 1970; Baker, 1970; Calman, 1976) which occurs when $\text{Ri} < (\text{Pr}+1)^2/4\text{Pr}$ and is absent at $\text{Pr} = 1$. One of the important distinctions between these two models of baroclinic front instability is that in the present model the disturbances are allowed to have a nonzero slope in the along-front direction while in the model of diffusive instability by McIntyre (1970) the slope is taken to be zero. Therefore, the McIntyre's model and other models in which the term $\partial p/\partial x$, where x is the along-front co-ordinate, in the equations of motions is ignored (McIntyre, 1970; Baker, 1971; Calman, 1977; Munro et al., 2010), can be referred to as the 2D models.

From the mathematical point of view, the models that take into account the along-front slope of disturbances, are much more complicated. Indeed, the analysis of instability in the 2D models ultimately reduces to finding the roots of a polynomial depending upon the wave-numbers and growth rate. The models, that take into account the along-front slope of disturbances, are reduced to the differential equations with variable, z -dependent coefficients, and such problems can be solved analytically only in rare cases.

Also, it is worth to note that the instability described by Eq. (11) is not the critical layer instability analyzed by Miles (Miles, 1965) for a geostrophic current with constant vertical shear based on similarity between the equation of potential vorticity and the Orr-Sommerfeld equation. Indeed, the phase velocity of the unstable disturbances in our model satisfies the inequality $U_1 + U_3 - \text{Re}c \neq 0$ for any value of the vertical coordinate z . As we can see from Eq. (16), the phase velocity of unstable disturbances is directed along the geostrophic current and exceeds the maximum velocity of the current. In such a case, in the author's opinion, the most likely is the conversion of the kinetic energy of the main flow into the kinetic energy of disturbances.

2.2 Stable solutions

Stable solutions of Eq. (13) are realized for $\text{Re}a > 0$. In this case $F_1(z)$ and all its derivatives vanish for $z \rightarrow \pm\infty$, but $F_3(z)$ and all its derivatives increase infinitely for $z \rightarrow \pm\infty$. To construct our own functions of the eigenvalue problem for the case 1 ($2H_0 \sim H \ll fL/N$), we have to take $B_3 = 0$.

The solutions describe slow time-decay, long waves that can move, in contrast to the Rossby waves, not only to the west but also to the east depending on the sign of s (see Eq. (7)). Moreover, if $|s| > \beta N_0^2 / f^2$ (which is quite possible especially in polar regions), the long-wave dynamics in the β -plane approximation is determined by the linear shear of geostrophic flow rather than the β -effect.

The real and imaginary parts of the growth rate of stable perturbations are

$$kc_1 = -2.5 \cdot \sqrt{|s|kK} + kU_3, \quad kc_2 = -2.5 \cdot \sqrt{|s|kK} \quad \text{for } s < 0, \quad (19 \text{ a})$$

$$kc_1 = 2.5 \sqrt{skK} + kU_3, \quad kc_2 = -2.5 \cdot \sqrt{skK} \quad \text{for } s > 0. \quad (19 \text{ b})$$

According to Eq. (19), the condition $U_1 + U_3 - \text{Re}c \neq 0$ is satisfied, if $2.5\sqrt{|s|kK}/k > |s|H_0^2/2$. Comparing Eqs. (16) and (19), we can conclude, that the phase velocity has different sign for the stable and unstable disturbances. That is, stable and unstable perturbations described by solutions Eq. (15) will move in opposite directions with respect to the flow and a fixed observer.

For the case 1 type 2 boundary conditions, a plot of the density disturbances $\text{Re} \rho = \text{Re}(dF/dz) = \tilde{\rho}$ corresponding to the stable solutions is presented in Fig. 3.

2.3 Obtained solutions: discussion

Our model does not allow to determine the maximum growth rate. Here again we can see an analogy with the work by Stern (1967). Indeed, in a well-known paper by Stern (1967), which was the first study of the double diffusion instability of the infinite thermohaline front, the magnitude of the fastest growing mode was not found. The reason is that the growth rate in Stern's model could indefinitely increase with the horizontal wave number due to the neglected vertical friction. A similar feature is typical for our model. The growth rate increases with the increase in the wave number k up to the limit \tilde{k} for which the constraint of $\tilde{k} \ll f/(2H_0N_0)$ is still valid. Nevertheless, for a rough estimate of the time of formation of unstable perturbations it is reasonable to use Eq. (16). It is also worth evaluating the relationship between the growth rate of unstable disturbances and the layer thickness (case 1) or the characteristic vertical scale of disturbances (case 2). Let us address Eq. (17), which follows from the boundary conditions for one of the problems of instability in a finite layer. The parameter $\chi = \text{Re}(-aH_0^2) = 0.5(ks/K)^{1/2}H_0^2$ governs Eq. (17). The higher the value of χ , the larger the wave number of the unstable mode for the given values of the problem parameters K , s , and H_0 , and therefore, the larger the growth rate. However, the applicability of our model imposes a constraint on the space of wave numbers, $k \ll f/(2N_0H_0)$. In order to satisfy these two conditions simultaneously in the wide range of variability of hydrological parameters in the ocean, it is reasonable to put $1 \leq \chi \leq 2$. For $\chi = 2$, taking into account Eq. (16), we obtain the following formula relating the growth rate of disturbances and the vertical scale of the layer:

$$kc_2 = 1/T = 10 \cdot K/H_0^2.$$

It is easy to understand the physical meaning of the parameter χ . This parameter characterizes the ratio of advection and vertical diffusion terms depending on the wave number k . Indeed, recalling that in our model $U = U_1 + U_3$ and the geostrophic velocity on the boundaries of the layer is zero, the maximum velocity at the midline of the layer shall be $U_{\max} = |s|H_0^2/2$. This allows the squared parameter χ to be presented as $\chi^2 = 0.25(ks/K)H_0^4 = 0.5 \cdot R_d k H_0$, where $R_d = 0.5sH_0^3/K = U_{\max}H_0/K$ is a diffusion analogue of the Reynolds number called the Peclet number.

To conclude this section, we note the following.

The instability of the weak geostrophic flow in the frame of the solutions Eq. (15) is an oscillatory instability (the growth rate has real and imaginary components). Generally, using interleaving models (Stern, 1967; Tool and Georgy, 1981; McDougal, 1985a, 1985b; Niino, 1986; Yoshida et al., 1989; Kuzmina and Rodionov, 1992; May and Kelley, 1997; Kuzmina and Zhurbas, 2000; Walsh and Ruddick, 2000; Merryfield, 2002), it is possible to obtain the monotonous unstable modes only (the phase velocity of the disturbances is equal to zero: $Re c = 0$). The exceptions to this rule are the interleaving models related to equatorial fronts. In accordance with the modelling efforts (Richards, 1991; Edwards and Richards, 1999; Kuzmina et al., 2004; Kuzmina and Lee, 2005), the instability of the equatorial fronts in the scale of intrusive layering is regarded as an oscillatory instability.

General solution Eq. (15) is one of the classes of solutions of Eq. (13). Thus, for example, at $ikc = 9a \cdot K + ikU_3$ it is also possible to find an analytical general solution of (13). This solution would have a more complex structure than the solution Eq. (15). The detailed analytical consideration of unstable modes based on the analysis of different classes of solutions of Eq. (13) taking into account the friction may be a subject for further research. In order to clearly define the range of applicability of our model, it would be worth solving the eigenvalue problem for Eq. (7) for small values of parameter δ^2 by means of numerical methods. This problem also may be a subject for further research. The analytical solutions found can be used to validate numerical solutions of the eigenvalue problems. Moreover, the analytical solutions obtained provide analytical expressions for eigenfunctions, phase velocities and growth/decay rates of disturbances that cannot, as a rule, be found exactly from numerical solutions.

3 Application to thermohaline intrusions in the Eurasian Basin of the Arctic Ocean

It is worth evaluating the time of formation of large-scale intrusions based on the results of the presented model. According to Kuzmina et al. (2011), in the upper layer of the Polar Deep Water (PDW) where the large-scale intrusions are observed in the Eurasian Basin at stable–stable stratification, the following estimates of N , f , and β are typical: $N \approx 2 \cdot 10^{-3} \text{ s}^{-1}$, $f = 1.4 \cdot 10^{-4} \text{ s}^{-1}$, and $\beta < 0.3 \cdot 10^{-11} \text{ m}^{-1} \text{ s}^{-1}$ (at latitude of 83°N and higher). Therefore, for disturbances, for example, with the vertical scale of $H = 100 \text{ m}$, the Rossby radius of deformation is only $HN/f \approx 1 \text{ km}$.

According to the derivation of Eq. (7), the value of the linear shear s is limited by the inequality of $|s|fL \ll N_0^2$. Given that the horizontal scale of the baroclinic fronts (along the cross-front axis y) in the upper layer of the PDW is approximately $L \approx 50\text{--}100 \text{ km}$ [see examples of transections across the fronts of different types observed in the PDW (Kuzmina et al., 2011)], the maximal linear shear can be estimated as $|s| \approx (1\text{--}2) \cdot 10^{-7} \text{ m}^{-1} \text{ s}^{-1}$. Such value of the linear shear is large enough to neglect the β -effect term relative to the linear shear term in Eq. (7): $\beta N_0^2 / f^2 s < 0.6 \cdot 10^{-2} \ll 1$. The vertical diffusivity K can be estimated in the range of $K = (1\text{--}3) \cdot 10^{-6} \text{ m}^2 \text{ s}^{-1}$ (Merryfield, 2002; Walsh and Carmack, 2003). We suggest a weak turbulence regime in the layer under consideration: $Pr > 1$, $Pr \cdot Bu \ll 1$. The typical vertical scale of intrusive layering in the fronts of PDW is approximately $30\text{--}40 \text{ m}$ (Merryfield, 2002; Kuzmina et al., 2014). Let us evaluate the time of formation of intrusions with the vertical scale of $\sim 40 \text{ m}$. Using formula $k_0 c = 10 \cdot K / H_0^2$ (see Subsection 2.3), we can estimate the time of formation of the unstable mode as $1/(k_0 c_2) \sim 5 \text{ years}$ for $K = 10^{-6} \text{ m}^2 \text{ s}^{-1}$ and approximately 2 years for $K = 3 \cdot 10^{-6} \text{ m}^2 \text{ s}^{-1}$.

To verify the applicability of our model, it is worth to estimate the wave number k_0 using the following formula (see Subsection 2.3):

$$k_0 = 16 \cdot K / (H_0^4 s). \quad (20)$$

Substituting $H_0 = 40$ m, $s = 2 \cdot 10^{-7} \text{ m}^{-1}\text{s}^{-1}$ and $K = 10^{-6} \text{ m}^2\text{s}^{-1}$ in Eq. (20), we find $k_0 = 3 \cdot 10^{-5} \text{ m}^{-1}$. The value of $k_0 \approx 10^{-4} \text{ m}^{-1}$ may be obtained at $s = 2 \cdot 10^{-7} \text{ m}^{-1}\text{s}^{-1}$, $K = 3 \cdot 10^{-6} \text{ m}^2\text{s}^{-1}$. These values of k_0 lie in the wave number range of applicability of our model, since $\delta^2 = N_0^2 H_0^2 k_0^2 / f^2 \sim 10^{-4} - 10^{-3}$.

The above-presented estimates of the time of formation of intrusions in PDW are evidently better than the evaluations from 2D modelling of baroclinic front instability (see Introduction).

In the closing of this section, let us justify the assumption that the circulations associated with changes in vorticity Δp are not essential in the description of the formation of intrusions in all considered cases. According to Eqs. (2) and (10), the characteristic scale of vertical velocity in such circulations can be written as $w_1 \sim U \cdot u \cdot H_0 \cdot k_0^2 / f$. In all the considered above cases of application of the model to the Arctic intrusions, the relation of $U \cdot k_0 < 10^{-8} \text{ s}^{-1}$ is satisfied. Given that small disturbances of horizontal velocity cannot exceed the value of geostrophic velocity U , we find $w_1 < 4 \cdot 10^{-11} \text{ m s}^{-1}$. A liquid particle with such vertical velocity travels less than 0.004 m over the period of formation of intrusion ($t_* = 1/(k_0 c_2) \approx 3$ years), while due to the vertical diffusion, the particle displacement is estimated as $\sqrt{K \cdot t_*} \approx 40$ m (i.e. four orders of magnitude larger). Note also, that the decreasing with $|z|$ solution $F_3(z)$ can be used for the description of generation of intrusions even if the vertical velocity is not negligibly small.

4 Conclusions

In this paper, we investigated analytically the instability of a baroclinic front in the quasi-geostrophic, long-wave approximation taking into account the vertical diffusion of buoyancy. Such instability has to be distinguished from the 2D McIntyre instability (McIntyre, 1970), the instability due to flow-dependent fluctuations in turbulent diffusivities (Smyth and Ruddick, 2010), and the 2D baroclinic instability due to the double diffusion (Kuzmina and Rodionov, 1992; May and Kelley, 1997; Kuzmina and Zhurbas, 2000; Kuzmina and Lee, 2005).

In contrast to the paper by Miles (Miles, 1965), who showed, that the vertical diffusion of buoyancy is not essential in comparison with the vorticity change in the destabilisation of zonal flow, we considered the opposite case, where the vertical diffusion of buoyancy can play an important role as a destabilizer of a very weak geostrophic current with linear shear and large cross-frontal scale.

The model we developed can be considered as a modification of Stern's model (Stern, 1967). However, instead of analysing the instability of a purely thermohaline front due to the double diffusion (Stern, 1967), in our case the instability of a weak baroclinic front is analysed taking into account the vertical diffusion of density. This model can be useful for describing stable and unstable disturbances of the planetary scale in the polar regions of the ocean under the stable–stable stratification, particularly in the deep water of the Arctic Ocean, where weak baroclinic fronts with a large horizontal (cross-frontal) scale and typical temporal variability period of the order of several years or more have been observed, and the β -effect is negligible.

The stable solutions are shown to describe long-wave disturbances, which, unlike Rossby waves, can move not only to the west but also to the east, depending on the magnitude and sign of the linear shear of geostrophic velocity. It is important to underline, that the linear shear of the mean flow (parabolic z -dependence of the mean velocity) affects the dynamics of disturbances likewise the β -effect.

5 The unstable solutions can contribute to better understanding of the formation of large-scale intrusions at baroclinic fronts of the Arctic Ocean in the layers characterized by absolutely stable thermohaline stratification, for example, in the upper layer of the PDW in the Eurasian Basin. It is important, that the vertical scale of the new modes of instability can reach tens of meters of magnitude, just in accordance with the observations. However, the model is so complex that obtaining the comprehensive results of modelling that can be fully comparable with the empirical data, would still remain a future task.

10 Acknowledgements

This work was supported by the Russian Science Foundation (Grant No 14-50-00095) and the Russian Foundation for Basic Research (Grant No 15-05-01479-a). The author is grateful to Victor Zhurbas for the constructive discussions. Comments by anonymous reviewers were very helpful in improving the manuscript.

References

- 15 Aagaard, K.: On the deep circulation of the Arctic Ocean, *Deep Sea Res. Part A*, 28, 251–268, 1981.
- Baker, D. J.: Density gradients in a rotating stratified fluid: experimental evidence for a new instability, *Science*, 172 (3987), 1029–1031, 1971.
- Calman, J.: Experiments on high Richardson number instability of a rotating stratified shear flow, *Dynamics of Atmospheres and Oceans*, 1 (4), 277–297, 1977.
- 20 Cushman-Roisin, B.: *Introduction to the Geophysical Fluid Dynamics*, Prentice Hall, Englewood Cliffs, New Jersey 07632, 320 pp., 1994.
- Eady, E. T.: Long waves and cyclone waves, *Tellus*, 1 (3), 33–52, 1949.
- Edwards, N. and Richards, K. J.: Linear double-diffusive-inertial instability at the equator, *J. Fluid. Mech.*, 395, 295–319, 1999.
- Kuzmina, N. P.: On the parameterization of interleaving and turbulent mixing using CTD data from the Azores Frontal Zone, *J. Mar. Syst.*, 23, 285–302, 2000.
- 25 Kuzmina, N. and Lee, J. H.: Driving Forces of Interleaving in the Baroclinic Front at the Equator, *J. Phys. Oceanogr.*, 35, 2501–2519, 2005.
- Kuzmina, N. P. and Rodionov, V. B.: Influence of baroclinicity on the formation of thermohaline intrusions in oceanic frontal zones, *Izvestia Akademii Nauk SSSR, Fizika Atmosferi i Okeana*, 28 (10–11), 1077–1086, 1992.
- 30 Kuzmina, N. P. and Zhurbas, V. M.: Effects of Double Diffusion and Turbulence on Interleaving at Baroclinic Oceanic Fronts, *J. Phys. Oceanogr.*, 30, 3025–3038, 2000.
- Kuzmina, N. P., Lee, J. H., and Zhurbas, V. M.: Effects of turbulent mixing and horizontal shear on double-diffusive interleaving in the Central and Western Equatorial Pacific, *J. Phys. Oceanogr.*, 34, 122–141, 2004.
- Kuzmina, N., Rudels, B., Stipa, T., and Zhurbas, V.: The Structure and Driving Mechanisms of the Baltic Intrusions, *J. Phys. Oceanogr.*, 35 (6), 1120–1137, 2005.
- 35

- Kuzmina, N., Rudels, B., Zhurbas, V., and Stipa, T.: On the structure and dynamical features of intrusive layering in the Eurasian Basin in the Arctic Ocean, *J. Geophys. Res.*, 116, C00D11, doi:10.1029/2010JC006920, 2011.
- Kuzmina, N. P., Zhurbas, N. V., Emelianov, M. V., and Pyzhevich, M. L.: Application of interleaving Models for the Description of intrusive Layering at the Fronts of Deep Polar Water in the Eurasian Basin (Arctic), *Oceanology* 54 (5), 557–566, 2014.
- Lin, C. C.: *The Theory of Hydrodynamic Stability*, Cambridge University Press, 155 pp., 1955.
- May, B. D. and Kelley, D. E.: Effect of baroclinicity on double-diffusive interleaving, *J. Phys. Oceanogr.*, **27**, 1997–2008, 1997.
- May, B. D. and Kelley, D. E.: Growth and steady stage of thermohaline intrusions in the Arctic Ocean, *J. Geophys. Res.*, 106 (C8), 16783–16794, 2001.
- McDougall, T. J.: Double-diffusive interleaving. Part I: Linear stability analysis, *J. Phys. Oceanogr.*, 15, 1532–1541, 1985a.
- McDougall, T. J.: Double-diffusive interleaving. Part II: Finite amplitude, steady state interleaving, *J. Phys. Oceanogr.*, **15**, 1542–1556, 1985b.
- McIntyre, M. E.: Diffusive destabilization of the baroclinic circular vortex, *Geophys. Fluid Dyn.*, 1 (1–2), 19–57, 1970.
- Merryfield, W. J.: Origin of thermohaline staircases, *J. Phys. Oceanogr.*, 30, 1046–1068, 2000.
- Merryfield, W. J.: Intrusions in Double-Diffusively Stable Arctic Waters: Evidence for Differential mixing? *J. Phys. Oceanogr.*, 32, 1452–1439, 2002.
- Miles, J. W.: Effect of Diffusion on Baroclinic Instability of the Zonal Wind, *J. Atmos. Sci.*, 22, 146–151, 1965.
- Munro, R. J., Foster, M. R., and Davies, P. A.: Instabilities in the spin-up of a rotating, stratified fluid, *Physics of Fluids* (1994-present), 22 (5), 054108, 2010.
- Niino, H. A.: Linear stability theory of double-diffusive horizontal intrusions in a temperature-salinity front, *J. Fluid. Mech.*, 171, 71–100, 1986.
- Polyain, A. D. and Zaitsev, V. F.: *Handbook: Ordinary differential equations*, Fizmatlit, Moscow, 576 pp., 2001.
- Pedlosky, J.: *Geophysical Fluid Dynamics*, Second Edition, Springer, New York, 710 pp., 1992.
- Radko, T.: A mechanism for layer formation in a double-diffusive fluid, *J. Fluid Mech.*, 497, 365–380, 2003.
- Radko, T.: Mechanics of merging events for a series of layers in a stratified turbulent fluid, *J. Fluid Mech.*, 251–273, 2007.
- Richards, K. J.: Double-diffusive interleaving at the equator, *J. Phys. Oceanogr.*, 21, 933–938, 1991.
- Richards, K. J. and Edwards, N. R.: Lateral mixing in the equatorial Pacific: The importance of inertial instability, *Geophys. Res. Lett.*, 30, (17), 1888, 2003.
- Ruddick, B. and Turner, J. S.: The vertical length scale of double-diffusive intrusions, *Deep Sea Res. Part A*, 903–913, 1979.
- Rudels, B., Bjork, G., Muench, R.D., and Schauer, U.: Double-diffusive layering in the Eurasian Basin of the Arctic Ocean, *J. Mar. Syst.*, 21 (1–4), 3–27, 1999.
- Rudels, B., Kuzmina, N., Schauer, U., Stipa, T., and Zhurbas, V.: Double-Diffusive Convection and Interleaving in the Arctic Ocean – Distribution and Importance, *Geophysika*, 45, 199–213, 2009.
- Smyth, W. D.: Instabilities of a Baroclinic, Double Diffusive Frontal Zone, *J. Phys. Oceanogr.*, 38, 840–861, 2008.
- Smyth, W. D. and Ruddick, B.: Effects of Ambient Turbulence on Interleaving at a Baroclinic Front. *J. Phys. Oceanogr.*, 40, 685–712, 2010.
- Stern, M. E.: Lateral mixing of water masses, *Deep Sea Res. Part A*, 14, 747–753, 1967.
- Stern, M.: *Ocean circulation physics*, International Geophysics Series, 19, Academic Press, New York, 246 pp., 1975.

- Toole, J. M. and Georgi, D. T.: On the dynamics and effects of double-diffusively driven intrusions, *Prog. Oceanog.*, 10 (2), 123–145, 1981.
- Walsh, D. and Carmack, E.: The nested structure of Arctic thermohaline intrusions, *Ocean Modelling*, 5, 267–289, 2003.
- Walsh, D. and Ruddick, B.: Double-diffusive interleaving in the presence of turbulence – the effect of a nonconstant flux ratio, *J. Phys. Oceanogr.*, 30, 2231–2245, 2000.
- 5
- Yoshida, J., Nagashima, H., and Niino, H.: The behavior of double-diffusive intrusion in a rotating system, *J. Geophys. Res.*, 94, 4923–4937, 1989.
- Zhurbas, V. M., Kuzmina, N. P., and Kul'sha, O. B.: Steplike stratification of the ocean thermocline resulting from transformations associated with thermohaline salt finger intrusions (numerical experiment), *Okeanologiya (Moscow)* 27 (3), 377–383, 1987.
- 10
- Zhurbas, V. M., Kuzmina, N. P., Ozmidov, R. V., Golenko, N. N., and Paka, V. T.: Manifestation of subduction in thermohaline fields of vertical fine structure and horizontal mesostructure in frontal zone of Azores Current, *Okeanologiya (Moscow)*, 33 (3), 321–326, 1993.

Figures




Figure 1. Modelled vertical profiles of density disturbances $\text{Re}(dF/dz) = \text{Re } \rho = \tilde{\rho}$ for case 1. Unstable (growing) solution for boundary conditions of type 1 (left) and type 2 (right) for $\chi = \text{Re}(-aH_0^2) = 0.5(ks/K)^{1/2}H_0^2 = 1.5$, $K = 10^{-5} \text{ m}^2 \text{ s}^{-1}$, $H_0 = 100 \text{ m}$, $s = 10^{-7} \text{ m}^{-1}\text{s}^{-1}$.

5




Figure 2. Modelled vertical profile of density disturbances $\text{Re}(dF/dz) = \text{Re } \rho = \tilde{\rho}$ for case 2. The function $\text{Re}(dF_3/dz)$ (left) and its stretched fragment (right) versus the dimensionless co-ordinate $\tilde{z} = z \cdot (ks/K)^{1/4}$ are presented for $k = 10^{-5} \text{ m}^{-1}$, $s = 10^{-7} \text{ m}^{-1}\text{s}^{-1}$, $K = 10^{-5} \text{ m}^2\text{s}^{-1}$.

10




Figure 3. Stable solution for case 1 and boundary conditions of type 2 for $\chi = \text{Re}(-aH_0^2) = 0.5(ks/K)^{1/2}H_0^2 = 2$, $K = 10^{-5} \text{ m}^2 \text{ s}^{-1}$, $H_0 = 100 \text{ m}$, $s = 10^{-7} \text{ m}^{-1}\text{s}^{-1}$.