## Freshening of Antarctic Intermediate Water in the South Atlantic Ocean in 2005 - 2014

2 Wenjun Yao<sup>a,\*</sup>, Jiuxin Shi<sup>a</sup>, Xiaolong Zhao<sup>a</sup>

<sup>3</sup> <sup>1</sup>Key Lab of Physical Oceanography (Ocean University of China), Ministry of Education,

4 Qingdao 266100, Shandong, China

5 *Correspondence to:* E-mail address: wjimyao@gmail.com (Wenjun Yao)

6 Abstract

7 Basin-scale freshening of Antarctic Intermediate Water (AAIW) is reported to have 8 occurred in the South Atlantic Ocean during the period from 2005 to 2014, as shown by the 9 gridded monthly means Argo (Array for Real-time Geostrophic Oceanography) data. This 10 phenomenon was also revealed by two repeated transects along a section at 30° S, performed 11 during the World Ocean Circulation Experiment Hydrographic Program. Freshening of the 12 AAIW was compensated by a salinity increase of thermocline water, indicating a hydrological 13 cycle intensification. This was supported by the precipitation less evaporation change in the 14 Southern Hemisphere from 2000 to 2014. Freshwater input from atmosphere to ocean surface 15 increased in the subpolar high-precipitation region and vice versa in the subtropical 16 high-evaporation region. Against the background of hydrological cycle changes, a decrease in 17 the transport of Agulhas Leakage (AL) which was revealed by the simulated velocity field, 18 was proposed to be a contributor to the associated freshening of AAIW. This estimated 19 variability of AL was inferred from a weakening of wind stress over the South Indian Ocean 20 since the beginning of the 2000s, which would facilitate freshwater input from the source 21 region and partly contribute to the observed freshening of AAIW. The mechanical analyses 22 used in this study are qualitative, but it is contended that this study would be helpful to 23 validate and test predictably coupled sea-air model simulations.

24

4 Keywords: Freshening; Antarctic Intermediate Water; South Atlantic; Agulhas Leakage;

25 Wind Stress

<sup>&</sup>lt;sup>a</sup> Key Lab of Physical Oceanography (Ocean University of China), Ministry of Education, Qingdao 266100, Shandong, China

 <sup>\*</sup> Corresponding author at: Key Lab of Physical Oceanography (Ocean University of China), Ministry of Education, 238 Shongling Road, Laoshan District, Qingdao 266100, Shandong, China. Mob: +86 151 5421 9251.
 E-mail address: wjimyao@gmail.com (Wenjun Yao)

# 26 1 Introduction

Thermocline and intermediate waters play an important part in global overturning
circulation by ventilating the subtropical gyres in different parts of the world oceans [*Sloyan and Rintoul*, 2001]. They also constitute the northern limb of the Southern Hemisphere
supergyre [*Ridgway and Dunn*, 2007; *Speich et al.*, 2002].

31 Previous studies have addressed the variability of intermediate water. Wong et al. [2001] 32 found that the intermediate water had freshened between the 1960s and the period 1985-94 in 33 the Pacific Ocean. Bindoff and McDougall [2000] reported that there had been freshening of 34 water between 500 and 1500 db from 1962 to 1987 along 32° S in the Indian Ocean. Curry et 35 al. [2003] showed a salinity reduction on the isopycnal surface of intermediate water for the 36 period 1950s - 1990s in the western Atlantic. The freshening variability can be traced back to 37 the signature of water in the formation regions [Church et al., 1991]. The freshening 38 examples given above are in agreement with changes in the hydrological cycle, in which the 39 wet (precipitation > evaporation, P > E dominance) subpolar regions have been getting wetter 40 and vice versa for the dry (P<E dominance) subtropical regions over the last 50 years [Held 41 and Soden, 2006; Skliris et al., 2014].

42 Antarctic Intermediate Water (AAIW) is characterized by a salinity minimum (core of 43 AAIW) and is centered at the depths of 600 m and 1000 m (Fig. 1), which lies within 44 potential density (reference to sea surface)  $\sigma_0 = 27.1 - 27.3 \text{ kg/m}^3$  [Piola and Georgi, 1982]. 45 The AAIW is found from just north of the Subantarctic Front (SAF) [Orsi et al., 1995] in the 46 Southern Ocean and can be traced as far as 20° N [Talley, 1996]. It is generally accepted that 47 the variability of AAIW is largely controlled by air-sea-ice interaction [Close et al., 2013; 48 Naveira Garabato et al., 2009; Santoso and England, 2004], but the argument about its origin 49 and formation process continues. For example, there is the circumpolar formation theory of 50 AAIW along the SAF, through mixing with Antarctic Surface Water (AASW) along 51 isopycnal [Fetter et al., 2010; Sverdrup et al., 1942]. Alternatively, it has been proposed that 52 there is a local formation of AAIW in specific regions, as a bi-product of Subantarctic Mode

Water (SAMW) relating to deep convection [*McCartney*, 1982; *Piola and Georgi*, 1982]. The
first standpoint states that the AAIW are primarily derived from entire subpolar sources,
meanwhile the second one emphasizes the role that air-sea interaction plays in the oceans
south of South America.

57 In the South Atlantic, AAIW constitutes the return branch of the Meridional Overturning 58 Circulation (MOC) [Donners and Drijfhout, 2004; Speich et al., 2007; Talley, 2013]. As an 59 open ocean basin, the South Atlantic is fed by two different sources of AAIW [Sun and Watts, 60 2002]. The first is younger, fresher and has a lower apparent oxygen utilization (AOU) and 61 originates from the Southeast Pacific [McCartney, 1977; Talley, 1996] and the winter waters 62 west of Antarctic Peninsula [Naveira Garabato et al., 2009; Santoso and England, 2004]. These source regions of AAIW are mostly dominated by the net surface freshwater flux from 63 64 atmosphere to ocean (P > E), which facilitates the freshening of AAIW with time. The second 65 is the older, saltier and higher AOU AAIW which comes from Indian Ocean, transported by 66 Agulhas Leakage (AL) as Agulhas rings (Fig. 2). The mixture of the above two types of 67 AAIW can lead to a transition of hydrographic properties across the subtropical South 68 Atlantic [Boebel et al., 1997].

The influence of AL on variability of AAIW in the South Atlantic has been demonstrated to be substantial [*Hummels et al.*, 2015; *Schmidtko and Johnson*, 2012], as 50 -60% of the Atlantic AAIW originates from the Indian Ocean [*Gordon et al.*, 1992; *G D McCarthy et al.*, 2012], with increased (decreased) transport of AL relating to salinification (freshening) of AAIW. AL has apparently increased during the period from 1950s to the early 2000s [*Durgadoo et al.*, 2013; *Lübbecke et al.*, 2015], but there have been no studies addressing the influence of AL on the AAIW in South Atlantic since 2000.

With the instigation of the Argo (Array for Real-time Geostrophic Oceanography) program, *in-situ* hydrographic observation has tremendously expanded since 2003 [*Roemmich et al.*, 2015], particularly in the Southern Ocean (SO) where historical data are sparse and intermittent. This decreases the uncertainty of estimates for the research on both seasonal and decadal variations of subsurface and intermediate waters. 81 The present work reports the freshening of AAIW in the South Atlantic over the
82 preceding decade (2005 - 2014). Against the background of an enhanced hydrological cycle,
83 decreased transport of AL contributes to such freshening and may be driven by a weakening
84 of wind stress in the South Indian Ocean during the same period.

#### 85 2 Data and Methods

Based on individual temperature (*T*) and salinity (*S*) profiles from Argo, International
Pacific Research Centre (IPRC) gridded monthly means data for period 2005 - 2014 are
produced using variational interpolation. The IPRC data have 27 levels from 0 to 2000 m
depth vertically, nominal 1°×1° grid globally and monthly temporal resolution.
(http://apdrc.soest.hawaii.edu/projects/Argo/data/gridded/On\_standard\_levels/index-1.html).

91 To reduce the error from low vertical resolution of data when computing the hydrographic 92 values on isopycnal surface, T and S profiles are first interpolated onto 1 m depth intervals 93 vertically using a spline instead of linear method. Because the IPRC data are interpolated 94 from randomly distributed Argo profiles, it is necessary to demonstrate the robust nature of 95 their signals by comparing with the other Argo gridded products. As a result, the Japan 96 Agency of Marine-Earth Science and Technology (JAMSTEC, [Hosoda et al., 2008]) T and S 97 data from 2005 to 2014 with 1° longitude and 1° latitude resolution are also collected for 98 comparison and verification. The number of Argo profiles is rapidly increasing year by year, 99 and part of their distribution has been outlined in previous studies, inter alia Hosoda et al. 100 [2008] and Roemmich et al. [2015].

101 Two hydrographic cruises of repeated transects along 30° S were conducted in the World 102 Ocean Circulation Experiment (WOCE) Hydrographic Program 103 (http://www.nodc.noaa.gov/woce/wdiu/diu summaries/whp/index.htm). Their locations are 104 presented in Fig. 2. The first transect consisted of 72 stations in 2003 by the R/V Mirai (Japan, 105 [Kawano et al., 2004]), the second was in 2011 with 81 stations sampled from the Ronald H. 106 Brown (United States, [Feely et al., 2011]) 107 (http://www.nodc.noaa.gov/woce/wdiu/diu summaries/whp/index.htm). These two transects

108 were not only performed in almost repeated positions in the subtropical South Atlantic, but 109 also conducted in the same season (Nov and Oct respectively). Furthermore, the investigation 110 time interval between the two sections from Nov 2003 to Oct 2011 is very similar to the IPRC 111 data (Jan 2005 - Dec 2014) and can therefore be used to confirm those results.

112 To reduce the effect of dynamic processes in the ocean interior (i.e. mesoscale eddies 113 and internal waves), the investigation of halocline variation should be along neutral density 114 surfaces [*G McCarthy et al.*, 2011; *McDougall*, 1987]. The layer of AAIW is defined using 115 neutral density ( $\gamma^n$ , unit: kg/m<sup>3</sup>) [*Jackett and McDougall*, 1997] instead of potential density, 116 with the upper and lower boundaries being 27.1 $\gamma^n$  and 27.6 $\gamma^n$  [*Goes et al.*, 2014], respectively.

117 Monthly 10m wind fields between years 1980 and 2014 from the ERA-Interim archive at 118 the European Centre for Medium Range Weather Forecasts (ECMWF) 119 (http://apps.ecmwf.int/datasets/data/interim-full-daily/levtype=sfc/) are used to display the 120 decadal variability of wind stress (WS) over the South Indian Ocean. Another reanalysis wind 121 product of NCEP2 (National Centers for Environmental Prediction-Department of Energy 122 Atmospheric Model Intercomparison Project reanalysis 2, NCEP-DOE AMIP Reanalysis-2, 123 http://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis2.html) is also used for the 124 period 1980-2014. Additionally, the satellite-derived wind products of QuikSCAT for 125 2000-2007 and ASCAT for 2008-2014 (Quick Scatterometer and Advanced Scatterometer, 126 both in ftp://ftp.ifremer.fr/ifremer/cersat/products/gridded/MWF/L3/) are used to compare and 127 verify the decadal variability of WS revealed by the ERA-Interim wind product. In this work, 128 the WS over open ocean is calculated from 10 m wind field data using the equation adopted in 129 Trenberth et al. [1989].

Reanalysis data including precipitation (P) and evaporation (E) from ERA-Interim are
used to reveal the freshwater input from the atmosphere to ocean surface in the preceding
decade.

133 The up-to-date SODA3.3.1 (Simple Ocean Data Assimilation,
134 http://www.atmos.umd.edu/~ocean/), which is forced by MERRA2 (The Modern-Era
135 Retrospective analysis for Research and Applications, Version 2), spans the 36-year period

136 1980-2015 ([*Carton et al.*, 2016]). The global simulated velocity field at specified depths
137 make it available to evaluate the transport of AL.

## 138 3 Freshening of Antarctic Intermediate Water

# 139 3.1 Freshening observed from Argo gridded products

140 The Argo gridded products provide a globally distributed and continuous time series of T141 and S profiles down to 2000 m ocean depth. The present work focuses on the AAIW in the 142 South Atlantic Basin (Fig. 2, Region A), which encompasses most of the subtropical gyre and 143 a part of the tropical regimes [Boebel et al., 1997; Talley, 1996]. Computed from the Argo 144 gridded data of IPRC, the biennial mean of  $\theta$ -S diagram (Fig. 3a) clearly shows that the 145 AAIW has experienced a process of progress of basin-scale freshening during the period from 146 Jan 2005 to Dec 2014. The linear trend of salinity (Fig. 3b) further reveals that the freshening 147 takes up most of the AAIW layer but with a little salinification in the deeper part of it. Except 148 around the 27.42 $\gamma^n$  neutral density surface, the AAIW variation is significant at 95% 149 confidence level, using the F-test criteria. In comparison with Fig. 3a, it was found that the 150 cut-off point of transformation from salinity decrease to increase is near the salinity minimum. 151 Above the salinity minimum, the shift of  $\theta$ -S trends towards cooler and fresher values along 152 density surface and seem to be a response to the warming and freshening of surface waters 153 where AAIW ventilates. Such thermohaline change has also been found in the Pacific and 154 Indian oceans over a different time period [Wong et al., 1999]. Church et al. [1991] and 155 Bindoff and Mcdougall [1994] has researched the counterintuitive cooling of AAIW 156 temperature induced by warming of surface water. They showed that a warming parcel at 157 mixed layer would subduct further equatorward, which would lead the S- $\theta$  curve to become 158 cooler and fresher at a given density. The salinity decrease of core of AAIW indicates that 159 such change can only be induced by freshwater input from the source region, as mixing with 160 more saline surrounding waters cannot give rise to a salt loss in the salinity minimum layer.

To demonstrate the robustness of AAIW variations revealed by IPRC data, re-plots of Fig. 3a-b using another Argo gridded product, the JAMSTEC, are also shown for comparison (see Supplementary 1, only AAIW layer shown). Not only the same variation along density surfaces in the AAIW layer found, but also a freshening of the salinity minimum. Both the isoneutral salinity increases of IPRC and JAMSTEC data below the salinity minimum are quite small. The main discrepancy between them is that the degree of freshening in the JAMSTEC data is somewhat less than IPRC and at a higher 95% confidence level.

168 The freshwater gain for the basin-scale salinity decrease of AAIW (mean salinity difference of 0.012 between  $27.1\gamma^n$  and  $27.6\gamma^n$  over a mean water mass thickness of 500 m) is 169 estimated at 15mm yr<sup>-1</sup> in its source region. However, the depth-integrated salinity change 170 171 over the water column (between  $26.6\gamma^n$  and  $27.6\gamma^n$ ) is 0.0014; since a salinity increase of 172 thermocline water balances the observed freshening of AAIW. This salinity budget implies 173 contemporary hydrological cycle intensification in the southern hemisphere, which is 174 illustrated by the P minus E change from 2000 to 2014, with P-E increasing in the subpolar 175 region and vice versa in the subtropical region (Fig. 4a). In these cases, the thermocline 176 (intermediate) water that ventilates in the high-evaporation (precipitation) subtropical 177 (subpolar) regions gets more saline (freshened), as shown by the hydrographic observations 178 (Fig. 3b).

Against the background of hydrological cycle augmentation, the annual freshwater input in AAIW ventilation region during the freshening period increased by 0.2 mm day<sup>-1</sup>, about 17% of the *P-E* in 2005 (Fig. 4b). It is considered that, the increase of *P-E* began in 1992, but there was a significant increase around 2003 (Fig. 4b, 5-yr running mean line), which means the observed freshened AAIW could be traced back to 2003. Though it was not possible to compute the direct freshwater input to the South Atlantic Basin in this study, the Argo era freshening of AAIW is qualitatively consistent with the freshwater gain in its source region.

186 3.2

#### Freshening in the quasi-synchronous WOCE CTD observations

187 Here two synoptic transatlantic sections from WOCE hydrographic program were used188 to explore the decadal freshening signal identified in the above subsection. Similar to Fig. 3a,

189 sectional mean  $\theta$ -S diagram (Fig. 5a) displays the same shift of thermohaline values, 190 including freshening of the salinity minimum, salinity reduction in the upper AAIW layer and 191 *vice versa* in the lower layer. Compared to the  $\theta$ -S curves of IPRC data (Fig. 3a), the curves of 192 WOCE (Fig. 5a) seem to be, in general, cooler  $\theta$  and fresher S. It is suggested that this is 193 because the IPRC mean is weighted towards the warmer and saltier waters in the north.

194 Unlike the Argo gridded product which has continuous time series of T and S data, there 195 are only two sections in the WOCE observations. Instead of calculating the linear trend of 196 salinity (as was done with the IPRC data), the difference of salinity observed in 2003 and 197 2011 is estimated (Fig. 5b). The light grey shading denotes the 95% confidence interval using 198 simple *t*-test criteria and having consider the number of degrees of freedom. Above the 199 salinity minimum, the freshening of AAIW revealed by IPRC and WOCE data are quite 200 similar, with the maximum appearing near  $27.2\gamma^n$ . Because the last WOCE observation 201 terminated in 2011 and the salinity reduction would continue at least up to 2014 as displayed 202 in Fig. 3a, the magnitude of the freshening in WOCE (Fig. 5b) is smaller than IPRC (Fig. 3b). 203 In the water layer below the salinity minimum (around  $27.41\gamma^n$ ), the salinity increase shown 204 in the WOCE data is relatively large (Fig. 5b). This is thought to be because the salinity rise 205 reached its maximum around 2011, which is shown in the time series of basinwide averaged 206 salinity on 27.45 $\gamma^n$  and 27.55 $\gamma^n$  density surfaces (see Supplementary 2).

For the salinification of thermocline water, there is a large discrepancy between IPRC and WOCE data, on neutral density surfaces  $26.6-26.7\gamma^n$  (Fig. 5b). It is considered that this would not affect the salinity budget over the water column (Fig. 5b), giver that the salt gain of thermocline water would balance the observed freshened AAIW. In conclusion, the general trend and consistency of the detail therein of the salinity change over the last ten-year time period revealed by IPRC and WOCE data leads us to state that the freshening of AAIW is a robust finding and valid.

# 214 4 Decrease of Agulhas Leakage transport

215 AAIW in the South Atlantic is largely influenced by AL through the intermittent pinching off of Agulhas rings (Fig. 2) [Beal et al., 2011], transferring salty thermocline and 216 217 intermediate water from the Indian Ocean to the South Atlantic [De Ruijter et al., 1999]. The 218 above discussion suggests that the freshening of AAIW is induced by the input of freshwater 219 from the source regions, which are consisted of the southeast Pacific Ocean and the 220 circumpolarly subpolar oceans (see introduction). As a result, if the transport of more saline 221 water from the Indian Ocean decreased, it would promote the effect of this freshwater 222 increase. In this section, the decrease of AL transport was evaluated by depth integration of 223 velocity field and further demonstrated by using an indirect indicator.

# 224 4.1 Evaluation from SODA velocity

225 In the study of modeling, it is widely acceptable to quantify the leakage follows a 226 Lagrangian approach [Biastoch et al., 2009; van Sebille et al., 2009]. Here a simplified 227 strategy was employed to compute the leakage by integrating the velocity within AAIW layer 228 (approximately between 610 and 1150m, according to Fig. 1), which was demonstrated to 229 result to a similar quantification to the Lagrangian one [Le Bars et al., 2014]. The depth 230 integration is along the GoodHope section (green line in Fig. 2), using the cross-component 231 velocity. Note that the leakage calculation is from the continent to the zero line of barotropic 232 streamfunction, which is the separation of the Agulhas regime and the Antarctic Circumpolar 233 Current [Biastoch et al., 2015].

Before showing the transport computed from the SODA velocity data, it is necessary to verify that the SODA hydrographic data could reveal the same freshening of AAIW as the other dataset done. And in consequence, the AAIW in the South Atlantic was also shown to have freshened during period 2005-2014, though with relatively small magnitude (Supplementary 3). Yearly leakage computation within AAIW layer was employed for the period 2000-2015 (Fig. 6). It shows that the leakage in the early 2010s is smaller than that in the middle and post 2000s, forming a decreased trend in a nearly ten-year period. This
estimation of leakage seems to be consistent with the below indirect estimation of AL
transport.

# 243 4.2 Weakening of the westerlies in the South Indian Ocean

244 Continuous measurements of the AL transport have never been realized so far. The 245 earlier study suggested that an increased AL transport correlates well with a poleward shift of 246 westerlies [Beal et al., 2011]. However, after using reanalysis and climate models, Swart and 247 Fyfe [2012] argued that strengthening of Southern Hemisphere surface westerlies has 248 occurred without major transgressions in its latitudinal position over the period 1979-2010, 249 during which period the AL has largely increased [Biastoch et al., 2009]. A more recent study 250 of Durgadoo et al. [2013] showed that the increase of AL is concomitant with equatorward 251 rather than poleward shift of westerlies in their simulation cases. They also concluded that the 252 intensity of westerlies is predominantly responsible in controlling this Indian-Atlantic 253 transport. Many relevant studies agreed on this relationship, that the enhancement of 254 westerlies intensity is related to the increase of AL [Goes et al., 2014; Lee et al., 2011; 255 Loveday et al., 2015].

256 The AL corresponds most significantly to westerlies strength averaged over the Indian 257 Ocean in contrast to that averaged circumpolarly or locally [Durgadoo et al., 2013]. 258 According to the work of Durgadoo et al. [2013], zonally averaged WS was calculated from 259 the wind product of ERA-Interim over the Indian Ocean (20-110° E) for every 5-vr period 260 since 1980 (Fig. 7a and d). Previous studies [Lee et al., 2011; Loveday et al., 2015] have 261 found that the WS has considerably increased from 1980s to the beginning of 2000s (Fig. 7d), 262 consistent with the contemporary increase of AL transport. Though there are oscillations 263 during 1990s, the WS reached its peak around the years 2000-2004 (Fig. 7d). then began to 264 decline. It can be concluded that the WS has weakened for period 2000 - 2014 (Fig. 7d), 265 which implies a concurrent decrease of AL transport.

In addition to the ERA-Interim wind data, we have further checked the zonally averaged
WS over the Indian Ocean (20-110° E), using another reanalysis product of NCEP2 (Fig. 7b

and e) and the combined QuikSCAT-ASCAT (Fig. 7c and f) satellite-derived wind products. The three zonally averaged WS agree that during the period 2000-2014, the westerlies reached a peak in the years 2000-2004, and then progressively subsided through 2005-2009 to 2010-2014. The process of gradual decline of WS is most pronounced in the NCEP2 data. It is noteworthy that none of the three products show a significant meridional shift of the latitude of maximum WS from 2000 to 2014, in corroboration with the conclusion of *Swart and Fyfe* [2012].

275 4.3 H

#### **3** Evidence from other works

276 Many efforts have been made to estimate AL transport, especially using model 277 simulations [Lübbecke et al., 2015; Loveday et al., 2015]. In recent years, Le Bars et al. [2014] 278 provided the time series of AL transport over the satellite altimeter era, computed from 279 absolute dynamic topography data, which can show the decadal variation of AL present. In 280 their result (Figure 8 in Le Bars et al. [2014]), the anomalies of AL from satellite altimetry 281 reached a peak around 2003 (annual average), and then began to subside, apart from a 282 mid-2011 increase. In addition, their negative trend of AL (Figure 9 in Le Bars et al. [2014]) 283 over the period from Oct 1992 to Dec 2012 indicates that the transport was reduced during the 284 2000s in contrast to the 1990s. Another study by Biastoch et al. [2015] should be of help in 285 the present discussion. Though the time series of AL obtained from models didn't show a 286 distinct decline of AL transport in the last decade, which seems partly due to the data filter 287 applied and the end of time series in 2010 (Figure 4 in *Biastoch et al.* [2015]), it displays a 288 maximum of salt transport around 2000 (Figure 5 in Biastoch et al. [2015]). This peak and the 289 subsequent decline of salt transport are consistent with the freshening of AAIW over the 290 similar time period considered here.

Thus, in addition to the freshwater input that gives rise to the salt loss of the AAIW in the South Atlantic Ocean, reduced transport of AL or salt will further enhance this signal. Unfortunately, the analysis of the contribution from both the source region and AL is qualitative. Future work should be focused on quantification of each factor based on model simulations.

### 297 5 Conclusions

The analysis of IPRC gridded data shows that AAIW in the South Atlantic has experienced basin-scale freshening for the period from Jan 2005 to Dec 2014 (Fig. 3a and b), with freshwater input estimated at 15 mm yr<sup>-1</sup> in its source region. Two transects of WOCE hydrographic program observed in 2003 and 2011 also reveal the above variation of AAIW in the last decade (Fig. 3c and d).

This freshening in the intermediate water layer is thought to be compensated by increased salinity in shallower thermocline water, indicating a contemporary intensification of hydrological cycle (Fig. 3b and Fig. 5b). In this case the freshwater input from atmosphere to ocean surface increased in the subpolar high-precipitation region and *vice versa* in the subtropical high-evaporation region (Fig. 4a). Over the last ten year time period, significant freshwater gain began around 2003 (Fig. 4b), suggesting the observed freshened AAIW could be traced back to this time.

310 Against the background of hydrological cycle intensification, the decrease of AL 311 transport is proposed to contribute to the freshening of AAIW in the South Atlantic, 312 associated with a weakening of westerlies over the South Indian Ocean. This decrease was 313 revealed by the leakage evaluation along the GoodHope section. The mechanical analysis 314 shows that the WS over the South Indian Ocean reached its peak around 2000-2004 and 315 began to subside through 2005-2009 to 2010-2014 (Fig. 7), reversing its increasing phase 316 from 1950s to the beginning of 2000s, during which period the AL had increased [Durgadoo 317 et al., 2013; Lübbecke et al., 2015]. This indirectly estimated variability of AL is consistent 318 with other studies covering a similar period [Biastoch et al., 2015; Le Bars et al., 2014]. As 319 the AAIW carried by AL is more saline relative to its counterpart in the South Atlantic Ocean, 320 its decrease would promote the effect of freshwater input from the source region, contributing 321 to the observed freshening.

Both the analyses of freshwater input and reduced transport in the AL reported here are qualitative but not quantitative. The purpose of this work is to reveal the decadal freshening of AAIW in the South Atlantic Ocean over the last ten-year time period, and suggest a contributing mechanism. Future work should be focused on the quantification of these two contributors, and the influence they have on the world ocean circulation.

- 327
- 328 Acknowledgements
- 329 This study is supported by the Chinese Polar Environment Comprehensive Investigation330 and Assessment Programs (Grant nos. CHINARE-04-04, CHINARE-04-01).
- 331



332 Captions of Figures

333

Fig. 1 WOCE salinity sections along 30° S in the South Atlantic Ocean (positions shown in Fig. 2) observed in (a) 2003 and (b) 2011. Overlaid white solid-dotted lines are  $\gamma^n$  surfaces ranging from 26.9 to 27.5 kg/m<sup>3</sup>, with 0.2 kg/m<sup>3</sup> interval.



338

Fig. 2 Bathymetry of the South Indian-Atlantic oceans. Color shading is ocean depth. Red box
delineates the area for the basinwide average of gridded data (hereafter refers to Region A).
The green line shows the GoodHope section which is used to calculate the leakage transport
to the South Atlantic. Magenta stars represent transatlantic CTD stations measured in 2003,
meanwhile blue dots in 2011. The Agulhas Current, Retroflection, Agulhas Return Current
and Agulhas Leakage (as eddies) are also shown and ticked.





Fig. 3 (a) Biennial mean  $\theta$ -S diagram averaged over Region A for IPRC data with  $\gamma^n$  surfaces superimposed (grey solid-dotted lines). The inserted figure is the magnification of the area delineated by cyan solid-dotted box. The corresponding time for each  $\theta$ -S curve is listed in their bottom-right corner (i.e. 05/06 for 2005-2006). (b) Salinity trend along  $\gamma^n$  surfaces for period Jan 2005 – Dec 2014 is displayed by the thick black line, and the 95% confidence intervals (*F*-test) are represented by the light grey shadings, calculated from IPRC data.

353



Fig. 4 Calculated from ERA-Interim precipitation and evaporation data: (a) Zonally mean (ocean areas only) of annually *P-E* (freshwater input, mm day<sup>-1</sup>), each line represents a 5-yr averaged result. The corresponding time period (i.e. 00-04 for 2000-2004) is listed in the bottom-left corner. (b) Time series of annually *P-E* averaged over the oceans in 45-65° S, 0-360° E band from 1979 to 2014 (blue star), and its 5-yr running mean (black).



Fig. 5 (a) The same as Fig. 3a but for sectional mean of WOCE hydrographic casts. The corresponding year for each  $\theta$ -S curve is listed in their bottom-right corner. (b) Sectional mean differences (thick black line) of WOCE hydrographic data along  $\gamma^n$  and their 95% confidence intervals (grey shadings, *t*-test).



Fig. 6 Computation of Agulhas Leakage transport anomaly from the SODA velocity field
along the Goodhoop Line. Note that the depth integration is only for the AAIW layer.

- 373
- 374



375

Fig. 7 Zonally averaged wind stress calculated from the wind product of (a) ERA-Interim, (b)
NCEP2 and (c) ASCAT/QSCAT over the Indian Ocean (20° E-110° E) for different periods
(i.e. 80-84 for Jan 1980 - Dec 1984; 00-04 for Jan 2000 - Dec 2004) listed in the top-right
corners. (d), (e) and (f) are the magnification of cyan boxes in (a), (b) and (c), respectively.

**380** REFERENCES

Beal, L. M., W. P. De Ruijter, A. Biastoch, and R. Zahn (2011), On the role of the Agulhas
system in ocean circulation and climate, *Nature.*, 472(7344), 429-436, doi:
10.1038/nature09983.

Biastoch, A., C. W. Böning, F. U. Schwarzkopf, and J. Lutjeharms (2009), Increase in
Agulhas leakage due to poleward shift of Southern Hemisphere westerlies, *Nature.*,
462(7272), 495-498, doi: 10.1038/nature08519.

Biastoch, A., J. V. Durgadoo, A. K. Morrison, E. van Sebille, W. Weijer, and S. M. Griffies
(2015), Atlantic multi-decadal oscillation covaries with Agulhas leakage, *Nature communications*, 6, doi: 10.1038/ncomms10082.

Bindoff, N. L., and T. J. Mcdougall (1994), Diagnosing climate change and ocean ventilation
using hydrographic data, *J. Phys. Oceanogr.*, 24(6), 1137-1152, doi:
10.1175/1520-0485(1994)024<1137:DCCAOV>2.0.CO;2.

Bindoff, N. L., and T. J. McDougall (2000), Decadal changes along an Indian Ocean section
at 32 S and their interpretation, J. Phys. Oceanogr., 30(6), 1207-1222, doi:

395 <u>http://dx.doi.org/10.1175/1520-0485(2000)030<1207:DCAAIO>2.0.CO;2</u>.

- Boebel, O., C. Schmid, and W. Zenk (1997), Flow and recirculation of Antarctic intermediate
  water across the Rio Grande rise, J. Geophys. Res. Oceans (1978–2012), 102(C9),
  20967-20986, doi: 10.1029/97JC00977.
- Carton, J. A., G. A. Chepurin, and L. Chen (2016), An updated reanalysis of ocean climateusing the Simple Ocean Data Assimilation version 3 (SODA3), *manuscript in preparation*.
- 401 Church, J. A., J. S. Godfrey, D. R. Jackett, and T. J. McDougall (1991), A model of sea level
  402 rise caused by ocean thermal expansion, *J. Climate.*, 4(4), 438-456, doi:
  403 http://dx.doi.org/10.1175/1520-0442(1991)004<0438:AMOSLR>2.0.CO:2.
- 404 Close, S. E., A. C. Naveira Garabato, E. L. McDonagh, B. A. King, M. Biuw, and L. Boehme
- 405 (2013), Control of mode and intermediate water mass properties in Drake Passage by the
  406 Amundsen Sea Low, J. Climate., 26(14), 5102-5123, doi: 10.1175/JCLI-D-12-00346.1.
- 407 Curry, R., B. Dickson, and I. Yashayaev (2003), A change in the freshwater balance of the
  408 Atlantic Ocean over the past four decades, *Nature.*, 426(6968), 826-829, doi:
  409 10.1038/nature02206.
- 410 De Ruijter, W., A. Biastoch, S. Drijfhout, J. Lutjeharms, R. Matano, T. Pichevin, P. Van
- 411 Leeuwen, and W. Weiger (1999), Indian-Atlantic interocean exchange: Dynamics, estimation
- 412 and impact, J. Geophys. Res., 104(C9), 20885-20910.
- 413 Donners, J., and S. S. Drijfhout (2004), The Lagrangian view of South Atlantic interocean
- 414 exchange in a global ocean model compared with inverse model results, J. Phys. Oceanogr.,
  415 34(5), 1019-1035, doi:
- 416 <u>http://dx.doi.org/10.1175/1520-0485(2004)034<1019:TLVOSA>2.0.CO;2</u>.
- 417 Durgadoo, J. V., B. R. Loveday, C. J. C. Reason, P. Penven, and A. Biastoch (2013), Agulhas
- 418 Leakage Predominantly Responds to the Southern Hemisphere Westerlies, J. Phys. Oceanogr.,
- **419** *43*(10), 2113-2131, doi: 10.1175/JPO-D-13-047.1.
- 420 Feely, R. A., R. Wanninkhof, S. Alin, M. Baringer, and J. Bullister (2011), Global Repeat
- 421 Hydrographic/CO2/Tracer surveys in Support of CLIVAR and Global Cycle objectives:422 Carbon Inventories and Fluxes.
- 423 Fetter, A., M. Schodlok, and V. Zlotnicki (2010), Antarctic Intermediate Water Formation in a
- 424 High-Resolution OGCM, paper presented at EGU General Assembly Conference Abstracts.
- Goes, M., I. Wainer, and N. Signorelli (2014), Investigation of the causes of historical
  changes in the subsurface salinity minimum of the South Atlantic, *J. Geophys. Res. Oceans*,
- **427** *119*(9), 5654-5675, doi: 10.1002/2014JC009812.
- 428 Gordon, A. L., R. Weiss, W. M. Smethie Jr, and M. J. Warner (1992), Thermociine and
- 429 Intermediate Water Communication, J. Geophys. Res., 97(C5), 7223-7240, doi:
- **430** 10.1029/92JC00485.
- Held, I. M., and B. J. Soden (2006), Robust responses of the hydrological cycle to global
- 432 warming, J. Climate., 19(21), 5686-5699, doi: <u>http://dx.doi.org/10.1175/JCLI3990.1</u>.
- 433 Hosoda, S., T. Ohira, and T. Nakamura (2008), A monthly mean dataset of global oceanic
- temperature and salinity derived from Argo float observations, *JAMSTEC Rep. Res. Dev.*, 8,
  435 47-59, doi: http://doi.org/10.5918/jamstecr.8.47.
- 436 Hummels, R., P. Brandt, M. Dengler, J. Fischer, M. Araujo, D. Veleda, and J. V. Durgadoo
- 437 (2015), Interannual to decadal changes in the western boundary circulation in the Atlantic at

- 438 11° S, Geophys. Res. Lett., 42(18), 7615-7622, doi: 10.1002/2015GL065254.
- 439 Jackett, D. R., and T. J. McDougall (1997), A neutral density variable for the world's oceans, *J.* 440 *Phys. Oceanogr.*, *27*(2), 237-263, doi: 441 http://dx.doi.org/10.1175/1520-0485(1997)027<0237:ANDVFT>2.0.CO;2.
- 442 Kawano, T., H. Uchida, W. Schneider, Y. Kumamoto, A. Nishina, M. Aoyama, A. Murata, K.
- Sasaki, Y. Yoshikawa, and S. Watanabe (2004), Cruise Summary of WHP P6, A10, I3 and I4
  Revisits in 2003, paper presented at AGU Fall Meeting Abstracts.
- 445 Lübbecke, J. F., J. V. Durgadoo, and A. Biastoch (2015), Contribution of increased Agulhas
  446 leakage to tropical Atlantic warming, *J. Climate.*, 28(24), 9697-9706, doi:
  447 http://dx.doi.org/10.1175/JCLI-D-15-0258.1.
- Le Bars, D., J. V. Durgadoo, H. A. Dijkstra, A. Biastoch, and W. P. M. De Ruijter (2014), An
  observed 20-year time series of Agulhas leakage, *Ocean. Sci.*, 10(4), 601-609, doi:
  10.5194/os-10-601-2014.
- 451 Lee, S. K., W. Park, E. van Sebille, M. O. Baringer, C. Wang, D. B. Enfield, S. G. Yeager, and
- 452 B. P. Kirtman (2011), What caused the significant increase in Atlantic Ocean heat content
- 453 since the mid 20th century?, *Geophys. Res. Lett.*, *38*(17), doi: 10.1029/2011GL048856.
- 454 Loveday, B., P. Penven, and C. Reason (2015), Southern Annular Mode and westerly -
- wind driven changes in Indian Atlantic exchange mechanisms, *Geophys. Res. Lett.*, 42(12),
  4912-4921, doi: 10.1002/2015GL064256.
- McCarthy, G., E. McDonagh, and B. King (2011), Decadal variability of thermocline and
  intermediate waters at 24 S in the South Atlantic, *J. Phys. Oceanogr.*, 41(1), 157-165, doi:
  10.1175/2010JPO4467.1.
- 460 McCarthy, G. D., B. A. King, P. Cipollini, E. L. McDonagh, J. R. Blundell, and A. Biastoch
  461 (2012), On the sub-decadal variability of South Atlantic Antarctic Intermediate Water,
  462 *Geophys. Res. Lett.*, 39, L10605, doi: 10.1029/2012GL051270.
- 463 McCartney, M. S. (1977), Subantarctic Mode Water, in Angel, M.V. (Ed.), A Voyage of
- 464 Discovery: George Deacon 70th Anniversary Volume (Suppl. To Deep-Sea Res.). Pergamon,
- 465 edited, pp. 103-119, Woods Hole Oceanographic Institution.
- 466 McCartney, M. S. (1982), The subtropical recirculation of mode waters, J. Mar. Res., 40,
  467 427-464.
- 468 McDougall, T. J. (1987), Neutral surfaces, J. Phys. Oceanogr., 17(11), 1950-1964, doi:
   469 <u>http://dx.doi.org/10.1175/1520-0485(1987)017<1950:NS>2.0.CO;2</u>.
- 470 Naveira Garabato, A. C., L. Jullion, D. P. Stevens, K. J. Heywood, and B. A. King (2009),
- 471 Variability of Subantarctic Mode Water and Antarctic Intermediate Water in the Drake
- 472 Passage during the Late-Twentieth and Early-Twenty-First Centuries, J. Climate., 22(13),
- **473** 3661-3688, doi: 10.1175/2009jcli2621.1.
- 474 Orsi, A. H., T. Whitworth Iii, and W. D. Nowlin Jr (1995), On the meridional extent and fronts
  475 of the Antarctic Circumpolar Current, *Deep-Sea Res. I.*, 42(5), 641-673, doi:
  476 <a href="http://dx.doi.org/10.1016/0967-0637(95)00021-W">http://dx.doi.org/10.1016/0967-0637(95)00021-W</a>.
- 477 Piola, A. R., and D. T. Georgi (1982), Circumpolar properties of Antarctic intermediate water
- **478** and Subantarctic Mode Water, *Deep-Sea Res. A.*, 29(6), 687-711, doi:
- **479** 10.1016/0198-0149(82)90002-4.
- 480 Ridgway, K. R., and J. R. Dunn (2007), Observational evidence for a Southern Hemisphere

- 481 oceanic supergyre, Geophys. Res. Lett., 34, L13612, doi: 10.1029/2007gl030392.
- 482 Roemmich, D., J. Church, J. Gilson, D. Monselesan, P. Sutton, and S. Wijffels (2015),
- 483 Unabated planetary warming and its ocean structure since 2006, Nature Clim. Change, 5(3), 484 240-245, doi: 10.1038/nclimate2513.
- 485 Santoso, A., and M. H. England (2004), Antarctic Intermediate Water circulation and 486 variability in a coupled climate model, J. Phys. Oceanogr., 34(10), 2160-2179, doi: 487 http://dx.doi.org/10.1175/1520-0485(2004)034<2160:AIWCAV>2.0.CO;2.
- 488 Schmidtko, S., and G. C. Johnson (2012), Multidecadal Warming and Shoaling of Antarctic 489 Intermediate Water\*, J. Climate., 25(1), 207-221, doi: 10.1175/jcli-d-11-00021.1.
- 490 Skliris, N., R. Marsh, S. A. Josey, S. A. Good, C. Liu, and R. P. Allan (2014), Salinity changes 491 in the World Ocean since 1950 in relation to changing surface freshwater fluxes, Climate 492 dynamics, 43(3-4), 709-736, doi: 10.1007/s00382-014-2131-7.
- 493 Sloyan, B. M., and S. R. Rintoul (2001), Circulation, Renewal, and Modification of Antarctic 494 Mode and Intermediate Water\*, J. Phys. Oceanogr., 31(4), 1005-1030, doi: 495 http://dx.doi.org/10.1175/1520-0485(2001)031<1005:CRAMOA>2.0.CO;2.
- 496 Speich, S., B. Blanke, and W. Cai (2007), Atlantic meridional overturning circulation and the
- 497 L23614, Southern Hemisphere supergyre, Geophys. Res. Lett., 34, doi:
- 498 10.1029/2007GL031583.
- 499 Speich, S., B. Blanke, P. de Vries, S. Drijfhout, K. Döös, A. Ganachaud, and R. Marsh (2002),
- 500 Tasman leakage: A new route in the global ocean conveyor belt, Geophys. Res. Lett., 29(10), 501 1416, doi: 10.1029/2001gl014586.
- 502 Sun, C., and D. R. Watts (2002), A view of ACC fronts in streamfunction space, Deep-Sea 503 Res. I., 49(7), 1141-1164, doi: 10.1016/S0967-0637(02)00027-4.
- 504 Sverdrup, H. U., M. W. Johnson, and R. H. Fleming (1942), The Oceans: Their physics, 505 chemistry, and general biology, Prentice-Hall New York.
- Swart, N., and J. Fyfe (2012), Observed and simulated changes in the Southern Hemisphere 506 507 surface westerly wind stress, Geophys. Res. Lett., *39*(L16711), doi: 508 10.1029/2012GL052810.
- 509 Talley, L. D. (1996), Antarctic intermediate water in the South Atlantic, in *The South Atlantic*, 510 edited, pp. 219-238, Springer.
- 511 Talley, L. D. (2013), Closure of the global overturning circulation through the Indian, Pacific,
- 512 and Southern Oceans: Schematics and transports, Oceanography, 26(1), 80-97, doi: 513 http://dx.doi.org/10.5670/oceanog.2013.07.
- 514 Trenberth, K. E., W. G. Large, and J. G. Olson (1989), The effective drag coefficient for 515 evaluating wind stress over the oceans, J. Climate., 2(12), 1507-1516, doi: 516
- http://dx.doi.org/10.1175/1520-0442(1989)002<1507:TEDCFE>2.0.CO;2.
- 517 van Sebille, E., A. Biastoch, P. Van Leeuwen, and W. De Ruijter (2009), A weaker Agulhas 518 Current leads to more Agulhas leakage, Geophys. Res. Lett., 36(3), doi: 519 10.1029/2008GL036614.
- 520 Wong, A. P., N. L. Bindoff, and J. A. Church (1999), Large-scale freshening of intermediate
- 521 waters in the Pacific and Indian Oceans, *Nature.*, 400(6743), 440-443, doi: 10.1038/22733.
- 522 Wong, A. P., N. L. Bindoff, and J. A. Church (2001), Freshwater and heat changes in the
- 523 North and South Pacific Oceans between the 1960s and 1985-94, J. Climate., 14(7),

524	1613-1633,	doi: htt	p://dx.doi.org	/10.1175/	1520-0442	(2001)	)014<1613:FA	HCIT>2.(	).CO;2.