

Interactive comment on “Wave spectral shapes in the coastal waters based on measured data off Karwar, west coast of India” by M. Anjali Nair and V. Sanil Kumar

Anonymous Referee #1

Received and published: 5 February 2017

General Comments

Generally, this work presents an impressive data set spanning 5 years. The data appear largely to be a new contribution to the literature, and if so, should qualify the work for publication as a case study. Having recognized that, there are several concerns with potential to change this assessment.

1) It appears that a significant portion of this data may have been published elsewhere. For example, here:

Glejin, J., Sanil Kumar, V., Amrutha, M.M. and Singh J., Characteristics of long-period swells measured in the in the near shore regions of eastern Arabian Sea, Int. J. Naval

Printer-friendly version

Discussion paper



Architecture and Ocean Engineering, 8, 312-319, 2016.

it seems clear that data from January 1 2011 through December 2012 were recorded at, or very near the location reported in this paper, with the same methods and perhaps instruments. The Glejin et al. paper present statistics that appear to overlap with statistics presented here.

2) Why is the broader-context not communicated? It seems peculiar that several existent publications by the authors use verbatim text describing the area, methods and motivation, as well as the same analysis techniques and products, but those works are referenced only narrowly in regard to specific details. For example, data collected over nearly the same period of record (March 23, 2010 to November 6, 2014) only a few kilometers up the coast, and with essentially the same analysis is reported here:

Anjali Nair, M. & Kumar, V.S., Spectral wave climatology off Ratnagiri, northeast Arabian Sea, Nat Hazards (2016) 82: 1565. doi:10.1007/s11069-016-2257-5

These publications share significant portions of text and techniques:

Glejin, J., Sanil Kumar, V., Sajiv, P.C., Singh, J., Pednekar, P., Ashok Kumar, K., Dora, G.U., and Gowthaman, R., Variations in swells along eastern Arabian Sea during the summer monsoon, Open J. Mar. Sci., 2 (2), 43–50, 2012.

Glejin, J., Sanil Kumar, V., Amrutha, M.M. and Singh J., Characteristics of long-period swells measured in the in the near shore regions of eastern Arabian Sea, Int. J. Naval Architecture and Ocean Engineering, 8, 312-319, 2016.

Indeed, there appears to be such a plethora of work in the area that a recent compendium was published:

P. V. Ethamony, R. Rashmi, S.V. Samiksha, V. M. Aboobackerm, Recent Studies on Wind Seas and Swells in the Indian Ocean: A Review, The International Journal of Ocean and Climate Systems, Vol 4, Issue 1, pp. 63 - 73.

Informing the reader of the broader context in which these measurements exist, and to clarify novel differences between published work and data presented in this paper is warranted.

3) The expressed objective of the work is: "This study addresses two main questions: (1) How the high-frequency tail of the wave spectrum varies in different months? (2) What are the spectral parameters for the best-fit theoretical spectra?"

The first objective is questionable since the method to assess slopes of the spectral tails is not revealed. The second objective is questionable since an ad hoc method consisting of a piecewise concatenation of different theoretical spectral functions is used. Specifics of how the concatenated frequency bands are determined is not provided, nor is an assessment of the physical assumptions inherent in concatenation of these different spectra.

The authors have attempted to clarify the data with statistical analysis, but the specific methods and algorithms are not provided, a prime deficiency. If there is desire to improve understanding and interpretation of the data, some effort to connect the data/statistics to underlying physical processes could improve the publication value. For example, you have argued that "Understanding of the wave spectral shapes is of primary importance for the design of marine facilities" yet little effort is made to quantify or relate physical parameters/forcings pertaining to the change in spectral slopes/peaks.

4) Please address the lack of consistency in precision of your statistical estimates and the lack of standard error or confidence limits. For example, wave frequencies are variously reported with precisions of 1, 2 or 3 decimal places, even in the same sentence. Slopes of the wave spectra are reported with 1 or 2 decimal point precision. Kindly adopt a uniform usage for expressing the precision of your estimates, ideally ones that reflect physically pertinent cutoffs. Please add error estimates or confidence bounds to your estimates, or address quantitatively why they are not germane.

5) You quote estimates of the slope of high-frequency portions of the wave spectra, but never define how those estimates are made.

6) You plot and describe "normalized" spectral densities, but never detail how this normalization is applied. One is then left to question whether the normalization is specific to each time period, or normalized across all periods so that direct comparisons of different years is meaningful.

7) Line 92 "The average monthly sea level at Karwar varies from 1.06 m (in September) to 1.3 m (in January)"

You are apparently quoting geodetic elevations here, but there is no reference datum specified.

8) Line 100 "The data for every 30 minutes from the continuous records at 1.28 Hz are processed as one record. From the time series data, the wave spectrum is obtained through fast Fourier transform (FFT)."

My understanding of waverider spectral processing is that motion samples are recorded at a sample rate of $f_s = 2.56$ Hz, not 1.28 Hz. The bandwidth of the spectral estimate is $f_s/2 = 1.28$ Hz. I do not believe that the data over 30 minutes are processed as one record. In standard Datawell processing, records of length 200 seconds are collected ($N = 512$ samples). 17 records are then FFT'd, windowed, and averaged with a 50% overlap to produce the 30 minute power spectral density estimate. Kindly verify your description, and if it differs from the standard Datawell spectral processing, detail and justify the differences.

9) Line 185 - 189 Here you describe dispersion relation impacts characterizing the wave spectrum of your data, you might consider moving this information to the beginning of the section.

10) Line 268 "Contour plots of spectral energy density (normalized) clearly show the predominance of wind-seas and swells during the non-monsoon period (Fig. 9)."

[Printer-friendly version](#)[Discussion paper](#)

Here you have apparently applied the unspecified 'normalization' to month long temporal windows of spectral density averaged over years. Is the normalization value specific to each month, or is it over the entire ensemble so that meaningful intercomparisons can be made? The normalization, as well as the averaging must be defined.

11) Line 281 "To study the characteristics of different wave systems, average mean wave direction and average wave spectral energy density grouped under different peak frequency bins are plotted in Fig. 10."

The meaning of "wave systems" is not clear. The meaning of "average mean" is not clear. Please see the following comment.

12) Line 281 - 294 It isn't clear to me that this section along with figures 10 and 11 provide meaningful analysis that isn't already discernible from the previous results, rather this seems like part of the exploratory data analysis and seems redundant. My recommendation is to remove this section and figures.

13) Line 296 "The behavior of the high-frequency part of the spectrum is governed by the energy balance of waves generated by the local wind fields. When the wind blows over a long fetch or for a long time, the wave energy for a given frequency reaches the equilibrium range and the energy input from the wind are balanced by energy loss to other frequencies and by wave breaking."

You have argued that the change in slope is indicative of a change from local wind-dominated to swell waves, a physical connection. And have recognized that the dynamic equilibrium between generation and dissipation processes are what control this slope... but fail to make any meaningful physical connection between the slope behavior and the underlying physics. This is not required in a paper that only presents data, but you have invested some effort in quantifying the spectra, it would seem natural to attempt a physical connection to sea state, wave height, wave age, wind speed or some physical parameter.

[Printer-friendly version](#)[Discussion paper](#)

14) Line 296 - 313 This section seems to try and convey in detail a very simple observation that more energetic wave spectra have steeper high-frequency tail slopes. If this needs explanation at all, it seems it should be much simpler. Again, how the slopes are computed is not defined.

Why are slopes in table 4 and figure 12 numerically negative, yet in the text are all positive? You refer to slopes as increasing, but are they not becoming more negative?

15) Line 307 "It is shown in Fig. 12 that the slope also increases as the mean wave period increases."

Figure 12 presents an interesting association between high frequency wave spectra slope and monthly mean wave height, but there is only one sentence referring to it with no exploration of its physical significance. It suggests a nonlinear saturation of slope as a function of wave height. You may wish to consider the suggestion following the comments.

16) Line 312 "The slope of the high-frequency end of the wave spectrum becomes milder when the wave nonlinearity increases. The study shows that the tail of the spectrum is influenced by the local wind conditions."

An attempt to physically connect your observations/statistics to physical forcing... Good. But this is unsatisfying. You are arguing that by virtue of large H_{m0} alone that nonlinearity increases? This seems speculative at best.

17) Line 315 4.3 Theoretical wave spectra

This section is very unsatisfying. How the fits to the theoretical spectra are determined is not provided, the only clue is: 'values for α and β were randomly varied within a range'. That this is not explained at all is a serious deficiency.

18) Line 317 "The monthly average wave spectra for the year 2015, is compared with JONSWAP and Donelan theoretical wave spectra."

[Printer-friendly version](#)[Discussion paper](#)

Why was the only the year 2015 used in the monthly average wave spectra fits? Is there reason to believe that interannual variability is insignificant and can be ignored, figure 6 suggests not. How then can monthly means from one year be considered representative of the wave climate? Year 2015 is reported to have 14772 observations, while years 2011, 2012 and 2014 have 17300. How do you justify extracting mean statistics on a data set missing 15% of the data?

19) Line 320 - 326 "For these months the first peak is fitted with JONSWAP spectrum and second with Donelan, and the fitted spectrum shows a good match with the measured one. In the monsoon period, the spectrum is single peaked with high spectral energy density and during this period JONSWAP spectrum is fitted up to the peak frequency and after that Donelan spectrum is used. During the months May, October and November, after the peak frequency, the measured spectrum is not smooth and hence for this part, Donelan spectrum is fitted in two parts in order to obtain the best fit."

The ad hoc piecewise fitting of spectra is unsatisfying. A uniform criteria from which spectra are fit to different portions of the data does not seem to exist. How can an ad hoc scheme be deemed useful to those engaged in the "design of marine facilities"? How does one interpret the physical basis of these piecewise spectra? Does it make physical sense to apply an ad hoc piecewise spectral concatenation? Are the essential assumptions inherent in the different piecewise spectra being satisfied?

In absence of answering these questions, an alternative approach would be to fit the different theoretical spectra over the entire frequency band, present the results, and perhaps hypothesize/speculate why the different inherent assumptions in the theoretical spectra result in different fit fidelity over the different frequency bands/environmental conditions.

20) Figure 3 The color bands need improvement. For example, in a) there are two yellow bands with only slightly different saturation, but widely different amplitudes. It is difficult to discern which amplitude corresponds to the sections in the plot.

[Printer-friendly version](#)[Discussion paper](#)

21) Figure 8 Instead of scaling the ordinal axis for each plot to maximize the dynamic range of the curves, it would be more informative to have a single uniform axis for all plots with the same angular range.

22) Figure 9 This is commonly referred to as a spectrogram. The amplitude scale has not been defined. No units are shown.

23) Figure 10 - recommend removing this figure and section To make figure 10 more informative for relative comparison across spectral bands, the ordinal axes should have uniform ranges, both for spectral amplitude and direction. This will allow the reader to immediately discern important differences between bands.

24) Figure 11 - recommend removing this figure and section Abcissal axes need labels and units.

25) Figure 12 Needs to explicitly label ordinates as slopes.

26) Figure 13 In contradiction to the text describing this figure, I'm unable to see where the Donelan spectrum was applied.

Suggested Analysis

As mentioned above, Figure 12 presents an interesting association between high frequency wave spectra slope and monthly mean wave height suggesting a nonlinear saturation of slope as a function of wave height.

The introduction contains a nice discussion of findings that quantify high-frequency slopes, both theoretically and experimentally, with substantial support that in your oceanographic setting that the expected high-frequency decay would be affine with f^{-4} . You mention in general terms how this decay represents an equilibrium between dissipation and energy input dominated by local winds, it may be useful to think about specific processes. For example Donelan et al. (2012) find that in addition to the k^{-4} dissipation that swells modulate the equilibrium in breaking waves dependent on the mean surface slope, while Melville (1994) also quantified a relation between wave

[Printer-friendly version](#)[Discussion paper](#)

packet slopes and dissipation rate. These results are specific to breaking waves, but one might expect similar relations between surface dynamics and dissipation rate for non breaking waves. If you do not find existing literature pertinent to non-breaking wave dissipation, then perhaps a functional representation of the data shown in figure 12 might be useful in revealing something about the physical connection, and at the very least would provide a predictive basis relating spectral slopes with mean wave heights as a basis for future research.

To that end, you might wish to fit a function of the form: $A * \exp(\lambda Hm0) + s0$, with initial parameters of $A = 8$, $\lambda = -2.4$, $s0 = -3.7$ to the data of figure 12. This is exemplified with a subset of your data below. With optimized parameter values you will then have a functional representation of your spectral slopes based on $Hm0$. Presumably, $Hm0$ along with other parameters (wavelength, wind...) may lead you or others to a hypothesis relating your spectral slopes to sea surface physics.

[Interactive comment on Ocean Sci. Discuss., doi:10.5194/os-2016-91, 2017.](#)

[Printer-friendly version](#)[Discussion paper](#)




Fig. 1. Slope vs H_{m0}

[Printer-friendly version](#)[Discussion paper](#)