# Interactive comment on "Bathymetry and oceanic flow structure at two deep passages crossing the Lomonosov Ridge" by Göran Björk et al.

5 Anonymous Referee #1

We thank the reviewer for a thorough reading of our manuscript and many insightful and constructive suggestions for revisions.

Summary

- 10 Referee #1 finds that the detailed bathymetric survey and collection of CTD casts presented in our paper from the Lomonosov Ridge are scientifically interesting and deserves publication, although first after a major revision. It is specifically the "methods", "results" and "discussion" sections Referee #1 comments on. We have decided to undertake the revision. Referee #1 provided some specific comments that we will address one by one in our revision (The comments by Referee #1 are written in
- 15 italics with blue text).

# **Methods**

Missing information: Detailed period of the cruise (days, months), explanation about collection and use of silicate and dissolved oxygen data (maybe coming from discrete water samples and laboratory analyses?)

This information will be added in our revision.

Moreover, if you compare your 2014 data with previous ones, especially if the latter are taken 10 years before, you should justify your choice, being aware that the comparison of thermohaline properties taken in period much different can raise some criticisms.

The choice for our data comparison is simple, it is it the only data we possess from this very sparsely investigated area of the Arctic Ocean. We strongly believe that a comparison with the older data

30 collected from previous years is important to show since they overlap spatially and support the notion that our observations not only represent anomalous snapshots. However, we agree with Referee #1 in that it is important to emphasize the time differences between the data sets. We will therefore add some clarifying words in the revised paper, and include more clearly from when over the year all data shown in the paper was collected, and not only at what year.

35

20

The following will be included:

"Additional evidence of water mass transport from the Makarov to the Amundsen Basin at the northern end of the southern gap is found in two zonal sections at 81°N acquired by RV Polarstern expeditions ARK-XI/1 and ARK-XII in 1995 and 1996 respectively. The 1995 section was acquired

- 40 August 20-27 and the 1996 section August 16-20. They coincide with the northern end of the SWERUS-C3 section and include deep stations in both basins. This implies that they may serve as reference stations representative for each year to exclude inter-annual variability and long-term trends that have been identified in the Arctic (Polyakov et al., 2012)..
- **45** *A minor comment: I suggest the Authors to write somewhere in the methods that they analyze thermohaline properties by using potential temperature, salinity and dissolved oxygen, and define those parameters and their symbols once and for all.*

This will be added as suggested.

# Results

5

In section 3.1 the Authors start describing the bathymetry of the region. I had to spend a time to compare figures 1 and 2: I would suggest the Authors to use the same criteria of orientation in panels of figure 2 (a, b) with respect to that in figure 1 (b). It would render easier for the reader to visualize regions and possible pathways of water masses in the study area discussed later within the ms.

Point taken. We will rotate the main map so 140° is down, implying that it fits with the insets in later figures.



15 Top: Original map, Bottom: Revised rotated map. Please refer to the original pdf manuscript for comparing original and revised figures for figures 2-9.

Page 5: After line 20, Authors describe the hydrological properties. I suggest adding a phrase that could connect the first part of the section 3.1 with its second part.

5

We suggest bridging the two sections with:

"Overall, the new multibeam bathymetry from SWERUS-C3 together with depth information from IBCAO 3.0 provides a spatial context for the north-south hydrographic section across the southern passage (along the ridge crest) close to the Siberian continental slope (Fig. 3)."

10

*Line 21: substitute "the section across the passage" with something like "the northsouth hydrological section across the passage..."* 

#### 15 This will be inferred in the bridging suggestion above.

Lines 27-29: Are you speaking about the surface warm core? If so, please move up this phrase within the text, where you are speaking about the surface layer, otherwise it is difficult to follow the description of the thermohaline properties.

20

No, this is referring to the Atlantic water, which is subsurface. We have removed "core" in the first sentence to avoid any confusion.

- Lines 31-32: explain why such properties of the bottom layer are "anomalous". From figure 3, which
  this part of the text is referred to, the mentioned temperature increase at the very thin bottom layer is not visible. Moreover, cold and salt waters, hence very dense (why you do not add also potential density data?), are normally trapped in the bottom layer of a basin, and lower dissolved oxygen values confirm that they are also pretty old (i.e. not ventilated since a relatively long time).
- 30 Here, "anomalous" simply refers to a slight change in near bottom layer water properties; we will use more clarifying wording. It should be noted that the changes we refer to are also seen in Figure 4, specifically salinity. For this reason, the reader pointed to Figure 4 in this section in addition to Figure 3.

We will also show potential density data. Yes, normally oxygen content reflects age. Here alsodifferences in source waters contribute to the oxygen contrast.

Line 34: remove double 'to be'.

Will be fixed.

40

Lines 35-37: please better explain this part and/or support with adequate references (e.g. Chelton et al 1997, JPO). How did you calculate the Rossby radius? Why did you not mentioned this calculation in the "methods"? Again, did you consider checking satellite images, sea level, or horizontal distribution of potential density to see if any eddies would be visible? It could support the discussion.

45

After some further analyze of the special characteristics of station 137 and 138 we have arrived at the conclusion that we probably see some type of interleaving structure between the Barents Sea branch and Fram Strait branch waters which is advected along with the boundary current. We will adjust the text according to this and thus exclude the discussion about eddy and the associated Rossby radius.

50

Lines 4-5: remove "matching the salinity and temperature data". Perhaps, you could discuss the origin of this bottom water masses, and how they accumulated in this part of the ridge, remaining likely isolated from the rest of the water column.

55 "Matching the salinity and temp......" will be removed.

Lines 6-7: please change this part with "Vertical profiles of  $\theta$  and S collected in the southern passage of the LR are compared with those collected in the Makarov Basin (Stn. 145 in fig. 2b) and Amundsen Basin (Stn. 148 in fig. ???) and shown in figure 4."

5

# This section will be revised for clarity.

Lines 7-19: this part need to be re-written to render it clearer. I was totally lost reading this part. Please write in more orderly manner about layers, water masses, possible pathways, explain why you chose two reference stations, and so on.

# This section will be revised for clarity.

Lines 20-36: the same comment as above: re-write this part to be more clear. Authors start
introducing silicate data without any previous explanation of them in the "methods". Additionally, they suddenly refer to mooring data gathered in 1995-1996 but this is not well inserted in the context.

This section will be revised for clarity and the measurement procedure of silica in water samples will be described in the methods.

20

Page 7: Line 2: "upwelling" is a specific oceanographic process, in this case is it driven by what? Maybe Authors mean "upward displacement"

Yes, it should be upward displacement.

25

*Line 5: "turbulent diffusion"? ok, but please justify or refer to appropriate bibliography.* 

This will be expanded on in a separate sentence with reference included:

30 "Using a value of  $5 \cdot 10^5 \text{ m}^2 \text{s}^{-1}$  for the turbulent diffusion coefficient according to observations at the Lomonosov Ridge (Rainville and Winsor, 2008), it would take about one year to reduce the maximal vertical salinity gradient with 50%."

Lines 8- end of the page: to be re-organized. Again, comparing 2014 thermohaline conditions with
those of 1995-1996, after you have written that the variability is large in this region, does not make any sense, unless you justify this approach. Moreover, remove conclusions from this section.

The approach is now justified and explained with more metadata, as described above.

40 *Page 8*.

Lines 1-6: move this part to the "methods".

# This will be moved to the methods."

**45** *Lines* 20. . .: *the same comment used for the previous page. The description of the thermohaline properties distribution is confusing, and need heavy improvements, in terms of language used and organization of the text.* 

#### This section will be revised for clarity.

50

Based on the comments I have provided for the results, I could say that the discussion has to be revised accordingly to the future changes required for the "results". However, I will provide here some comments: Page 10, Lines 34-35: how do you define the flow "largely barotropic"? Is it reported in literature?

Yes, the batropic flow is reported in literature, which we will include in the revised version (Noest and Isachsen, 2003).

Page 11, line 7: I do not think that "water streaming" is appropriate, please check it.

5

15

Will be changed to "water flowing".

Page 11, Line 11: indicate "Gakkel Ridge" in figure 1.

# 10 To be changed in revised figure.

In general, it seems to me that parts of the discussion could be moved to the introduction, while here the Authors should discuss their own data with more detail. Doing so, they could provide some nice conclusions (now they are not clear) on water masses distribution (as they did) and some speculations or hypotheses on the evolution of the thermohaline properties according with previous already published data.

The discussion will be expanded considering the broader implications of our results.

- 20 FIGURES: Figure 3: To use different color scales in each layer can be useful to see the variability within each of them, but can confuse the reader because it seems that (e.g.) intermediate and deep layers have different values while they are almost similar with the exception of the very bottom layer. Try to use the same scale in each layer. Figure 4: respect always the same order, θ first, S second, Dissolved oxygen third (if you want to show), and then θ/S diagram. In general, figure is not clear, all
- **25** profiles seem bold, and colors between st. 145 and 148 are not clearly distinguishable. Finally, for the exact location of the stations, it is better to refer to figure 2, not figure 1. Figure 5: from this figure, it seems that silicate and dissolved oxygen data comes from discrete water samples. Why the Authors did not described this aspect in the "methods"?
- 30 It is actually necessary with different color scales for each depth interval since it otherwise is impossible to see the variability. We will add a note in the figure text to alert the reader about the different scales. We will revise figure 4 in order to make it clearer and will refer to figure 2 for locations. The discrete water sampling data will be included in methods.

# Interactive comment on "Bathymetry and oceanic flow structure at two deep passages crossing the Lomonosov Ridge" by Göran Björk et al.

5

#### Anonymous Referee #2

We thank the reviewer for a thorough reading of our manuscript and many insightful and constructive suggestions for revisions (The comments by Referee #2 are written in italics with blue text).

#### 10

# **General comments:**

We think that the reviewer points to important general improvements concerning language, clarity and readability of figures, and the benefits of extending discussions on the broader implications of our

study; in particular pointing to potential new questions emerging from our work. We will revise the paper based on these overarching suggestions, as will be detailed below.

## **Specific comments:**

P6, Line 11: Could you colour stations 137 and 138 differently so that this statement is clearer on Fig
4? I can just about see what you mean, lower salinities c 500 m, but all profiles being blue it is hard to distinguish from the other profiles for the remaining depths.

We will introduce different line types for these two stations (solid and dashed lines) to facilitate distinguishing between them.

25

P7, line 14 & figures: Please make all multi-panel figures in the paper a,b,c etc. You have done this for some figs but not all. Make consistent across the figures, and also how you label them and the fonts used (some bold with a bracket, some not bold no bracket...). You can then refer more easily to the salinity/potential temperature plots in the text.

#### 30

We will edit the figures concerning panel labels and fonts as suggested.

*P7*, line 27: Any comment on this westernmost station? Makarov water just not reached here or Amundsen signal overwhelming, barrier to W transport/mixing?

35

Most likely, the Makarov Basin water in this depth range is deflected along the Ridge bathymetry (in the north-south direction; see Woodgate and Nøst and Isachsen in the reference list) slightly to the east of the westernmost station, where the topography becomes steeper. We will add the following comment in the revised ms.:

40 "The absence of a Makarov Basin hydrographic signature at the westernmost station indicates that the Makarov Basin waters are deflected along the steep topography on the western side of the Lomonosov Ridge."

P9 line 9: Interleaving motions = or could it by a gyre/circulation within the intra basin? Perhaps
explain the origin of "interleaving motions" if this is the correct oceanographic term (not my speciality)

We will rewrite the sentence related to interleaving, providing a reference to Rudels et al. (1999), who in detail review interleaving processes in the Arctic Ocean. The revised sentence reads:

<sup>50</sup> "The profiles suggest that this exchange is not a unidirectional organized flow, but more likely interleaving motions resulting from double-diffusive mixing process, which can create relatively distinct layers with variable flow directions that are intermittent in nature (Rudels et al., 1999)."

P10, line 25: Label disturbed bottom sediments and transparent lenses on Fig 9

# We will do this.

P12 line 4-5: I would prefer to see a comment on what kind of data could be used to elucidate the flow exchange rather than a negative comment on the data presented in this study! Otherwise why are we publishing it?

We will replace the sentence "It is hard to say anything more specific from this type of data." with "Information on the flow speeds, ideally from moored current meters, would be needed to decide the significance of the cross-ridge exchanges of volume, heat, and salt in this northern passage."

10

5

P12 line 20-27: The discussion lacks discussion on larger implications of this work plus any comments on unresolved/new questions raised by the study. What is the implication, if any, on seabed temp, for example.

15 We will replace the last paragraph in the discussion section with a new one pointing to a few new open research questions (see examples below) which are highlighted by the present study.

1) Our study suggests that the exchange flows across the Lomonosov Ridge have rather different character in the southern and the northern passages. In the southern one, the flow has a more coherent structure, comprised by a few vertical layers with unidirectional velocities. In the northern passage, on the other hand, the flow is broken up into multiple shallow vertical layers with alternating velocity directions. Thus, our results highlight the question of what physical and bathymetrical features that control the nature of the exchange flow through the deep passages cutting across the Lomonosov Ridge.

25

30

20

2) Our survey of the bathymetry and hydrography in the southern passage shows that the detailed saddle bathymetry can have a strong influence on the flow of most dense Makarov Basin waters crossing the LR. Although, our survey does not fully resolve the hydrography and bathymetry over the southern saddle, it indicate that relatively minor topographical features may control if the dense waters reaching the saddle cross over the ridge or return back to the Makarov Basin.

# **Technical corrections:**

We appreciate these detailed comments. We will go through the manuscript and address all suggested technical corrections.

#### Bathymetry and oceanic flow structure at two deep passages crossing the Lomonosov Ridge

Göran Björk<sup>1</sup>, Martin Jakobsson<sup>2</sup>, Karen Assmann<sup>1</sup>, Leif <u>G.</u> Andersson<sup>1</sup>, Johan Nilsson<sup>3,4</sup>, Christian Stranne<sup>2,5</sup>, and Larry Mayer<sup>5</sup>

<sup>1</sup>Department of Marine Sciences, University of Gothenburg, Gothenburg 405 30, Sweden
 <sup>2</sup>Department of Geological Sciences, Stockholm University, Stockholm 106 91, Sweden.
 <sup>3</sup>Bolin Centre for Climate Research, Stockholm University, Stockholm106 91, Sweden.
 <sup>4</sup>Department of Meteorology, Stockholm University, Stockholm 106 91, Sweden.

- <sup>5</sup>Center for Coastal and Ocean Mapping, University of New Hampshire, New Hampshire 03824, USA *Correspondence to:* Göran Björk (goran.bjork@marine.gu.se)
- 15 Abstract. The Lomonosov Ridge represents a major topographical feature in the Arctic Ocean which has a large effect on the water circulation and the distribution of water properties. This study presents detailed bathymetric survey data along with hydrographic data at two deep passages across the ridge: A southern passage (80-81 °N) where the ridge crest meets the Siberian continental slope and a northern passage around 84.5 °N. The southern channel is characterized by smooth and flat bathymetry
- 20 around 1600-1700 m with a sill depth slightly shallower than 1700 m. A hydrographic section across the channel reveals an eastward flow with Amundsen Basin properties in the southern part and a westward flow of Makarov Basin properties in the northern part. The northern passage includes an approximately 72 km long and 33 km wide trough which forms an intra basin in the Lomonosov Ridge morphology (the Oden Trough). The eastern side of Oden Trough is enclosed by a narrow and steep
- 25 ridge rising 500-600 m above a generally 1600 m deep trough bottom. The deepest passage (the sill) is 1470 m deep and located on this ridge. Hydrographic data show irregular temperature and salinity profiles indicating that water exchange occurs as midwater intrusions bringing water properties from each side of the ridge in well-defined but irregular layers. There is also morphological evidence that some rather energetic flows may occur in the vicinity of the sill. A well expressed deepening near the
- 30 sill may be the result of seabed erosion by bottom currents.

# **1** Introduction

The Arctic Ocean includes the northernmost loop of the global ocean circulation system. Warm water from the North Atlantic flows through Fram Strait between Svalbard and Greenland and continues into the central Arctic Ocean basin as a warm core along the northern Barents Sea shelf slope (Fig. 1).

35 Another branch of Atlantic water flows across the Barents Sea where it undergoes significant cooling

by heat loss to the atmosphere before entering the central Arctic Ocean further east in the St. Anna Trough (Schauer et al., 2002). The two branches of Atlantic inflow meet to partially mix together and continue further eastward along the continental shelf slopes of Kara, Laptev and East Siberia seas (Rudels et al., 2000). The inflow of Atlantic water implies a net heat transport from low to high

5 latitudes and results in the entire central Arctic Ocean basin filling up with a relatively warm (>0  ${}^{\bullet}C^{\circ}C$ ) layer at depths between about 100 and 600 m, the Atlantic Water L-layer. The maximum temperature core is located at around 200-300 m depth. Further on, the water mass definition AWL will be used for the warm core of Atlantic water >0 °C. When referring more generally to waters of Atlantic origin we use the term Atlantic water. The temperature of the Atlantic water inflow has not

10

been steady, not even over short time-scales. Several warm pulses on decadal scales have been observed as well as sudden changes in the core temperature (Dmitrenko et al., 2009; Woodgate et al., 2001).

The currents in the central Arctic Ocean are generalized by a weak interior circulation and intensified boundary currents following the slopes of the shallow continental shelf seas that nearly enclose the

- central basin. The boundary currents carry not only the warm core of the Atlantic layer (>0 $^{-\theta}$ C)-WL 15 but also intermediate and deep water with lower temperature that generally decreases with depth. The system of boundary currents including containing water from the two branches of Atlantic inflow is referred to as the Arctic Circumpolar Boundary Current (Rudels et al., 1999a). However, the detailed circulation is far from one simple flow along the continental slopes of the shelves which emerges from spatial contrasts of the Atlantic layer and deep water properties (Rudels et al., 20122013). The 20
- complex seafloor landscape consists of extensive submarine ridges with morphologies that influence and steer the currents (Fig. 1).

As is the general case for the mean circulation in the weakly stratified high-latitude oceans, the circulation in the Arctic is strongly guided by bathymetry with currents tending to follow isobaths and giving rise to several internal circulation loops associated with deep sea ridges (see e.g., Nøst and

- Isachsen, 2003). At places where the deep sea ridges meet the continental slope they may deflect a part of the boundary current from the continental slope and make it follow the ridge instead. The largest deep sea ridge system in the Arctic Ocean is the Lomonosov Ridge (LR) rising up several thousand meters above the abyssal plains and stretching all the way from the Siberian slope to the continental
- 30 slope north of Greenland (Fig. 1). The LR also defines the border between the two major ocean basins in the Arctic: the Eurasian Basin, with sub-basins Nansen Basin and Amundsen Basin and the Amerasian Basin, with sub-basins Makarov Basin and Canada Basin. The impact of the LR as a major obstacle for the boundary current's passage along the Siberian slope from the Eurasian Basin to the Amerasian Basin side is clearly seen in hydrographic data (Anderson et al., 1994), current
- 35 measurements (Woodgate et al., 2001) and ocean modelling studies (Aksenov et al., 2010). The boundary current splits up in two branches where the LR meets the Siberian shelf slope (Fig. 1). One

9

branch turns north and flows along the ridge towards Greenland and one part passes the ridge and continues along the shelf slope in the Amerasian Basin. In addition to the passage of warm Atlantic water flowing across the LR from the Amundsen Basin to the Makarov Basin, there is also evidence based on mooring data that deep water from the Makarov Basin passes the ridge in the opposite direction in this area (Woodgate et al., -2001).

5

10

15

Mapping the detail pathways of the warmer water of Atlantic origin is of importance as the heat it brings influences the Arctic Ocean environment, specifically the marine cryosphere, including for example gas hydrates stored in bottom sediments, sea ice and outlet glaciers. More specifically, the inflowing warm Atlantic water controls the temperature near the seabed along the continental shelf slope, which affects the location of the Gas Hydrate Stability Zone and thus impacts the storage and release of methane (Biastoch et al., 2011; Stranne et al., 2016; Westbrook et al., 2009). Furthermore, the heat carried with the inflow has a large potential to melt the perennial sea ice cover (Polyakov et al., 2010), although mostly in areas with weak salinity stratification such as north of Svalbard. In most parts of the Arctic, the Atlantic layerAWL is shielded from the sea ice by a strong cold halocline resulting from freshwater supply by rivers and low salinity water coming from the north Pacific through Bering Strait (Sirevaag and Fer, 2009; Linders and Björk, 2013). Another example illustrating the complex pathways of Atlantic water is that it finds its way across Lincoln Sea to reach the Petermann Glacier's floating ice tongue (Münchow et al., 2007).

In a larger scale context it is clear that the LR not only plays a critical role by shaping the property distribution of the intermediate and deep waters among the major Arctic Ocean basins but also by controlling the mixing and water mass transformation from the inflowing Atlantic water contained in the two branches to the outflow of colder and fresher modified water through the western Fram Strait (Rudels et al., 1999a). This outflow then contributes to the overflow across the Greenland-Scotland
 ridge and the Nnorth Atlantic Ddeep Wwater formation.

The bathymetric portrayal of the LR in the latest version 3.0 of the International Bathymetric Chart of the Arctic Ocean (IBCAO) is mainly based on sparse single beam echo soundings from icebreakers and submarines and digitized depth contours from published maps, apart from a few areas mapped with multibeam echo sounder (Jakobsson et al., 2012). The sparse source data implies that bathymetric

- details of importance from an oceanographic perspective may be missed in some areas, such as the location of bathymetric passages or saddles in LR, which are critical as this is where a large part of the water exchange between the basins can occur. The (presumably) deepest passage across the LR, with a sill depth of 1870 m, has been located at 88°88° N. This passage was mapped in detail with multibeam together with hydrographic observations in 2005 and it was shown that a substantial flow of Canadian
- Basin Deep Water passes through it (Björk et al., 2007; Björk et al., 2010).

Here we present a detailed bathymetry of two other deep passages in the LR morphology together with hydrographic data acquired during the SWERUS-C3 (Swedish-Russian-US Arctic Ocean Investigation of Climate-Cryosphere-Carbon Interactions) expedition in 2014 with Swedish icebreaker (IB) Oden. The southernmost of these passages is located close to the Siberian shelf slope between about 80°N

5 and 81°N where it can be expected that most of the Atlantic water passes the LR towards the Amerasian Basin. The other passage is located further north at  $\sim$ 85°N in an area that has not been studied before but has been suggested as a possible location for exchange of Atlantic water (Woodgate et al.-, 2001). Our study shows the critical importance of knowing the detailed shape of the seafloor in order to map the deep bottom currents, which in the Arctic Ocean are critical for the distribution of heat and other properties and have implications for the marine environment, including the vulnerable

#### 10

#### 2 Methods

marine cryosphere.

15

- The entire water sampling program during the SWERUS-C3 expedition took place August 15 September 25, 2014. Collection of water samples from the southern LR area, in focus here, was performed between September 14 – and September 22. A rosette equipped with 24 Niskin bottles was used for the water sampling. CTD observations were made using a SeaBird 911 CTD equipped with dual SeaBird temperature (SBE 3), conductivity (SBE 04C) and oxygen sensors (SBE 43). Salinity samples were analyzed using a Guildline Autosal instrument which was kept in a well isolated lab container with relativelyin order to keep constant temperature as constant as possible. The salinometer 20 was calibrated using one standard sea water ampule (IAPSO standard sea water from OSIL Environmental Instruments and Systems) before each batch of 24 samples. The CTD data files were post processed with standard SeaBird data processing software. Removal of spikes in the CTD data They were was done despiked manually by first visually identifying spikes and then interpolating across the spike in the file with full time resolution. The alignment parameter was tuned for each 25 sensor pack following the suggested method described in the SeaBird Data Processing manual. Salinities were compared with bottle salinities analyzed with the Autosal lab salinometer. This analysis revealed a systematic offset of conductivity sensor 1, from station 16 and onward, corresponding to about 0.002 PSU-psu higher salinities compared to the salinometer and to conductivity sensor 2. A conductivity slope correction was calculated and the raw data conversion was
- 30 redone with the slope correction. Water column data is mainly analyzed using the potential temperature  $\theta$  (°C), salinity S (psu) and dissolved oxygen O<sub>2</sub> (µmol kg<sup>-1</sup>)

We also make use of bottle data of oxygen and silicate sampled from the rosette. Dissolved silicate was measured on board using a continuous flow analyser (QuAAtro system, SEAL Analytical). Within each analysis run a calibration was done using a commercial stable silica-compound solution.

Analysis quality was further assured by automatic drift control using standard solutions, with the precision being 1.3 % based on 28 determinations of standards. An automated Winkler titration system was used for the oxygen measurements with a precision of 1  $\mu$ mol kg<sup>-1</sup>. The accuracy was set by titrating known amounts of KIO<sub>3</sub> salts that were dissolved in sulfuric acid. As the amount was known to better than 0.1% the accuracy should be significantly less than the precision.

5

10

25

A description of the geophysical mapping program onboard IB *Oden* during the SWERUS-C3 2014 expedition is included in Jakobsson et al. (2016). Here we provide a methods summary with emphasis on the mapping of the areas of the LR that are the focus for this study. Multibeam swath bathymetry was acquired using the Kongsberg EM 122 (12 kHz, 1°x1°) multibeam system installed in IB *Oden*. The CTD observations gathered for the oceanographic program were used for sound velocity control and supplemented with XBT (Expendable Bathy Thermograph) casts. The target areas of the LR were bathymetrically mapped with between 50 and 100 % overlapping multibeam swaths to acquire a high

quality imagery of the ridge morphology. All multibeam bathymetric data were processed using a combination of the software Caris and Fledermaus-QPS. Grids with horizontal resolutions ranging between 15x15 m and 30x30 m were produced and used for the final analyzes in the 3D environment of Fledermaus and in the GIS software ArcMap. In addition to multibeam bathymetry, sub-bottom profiles were collected using the Kongsberg SBP 120 3°x3° chirp sonar installed in IB *Oden*. The
 chirp sonar was operated continuously using a 2.5-7 kHz pulse.

The high quality of the bathymetric data collected was due, in large part, to the exceptionally ice free conditions along the LR during summer 2014. <u>The northern section of the LR shown in Figure 2b was</u> the last area to be investigated in detail during the SWERUS-C3 2014 expedition. The limited time available before IB *Oden* had to begin the return transit implied that all oceanographic stations were carried out concurrently with the multibeam mapping program. This excluded the possibility to strategically place hydrographic stations along section lines based on a detailed <u>multibeam</u>bathymetry map of the area.

# 30 **3 Results and discussion**

# 3.1 Southern passage

The LR meets the continental slope of the Siberian shelf approximately at the defined border between the Laptev and East Siberian Seas is (Fig. 1). This section of the LR, here referred to as the southern passage, is characterized by smooth and flat bathymetry with depths around 1600–1700 m in the

deeper central parts where the hydrographic section is located (Fig. 2 a, c). North of the deepest flat part, the ridge rises rather abruptly to reach depths shallower than 800 m. The southern end of the southern passage is confined by the continental slope that rises to the continental shelf. The single multibeam track on the western side of the surveyed section reveals a stripe of rugged bottom

- 5 topography with generally shallower depths than in the central parts, <u>undulating roughly between 1300</u> and 1500 m. Increasing depths at the southern end of this western section suggest that the bathymetric sill of the southern gap is likely located at the south-western rim of the central plateau. However, since the southern passage in the LR not was completely covered by the SWERUS-C3 multibeam survey, it is not possible to exactly pinpoint the location of the sill. We cannot exclude that there might be a
- shallower area forming a sill where there is no multibeam coverage between the hydrographic section and the western single track of multibeam data. However, the multibeam data we have at hand together with the general morphology of the LR provided by Version 3.0 of IBCAO, suggest a sill shallower than 1700 m somewhere south of 81° 15'N 141° 30'E (Fig. 2a). IBCAO is generally in good agreement with the newly acquired multibeam bathymetry, but approximately 50 m shallower in the
- deeper parts of the southern passage than our new multibeam bathymetry. IBCAO Version 3.0 is in this area primarily based on digitized contours from the Russian bottom relief map of the Arctic Ocean (Naryshkin, 2001), which do not include information on the bathymetric source data used. <u>Overall, the new multibeam bathymetry from SWERUS-C3 together with depth information from IBCAO 3.0</u>
  provides a spatial context for the north-south hydrographic section across the southern passage (along the ridge crest) close to the Siberian continental slope (Fig. 3).

The section across the <u>southern</u> passage (along the ridge crest) close to the Siberian continental slope (Fig. 3) shows a low salinity surface layer down to about 20 m with high temperature (up to 3 °C) above a cold layer close to the freezing point (Fig. 3). The low salinity is likely a result of ice melting and as well as possibly also fresh water input from the north Siberian rivers. The high surface temperature is a result of the open water conditions which allowed for heat supply by absorption of solar radiation and air-sea heat exchange during the summer. The interval between about 100 m and 900 m contains the A<u>Wtlantic water layer L</u> with temperature above 0 °C. The highest temperature in the Atlantic <u>AWL</u> water core is above 1.5 °C and is seen in the southern half of the section indicating that Atlantic water from the Amundsen Basin passes the ridge through this channel with a high temperature/salinity core in the southern part. The deep waters (>1000 m) are characterized by decreasing temperature and increasing salinity towards the bottom. The deepest part of the section shows anomalous deviating properties in a thin bottom layer with significantly higher salinity while theand temperature deviation is less pronounced. The density (Fig. 3c) (not shown) is also also higher

35

25

30

in this layer since the salinity dominates the effect on the density. Another noteworthy feature is at

station 137 with higher temperature and salinity at mid depth in the range 600-1300 m. <u>There are also</u> cores with lower salinity in the AWL around 450 m as seen in profiles from stations 137 and 138 as

well as at the southernmost station 143. These are barely visible in the section plots but stands out in the  $\theta$ S profiles (Fig. 4). This appears to be to be a small anticyclonic eddy which is not fully resolved with the present station spacing which is larger than the Rossby radius of ~7.5 km based on the first baroclinic mode phase speed of station 137. The oxygen section (Fig. 3d) shows generally high

5

concentration in the surface layer roughly matching the salinity distribution. It is relatively constant in most of the depth interval 200-1000 m with concentration around 295 µmol kg<sup>-1</sup>. The deepest part shows lower oxygen concentrations at the northern stations below 1200 m. The concentration is also significantly lower in the thin bottom layer-matching the salinity and temperature data.

Vertical profiles of  $\theta$  and S collected in the southern passage The TS profiles are compared with 10 reference stations in the Makarov Basin (Stn. 145) and Amundsen Basin (Stn. 148) in figure 4. The reference stations were chosen in order to represent the water properties, at each side of the ridge, away from the passages. We focus first on the observed properties above the thin bottom layer. In the depth interval 700-1300 m, Fit can be seen that the depth profiles of the three northernmost stations are are shifted towards the dominated by Makarov Basin reference profile while the other stations are shifted towards the Amundsen Basin reference profile. This difference is seen even clearer in the  $\theta S$ 15 plot (Fig. 4c). TS properties (red curves) with less saline and colder water in the depth interval 700-1300 m. The other stations in the section follow more the Amundsen Basin reference profile. The situation is more complicated in the A<u>Wtlantic LayerL (300-700 m)</u>, because of the eddy-likeisolated <u>core</u> feature at stations 137 and 138 around 500 m depth with significantly lower salinity and slightly 20 colder temperature. Also the southernmost station (Stnstn, 143) has a similar pattern in the AWtlantie layerL. Apart from these three stationsspecial features it is clear that the Makarov Basin properties dominate the three northern stations above a depth of about 1500-1300 m and well into the Atlantic layerWL. This is also seen in the oxygen section data with lower concentrations at depth in the northern deep-part of the section indicating a flow from the Makarov Basin having generally lower 25 oxygen concentration in this depth interval (see also Fig. 5). These observed profiles along the section observations thus suggest a flow from the Makarov Basin to the Amundsen Basin towards thein the north and a flow from the Amundsen Basin to the -Makarov Basin - in the depth interval 400-1300 m over the major southern part of the channel, in the depth interval 400-1300 m. but there is also clear evidence of a flow in the opposite direction in the northern part.

A possible explanation to the special features at station 137 and 138 is that they are formed between 30 the warmer and saltier Fram Strait branch of the Atlantic inflow and fresher and colder Barents Sea branch. The differences in  $\theta$ S properties results in highly variable vertical structures at the confluence when these two water masses meet east of the St. Anna Trough. The structures are formed by interleaving motions bringing layers of each water mass in under and above each other, or by isolated 35 parcels/eddies of Barents Sea branch water which are brought into the Fram Strait branch and vice

versa (Rudels et al., 2000). The water from the two branches then moves together in the boundary

current along the Laptev Sea slope, with the Barents Sea branch closest to the continental slope. Evidence of interleaving/eddy structures are also seen downstream at long distances from the confluence area indicating that these are carried with the flow while diminishing in amplitude due to mixing processes (Rudels et al., 2000). These structures will then also likely exist at the southern section where some part of the boundary current passes the LR, as seen at stations 137 and 138,.

5

<u>The deviating properties near bottom are clearly seen in the vertical profiles</u> Deeper down in water column, below 1500 m<sub>2</sub>, the <u>The</u> salinity and temperature increases sharply in a bottom layer with a thickness of about 50 m thickness. The salinity anomalies in this layer are typically 0.02 psu while the temperature is higher by about 0.05 °C at the deepest three stations. The salinity anomaly dominates the density resulting in enhanced potential density (Fig. 2c). This bottom layer is likely associated with

10

the density resulting in enhanced potential density <u>(Fig. 2c)</u>. This bottom layer is likely associated with the of-bowl shape of the central plateau and we surmise that this is water that has become trapped in the depression. This appears clear for the deepest part below 1700 m, but is not so obvious for the shallower signals at stations 134 and 138-140 which lie at about 1500 m and are thus well above the sill depth of the local depression. The properties near the bottom are similar to Makarov Basin

- 15 properties, but the salinity is higher than the reference profile. The salinity and density match the Makarov Basin profiles 400 m deeper for the deepest stations along the section and about 200 m deeper for the more shallow stations. The bottom layer has also anomalous concentrations of chemical constituents with significantly higher silicate concentration and lower oxygen concentration near the bottom (Fig. 5). These values also match the Makarov Basin properties 200-400 m deeper down. This
- 20 suggests that Makarov Basin water has been uplifted 200-400 meter and filled up a bottom layer with relatively deep Makarov Basin characteristics across the high plateau of the passage. Evidence that such vertical motions actually occur can be found from deep sea mooring data nearby in the Makarov Basin. The salinity data from a nearby moored instrument at 1700 m depth during 1995-1996 (Woodgate et al., 2001) show relatively large salinity variability, both short and long term, of about
- 0.02 psu which likely is associated with vertical motions in combination with a vertical property
   gradient. The bottom-layer water characteristics can thus be explained by upwelling upward
   <u>displacement</u> of water from deeper levels in the Makarov Basin that subsequently have become
   trapped on the plateau. The small vertical extent and sharp halocline at top of the layer is still puzzling,
   however. It indicates that the structure was formed relatively recently since vertical turbulent diffusion
- 30 would smear it out relatively quickly. Using a value of  $5 \cdot 10^{-5} \text{ m}^2 \text{s}^{-1}$  for the turbulent diffusion coefficient according to observations at the LR (Rainville and Winsor, 2008), it would take about one year to reduce the maximal vertical salinity gradient with 50%. The layer is also located at more shallow depth at the northern and southern stations which indicates that it is takes part<u>contributes to in</u> the across\_ridge flows giving rise to sloping isopycnals.

Additional evidence of water mass transport from the Makarov to the Amundsen Basin at the northern end of the southern gap is found in two zonal sections at 81°N acquired by RV Polarstern expeditions ARK-XI/1 and ARK-XII in 1995 and 1996 respectively. The 1995 section was acquired August 20-27 and the 1996 section August 16-20. They coincide with the northern end of the SWERUS-C3 section 5 and include deep stations in both basins. This implies that they may serve as reference stations representative for each year to exclude inter-annual variability and long-term trends that have been identified in the Arctic (Polyakov et al., 2012). To provide additional evidence of water mass transport from the Makarov to the Amundsen Basin at the northern end of the southern gap we use two zonal sections at 81°N taken by RV Polarstern in 1995 and 1996 that coincide with the northern end of the 10 SWERUS-C3 section. These sections include deep stations in both basins that can be used as reference stations to account for the inter annual variability and long term trends that have been identified in the Arctic (Polyakov et al., 2012). Both the 1995 and 1996 the AWL tlantic Water layers areis around 0.03 psu fresher, but have a similar core temperature as SWERUS-C3 data from 2014 (Fig. 6 c, d, e, f). In 1995, there is with a salinity minimum between 500 and 700 m (Fig. 66 c and g), originating from the Barents Sea branch inflow (Schauer et al., 2002+). In the 1995 section, both S and 0 between 15 400-1000 m depthconditions show remarkably little difference between 400-1000 m depth-west and east of the ridge-with a salinity minimum between 500 and 700 m (Fig. 6 c and g), originating from the Barents Sea branch inflow (Schauer et al., 2001). Deep water properties below 1500 m are similar to those observed during SWERUS-C3, so that we can use the latter to identify the origin of these water masses. The profiles start to divide between Makarov and Amundsen Basin properties below 1200 m 20 with the three easternmost profiles (red) being warmer and saltier than the westernmost ones. There is however signs of Makarov Basin water at the western side of the ridge in one of the blue profiles between 1100-1700 m and also in a relatively thin bottom layer with higher salinity below 1700 m (light blue). In the 1995 section, the three stations furthest west (Fig. 6 g, blue) and east (Fig. 6 g, red) show clear Amundsen and Makarov deep water characteristics, respectively. Two stations west of the 25 ridge around 138°E (Fig. 6, a, c and g, light blue) show Amundsen Basin characteristics down to 1700 m and saltier Makarov characteristics below 1700 m. The deepest part includes a thin bottom layer of fresher, but colder Amundsen water. A station taken in the same position in 1996 (Fig. 6, b, d and h, magenta) shows a similar behavior. This suggests that the thin, salty bottom layer we observed in the SWERUS-C3 data is not an isolated phenomenon in time and that the Makarov deep water that has 30 been uplifted onto the ridge may spread westwards across the rugged western edge of the bowl in the southern passage.- In 1996, all but the westernmost station along 81° N (Fig. 6, b, d and h, blue), show fresher Makarov Basin water in the depth interval 500-1500 m, as referenced to the two easternmost stations (Fig. 6, b, d and h, red). The absence of a Makarov Basin hydrographic signature at the 35 westernmost station indicates that the Makarov Basin waters are deflected along the steep topography on the western side of the LR. This confirms our conclusion from the SWERUS C3 data that there is indeed a westward transport of water at the northern end of the southern passage in this depth interval.

Data from both years show evidence of a thin salty bottom layer at the western side of the ridge (light blue stations 1995 and magenta 1996). The deepest part includes a thin bottom layer of fresher, but colder Amundsen water. A station taken in the same position in 1996 (Fig. 6, b, d and h, magenta) shows a similar behavior. This suggests that the thin, salty bottom layer we observed in the SWERUS-C3 data is not an isolated phenomenon in time and that the Makarov Basin deep water that has been uplifted onto the ridge may spread westwards across the rugged western edge of the bowl in the southern passage.

Both the SWERUS-C3 data and historical sections thus show signals of a flow of Makarov Basin water towards the Amundsen Basin crossing the LR through the southern passage. This type of flow was also inferred from mooring data in 1995-1996 showing clear signals of Makarov Basin water at a mooring site (located at  $81^{\circ}$  34.5<sup>2</sup> N,  $138^{\circ}$  54.0<sup>2</sup> E) at the western flank of the LR and just north of the

10

5

# southern passage (Woodgate et al., 2001). **3.2** Northern passage

#### 15

The northern section of the LR shown in Figure 2b was the last area to be investigated in detail during the SWERUS-C3 2014 expedition. The limited time available before IB Oden had to begin the return transit implied that all oceanographic stations were carried out concurrently with the multibeam mapping program. This excluded the possibility to strategically place hydrographic stations along section lines based on a detailed multibeam map of the area.

The multibeam mapping at the northern passage reveals a trough-like structure in the LR extending 20 from about 84°24'N - 85°3'N in north-south direction and 148°E -151°E in east-west (Fig. 2b). The approximately 72--km long and 33--km wide trough forms an intra basin in the LR morphology, which not is well expressed in IBCAO Version 3.0. The name *Oden Trough* for this feature has been was formally accepted by GEBCO's (General Bathymetric Chart of the Oceans) Sub-Committee for Undersea Feature Names (SCUFN) in 2015 (Fig. 2e). The eastern side of Oden Trough is enclosed by 25 rather steep walls rising 500-600 m above a generally 1600 m deep trough bottom. The deepest passage, i.e. the sill, between the Makarov and Amundsen basins is 1470 m and located in the eastern wall at about 84°43'N 151°15'E (Fig. 2b, e). The western opening towards Amundsen Basin is located at the northern end of the trough and is deeper as well as substantially wider than the eastern sill. The deepest part of Oden Trough is about 1704 m deep and located just northwest of the sill along the foot of the eastern wall, close to hydrographic station 152 (Fig. 2b). This deepening extends  $\sim 10$  km along the foot of the steep wall north of the sill and is well pronounced in the bottom morphology.

30

The  $\theta$ TS profiles in this area (Fig. 7) show a quite irregular behavior with influences of water mass properties from each side of the ridge at different depth intervals. Station 153, located in the Makarov Basin near the sill, closely follows the reference profile in the Makarov Basin below 1400 m, but in

the depth interval between 1400-1100 - 1100-1400 m it has more Amundsen Basin properties. At 1100 m there is a sharp transition towards more Makarov Basin properties up to 800 m. Further up the profile, while at 700 m depth there is again is a transition towards Amundsen Basin properties. In the Atlantic AWL layer (300-600 m) the water mass has Makarov Basin properties as demonstrated by the salinity and temperature maximum at  $\sigma_{\theta} = 27.94$  kgm<sup>-3</sup>. The nearby station 152 on the Amundsen Basin side of the ridge shows more Amundsen Basin characteristics below the 1470 m sill depth as indicated by the salinity profile but is similar to the Makarov Basin in the  $\theta$ TS plot. Further up in the column it follows the Amundsen Basin properties relatively closely, up to about 800 m, where it joins the Makarov Basin profile and has Makarov Basin properties all the way up to 300 m.

- 10 The station located at the western end of the of the passage close to the Amundsen Basin (stn. 155) has a structure more like the Amundsen Basin characteristics in the deepest part below 1400 m. In the interval 900-1400 m it has a mixture of properties. Above 900 m it undulates between clear Makarov and Amundsen Basin characteristics up to 700 m and Makarov properties dominates above this depth. Station 154 is located in a more central part of the trough where there is a local depression. This
- 15 station also undulates between different water mass properties but is clearly more dominated by Amundsen Basin characteristics especially in the interval 400-800 m where it is has significantly higher salinity.

It is difficult to interpret this complicated water mass structure based on a few semi-synoptic CTD stations but it is possible to deduce some general aspects. It appears that this is an area of active water

- 20 exchange across the LR presenting properties from the basins on either side. <u>The profiles suggest that</u> this exchange is not a unidirectional organized flow, but more likely interleaving motions resulting from double-diffusive mixing processes, which can create relatively distinct layers with variable flow directions, that are intermittent in nature (Rudels et al., 1999b). The profiles imply that this exchange is not an organized flow in one direction but more likely interleaving motions in relatively distinct
  25 layers that can go in both directions and are probably intermittent in nature. An indication of the horizontal spatial structure of these flows is that Makarov Basin properties appear to dominate for stations 152 and 155 in the depth interval 300-900 m. Both of these stations are located near the northern high plateau suggesting a general flow from the Makarov Basin along the steep slope, but most of this flow appears to be occurring above 1000 m. On the other hand we see Amundsen Basin
- 30 characteristic's around 700 m in the profile on the Makarov Basin side of the ridge indicating a flow from the Amundsen Basin. Additionally, the profiles from the Amundsen Basin side of the ridge show Makarov Basin properties below the sill depth indicating spillover of denser water from the Makarov Basin that is stored in the flat central area.

A 2011 Polarstern section (Schauer et al., 2012) crosses the LR at 84°30'N and provides a useful complement to the SWERUS-C3 stations (Fig. 8d). It shows an AWL -water core that is cooler and

35

5

fresher than in 2014 (Fig. 8). The two westernmost (Fig. 8, dark blue) and easternmost (Fig. 8, red) stations show coherent water mass properties of the Amundsen and Makarov Basins, respectively and we will use them as reference stations for the two basins. Similarly to the SWERUS-C3 data, the 4 stations over the ridge show evidence of water mass exchange across the ridge. A station at 140°E

- 5 (Fig. 8, light blue), west of the ridge, has the coolest and narrowest AWW layer L of the section with salinities in the 300-700 m depth range similar to those on the Makarov side, but follows Amundsen characteristics below 800 m. A station east of the ridge at 156°E (Fig. 8, magenta) has mostly Makarov Makarov AW and Basin characteristics deep water characteristics, but shows signs of an intrusion of water from the Amundsen side and interleaving between 600 and 1200 m. Two stations at 10 145°E (Fig. 8, green) and 150°E (Fig. 8, yellow) show further indication of water mass exchange across the ridge. Notable is that the more western (Fig. 8, green) of those stations has the cold and
  - fresh Makarov Basin AW-characteristics down to 1300 m and follows the Amundsen Basin deep water below.
- Both the SWERUS-C3 and historical data show that the northern passage is an area of active water 15 exchange across the LR. It is indicated from <u>T</u>the  $\theta$ S profiles indicates that this exchange occurs as midwater intrusions bringing water properties from each side of the ridge in well-defined but irregular layers. Similar irregular intrusions are found at the Kara Sea slope east of the St. Anna Trough where the colder and fresher-Barents Sea branch of the Atlantic inflow enters the Arctic Ocean and meets the warmer and saltier Fram Strait branch as described above(Rudels et al., 2000,; Rudels et al., 2013). A 20 narrow front is formed in the confluence zone which is highly unstable for-to horizontal perturbations due to double diffusive fluxes starting to act when colder and fresher water comes on top of a layer with warmer and saltier water (and vice versa). The perturbations then grow to large amplitudes and form the irregular layers (Rudels et al. 1999b). A similar situation, with a narrow front between water masses of different 0 properties and near equal density, seems to be the case at the eastern wall of the Oden Through where water with Amundsen Basin properties comes into close contact with water 25 having Makarov Basin properties. Since the LR has a strong steering effect, with flows generally along the ridge, it reduces the water exchange between the major ocean basins and maintains a frontal structure between the water mass properties on each side which is typically seen on hydrographic sections crossing the ridge (Rudels et al., 2013). Note that the flows on each side go in the opposite 30 directions and therefore typically bring different water mass properties. At locations where the ridge morphology is narrow it can be expected that this front will be sharper. The narrow eastern wall of the northern passage (including the sill) appears to be such a feature which can guide the water masses streaming at on either each side of the ridge to come close together, sharpening the horizontal property gradient and forming irregular intrusions. It can beis expected that these intrusions should have a very small-low across frontal speed since they are driven by the minute density gradients generated by the 35 double diffusive fluxes (McDougall, 1985).

On the other hand there are morphological expressions in the ridge seafloor that may result from more energetic flows in the vicinity of the sill. There is a nearly 100 m deepening, beginning at from the sill to continue northward along the northern footwall of Oden Trough, possibly formed by long-term seabed erosion of bottom currents (Fig. 9a). The sub-bottom chirp sonar data collected along with the multibeam bathymetry show disturbed bottom sediments closest to the northern footwall as well as transparent lenses with an acoustic appearance and stratigraphic location suggestions that they may be comprised of sediments redistributed by bottom currents (Fig. 9b).

# 4-3.3 DiscussionImpact by Lomonosov Ridge

5

10 distribution in the Arctic Ocean. The deepest waters are directly isolated from each other by the ridge and there is a clear contrast in properties between the opposite sides with deep waters in the Amerasian Basin warmer and saltier water than those in the Eurasian Basin. As noted above, the ridge also affects the property distribution further up in the water column as seen in many hydrographic sections crossing the ridge (Rudels et al., 20132). The main reason for this property contrast is that the flow is largely barotrophic with small vertical shear and tends to follow the bathymetry even far away 15 from the bottom. It is therefore difficult for the flow to pass the ridge except at places where isobaths cross the ridge. The effect of the ridge is much less pronounced in the upper layers above 100 m, including the halocline and the surface mixed layer, where the circulation is more controlled by wind forcing and strong horizontal density gradients associated with the supply and spreading of freshwater

We finally discuss some general aspects on how the LR effects the water circulation and property

20 from rivers and inflowing low salinity water through Bering Strait (Morison et al., 2012). The horizontal density gradients in the upper layers generate baroclinic motions which are decoupled from the topographical steering.

The southern passage provides a direct pathway above 1700 m for the warm Atlantic water streaming along the continental slope to enter the Amerasian Basin. It is mostly water from the colder Barents

25 Sea branch that passes the ridge while the warmer Fram Strait branch water turns and follows the ridge towards north (Rudels et al., 2013). A substantial part of the Fram Strait branch can also turn earlier and follow the Gakkel Ridge back towards the Fram Strait (Rudels et al., 2013). This means that most of the hydrographic sections crossing the ridge show a warmer Atlantic water core on the Amundsen Basin side than on the Makarov Basin side. The dynamics that actually determine the properties of the

water passing the ridge, of which the temperature is a critical quantity, appears to be an open question. 30 The temperature of the Atlantic water, including the seabed temperatures in the entire Amerasian Basin, should be highly dependent on this control. Current meter data from three moorings located at the Siberian shelf slope (one at each side of the L-R) and one at the L-R slope (Woodgate et al., 2001) show that the mean velocities are small and follow the depth contours. <u>but T</u>there are also numerous and strong cross-isobath flow events which are mostly related to mesoscale eddies but also likely due

35

to wind forced up and down-welling. This means that not only topographic steering, but also other types of dynamics will control the passage of water masses across the ridge. Also, the mixing between the Barents Sea and Fram Strait branches along the continental slope upstream from the LR will be critical.

- 5 The southern passage appears also to guide a westward flow of Makarov Basin water towards the Amundsen Basin as seen here and in earlier investigations. This flow is likely of intermittent nature which is evident from the 1700 m mooring data in the Amundsen Basin side where the 0TS properties undulate between Makarov and Amundsen basin signatures (Woodgate et al., 2001). Our section shows that this flow is present over a quite large depth range from the bottom and upseabed upwards to at least 500 m depth.
- 10

The northern passage seems to provide a different type of exchange in terms of interleaving motions above the 1470 meters sill depth, with flow in opposite directions over different depth intervals. It is likely that this flow structure is less important for the exchange of properties across the ridge since velocities associated with double diffusive interleavings should be small ~5 mm s<sup>-1</sup> (McDougall,

- 1985). Combining this with the relatively small horizontal scale of the passage of ~50 km and a 15 vertical scale -100 m results in a rather small volume flow -0.05 Sv compared with the 2 Sv estimated to pass eastward through the southern channel (Woodgate et al., 2001). However, the interleaving signals of Makarov Basin water are seen at quite large distances from the sill which means that they actually carry properties across the entire LR and should therefore contribute to the
- 20

overall water exchange across the ridge, but the question is to what extent. Information on the flow speeds, ideally from moored current meters, would be needed to decide the significance of the crossridge exchanges of volume, heat, and salt in this northern passage.

# It is hard to say anything more specific from this type of data.

There are also other deep channels which are pathways for flows crossing the ridge. Earlier 25 investigations show that Canadian Basin Deep Water crosses the ridge through the 1870 m deep passage (the deepest passage) near the North Pole (Björk et al., 2007) and can be traced along the Amundsen Basin slope towards Greenland and further along the northern Greenland continental shelf slope (Björk et al., 2010). Signs of this water mass are also seen as a salinity maximum around 2000 m over a large part of the Amundsen Basin. Closer to Greenland where the ridge meets the continental slope there is a passage with sill depth of around 1200 m. The overall bathymetrical structureseafloor 30 morphology in this area is similar to the passage near the Siberian continental slope at the other end of the ridge. This will cause a bifurcation of the westward rim current along the northern Canada continental slope, as also noted by Rudels et al. (2000). The shallow part crosses the ridge while the deeper part turns northward and flow towards Siberia along the LR flank on the Makarov Basin side.

There are also indications that a part of the northward flowing branch is guided to cross the ridge 35

hugging the slope around a 500 m depth shoal or at some deeper passages between the southern 1200 m passage near Greenland and the deepest 1870 m passage (see Björk et al., 2010 for details).

In a larger scale context it is clear that the LR not only plays a critical role by shaping the property distribution of the intermediate and deep waters among the major Arctic Ocean basins but also by controlling the mixing and water mass transformation from the inflowing Atlantic water contained in the two branches to the outflow of colder and fresher modified water through the western Fram Strait (Rudels et al., 1999). This outflow then contributes to the overflow across the Greenland-Scotland ridge and the north Atlantic deep water formation. It should therefore be valuable to make further and more detailed studies of the different deep passages in the LR including moored instruments to obtain time series.

#### 4. Summary and conclusion

Our study suggests that the exchange flows across the Lomonosov Ridge have rather different character in the southern and the northern passages. In the southern one, the flow has a coherent vertical structure, with unidirectional velocities over a large depth interval, but there is a horizontal division with flow directed towards the Makarov Basin in the southern part and the opposite in the northern part. In the northern passage, on the other hand, the flow is broken up into multiple shallow vertical layers with alternating velocity directions. Thus, our results highlight the question of what physical and bathymetrical features that control the nature of the exchange flow through the deep passages cutting across the Lomonosov Ridge.

- 20 Our survey of the bathymetry and hydrography in the southern passage shows that the detailed saddle bathymetry can have a strong influence on the flow of most dense Makarov Basin waters crossing the Lomonosov Ridge. Although, our survey does not fully resolve the hydrography and bathymetry over the southern saddle, it indicate that relatively minor topographical features may control if the dense waters reaching the saddle cross over the ridge or return back to the Makarov Basin.
- 25 <u>The possibility of more energetic flows in the vicinity of the sill at the northern passage, as inferred</u> from morphological expressions in the ridge seafloor, is of interest to study further from a dynamical perspective in order to estimate expected flow speeds under different conditions. <u>It should therefore be</u> valuable to make further and more detailed studies of the different deep passages in the LR including moored instruments to obtain time series.

#### 30

5

10

15

# Acknowledgments

The SWERUS-C3 expedition was financed by Knut and Alice Wallenberg Foundation, Swedish Polar Research Secretariat and Stockholm University. Research grants to Larry Mayer

was provided by the U.S. National Science Foundation Grant No. PLR-1417789. We also thank the crew of *I/B Oden* for all their professional help during the cruise. The 1995 and 1996 Polarstern data were received via the National Oceanographic Data Center.

#### 5

#### References

- Aksenov, Y., Ivanov, V. V., Nurser, A. J. G., Bacon, S., Polyakov, I. V., Coward, A. C., Naveira-Garabato, A. C., and Beszczynska-Moeller, A.: The Arctic Circumpolar Boundary Current, J. Geophys. Res., 116, C09017, doi:10.1029/2010JC006637, 2011.
- 10 Anderson, L. G., Björk, G., Holby, O., Jones, E. P., Kattner, G., Koltermann, K. P., Liljebladh, B., Lindergren, R., Rudels, B., and Swift, J.: Water masses and circulation in the Eurasian basin: results from the Oden 91 expedition, J. Geophys. Res., 99, 3273-3283, doi:10.1029/93JC02977, 1994.

Biastoch, A., et al.: Rising Arctic Ocean temperatures cause gas hydrate destabilization and ocean acidification, Geophys. Res. Lett., 38, L08602, doi:10.1029/2011GL047222, 2011.

Björk, G., Jakobsson, M., Rudels, B., Swift, J. H., Anderson, L., Darby, D. A., Backman, J., Coakley,
B., Winsor, P., Polyak, L. and Edwards, M.: Bathymetry and deep-water exchange across the central Lomonosov Ridge at 88°-89°N, Deep-Sea Res.- pt I, 54, 1197–1208, 2007.

Björk, G., Anderson, L., Jakobsson, M., Antony, D., Eriksson, P. B., Eriksson, B., Hell, B.,

- Hjalmarsson, S., Janzen, T., Jutterström, S., Linders, J., Löwemark, L., Marcussen, C., Olsson, K. A., Rudels, B., Sellén, E., and Sølvsten, M.: Flow of Canadian Basin Deep Water in the Western Eurasian Basin of the Arctic Ocean, Deep-Sea Res. pt-I, 57, 577-586, doi:10.1016/j.dsr.2010.01.006, 2010.
- Dmitrenko, I. A., Bauch, D., Kirillov, S.A., Koldunov, N., Minnett, P. J., Ivanov, V. V., Hölemann, J.
   A., and Timokhov, L. A.: Barents Sea upstream events impact the properties of Atlantic water inflow into the Arctic Ocean: Evidence from 2005 to 2006 downstream observations, Deep-Sea Res. pt-I, 56, 513–527, doi:10.1016/j.dsr.2008.11.005, 2009.
  - Jakobsson, M., Nilsson, J., Anderson, L., Backman, J., Bjork, G., Cronin, T. M., Kirchner, N.,
    Koshurnikov, A., Mayer, L., Noormets, R., O/Regan, M., Stranne, C., Ananiev, R., Barrientos Macho, N., Cherniykh, D., Coxall, H., Eriksson, B., Floden, T., Gemery, L., Gustafsson, O., Jerram, K., Johansson, C., Khortov, A., Mohammad, R., and Semiletov, I.: Evidence for an ice shelf covering the central Arctic Ocean during the penultimate glaciation, Nature Communication, v.-7, p. 1-10. doi:-10.1038/ncomms10365, 2016.

Linders, J., and Björk, G.: The melt-freeze cycle of the Arctic Ocean ice cover and its dependence on ocean stratification: J. Geophys. Res., 118 (11), 5963-5976, -doi:10.1002/jgrc.20409, 2013.

15

30

- McDougall, T. J.: Double-diffusive interleaving. Part II: Finite amplitude, steady state interleaving, J. Phys. Oceanogr., 15(11), 1542-1556, 1985.
- Morison, J., Kwok, R., Peralta-Ferriz, C., Alkire M., Rigor, I., Andersen, R. and Steele, M.: Changing Arctic Ocean freshwater pathways, Nature, 481, 66–70, doi:10.1038/nature10705, 2012.
- Münchow, A., Falkner, K. K., and Melling, H.: Spatial continuity of measured seawater and tracer fluxes through Nares Strait, a dynamically wide channel bordering the Canadian Archipelago, J. Mar. Res., v. 65, no.( 6), p. 759-788, 2007.

Naryshkin, G.: Bottom relief of the Arctic Ocean, bathymetric contour map, scale 1:2,500000, Russ. Acad. of Sci., St. Petersburg, Russia, 2001.

- 10 Nøst, O.A. and Isachsen, P.E: The large-scale time-mean ocean circulation in the Nordic Seas and Arctic Ocean estimated from simplified dynamics, J. Mar. Res., 61(2), 175-210, doi:10.1357/002224003322005069, 2003.
  - Polyakov I.V., Timokhov, L. A., Alexeev, V. A., Bacon, S., Dmitrenko, I. A., Fortier, L., Frolov, I. E., Gascard, J.-C., Hansen, E., Ivanov, V.\_V., Laxon, S., Mauritzen, C., Perovich, D., Shimada, K.,

Simmons, H. L., Sokolov, V. T., Steele, M. and Toole, J.: Arctic ocean warming contributes to reduced polar ice cap, J. Phys. Oceanogr., 40, 2743–2756. doi: 10.1175/2010JPO4339.1, -2010.
Polyakov, I. V., Pnyushkov, A. V. and Timokhov, L. A.: Warming of the intermediate Atlantic Water

of the Arctic Ocean in the 2000s, J. Climate., 25(23), 8362–8370, doi:10.1175/JCLI-D-12-00266.1, 2012.

20

25

15

- Rainville, L., and Winsor, P.: Mixing across the Arctic Ocean: Microstructure observations during the Beringia 2005 Expedition, Geophys. Res. Lett., 35, L08606, doi:10.1029/2008GL033532, 2008.
- Polyakov, I. V., Pnyushkov, A. V. and Timokhov, L. A.: Warming of the intermediate Atlantic Water of the Aretic Ocean in the 2000s, J. Climate., 25(23), 8362–8370, doi:10.1175/JCLI-D-12-00266.1, 2012.

Rudels, B., Anderson, L., Eriksson, P., Fahrbach, E., Jakobsson, M., Jones, E. P., Melling, H.,
 Prinsenberg, S., Schauer, U., and Yao, T.: Observations in the Ocean, in Lemke, P., ed., Arctic
 Climate Change: The ACSYS Decade and Beyond, Volume 43, Springer, p. 117–198, doi:
 10.1007/978-94-007-2027-5, 2012.

- Rudels, B., Friedrich, H. J., and Quadfasel, D.: The Arctic Circumpolar Boundary Current, Deep-Sea Res. pt II, 46, 6–7, 1023-1062, ISSN 0967-0645, doi:10.1016/S0967-0645(99)00015-6, 1999<u>a</u>.
  Rudels, B., Björk, G., Muench, R. D., and Schauer, U.: Double-Diffusive layering in the Eurasian
- basin of the Arctic Ocean, J. of Marine Syst., 21(1-4), 3-27, 1999b.
- 35 Rudels, B., Muench, R. D., Gunn, J., Schauer, U., and Friedrich, H. J.: Evolution of the Arctic Ocean boundary current north of the Siberian shelves, J. Marine Syst. I, 25, 77-99, ISSN 0924-7963, doi:1016/S0924-7963(00)00009-9, 2000.

Rudels, B., Schauer, U., Björk, G., Korhonen, M., Pisarev, S., Rabe, B., and Wisotzki, A.:
Observations of water masses and circulation in the Eurasian Basin of the Arctic Ocean from the 1990s to the late 2000s, OS Special Issue: Ice-Atmosphere-Ocean interactions in the Arctic Ocean during IPY: the Damocles project, 9(1), 147-169, 2013.

- Sirevaag, A., and Fer, I.: Early spring oceanic heat fluxes and mixing observed from drift stations north of Svalbard, J. Phys. Oceanogr., 39, 3049-3069, doi:10.1175/2009JPO4172.1-,2009.
  - Schauer U., Rudels, B., Jones, P., Anderson, L. G., Muench, R. D., Björk, G., Swift, J. H., Ivanov, V. and Larsson, A.-M.: Confluence and redistribution of Atlantic waters in the Nansen, Amundsen and Makarov basins, Annales Geophysicae, 20-(2), 257-273, 2002.
- Schauer, U., Rabe, B., and Wisotzki, A.: Physical oceanography during POLARSTERN cruise ARK-XXVI/3 (TransArc). Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Bremerhaven, doi:10.1594/PANGAEA.774181, 2012.
  - Stranne, C., O'Regan, M., Dickens, G. R., Crill, P., Miller, C., Preto, P., and Jakobsson, M.: Dynamic simulations of potential methane release from East Siberian continental slope sediments,
- 15 Geochemistry, Geophysics, Geosystems, 1<u>7</u>-(13), 872-886, doi: 10.1002/2015GC006119, 2016.
  - Westbrook, G. K., Thatcher, K. E., Rohling, E. J., Piotrowski, A. M., Pälike, H., Osborne, A. H.,
    Nisbet, E. G., Minshull, T. A., Lanoisellé, M., James, R. H., Hühnerbach, V., Green, D., Fisher,
    R. E., Crocker, A. J., Chabert, A., Bolton, C., Beszczynska-Möller, A., Berndt, C., and
  - Aquilina, A.: Escape of methane gas from the seabed along the West Spitsbergen continental margin, Geophys. Res. Lett., 36, 15, p. L15608, doi:10.1029/2009gl039191,-2009.
    - Woodgate, R. A., Aaagard, K., Muench, R.D., Gunn, J., Björk, G., Rudels, B., Roach, A. T., and Schauer, U.: The Arctic Ocean boundary current along the Eurasian slope and the adjacent Lomonosov Ridge: Water mass properties, transports and transformation from moored instruments, Deep-Sea Res.-pt I, 48, 1757-1792, 2001.



Figure 1: a) Map of the Arctic Ocean with the major ocean currents. Red arrows: flow of warm
Atlantic water and inflow Pacific water through Bering Strait. Purple: flow of colder Barents Sea
Branch of the Atlantic water inflow which has been modified when crossing the Barents Sea. Blue: flow of colder and fresher modified Atlantic water. The red and blue arrows also represent the deep water circulation below the warm core of Atlantic water. White hatched arrows: Flow of low salinity water in the surface layer and halocline. Abbreviations: Lomonosov Ridge (LR), Fram Strait (FS),

Bering Strait (BS), Nansen Basin (NB), Amundsen Basin (AB), Makarov Basin (MB), and Canada Basin (CB). Yellow dots show the positions for all CTD station during the SWERUS-C3 expedition.
b) Details of the special study area of the LR including the CTD stations. Note the position of station 148 which is referred in the text. For further information of station numbers see Figure 2.



**Figure 2:** Detailed maps of the southern passage (a, c) and northern passage (b, e). The likely position of the sill for the southern passage is within the white circle. A semitransparent plane visualize the aproximate sill level of 1700 m for the southern passage (c) and the actual sill level of 1470 m for the northern passage (e).



**Figure 3:** Sections across the southern passage showing: Potential temperature <u>a</u>) Salinity S (psu), b) potential temperature  $\theta$  (°C), salinity S (psu) c) potential density  $\sigma_{\theta}$ -(kg m<sup>-3</sup>) and <u>d</u>) dissolved oxygen O<sub>2</sub> (µmol kg<sup>-1</sup>) sections across the southern passage. Please note the different color scales for the separate depth intervals. Note also that stations 141 and 142 are labeled as 41 and 42 in order to save space.





**Figure 4:** Salinity and potential temperature  $(\Theta)$  profiles in the southern passage. The bold profiles show reference stations in the Makarov Basin (<u>Stn-station</u> 145, black) and Amundsen Basin (<u>sStntation</u> 148, gray). Blue, <u>magenta and cyan</u>-profiles are those at the southern end of the section, the three northernmost are red. For <u>exact locationspositions</u> of the stations, please refer to Fig. <u>1 and 12</u>.



**Figure 5:** Silicate (a) and dissolved oxygen (b) from bottle data (circles show sampling depth) in the southern passage (stations 135, 138 and 142) compared with reference stations in the Makarov Basin (MB) and Amundsen Basin (AB). For positions of the stations, please refer to Figs. 1 and 2. Station 149 is located in the westward direction next to station 148 as shown in Fig. 1b.



**Figure 6:** Historical water mass distributions across the Lomonossov Ridge. Potential temperature  $\underline{\theta}$  and salinity profiles and potential temperature-salinity diagrams for 2-two sections taken at 81°N in 1995 (a, c, e, g) and 1996 (b, d, f, h). The black and gray profiles mark the SWERUS-C3 reference stations for the Makarov <u>Basin</u> (station 145) and Amundsen <u>Basin</u> (station 148) basins, respectively. The historical stations are colour-coded according to longitude and water mass characteristics and are marked in the same colour in the maps in panels a and b. <u>Black dots show the SWERUS-C3 stations</u>.



**Figure 7:** Salinity and potential temperature ( $\Theta$ ) profiles at the northern passage. The bold curves show reference stations in the Makarov Basin (<u>Stn-station</u> 145) and Amundsen Basin (<u>Stn-station</u> 148). For positions of the stations, please refer to Fig. 1 and 2.





**Figure 8:** Water mass characteristics across Lomonosov Ridge at the northern passage from a 2011 Polarstern cruise. The black and gray profiles mark the SWERUS-<u>C3</u> reference stations for the Makarov (station 145) and Amundsen basins (station 148), respectively. The historical stations are colour-coded according to longitude and water mass characteristics. Please refer to the map for their exact-positions. Black dots show the SWERUS-C3 stations.

10



**Figure 9:** a) Detail of the northern passage in the vicinity of the sill showing a deepening which is possibly formed by long term seabed erosion of bottom currents. b) Sub-bottom chirp sonar data along the Y-Y' section.