

Response to Referee #1

The paper has been revised according to the comments from the reviewers and we thank both reviewers for their very helpful comments and suggestions. Our point-by-point response is inserted in the reviewer's comments.

Reviewer's Comment (1): The authors are presenting a novel study assessing the ability of a regional ocean atmosphere coupled system to correctly represent ocean convection, especially the sensitivity of the system to the parameterization of turbulent fluxes. First, the authors assessed the model results through detailed comparisons with different observational datasets and show that the coupled system satisfactorily simulates the formation of deep water. After evaluating the uncertainties associated with the different turbulent fluxes parameterizations, the authors carried out several simulations based on 3 commonly used turbulent flux parameterizations. Their results highlight that the choice of the turbulent flux parameterization strongly influences the simulation of open ocean deep convection, especially in terms of volume of newly-formed deep water that can be different from one order of magnitude according to the parameterization choices. Open ocean deep convection plays a key role in the ocean circulation and the results found by the authors are important and will be certainly useful not only for the research groups working on the Mediterranean Sea but also in the North Atlantic, Nordic Seas, and/or Antarctic Seas. From my point of view, the manuscript represents an important contribution to our understanding of modeling deep water formation.

15 If the scientific and presentation quality of this article are good in general, I have some (minor) comments for the authors:

(1): p5,116 & p8,131: What do you mean by "departure"?

Authors' Answer (2): These two sentences have been rephrased.

Author's changes (3): "However, the difference between COARE3.0 and ANDREAS only occurred from wind speeds greater than 16 m s^{-1} (as compared to 8 m s^{-1} for the sensible heat flux)." "As the results were found in good agreement with the observations and did not reveal significant difference between the simulations, they are not presented here. "

(1): p6,111: the "MAW" are not introduced before. What is the difference with the AW?

(2): The Atlantic Water corresponds to the recent surface water which enters the Mediterranean at the Gibraltar strait and spreads in the southern basin. Along its pathway into the Mediterranean basin, this Atlantic water is modified under the effect of surface heat fluxes and of vertical mixing in the convective regions. It becomes the Modified Atlantic water (MAW).

25 (3): "In summer (Fig. 3a), the most stratified water ($SI(1000 \text{ m}) > 120 \text{ kg m}^{-2}$) present in the south corresponds to recent Atlantic Water (AW), while the less stratified water ($SI(1000 \text{ m}) < 80 \text{ kg m}^{-2}$), confined to the north of the deep basin (above 42°N), corresponds to an older Atlantic water mass, which along its pathway into the Mediterranean basin has been modified under the effect of surface heat fluxes and of vertical mixing in the convective regions."

(1): P6,127: "(ECMWF) with a horizontal resolution of $\frac{1}{8}$ " is it the horizontal resolution of the grid choose to export the reanalysis, or is it the resolution of the atmospheric model ?

(2): It is the resolution of the ECMWF atmospheric analysis (16 km)

(1): P8,18: What do you mean by "sound interpretation"?

(2-3): "sound interpretation" has been replaced by "rigorous analysis"

(1): p8,110 : "two simulations do".two simulations.

35 (2): Seems correct with do.

(1): P9,11 : "give good agreement" are in good agreement

(2): Done

(1): p9, 124: Maybe you could had that if the SST does'nt decrease at that time, it's because the mixed layer is deepening and SI continue to decrease.

40 (2): Done

- (1): p10,13 : “... the water column experienced ...” the water column could experience [...], in absence of horizontal advection
 (2): Done
- (1): p11,112-15: Maybe you could quantify this , for example by calculating RMSE for each case and add them to the table. What about the excess of mixing outside the deep-mixing area, is it due to the ocean model or the air-sea flux? Is it more important using the MOON simulation? To answer these questions, it might be interesting to look at the bias in different sub-regions (Northern Current, Deep Mixing, NBF-South) instead of a single one.
 (2): RMSEs have been included in Table 3 and the following text added in the manuscript:
 (3): “The slightly better performance of MOON is confirmed by the RMSEs which are weaker for MOON than for COARE and ANDREAS, including for DEWEX Leg 2 at 1500 m and 2000 m depths.”
- (2): The excess of mixing outside the deep-mixing area, for all simulations, is probably due to the inaccurate initial conditions in the Algerian basin. In the northern part of the domain and south of the NBF, the initial conditions have been corrected following Estournel et al, 2016, but due to the lack of observations initial conditions remained unchanged over the Algerian basin. By advection, any excess of mixing in the Algerian basin will propagate and eventually contaminate the NBF-south region. This issue is currently under investigation and has to be addressed before running multi-year simulations.
- (2): The reviewer’s suggestion regarding sub-domain analysis is interesting and was explored. However, for Dewex Leg 2, due to the limited number of observations, the results were found very sensitive to the definition of the sub-domains. These results are not included in the paper.
- (1): P12, section 5.3.2 Are you expecting to really be able to simulate the exact timing of convective mixing as in the obs? Is the too early mixing in MOON due to too important BMF? By looking figure 11, the stratification at the end of January in the obs seem to be small. Maybe adding the MLD superimposed to the 4 different sections and a 5th sub-panel with a comparison of IS calculated from the simulations and the observation would be clearer for the reader to appreciate the time evolution of the mixing and the difference between simulations and observations.
 (2): To simulate the exact timing of deep convection is a tricky issue due to the weak stratification prevailing before the convective events (end of January). Small differences in the BMF in January can delay (or advance) the triggering of convection from one strong wind episode to the next (or previous) one. The too early mixing in MOON is probably induced by a too large BMF. However, this error on timing is only of a few days and the MOON simulation reproduces fairly well the succession of mixing events. As suggested by the reviewer, the MLD has been superimposed on the 3 simulated sections to ease the comparison. Unfortunately, the lack of observations in the surface boundary layer did not allow the provision of a similar figure for the observations.
- (1): P13, 112-14: “ Our study demonstrates that are strongly sensitive to the turbulent flux parameterizations, not only air surface temperature and moisture but also sea surface temperature” you should simplify this sentence. For example: In addition to air surface temperature and moisture , sea surface temperature is also strongly sensitive to the turbulent flux parameterizations.
 (2): Done
- (1): P13: 126: “ In terms of stratification, the effect of MOON was also found to be positive, with again a general reduction of the bias between observed and computed parameters.” you should simplify this sentence to make it clearer (e.g.: In terms of stratification, the use of MOON also led to a general reduction of the bias between observed and computed parameters.
 (2): Done
- (1): Table2: It seems that there are extra zero in front of some first digits.
 (2): Corrected
- (1): Figures: A lot of the figures are very low resolution (label difficult or impossible to read) and should be of better quality before being published (increase dpi or export in pdf) -
 (2): The quality of the figures has been improved. The new figures have been uploaded.

(1): Figures 4,6,7: you should add in the legend that the grey shaded areas correspond to strong wind periods. The grey areas are also very difficult/impossible to see and should be darker, or you could replace the grey shade by an horizontal line on the upper and lower part of each panel.

(2): The grey areas have been made darker and their meaning added to the caption.

5 (1): Figure 11: During the mixing period in mid-February The instrument at 500m and 700m seem to give a lower potential density than the upper and lower instruments (> 29.12). Is it a calibration issue or a colorbar effect ?

(2): Unfortunately, this is due to a calibration issue. The 500 m observations have been removed from the plot to avoid confusion.

Response to Referee #2

10 The paper has been revised according to the comments from the reviewers and we thank both reviewers for their very helpful comments and suggestions. Our point-by-point response is inserted in the reviewer's comments.

Reviewer's Comment (1): This paper presents a new modelling system consisting in a coupled ocean-atmosphere model and show some results regarding deep-water formation events in the North Western Mediterranean. Additionally the authors run some sensitivity experiments to show the impact of the choice of flux bulk formulas. I think the paper addresses an interesting topic, is well written and the results are interesting. Therefore I recommend it for publication after some issues are addressed. I have recommended a major review because there are many small issues to address, even if none of them are critical.

In general I think that some more details should be provided in what regards the modelling system description and the different bulk formula that are used in the paper, as these are key aspects to understand the results.

20 Authors' Answer (2): Additional information on the coupling platform is now given in the introduction (see detailed comments below). In section 2, the differences between the 3 parameterisations are better presented and discussed (see detailed comments below).

(1): Another issue is that I think the results are not discussed in depth. For instance, an important question that is now present in the modelling community is what is the role of high resolution on the modelling of these type of processes. In this sense, it is not clear to me what part of the improvement brought in this modelling system is due to the high resolution and which part due to the air-sea coupling. Some discussion on this aspect would be appreciated.

30 (2): We agree that these issues are really important for the modelling community. However, the present study was not designed to study to role of the coupling and can not provide very relevant answers with that respect. A preliminary uncoupled experiment (not included in the paper) suggests that in absence of coupling the heat fluxes could be strongly overestimated during the preconditioning of deep ocean convection. These results are consistent with previous findings (eg. Lebeaupin Brossier and Drobinski (2009); Small et al. (2012); Renault et al. (2012)). However, it is not straightforward to distinguish the improvement resulting from the coupling itself (ie from its different feedbacks) from the one resulting from a more accurate atmospheric forcing or a more accurate flux parameterisation. This would require a series of carefully-designed experiments in which the current coupled system would be step-by-step downgraded into an uncoupled system till it would exactly mimic the behaviour of the atmospheric and ocean models in their stand-alone configuration. Currently, this type of work is hampered by the fact the surface fluxes are computed on the atmospheric grid and not on the oceanic grid.

Author's changes (3): Most of these arguments are now given in the conclusion:

"However, this conclusion regarding the good performance of the MOON flux parameterization needs to be further consolidated.

40 First the present results were obtained with a coupled system. They could probably be different with uncoupled simulations. In air-sea coupled simulations, the interactive evolution of ocean and atmosphere influences the turbulent heat fluxes, which themselves modify the atmospheric and oceanic surface fields involved in the flux calculation. In statically unstable Mistral and Tramontane conditions, if the sensible (or latent) heat flux increases, the vertical temperature (or humidity) gradient is reduced, which in turn limits the increase in the sensible (latent) heat flux. It is likely that these feedback loop effects tend to limit the discrepancies induced differences between by the different parameterizations. The results of partial and preliminary uncoupled simulations (not shown) suggest that these discrepancies could be larger than in the coupled simulations. It would be therefore

of great interest to disentangle the effect of the flux formulation from the effect of the air-sea coupling and to check whether the MOON parameterization still improves the results in uncoupled conditions. However, it is not straightforward to isolate the coupling effect in a clean and rigorous way. This requires a series of carefully-designed experiments in which the current coupled system is step-by-step downgraded into an uncoupled system till it exactly mimics the behavior of the atmospheric and ocean models in their stand-alone configuration. In our current system, this type of study is hampered by the fact that the surface fluxes are computed on the atmospheric grid, ie at a coarser resolution than the one used by the ocean model.

The differences in resolution between the atmospheric and ocean models (10 and 1 km, respectively), though partly justified by scale considerations, is also a debatable question. A further development will thus investigate the sensitivity to the resolution of the atmospheric model. In the present configuration, the atmospheric model does not have the possibility of representing scales fully adjusted to that of the oceanic model. In particular, with a 10 km resolution, the local maxima and horizontal gradients of the surface parameters are probably too smooth, which may affect the air-sea interactions especially in the vicinity of the oceanic front (Small et al., 2008) and could also modify the response of the coupled system to the different parameterizations.

In addition, the role of the waves necessitates further investigation. In our study, the waves are not considered in COARE and MOON and only indirectly accounted for in ANDREAS. In ANDREAS, the depth of the spray layer is computed as a function of the significant wave height (Andreas et al., 1995). The latter is rather roughly estimated from a simplified parameterization based on wind speed (Andreas and Wang, 2007). Similar crude relationships are used in COARE3.0 for the wave height and wave period. Another envisaged development will couple the current system with a wave model (Michaud et al, 2012) and revisit the results obtained with the ANDREAS and COARE3.0 parameterizations.

(1): Also, the atmospheric domain looks relatively small so I wonder if the good results of the atmospheric parameters aren't induced by the lateral boundary conditions. Again, what is the role of the coupling in the good quality of the results? Could one obtain similar quality using uncoupled models?

(2): The size of the atmospheric domain is typical of the one used in the field of Numerical Weather Prediction. The fact that the model is forced at its lateral boundaries with an analysis (as opposed to a larger-scale forecast) certainly contributes to improve the results. However, most of the improvement (as compared to regional climate results) is likely due to a better resolution of the topography and thus of the regional winds such as tramontana and mistral which are strongly controlled by the terrain. Further improvement is even expected from a higher resolution of the atmospheric model (on going work) since a 10 km resolution is still insufficient to accurately represent the atmospheric deep convective systems.

(1): Finally, you have shown that the choice of bulk formula have small impacts on the evolution of each parameter but a huge impact on the dense water volume formed (for instance). In your opinion, what should be done to improve the parameterizations? What kind of observations would help to improve them?

(2): The development of accurate flux parameterization in strong wind conditions is an area of research in itself. First, measurements in severe weather are difficult, often inaccurate and/or incomplete (e.g. simultaneous sea state observations are missing). Additional dedicated field campaigns together with wind-water tunnel experiments (Andreas et al. 2016) would certainly help to go one step further. Second, the physical processes taking place in the diphasic surface layer are complex and may be not fully understood yet. Third, there is still a large gap between our understanding of these processes and our ability to represent them in numerical weather predictions models. This is why, as atmosphere-ocean modelers, we should remain particularly attentive to the most recent developments and multiply the experiments to test and evaluate new propositions, would they be fairly pragmatic and model-based (as Moon's) or more sophisticated and physically-based (as Andreas').

(1): Page3 L4-7. As the paper has an important technical component it would be good to provide more details on the platform.

(2): Added/modified text in the introduction

(3): "These issues, among others, have motivated the recent development of a new coupling platform (SURFEX OASIS3-MCT) providing better numerical tools to address the scientific and technical questions related to ocean-wave-atmosphere coupling (Voldoire et al., 2017). This coupling platform is based on an external multi-surface model SURFEX (Masson et al., 2012) and on the OASIS3-MCT (Valcke et al., 2015) coupling interface. SURFEX computes the surface-atmosphere fluxes over four surface types (land, town, ocean and inland waters) and can be used in a stand-alone version with prescribed atmospheric forcing or embedded in an atmospheric models. The use of OASIS3-MCT allows SURFEX to be linked to various other

models including ocean, atmosphere, hydrology, waves and sea-ice models. This generic coupling strategy based upon an externalized surface model ensures that the surface flux computations are done in a consistent way, independently of the models to be coupled. As illustrated in Voldoire et al. (2017), this strategy has greatly facilitated the coupling of the different models developed in the French community, including the coupling of the MESONH atmospheric model (Lafore et al., 1997) and the SYMPHONIE ocean model (Marsaleix et al., 2008, 2009, 2012)"

(1): P3 L7-9. Please, provide more details on what are the conclusions of those studies. Why the air-sea coupling is beneficial? What is it providing? Introduction. I think that the interest of using a coupled system to analyse DWF should be better presented.

(2): Added text in the introduction:

(3)"Regarding the atmospheric forcing, the benefit of using a fully coupled system to study air-sea interactions in the numerical weather prediction models was already illustrated in previous studies based upon different air-sea coupled systems (eg Lebeaupin Brossier and Drobinski, 2009; Small et al., 2012; Renault et al., 2012). These studies have shown that coupled simulations provide a better representation of atmospheric and oceanic surface parameters compared to uncoupled simulations. In particular during strong wind events coupled simulations capture the rapid SST cooling more accurately, which makes the atmospheric boundary layer more stable and reduces the heat and moistures exchanges. It is likely that this improved representation of the atmospheric forcing could also lead to an improved representation of the deep water formation.

Besides the question related to coupling, there is still significant uncertainty as to the choice of a relevant parameterization to compute the turbulent fluxes for strong wind conditions such as the Mistral and Tramontane. Current parameterizations have been carefully assessed and validated against large data sets. However, due to the limited number of available observations in strong wind conditions, they are known to be inaccurate for wind speeds exceeding 20 m s^{-1} (e.g. Hauser et al, 2003). The sensitivity tests performed by Estournel et al., 2016b, suggest that the uncertainty associated with the turbulent flux computations could have a strong impact on the deep water formation process in the NWMS."

(1): P3 L30. How many levels are close to the surface?.

(2): 52 terrain-following vertical levels stretched from 15 m to 15000m, with 16 of them in the first km. This information is now given in the text.

(1): P3 L32. The modelling of convection is of paramount importance in this paper. Thus, more details on how this is parameterized should be included.

(3): "In the case of the ocean, the vertical diffusion is parameterized following Gaspar et al. (1990) with a prognostic equation for the turbulent kinetic energy and a diagnostic relation for the mixing and dissipation lengths. A 1-km resolution is still too coarse to explicitly resolve convective plumes, which thus need to be parameterized. Different parameterizations have been proposed (e.g., Marsland et al. (2003)). The most common and basic one consists in artificially increasing the vertical diffusion coefficient in statically unstable layers (eg Waldman et al. (2016a)). In our case, the heat and water fluxes are linearly distributed over the whole mixed layer, the depth of which is given by the depth at which the vertical density gradient becomes negative. By doing so the first level under the surface does not support the entire amount of heat loss by itself, which prevents the development of static instabilities at the surface. Furthermore, this parameterization is consistent with the nearly linear vertical variation of the buoyancy flux in the convective layer (Deardorff et al., 1969).

(1): Section 2.1. Please, give more details about SURFEX. For non-expert readers its role in the modelling system is confusing.

(2): More details about the role of SURFEX are given in the introduction (see above)

(1): P4 L5. A "3" is missed in OASIS-MCT

(2): Corrected

(1): Section 2.2. P4. L23. The different parameterizations used are for Cd, Ch and Ce? Please, be more clear in the description of the parameterizations and include more details. This is also a very relevant part of the paper and the reader needs to know what are the differences between the different options.

(3): Modified text in section 2. Turbulent air fluxes at the air/sea interface are computed from bulk type parameterizations based on the Monin-Obukhov similarity theory (Foken, 2006). These parameterizations compute the turbulent fluxes as

$$|\tau| = \rho_a u^{*2}$$

$$H = -\rho_a C_p u^* \theta^*$$

$$LE = -\rho_a L_e u^* q^*$$

where τ is the momentum flux, H the sensible heat flux, LE the latent heat flux; ρ_a the air surface density and where u^*, θ^*, q^* are scaling parameters for momentum, potential temperature, and humidity, respectively. The momentum scale, u^* , is referred to as friction velocity.

Classically, the scale parameters are expressed as a function of the vertical gradients of the mean fields at the air-sea interface, the surface roughness and the atmospheric stability. Although based upon the same formalism, the turbulent flux parameterizations differ in the way they specify the different roughness lengths and the so-called stability functions. In particular, the validity of the Charnok's formulation (Charnock, 1955) which is generally used to relate u^* to the dynamic roughness length has often been questioned for strong wind conditions.

In this study, three well-established parameterizations have been used:

- The COARE3.0 parameterization (Fairall et al., 2003) is one of the most widely used in the modelling community. This parameterization derives from the COARE 2.6 algorithm (Fairall et al., 1996) originally developed from the observations performed during the TOGA-COARE experiment (Webster and Lukas, 1992) in the North Pacific. An important upgrade in COARE3.0 is a new formulation of the surface (dynamic and scalar) roughness lengths which slightly increases the fluxes for wind speeds exceeding 10 m s^{-1} . Although COARE3.0 has been validated against a much larger data set (~ 7000 observations) than the one used for COARE2.6, COARE3.0 remains mostly reliable for wind speeds below 20 m s^{-1} due to the limited number of observations available in strong wind conditions. It is worth noting that the influence of waves (available as two possible options in COARE3.0 but not extensively validated) was not activated in our study.
- The ANDREAS parameterization (Andreas et al., 2015) is a novel and more physically-based approach which distinguishes two different contributions to the turbulent heat fluxes: the standard air-sea interfacial fluxes controlled by molecular processes right at the air-sea interface on the one hand, and the sea spray fluxes controlled by microphysical processes around sea spray droplets on the other hand. As opposed to the COARE-type algorithm, the friction velocity used to compute the interfacial fluxes is parameterised as a function of the 10 m wind speed at neutral stability, eliminating thereby the uncertainty associated with the definition of the dynamic roughness length and the use of the (Charnock, 1955) expression. The sea-spray contribution becomes notable only for wind speeds exceeding 13 m s^{-1} . Small droplets are then ejected by surface waves into the atmospheric surface layer. They cool, evaporate and can significantly contribute to the air-sea exchanges of heat and water. The sea spray fluxes are computed using the fast microphysical algorithm described in (Andreas, 2005). ANDREAS parameterization has been established with a data set of ~ 4000 observations with wind speeds up to almost 25 m s^{-1} .
- As opposed to COARE and ANDREAS, the MOON parameterization (Moon et al., 2007) mainly relies upon model results. It has been developed based upon the results of a coupled wave-wind model. The simulations of 10 idealized tropical cyclones have been used to derive a new expression of the dynamic roughness length, which limits the increase of friction velocity with wind for wind speeds exceeding 12.5 m s^{-1} . This new formulation was indirectly validated using the Geophysical Fluid Dynamics Laboratory coupled hurricane-ocean prediction model (Kurihara et al., 1998). For 5 hurricanes observed in the Atlantic Ocean, the new formulation lead to better results than the former one (based upon the Charnock's formulation) with a clear improvement of the cyclone intensity and no degradation of its track and central pressure.

(1): P5 L8 "They also allow the impact of the sea spray in ANDREAS to be distinguished". I don't understand this sentence. Could you please clarify the text here?

(3): Text replaced by "Although not used further in the following, the results of ANDREAS without sea spray effect (ANDREAS no-spray) have been added to assess its impact"

- (1): P7.L6. Please, summarize the conclusions of Estournel et al. (2016a).
- (2-3): This comment was referring to the following sentence : “Sensitivity to the initial state is not discussed here as it has been the subject of a thorough study by Estournel et al. (2016a).” which has been now removed as the conclusion of Estournel et al. (2016a) regarding the sensitivity of model results to the oceanic initial state was already summarized P7.L1 “. Estournel et al. (2016a) show that this correction is necessary to properly simulate the preconditioning phase and the triggering of the convective phase.”.
- (1): P7.L28-32. I think this paragraph is too pessimistic. The agreement between different time series is very high and differences are not so large.
- (3): Rephrased sentences: “For the other surface atmospheric parameters (2 m air temperature and relative humidity), slightly larger discrepancies are found from one simulation to another. Air temperature and humidity remain relatively close to observations in terms of correlation (respectively 0.98 and 0.85, Table 2). Bias and root mean square error exhibit larger but still weak differences between simulations. The largest difference is found for humidity. In particular, it is clear from Fig. 4 c that the moisture drops associated with the strong wind episodes are more pronounced in COARE and ANDREAS than in MOON”
- (1): P8 L1-2. Conversely I think that the extremely high correlations in the SST are over optimistic and due to the seasonal cycle.
- (3): Rephrased sentences: “This correlation is mainly due to the representation of the seasonal cycle and to the weak variability of the SST during the winter period when the SST ceases to evolve. The drops of SST associated with the events of Tramontane and Mistral in autumn are well captured by the three simulations.”
- (1): P8.L5-L7. Can you do a rough estimate of what is the relative importance of each mechanism (local process vs advection) ?
- (2): A rough estimate is provided by Estournel et al 2016a. Integrated during the autumn period the advection process in mass budget represent about 40% compared to local process. This information is now given in the text.
- (1): P8.L9-10. I think ANDREAS shows at least comparable skills with respect to MOON.
- (3): Rephrased sentences: “Nevertheless, it can be concluded from Figs 4 and 5 and Table 2 that in general the results of the MOON and ANDREAS appear to agree with the Lion buoy better than the results of the COARE do and that MOON slightly outperforms ANDREAS .”
- (1): P8. L23-25. I don’t understand this. It looks from the figures that differences between simulations are larger during the peaks. How can you deduce that the feedback mechanism is playing a significant role?
- (2): This discussion has been moved to the conclusion where the potential impact of coupling is now discussed in more details.
- (1): P9. L5. I agree MOON provides the best agreement, but it is just slightly better. Considering the simulation period is relatively small I think you should moderate that statement.
- (3): Rephrased sentence: “Although the differences remain fairly weak, as reflected by the statistical analysis, in our coupled system, the MOON parameterization gives the best agreement with the available observations”
- (1): P13. L13. “.. demonstrates that XXX are strongly ...” . XXX - Something is missed.
- (3): Replaced sentence: “In addition to air surface temperature and moisture, sea surface temperature is also strongly sensitive to the turbulent flux parameterizations.”
- (1): Conclusions. I don’t see that MOON is really outperforming the other parameterizations. For instance, for the SI on Leg-2 COADS seems to produce better results.
- (2): It is true that the MOON bias are not the best ones for DEWEX-Leg2. The conclusion regarding the good performance of MOON has been soften and is also now supported by the analysis of the root mean square errors (which have been added in Table 3).

- (1): Figure 1. Define in the caption what is DWF and NBF.
 (2): Done
- (1): Figure 2 . What is each subplot? What is the x-axis?
 (2): This figure has been redrawn with axis labels.
- 5 (1): Figure 3. What are the colours in (e) ? Isn't it redundant to use them in a time-depth plot?
 (2): The colours in Fig. 3(e) are similar to the ones used in Figs 3a-d and are defined with the colorbar, This color information is redundant in a time-SI plot, but in our opinion help the visualization.
- (1): Figure 4. What are the grey bars in the plots?
 (2): The grey bars correspond to the strong wind periods (hourly wind speed $> 15 \text{ m s}^{-1}$). This information is now given
 10 in the caption.
- (1): Table 1 "sigMa"
 (2): Corrected
- (1): Table 3. Include the averaged SI index obtained from observations, so the biases can be better interpreted.
 (2): Done
- 15 **Summary of all changes made in the manuscript**
- Title: Unchanged
- Abstract: Unchanged
- Introduction: We have added information on the benefit of using a fully coupled system and on the coupled system.
- Section 2.1: We have added information on the oceanic convection parameterization.
- 20 Section 2.1: We have added information on the turbulent fluxes parameterizations.
- Section 3: few rewording, mainly unchanged.
- Section 4: few rewording, mainly unchanged.
- Section 5: few rewording, mainly unchanged.
- Conclusions : We have added discussion on air-sea coupling, atmospheric resolution and waves impact on the turbulent
 25 fluxes.

Modelling deep-water formation in the North-West Mediterranean Sea with a new air-sea coupled model: sensitivity to turbulent flux parameterizations

Léo Seyfried¹, Patrick Marsaleix¹, Evelyne Richard¹, and Claude Estournel¹

¹Université de Toulouse, CNRS, UPS, Laboratoire d'Aérodynamique, Toulouse, France

Correspondence to: Léo Seyfried (leo.seyfried@aero.obs-mip.fr)

Abstract. In the north western Mediterranean, the strong, dry, cold winds, the Tramontane and Mistral, produce intense heat and moisture exchange at the interface between the ocean and the atmosphere leading to the formation of deep dense waters, a process that occurs only in certain regions of the world. The purpose of this study is to demonstrate the ability of a new coupled ocean-atmosphere modelling system based on MESONH-SURFEX-SYMPHONIE to simulate a deep-water formation event in real conditions. The study focuses on summer 2012 to spring 2013, a favourable period that is well documented by previous studies and for which many observations are available. Model results are assessed through detailed comparisons with different observation data sets, including measurements from buoys, moorings and floats. The good overall agreement between observations and model results shows that the new coupled system satisfactorily simulates the formation of deep dense water and can be used with confidence to study ocean-atmosphere coupling in the north-western Mediterranean. In addition, to evaluate the uncertainty associated with the representation of turbulent fluxes in strong wind conditions, several simulations were carried out based on different parameterizations of the flux bulk formulas. The results point out that the choice of turbulent flux parameterization strongly influences the simulation of the deep water convection and can modify the calculated volume of the deep water formed by up to one order of magnitude.

1 Introduction

The North-West Mediterranean Sea (NWMS) is one of the few regions in the world where the deep open ocean convection process is regularly observed (Marshall and Schott, 1999). The strong, dry, cold winds, the Tramontane (north-westerly) and the Mistral (northerly), play a major role in this process. These winds induce intense exchanges between the atmosphere and the sea (Flamant, 2003; Hauser et al., 2003), with a marked loss of surface buoyancy (Schott and Leaman, 1991). The oceanic deep convection can be separated into three phases. In autumn, a cyclonic gyre, bounded to the north by the Northern Current (NC) (Millot, 1999) and to the south by the North Balearic Front (NBF) (Millot and Taupier-Letage, 2005), isolates a weakly stratified water mass whose stratification is progressively eroded by northerly winds (preconditioning phase). In winter, in some years, vertical mixing induced by strong winds leads to the formation of a vertically homogeneous water mass (convective phase) identifiable by its temperature and salinity properties, and generally referred to as new Western Mediterranean Deep Water (nWMDW). After the convective phase, the mixed zone undergoes a restratification while the nWMDW is spread throughout the basin by the general circulation (Schott et al., 1996) and submesoscale eddies (Testor and Gascard, 2006) (restratification/spreading phase).

Two components of the MISTRALS programme (<http://www.mistrals-home.org/>) focused on the study of deep convection in the NWMS and made a major effort in collecting observations during 2012 and 2013. The first component, HyMeX (Drobinski et al., 2013), studied the atmosphere-land-ocean coupled system. In this context, two Special Observation Periods (SOPs) were organized; SOP1 in autumn 2012, during the preconditioning phase (Ducrocq et al., 2013), and SOP2 in winter 2013, during the convection phase (Estournel et al., 2016b). The second component, MERMEX (Sempéré et al., 2010) studied the impact of oceanic deep convection on the nutrient balance and the planktonic ecosystem. Three oceanographic cruises were organized by MERMEX: DOWEX in autumn 2012, DEWEX leg 1 in February 2013 and DEWEX leg 2 in April 2013. Combining all these different surveys provided a good description of the spatial distribution and temporal evolution of ocean stratification. Furthermore, the volume of dense water formed by deep convection could be evaluated thanks to the optimal interpolation of the many Conductivity-Temperature-Density (CTD) profiles available (Waldman et al., 2016b). This series of campaigns provided a unique opportunity to test the ability of models to simulate the different phases of the dense water formation process in the western Mediterranean.

Several modelling studies of the formation of deep water in the NWMS, have been carried out over different periods and with different models Herrmann et al. (2008); Herrmann and Somot (2008); Herrmann et al. (2010); Léger et al. (2016); Estournel et al. (2016a); Waldman et al. (2016a). Their results show high sensitivity to the initial stratification of the ocean model and to the accuracy of the atmospheric forcing. An attempt to progress on these two issues was proposed by the HYMEX and MERMEX groups. Concerning the first point, since 2010, oceanographic cruises have been organized in the NWMS each summer by the Mediterranean Ocean Observing System for the Environment (MOOSE; www.moose-network.fr/). These cruises provide a sample of the different water masses of the NWMS based on about 70 CTD profiles. In particular, the observations collected in summer 2012 have been assimilated to provide a more realistic initial state for ocean models (Estournel et al., 2016a; Léger et al., 2016) which has been shown to be crucial for the simulation of the winter convection 6 months later. Regarding the

atmospheric forcing, ~~besides the questions related to its space and time resolution, the benefit of using a fully coupled system to study air-sea interactions in numerical weather prediction models was already illustrated in previous studies based upon different air-sea coupled systems (e.g., Lebeaupin Brossier and Drobinski (2009); Small et al. (2012); Renault et al. (2012)).~~ These studies have shown that coupled simulations provide a better representation of atmospheric and oceanic surface parameters compared to uncoupled simulations. In particular, during strong wind events coupled simulations capture the rapid SST cooling more accurately, which makes the atmospheric boundary layer more stable and reduces the heat and moistures exchanges. It is likely that this improved representation of the atmospheric forcing could also lead to an improved representation of the deep water formation.

Besides the question related to coupling, there is still significant uncertainty as to the choice of a relevant parameterization to compute the turbulent fluxes for strong wind conditions such as the Mistral and Tramontane (Estournel et al., 2016b). Current parameterizations have been carefully assessed and validated against large data sets. However, due to the limited number of available observations in strong wind conditions, they are known to be inaccurate for wind speeds exceeding 20 ms^{-1} (e.g., Hauser et al. (2003)). The sensitivity tests performed by Estournel et al. (2016a) suggest that the uncertainty associated with the turbulent flux computations could have a strong impact on the deep water formation process in the NWMS.

These issues, among others, have motivated the recent development of a new modelling platform coupling platform (SURFEX OASIS3-MCT) providing better numerical tools to address the scientific and technical questions related to ocean-wave-atmosphere coupling (Voldoire et al., 2017). ~~In particular, this new platform greatly facilitates~~ This coupling platform is based on a multi-surface model SURFEX ((Masson et al., 2012) and on the OASIS3-MCT (Valcke et al., 2015) code coupler. SURFEX computes the surface-atmosphere fluxes over four surface types (land, town, ocean and inland waters) and can be used in a stand-alone version with prescribed atmospheric forcing or embedded in an atmospheric models. The use of OASIS3-MCT allows SURFEX to be linked to various other models including ocean, land, atmosphere, hydrology, waves and sea-ice models. This generic coupling strategy based upon an externalized surface model ensures that the surface flux computations are done in a consistent way, independently of the models to be coupled. As illustrated in Voldoire et al. (2017) , this strategy has greatly facilitated the coupling of the different ~~atmospheric and ocean~~ models developed in the French community. ~~The benefit of using a fully coupled system to study air-sea interactions in the NWMS was already illustrated in previous studies (based upon different air-sea coupling platforms) (Lebeaupin Brossier and Drobinski, 2009; Small et al., 2012; Renault et al., 2012), including the coupling of the MESONH atmospheric model (Lafore et al., 1997) and the SYMPHONIE ocean model (Marsaleix et al., 2008, 2009, 2012) .~~

A first objective of the present study is to show the capacity of the new coupled regional ocean-atmosphere system MESONH-SURFEX-SYMPHONIE to reproduce the formation of deep dense waters during the winter of 2013, in the NWMS. A second objective is to study the sensitivity of the simulations to the parameterization of the turbulent surface fluxes by testing three different parameterizations (Fairall et al., 2003; Andreas et al., 2015; Moon et al., 2007), all based on bulk formulas.

This paper is organized as follows. Section 2 describes the coupled modelling system, and the three parameterizations mentioned above that were used for the computation of turbulent fluxes. Section 3 presents the different observation data sets

used to evaluate the model results while section 4 details the setup of the numerical experiments. Results are analysed and discussed in section 5. Some conclusions and perspectives are presented in section 6.

2 Model description

2.1 Air-Sea Coupled model

5 The modelling system is based on the non-hydrostatic atmospheric model MESONH [Lafore et al. (1997); <http://mesonh.aero.obs-mip.fr/mesonh52/>] developed at the Laboratoire d’Aérodynamique (LA) and Centre National de Recherches Météorologiques (CNRM), the multi-surface model SURFEX [Masson et al. (2012); <http://www.cnrm-game-meteo.fr/surfex/>] developed at CNRM, and the Boussinesq hydrostatic ocean model SYMPHONIE [Marsaleix et al. (2008, 2009, 2012); http://sirocco.omp.obs-mip.fr/ocean_models/S-model] developed at LA. These models are coupled through the [SURFEX-OASIS interface \(Voldoire et al., 2017\)](#) using the [OASIS3-MCT coupler \(Valeke et al., 2015\)](#) [SURFEX-OASIS3-MCT interface \(Voldoire et al., 2017\)](#).

10 The main characteristics of the atmospheric and oceanic model configurations are listed in Table 1. The oceanic model uses a horizontal resolution of 1 km. Given the value of the Rossby radius (5-10 km in the NWMS), 1-km grid spacing appears to constitute a reasonable compromise between the computing cost and the necessary resolution. In the vertical, 40 generalized sigma levels are used; 10 of them in the first hundred meters (above the abyssal plain). The resolution just below the

15 sea surface is 1.5 m. The atmospheric model is run with a 10-km horizontal grid spacing and 52 terrain-following vertical levels ranging from 15 m to 15000 m, [with 16 of them in the first km](#). With such resolutions, both atmospheric and oceanic convection must be parameterized. In the case of the ocean, ~~as described in Estournel et al. (2016a)~~, [the vertical diffusion is parameterized following Gaspar et al. \(1990\) with a prognostic equation for the turbulent kinetic energy and a diagnostic relation for the mixing and dissipation lengths. A 1-km resolution is still too coarse to explicitly resolve convective plumes, which thus need to be parameterized. Different parameterizations have been proposed \(e.g., Marsland et al. \(2003\) \). The most common and basic one consists in artificially increasing the vertical diffusion coefficient in statically unstable layers \(e.g., Waldman et al. \(2016a\) \). In our case, the heat and water fluxes are re-distributed-linearly distributed over the whole mixed layer. This, the depth of which is given by the depth at which the vertical density gradient becomes negative. By doing so the first level under the surface does not support the entire amount of heat loss by itself, which prevents the development of static instabilities at the surface. Furthermore, this](#) parameterization is consistent with the ~~nearly-linear~~ [nearly linear](#) vertical variation of the buoyancy flux in the convective layer (Deardorff et al., 1969). Regarding the atmosphere, shallow and deep convection are parameterized with mass-flux schemes according to Bechtold et al. (2001) and Pergaud et al. (2009), respectively.

25

In the coupled system, the surface fluxes are computed by SURFEX on the atmospheric model grid. They are sent to the ocean model by the OASIS3-MCT coupler, which also performs their interpolation on the ocean model grid. Conversely, the

30 ocean model computes the sea surface temperature and sends it to SURFEX using the ~~OASIS-MCT~~ [OASIS3-MCT](#) coupler, which takes care of its interpolation on the atmospheric model grid. The coupling frequency is set to 10 minutes and the interpolation between the two model grids is bilinear.

The computational domains used for this study are presented in Fig. 1. The atmospheric grid covers the whole western Mediterranean basin while only part of it is covered by the ocean grid (blue area in Fig. 1) since the Alboran Sea and part of the Tyrrhenian Sea were excluded to avoid straits issues. Outside the ocean grid, the air-sea fluxes are computed using the sea surface temperature provided by the OSTIA data base (Donlon et al., 2012), the horizontal resolution of which is about 6 km.

5 2.2 Turbulent flux parameterizations

~~The turbulent~~ Turbulent air fluxes at the air/sea interface are ~~calculated from bulk-type~~ computed from bulk type parameterizations based on the Monin-Obukhov similarity theory (Foken, 2006). These parameterizations compute the turbulent fluxes as

$$\underline{|\tau| = \rho_a u^{*2}} \quad (1)$$

$$10 \quad \underline{H = -\rho_a C_p u^* \theta^*} \quad (2)$$

$$\underline{LE = -\rho_a L_e u^* q^*} \quad (3)$$

where τ is the momentum flux, H the sensible heat flux, LE the latent heat flux; ρ_a the air surface density and where u^ , θ^* , q^* are scaling parameters for momentum, potential temperature, and humidity, respectively. The momentum scale, u^* , is referred to as friction velocity.*

15 Classically, the scale parameters are expressed as a function of the vertical gradient-gradients of the mean fields at the air-sea interface (~~Eqs. 4, 4 and 4~~)

$$\underline{|\tau| = \rho_a C_d U^2}$$

$$\underline{H = -\rho_a C_p C_h (\theta - SST) U}$$

$$\underline{LE = -\rho_a L_e C_e (q - q_{sat}(SST)) U}$$

20 *where τ is the momentum flux, H the sensible heat flux, LE the latent heat flux; ρ_a the air surface density, C_p the specific heat of air, L_e the latent heat of vaporization, U the wind speed at 10, θ the air potential temperature at 10, SST the sea surface temperature, q the specific humidity at 10, $q_{sat}(SST)$ the specific humidity at saturation for air in contact with the sea surface, and C_d , C_h and C_e the transfer coefficients for momentum, heat and moisture. In this study three different parameterizations are tested* the surface roughness and the atmospheric stability. Although based upon the same formalism,
 25 the turbulent flux parameterizations differ in the way they specify the different roughness lengths and the so-called stability functions. In particular, the validity of the Charnok's formulation (Charnock, 1955) which is generally used to relate u^* to the dynamic roughness length has often been questioned for strong wind conditions.

In this study, three well-established parameterizations have been used:

- The COARE3.0 parameterization (Fairall et al., 2003) is one of the most widely used in the modelling community ~~but was~~
 30 ~~established for winds weaker than~~ This parameterization derives from the COARE 2.6 algorithm (Fairall et al., 1996) originally

developed from the observations performed during the TOGA-COARE experiment (Webster and Lukas, 1992) in the North Pacific. An important upgrade in COARE3.0 is a new formulation of the surface (dynamic and scalar) roughness lengths which slightly increases the fluxes for wind speeds exceeding 10 m s^{-1} . Although COARE3.0 has been validated against a much larger data set (~ 7000 observations) than the one used for COARE2.6, COARE3.0 remains mostly reliable for wind speeds below 20 m s^{-1} -due to the limited number of observations available in strong wind conditions. It is worth noting that the influence of waves (available as two possible options in COARE3.0 but not extensively validated) was not activated in our study.

- The ANDREAS parameterization (Andreas et al., 2015) is a novel and more physically-based approach distinguishing the fluxes at the which distinguishes two different contributions to the turbulent heat fluxes: the standard air-sea interface (computed with COARE2.6 (Fairall et al., 1996)) -interfacial fluxes controlled by molecular processes right at the air-sea interface on the one hand, and the sea spray fluxes (calculated with a fast microphysical algorithm) controlled by microphysical processes around sea spray droplets on the other hand. This parameterization was established with observations for As opposed to the COARE-type algorithm, the friction velocity used to compute the interfacial fluxes is parameterised as a function of the 10 m wind speed at neutral stability, eliminating thereby the uncertainty associated with the definition of the dynamic roughness length and the use of the Charnock (1955) expression. The sea-spray contribution becomes notable only for wind speeds exceeding 13 m s^{-1} . Small droplets are then ejected by surface waves into the atmospheric surface layer. They cool, evaporate and can significantly contribute to the air-sea exchanges of heat and water. The sea spray fluxes are computed using the fast microphysical algorithm described in Andreas (2005). ANDREAS parameterization has been established with a data set of ~ 4000 observations with wind speeds up to almost 25 m s^{-1} .
- The As opposed to COARE and ANDREAS, the MOON parameterization (Moon et al., 2007) proposes an alternative parameterization of mainly relies upon model results. It has been developed based upon the results of a coupled wave-wind model. The simulations of 10 idealized tropical cyclones have been used to derive a new expression of the dynamic roughness length, which limits the increase of friction velocity with wind for wind speeds exceeding 12.5 m s^{-1} . This new formulation was indirectly validated using the momentum transfer coefficient (C_d) which was developed and validated for strong winds and cyclones from wave-wind coupled simulations and applied to the Geophysical Fluid Dynamics Laboratory coupled hurricane-ocean prediction model (Kurihara et al., 1998). It is worth noting that the new formulation of the momentum transfer coefficient also impacts the turbulent heat fluxes since, in MOON, C_h and C_e are given as a functions of C_d . For 5 hurricanes observed in the Atlantic Ocean, the new formulation lead to better results than the former one (based upon the Charnock's formulation) with a clear improvement of the cyclone intensity and no degradation of its track and central pressure.

COARE3.0 is the standard parameterization used in SURFEX, while ANDREAS and MOON were implemented in SURFEX specifically for the purposes of the study.

To illustrate the discrepancies between these parameterizations, Fig. 2 shows the turbulent fluxes plotted as a function of the 10 m wind speed and computed with the different parameterizations. Although not used further in the following, the results of ~~COARE2.6-ANDREAS without sea spray effect (ANDREAS no-spray)~~ have been added for completeness. ~~They also allow the impact of the sea spray in ANDREAS to be distinguished.~~ The computations were carried out under unstable conditions typical of Mistral and Tramontane events : the 10 m air temperature fixed at 10 °C, the sea surface temperature at 15 °C and the 10 m relative humidity at 70 %. In these conditions, COARE3.0 and ANDREAS produced very similar momentum fluxes while the MOON momentum flux was found to be slightly weaker but only for the strongest winds ($>15 \text{ m s}^{-1}$). For the sensible heat flux, COARE3.0 and ANDREAS provided close results up to a wind speed 8 m s^{-1} but, for higher wind speeds, the impact of the sea spray considerably enhanced the ANDREAS sensible heat flux. Among the three parameterizations, MOON was the one that produced the strongest sensible heat flux over a wide range of wind speeds (from 6 m s^{-1} to 23 m s^{-1}). The latent heat fluxes presented the same hierarchy in intensity as the sensible heat fluxes. However, the ~~departure-difference~~ between COARE3.0 and ANDREAS only occurred from wind speeds greater than 16 m s^{-1} (as compared to 8 m s^{-1} for the sensible heat flux). To summarize, the three parameterizations gave significantly different results, particularly MOON, which produced the largest turbulent heat fluxes in a wide range of wind speeds. It should also be noted that the ~~departure-differences~~ between the parameterizations may become noticeable from wind speeds as low as 8 m s^{-1} and can dramatically increase with the wind speed, especially for the heat fluxes.

3 Available Observations

Several in situ datasets, collected by the MOOSE, HyMeX and MerMeX programmes, are used in the present study and are briefly described below. More details are available in Estournel et al. (2016b). First, the Lion meteorological buoy provides hourly measurements of the air temperature and humidity at 2 m above the sea surface, the wind speed at 10 m, and the sea surface temperature. A mooring named LION, part of the MOOSE network, and located only 5 km away from the Lion buoy is also available. It provides vertical profiles of temperature and salinity, sampled over 21 levels for temperature and 15 for salinity, between 150 and 2300 m deep (Houpert et al., 2016).

In addition, several oceanographic cruises were carried out in the NWMS during 2012-2013 (Waldman et al., 2016b). In August 2012, the annual cruise organized by MOOSE for the monitoring of the NWMS, provided 69 vertical surface-to-bottom profiles of temperature and salinity. The same network of CTD stations was deployed in February 2013 and April 2013, during the DEWEX leg 1 and leg 2 cruises. A smaller network was implemented in autumn 2012 during the DOWEX cruise, with only 41 profiles. Furthermore, 399 temperature and salinity profiles measured by 14 Argo floats, 4 of them specifically deployed by HyMeX with a daily cycle, are also taken into consideration in this study. The above-mentioned cruises allowed the spatial distribution and temporal evolution of ocean stratification to be well described.

Ocean stratification is commonly assessed using the Stratification Index (SI Eq. 4).

$$SI(Z) = \int_Z^0 (\rho(Z) - \rho(z)) dz \quad (4)$$

where ρ is the potential density and Z the reference level. SI is expressed in kg m^{-2} .

Note : that $SI(Z) = 0$, means that the water column is mixed at least to depth Z or, in other words, that the mixed layer depth is greater than or equal to Z .

5

Figures 3 (a-d) show the Stratification Index SI at 1000 m (Eq. 4) for each seasonal cruise and Fig. 3 e its time evolution from summer 2012 to spring 2013. In summer (Fig. 3 a), the most stratified water ($SI(1000\text{ m}) > 120\text{ kg m}^{-2}$) present in the south corresponds to recent Atlantic Water (AW), while the less stratified water ($SI(1000\text{ m}) < 80\text{ kg m}^{-2}$) is, confined to the north of the deep basin (above 42°N)~~and corresponds to the Modified Atlantic Water (MAW), while the more stratified water~~
 10 ~~($SI(1000\text{ m}) > 120$) in the south corresponds to the Atlantic Water (AW), corresponds to an older Atlantic water mass, which~~
along its pathway in the Mediterranean basin has been modified under the effect of surface heat fluxes and of vertical mixing in the convective regions. These two water masses are separated by the North Balearic Front. In autumn (Fig. 3 b) the northward displacement of this front brings AW toward the north of the basin, whereas the oceanic stratification increases in the south of the domain. In winter (Fig. 3 c), the oceanic deep convection is revealed by many profiles with $SI(1000\text{ m}) < 1\text{ kg m}^{-2}$,
 15 meaning that these profiles are mixed down to 1000 m depth. When the vertical mixing reaches the seafloor, it can then lead to the formation of dense deep water. Finally, in spring (Fig. 3 d), restratification occurs with an increase of stratification in the convective area ($SI(1000\text{ m}) > 10\text{ kg m}^{-2}$).

Combining all these available observations, Waldman et al. (2016a) estimated the volume of the dense deep water formed at $4.5(\pm 1.1) \times 10^{13}\text{ m}^3$ (for density greater than 29.11 kg m^{-3}) between summer 2012 (MOOSE 2012 cruise) and spring 2013
 20 (DEWEX leg 2 cruise) in the convective area defined between $[2.5^\circ\text{E}; 9^\circ\text{E}]$ and $[40^\circ\text{N}; 44^\circ\text{N}]$ and limited by the 2000 m bathymetry contour (red line in Figs 3 a-d)

4 Numerical Experiments

To model this well documented event of dense water formation in the NWMS, three coupled simulations are reported here. They differ only in their turbulent flux formulation and are named COARE, ANDREAS, and MOON according to their pa-
 25 rameterizations. These simulations start on 16 August 2012 with a realistic initial oceanic state and end on 1 May 2013. Initial and boundary conditions of the atmospheric model are provided by the 6-hour analyses produced by the European Centre for Medium-range Weather Forecasts (ECMWF) with a horizontal resolution of $1/8^\circ$. Initial and boundary conditions of the ocean model are provided by the analyses of the MERCATOR-OCEAN operational centre (Lellouche et al., 2013) based on the NEMO ocean model (Maraldi et al., 2013). These ocean fields are corrected according to the method proposed by Estour-
 30 nel et al. (2016a). Practically, the vertical background stratification is corrected using the CTD profiles of the MOOSE 2012

oceanographic cruise. Estournel et al. (2016a) show that this correction is necessary to properly simulate the preconditioning phase and the triggering of the convective phase. In this study, to also improve the sea surface temperature initial state, the surface temperature is restored toward the SST satellite data (Buongiorno Nardelli et al., 2013) during the 14 days preceding the beginning of the simulations using a restoring time scale of 3 days. At the end of this process, we obtain a surface temperature
5 that closely matches the satellite observation of 16 August, 2012. ~~Sensitivity to the initial state is not discussed here as it has been the subject of a thorough study by Estournel et al. (2016a).~~ During the rest of the simulation, the coupled model evolves freely, without further data assimilation and without any restoring or nudging procedure.

5 Results

5.1 Air-sea exchanges at Lion meteorological buoy

10 5.1.1 Atmospheric and oceanic surface parameters

A first evaluation of the simulations is carried out by comparing the computed fields with the corresponding measurements at the Lion buoy. This buoy has already been used in various previous studies to validate atmospheric and/or oceanic simulations (Lebeaupin-Brossier et al., 2014; Léger et al., 2016; Rainaud et al., 2016). Figure 4 shows the temporal evolution of surface atmospheric and oceanic parameters (10 m wind speed, 2 m temperature and humidity and sea surface temperature) computed
15 at the buoy, together with the corresponding observations. The same surface parameters are also presented in Fig. 5 in the form of scatter plots (simulations versus observations) while the associated statistics (bias, root mean square error and correlation coefficient) are given in Table 2.

Figure 4 a allows the strong Tramontane and Mistral events to be identified (with hourly wind speeds exceeding 15m/s outlined in grey), alternating with calm wind situations. The Mistral and Tramontane episodes are systematically accompanied
20 by marked drops in temperature and moisture (Figs 4 b and 4 c). By a cumulative effect, the succession of strong wind events in autumn leads to a progressive decrease of the sea surface temperature (Fig. 4 d), which reaches its minimum value of 12.9 °C (i.e. the temperature of the deep water) in early winter and then remains nearly constant during the convective period.

All three simulations accurately reproduce the time evolution of the wind speed at the buoy throughout the 8-month period, with a correlation of 0.9 and a bias lower than 0.2 m s^{-1} . In particular, the timing of the strong wind events is well captured.
25 Moreover, the wind speed maxima are well represented (Fig. 5 a), which is essential to correctly reproduce the intense air-sea exchanges associated with convection (Herrmann and Somot, 2008). Finally, there is no significant difference in wind speed between the different simulations (Figs 4 a, 5 a and Table 2).

For the other surface atmospheric parameters (2 m air temperature and relative humidity), slightly larger discrepancies are found from one simulation to another. Air temperature and humidity remain relatively close to observations in terms of
30 correlation (respectively 0.98 and 0.85, Table 2). ~~In contrast, bias~~ Bias and root mean square error exhibit larger but still weak differences between simulations, ~~especially~~. The largest difference is found for humidity. In particular, it is clear from Fig. 4

c that the moisture drops associated with the strong wind episodes are more pronounced in COARE and ANDREAS than in MOON.

The calculated sea surface temperature is remarkably well correlated with the observations for all simulations (> 0.98 , Table 2). This ~~is partly correlation is mainly~~ due to the ~~correct~~-representation of the ~~drop of SST during the events of Tramontane and Mistral in autumn but is also influenced by the very seasonal cycle and to the~~ weak variability of the SST during the winter period when the SST ceases to evolve. The drops of SST associated with the events of Tramontane and Mistral in autumn are well captured by the three simulations. However, there are significant differences between simulations in autumn. In particular, the COARE SSTs appear overestimated (Figs 4 d and 5 d). During this period, the Tramontane and Mistral events produce a cooling in response to the enhanced turbulent heat fluxes on the one hand and to the temperature advection associated with the northward displacement of the NBF on the other hand (Estournel et al., 2016a). Integrated during the autumn period the advection processes in mass budget represent about 40 % compared to local processes. In such conditions, it is clear that a reasoning limited to the local vertical exchanges is insufficient to provide a ~~sound interpretation rigorous analysis~~ of the results. Nevertheless, it can be concluded from Figs 4 and 5 and Table 2 that in general the results of ~~the MOON simulation MOON and ANDREAS~~ appear to agree with the Lion buoy ~~observations~~ better than the results of ~~the other two simulations~~ ~~do COARE do and that MOON slightly outperforms ANDREAS.~~

5.1.2 Air-sea fluxes

The time evolution of the turbulent fluxes computed at the buoy is shown in Fig. 6. Unfortunately, as there was no flux measurement at the buoy, this figure is limited to model-model comparison. During the strong wind episodes, all the turbulent fluxes are strongly enhanced and the air-sea exchanges are intensified. As suggested by Fig. 2, the wind stress is very similar for all three simulations (Fig. 6 a), while the sensible and latent heat fluxes (Figs 6 b and c) can differ significantly from one simulation to another, especially during the strong Mistral and Tramontane wind events. During these events, in accordance with Fig. 2, the strongest sensible and latent heat fluxes are obtained with MOON and the weakest with COARE. ANDREAS produces a sensible heat flux similar to that of MOON, and a weaker latent heat flux more similar to that of COARE. For example, on 28 November (corresponding to one of the strongest wind episodes), the daily average latent heat flux reaches 1100 W m^{-2} in MOON versus only 780 W m^{-2} in COARE and ANDREAS. On the same day, the sensible heat flux is 390 W m^{-2} in MOON and ANDREAS compared to only 300 W m^{-2} in COARE. ~~In air-sea coupled simulations, the interactive evolution of ocean and atmosphere influences the turbulent heat fluxes, which themselves modify the atmospheric and oceanic surface fields (Fig. 4) involved in the flux calculation. In statically unstable Mistral and Tramontane conditions, if the sensible (or latent) heat flux increases, the vertical temperature (or humidity) gradient is reduced, which in turn limits the increase in the sensible (latent) heat flux. These feedback effects tend to reduce the differences between the simulations.~~

Furthermore, during the autumn, as shown in Fig. 4 d, the sea surface temperature evolves differently in each simulation. This is particularly the case for the COARE simulation, which presents significantly warmer SSTs than the other two simulations and than observations. This results in larger turbulent heat fluxes during low wind periods in autumn for the COARE simulation than for ANDREAS and MOON (Fig. 6).

To complement the analysis, the radiative fluxes and precipitation, which also contribute to the air/sea exchanges, have been analysed. ~~As neither any major departures from the observations nor any significant differences~~ The results are in good agreement with the observations and do not reveal any significant difference between the simulations ~~were found~~, they are not ~~shown~~ presented here.

- 5 In summary, all the simulations generally ~~give are in~~ give are in good agreement with the surface parameters observed at the Lion buoy throughout the eight months considered, although significant punctual differences may appear between the different simulations, especially during the Tramontane and Mistral events. These differences mainly affect the heat and moisture exchanges whereas the momentum exchanges are very weakly impacted. ~~As~~ Although the differences remain fairly weak and as reflected by the statistical analysis, in our coupled system, the MOON parameterization gives the best agreement with the available
10 observations. However, considering the impossibility of directly validating the air-sea fluxes and also the multiple sources of uncertainty in such a complex modelling system (and their possible compensations) it is not clear whether the MOON flux parameterization is better per se or whether it is simply the most suitable parameterization for our modelling system.

5.2 Impact of the air-sea exchanges on the oceanic stratification

5.2.1 Buoyancy Mass fluxes and oceanic stratification

- 15 The air-sea exchanges are now assessed through the Buoyancy Mass Flux (BMF). This flux, directly linked to turbulent fluxes but also to radiative fluxes and precipitation, is formulated as follows:

$$BMF = \alpha \frac{Q_{net}}{C_p} + \beta SSS \rho_0 (E - P) \quad (5)$$

- 20 where α is the thermal expansion coefficient in K^{-1} , Q_{net} is the net heat flux (sum of the net radiative flux, and turbulent heat fluxes) in $W m^{-2}$, C_p is the specific heat capacity in $J kg^{-1} K^{-1}$, β is the saline contraction coefficient in psu^{-1} , SSS is the surface salinity in psu , ρ_0 is a reference density in $kg m^{-3}$, E is the evaporation and P the precipitations in $m s^{-1}$. BMF is expressed in $kg m^{-2} s^{-1}$.

- The time evolution of the buoyancy mass flux computed at the Lion buoy and for each simulation is shown in Fig. 7 a (instantaneous values) and 7 b (cumulative values). Its evolution closely follows (with opposite sign) the evolution of the
25 turbulent fluxes. The maximum buoyancy losses appear during Tramontane and Mistral events, which account for about 70 % of the total buoyancy loss. The simulations can be segmented into three periods; first the preconditioning period (16 August to 15 January) with a decrease in the cumulated buoyancy mass flux and in SST, then the convective period (15 January to 21 March) with a decrease in the cumulated buoyancy mass flux without significant decrease in SST (since the water column is nearly mixed) and, finally, the start of the restratification period (21 March to 30 April) when the buoyancy mass flux starts to
30 increase. As can be seen in Fig. 7 b, the buoyancy loss during the preconditioning period is nearly equivalent to the buoyancy

loss during the convective period. The largest cumulative buoyancy loss at the end of the simulations is obtained with MOON (195 kg m^{-2}) followed by COARE (170 kg m^{-2}) and ANDREAS (165 kg m^{-2}). By the end of the simulations, COARE has produced a slightly larger loss of buoyancy than ANDREAS while turbulent fluxes for the Tramontane and Mistral events are greater in ANDREAS. This is due to the larger buoyancy mass flux during the calm wind periods of the preconditioning phase in COARE.

Figure 7 c shows the time evolution of the stratification index (Eq. 4) relative to the 2000 m depth at the location of the buoy. At the beginning of the simulations the stratification is 160 kg m^{-2} , i.e. less than the total buoyancy loss produced by any of the simulations, suggesting, that in all three simulations, in absence of horizontal advection, the water column **experienced could experience** full mixing down to a depth of 2000 m at the buoy location. The decrease of the stratification index with time is not continuous and its evolution is not strictly directly correlated with the evolution the cumulated buoyancy mass flux (as it would be in a one-dimensional system). Even if most of the stratification losses occurs mainly during the Tramontane and Mistral, during the weak wind periods, despite the slow decrease of cumulated buoyancy mass flux, the stratification index may increase.

Figure 8 places these local results in a wider context and shows the spatial distribution of the cumulated buoyancy mass fluxes during the whole simulation period, the preconditioning period, and the convective period for each simulation. During the whole simulation period, three distinct maxima of buoyancy loss appear, two in the Gulf of Lion located within Tramontane and Mistral corridors, and a third one located further south. Whereas the first two stem from local wind maxima, the southern maximum, only present during the preconditioning period, is more related to the warm waters brought over by the seasonal northward displacement of the NBF. When the Tramontane or Mistral blows over this warm patch, the event is associated with strong air-sea temperature gradients and enhanced turbulent heat fluxes. This explains the very strong buoyancy mass flux losses observed during the autumnal Tramontane and Mistral events at the buoy location (Fig. 7 a), which is situated in the vicinity of the frontal zone. This suggests that, during the preconditioning period, the dynamics of the NBF plays a major role in the loss of surface buoyancy in the deep water zone. Furthermore, the front displacement is modulated by the wind intensity. The increase (decrease) of stratification during the period of weak (strong) winds (Fig. 7 c) is due to the lateral advection of light (heavy) water by the northward (southward) displacement of NBF (Estournel et al., 2016a). Because of these horizontal processes and their feedback on the buoyancy mass flux, the three simulations experience different time evolutions. The large discrepancy seen between the COARE and MOON buoyancy mass fluxes and stratification indices may also result from differences in the NBF progression. In winter, during the convective phase, only the two maxima of the Gulf of Lion remain. In the deep water zone, as the SST has reached a nearly-constant value, there is no significant effect of the SST structures on the buoyancy mass flux. At the Lion buoy, the buoyancy mass flux is not affected by the SST and the local increase in stratification during the period of weak winds (Fig. 7 c) is principally due to the advection of light water into the mixing zone by baroclinic instability (Marshall and Schott, 1999).

The comparison of the three simulations highlights the impact of the surface flux parameterization. In MOON, the buoyancy mass flux losses are stronger than in COARE and affect a much wider area.

5.2.2 Validation of oceanic stratification with oceanographic cruises

To assess the evolution of the oceanic stratification, Stratification Indices collocated in space and time with all CTD and Argo profiles were calculated for each simulation and compared with the corresponding values deduced from the observations. Results were analysed in terms of bias (observations minus simulations). Figure 9 shows the spatial distribution of the SI (1000 m) bias obtained for each simulation and for the two DEWEX oceanographic cruises (leg 1 in winter 2013 and leg 2 in spring 2013) while Table 3 gives the values of the averaged SI [bias for observations and the SI simulations bias and root mean square error \(RMSE\)](#) computed in the convective area for different depths (1000, 1500 and 2000 m).

During the convective phase (winter 2013), as shown in Figs 3 c and e, an area of mixed profiles (with $SI(1000m) < 10kg\ m^{-2}$) induced by the oceanic deep convection is present in the centre of the basin. This area is surrounded by more stratified waters corresponding to the NC to the North and to the NBF to the South. From Figs 9 a-c, it is clear that all the simulations present an excess of stratification in the convective area (negative bias) and an excess of mixing south of the NBF (positive bias). However, it is noteworthy that the SI (1000 m) bias is significantly reduced in the MOON simulation (Fig. 9 c). The good performance of MOON with respect to the stratification is further confirmed by Table 3: at 1500 and 2000 m, the SI bias reduction obtained with MOON is even more spectacular. During spring 2013, as seen in Figs 3 d and e, the stratification index starts to slowly increase in the convection area. For all simulations, as in winter, the negative $SI(1000m)$ bias is large to the south-east and the water column is too mixed while, in the convection area, the bias is fairly low, especially for the MOON simulation (Fig. 9 f). On average (Table 3) the bias is now positive and also greater with MOON than for ANDREAS and COARE. However, this is mainly due to the fact that the large positive biases present in the South of the domain in all the simulations are in ANDREAS and COARE partly compensated by large negative biases found in the centre of the convective area (Figs 9 d-f). [The slightly better performance of MOON is confirmed by the RMSEs which are lower for MOON than for COARE and ANDREAS, including for DEWEX Leg 2 at 1500 m and 2000 m depths.](#)

In summary, the evolution of stratification is not simply related to the buoyancy mass flux but results from the complex interaction between buoyancy mass fluxes and advective processes. In this study, the MOON simulation significantly reduces the negative bias of the stratification, in the convection area, during the convective and restratification periods, again indirectly suggesting that the air-sea fluxes are most realistic in the MOON simulation.

5.3 Impact on the air-sea exchanges on the deep water formation

In this section observations provided by the Lion buoy and the DEWEX cruises are compared with the results of the different simulations in terms of mixed patch, timing of convective process and volume of deep dense water formed.

5.3.1 Mixed patch

The spatial distribution of the convection is first examined. Figure 10 shows the position of the stratified and mixed CTD and Argo profiles and the extent of the mixed patch calculated by the model for the different simulations. The mixed patch is defined here as the grid points where the stratification index relative to 1000 m (or 2000 m) reaches zero at least once during the winter.

In other words, this corresponds to the area where deep convection reaches the 1000 m (or 2000 m) depth. The CTD and Argo profiles indicate that deep convection occurs between 41 °N and 43 °N, and between 3.5 °E and 6.5 °E. Obviously the CTD and Argo profiles only show a partial view of the mixed patch as they are punctual observations, whereas the simulated mixed patch corresponds to the area where deep convection occurred at least once during winter. The deep convection zone at 1000 m clearly appears in all the simulations but its eastern (and to a lesser extent southern) extension varies. The COARE and MOON simulations lead to the smallest and the largest mixed patch, respectively. However the observed profiles do not allow us to conclude on the most realistic extension at 1000 m. The deep convection zone at 2000 m is much smaller than at 1000 m but for this depth, observations clearly indicate that the extent of the mixed patch is better depicted by MOON than by COARE or ANDREAS.

5.3.2 Timing of convective process

The development of deep convection is examined using the observations produced by the LION mooring positioned in the centre of the deep water zone. Figure 11 shows the time evolution of the observed and simulated sea water density at this point. During the preconditioning period, the surface density anomaly is less than 29.0 kg m^{-3} and the bottom density anomaly is 29.11 kg m^{-3} . In the observations, the mixing reaches the first upper level of the LION mooring (150 m) in mid-January and the first complete mixing of the water column is achieved at the beginning of February. All the simulations produce deep convection but with different timing and intensity. In the COARE and ANDREAS simulations, complete mixing occurs too late (after the second event of Mistral and Tramontane of February) while for the MOON simulation, the process occurs too early (after the second event of Mistral and Tramontane of January). After the complete mixing, a sequence of Tramontane and Mistral events led to the densification of the water column (to a density between 29.12 and 29.13 kg m^{-3}). This densification in February is represented by all simulations. However, it is too weak in COARE and ANDREAS. Finally, at the beginning of March, a period of weak wind allows surface restratification. The last event of strong Tramontane and Mistral, around 15 March, destroys this restratification and again leads to the full mixing of the water column and to an increase of the seawater density up to 29.12 kg m^{-3} . This mixing event is not well represented by COARE and ANDREAS. Overall, despite the too-early full mixing, the MOON simulation gives the best representation of the deep convection at the LION mooring and correctly captures its three densification events.

5.3.3 Volume of deep water

The volume of the water mass created by deep convection is evaluated using the method proposed by Waldman et al. (2016b) for the same area and same period. Figure 12 shows the time evolution of the dense water formation rate computed in the simulations and estimated by Waldman et al. (2016b) with its error bar. During the autumn, the volume decreases slowly, whatever the simulation. This decrease is due to the dense water advection outside the study zone. Then, in winter, it increases rapidly, especially during the Tramontane and Mistral events, with the development of deep convection. After the convective events, the dense water volume decreases again, due to the restratification and export processes. The timing of deep water formation and the volume created are very different according to the simulations. The timing of deep water formation is the

same as that discussed for the LION mooring. The deep water volume created depends on the timing, spatial extent and intensity of deep convection processes. The COARE, ANDREAS and MOON simulations produce respectively $0.3 \times 10^{13} \text{ m}^3$, $1.5 \times 10^{13} \text{ m}^3$ and $3.4 \times 10^{13} \text{ m}^3$ between winter 2012 and spring 2013. For the three simulations, the amount of dense water formed is lower than the one calculated from the observations ($4.5(\pm 1.1) \times 10^{13} \text{ m}^3$). However, the most realistic volume is obtained with MOON. The differences between the simulations highlight the great sensitivity of deep water formation to turbulent flux parameterization.

6 Conclusions

This study focused on assessing the ability of a regional ocean atmosphere coupled system, based on the SYMPHONIE, SURFEX and MESONH models, to correctly represent ocean convection and deep water formation in the NWMS. Several realistic simulations, were carried out over a period of 8 months, from summer 2012 to spring 2013, and were used to investigate the sensitivity of the system to the parameterization of turbulent fluxes.

First, this study shows the ability of the air-sea coupled system to reproduce the evolution of the ocean and the atmosphere for several months by relying only on realistic initial and boundary conditions and without resorting to data assimilation or nudging. For all simulations, a good correlation is obtained between the observed and computed surface parameters at the Lion buoy. During the Tramontane and Mistral events, the turbulent heat fluxes differ significantly from one simulation to another, directly impacting the atmospheric and oceanic surface parameters. In a previous study devoted to the same case study, Rainaud et al. (2016) underlined the difficulty of reproducing air surface temperature and moisture during the Mistral and Tramontane events and advocated the use of an air-sea coupled model and a purposely adjusted turbulent flux parameterization. ~~Our study demonstrates that are strongly sensitive to the turbulent flux parameterizations, not only~~ In addition to air surface temperature and moisture ~~but also,~~ sea surface temperature is also strongly sensitive to the turbulent flux parameterizations. Our results also suggest that coupling plays a key role during the autumn storms when the rapid drops of SST reduce the turbulent heat fluxes. In winter, the impact of the coupling is likely to be weaker since the SST does not vary much anymore. Among the three parameterizations, we found that MOON (i.e. the parameterization yielding the strongest heat turbulent fluxes) significantly reduced the bias between the observed and computed surface parameters. Unfortunately, due to the lack of flux measurements at the buoy, it was not possible to validate the computed turbulent surface fluxes directly.

Then, the buoyancy mass flux was calculated and compared to the evolution of the stratification for each simulation. The stratification evolution is directly impacted by the buoyancy mass flux loss but there is no strict correlation between stratification and buoyancy mass flux. As already shown in Estournel et al. (2016a), this confirms the importance of the advective processes on the evolution of the stratification in the deep water area. Moreover, these advective processes also directly impact the surface buoyancy mass fluxes, particularly during preconditioning period, when the position and dynamics of the North Balearic Front clearly affect these fluxes. This interaction between the buoyancy mass fluxes and the advective processes is clearly an air-sea coupled process, which deserves to be analysed in greater depth. In terms of stratification, the ~~effect of MOON was also found to be positive, with again~~ use of MOON also led to a general reduction of the bias between observed and computed parameters.

Finally, the timing and the spatial extent of the convection process are very sensitive to the flux parameterization. This impacts the volume of deep water created, which varies from $0.3 \times 10^{13} \text{ m}^{-3}$ (with COARE) to $3.4 \times 10^{13} \text{ m}^{-3}$ (with MOON), i.e. by a factor of 11.3. For the same case study, Léger et al. (2016) studied the sensitivity to initial conditions and found a factor of 4.4 between two different sets of initial conditions. Our results suggest that the deep water formation process in the NWMS might be even more sensitive to atmospheric forcing. Here again, in spite of convection being triggered too early and the volume of dense water formed being slightly underestimated, MOON appears to give the most satisfactory results.

However, this conclusion regarding the good performance of the MOON flux parameterization needs to be further consolidated. ~~Future work will investigate the sensitivity of the present results to the resolution of the atmospheric model. In the present configuration (10 and 1 grid-spacing for the atmosphere and ocean, respectively), the local maxima and horizontal gradients of the surface atmospheric parameters are probably too smooth, which may affect the representation of the oceanic front, known to play a major role in~~

~~First the present results were obtained with a coupled system. They could probably be different with uncoupled simulations. In air-sea coupling (Small et al., 2008). A second point of interest, will be coupled simulations, the interactive evolution of ocean and atmosphere influences the turbulent heat fluxes, which themselves modify the atmospheric and oceanic surface fields involved in the flux calculation. In statically unstable Mistral and Tramontane conditions, if the sensible (or latent) heat flux increases, the vertical temperature (or humidity) gradient is reduced, which in turn limits the increase in the sensible (latent) heat flux. It is likely that these feedback loop effects tend to limit the discrepancies induced differences between by the different parameterizations. The results of partial and preliminary uncoupled simulations (not shown) suggest that these discrepancies could be larger than in the coupled simulations. It would be therefore of great interest to disentangle the effect of the flux formulation from the effect of the air-sea coupling and to check whether the MOON parameterization still improves the results in the absence of coupling uncoupled conditions. However, it is not straightforward to isolate the coupling effect in a clean and rigorous way. This requires a series of carefully-designed experiments in which the current coupled system is step-by-step downgraded into an uncoupled system till it exactly mimics the behavior of the atmospheric and ocean models in their stand-alone configuration. In our current system, this type of study is hampered by the fact that the surface fluxes are computed on the atmospheric grid, ie at a coarser resolution that the one used by the ocean model. A further step will be to couple the current system with a wave model~~

~~The differences in resolution between the atmospheric and ocean models (10 and 1 km, respectively), though partly justified by scale considerations, is also a debatable question. A further development will thus investigate the sensitivity to the ANDREAS parameterization, which takes the significant wave height into account to calculate the depth of the sea-spray layer (Andreas et al., 1995). In resolution of the atmospheric model. In the present configuration, the atmospheric model does not have the possibility of representing scales fully adjusted to those of the oceanic model. In particular, with a 10 km resolution, the local maxima and horizontal gradients of the surface parameters are probably too smooth, which may affect the air-sea interactions especially in the vicinity of the oceanic front (Small et al., 2008) and could also modify the response of the coupled system to the configuration used in this study, the wave height was only very roughly estimated as a function of wind (Andreas and Wang, 2007) different parameterizations.~~

- In addition, the role of the waves necessitates further investigation. In our study, the waves are not considered in COARE and MOON and only indirectly accounted for in ANDREAS. In ANDREAS, the depth of the spray layer is computed as a function of the significant wave height (Andreas et al., 1995) . The latter is rather roughly estimated from a simplified parameterization based on wind speed (Andreas and Wang, 2007) . Similar crude relationships are used in COARE3.0 for the wave height and wave period when the wave option is turned on. Another envisaged development will couple the current system with a wave model (Michaud et al. (2012) and the references cited therein) and revisit the results obtained with the ANDREAS and COARE3.0 parameterizations. Finally, the good overall performance of the newly developed coupled system encourages us to tackle future studies that necessitate an accurate description of the air/sea interface such as the study of the NBF dynamics during stormy events or the Mediterranean tropical-like cyclones (Miglietta et al., 2013) .
- 10 *Acknowledgements.* This work is a contribution to the MISTRALS/HyMeX programme through the ASICS-MED (ANR-12-BS06-0003) project funded by the French National Agency for Research (ANR). Data were obtained from the HyMeX programme, sponsored by grants from MISTRALS/HyMeX and Météo-France. The authors acknowledge the international ARGO programme, the LEFE/GMMC programme and the French NAOS project for supporting the deployment of profilers. Argo and CTD data were collected and made freely available by the CORIOLIS project (<http://www.coriolis.eu.org>) and programmes that contribute to it. We acknowledge the crews of R/V Suroit and Tethys II
- 15 and the scientists involved in the different cruises mentioned in this paper. Numerical simulations were performed using HPC resources from CALMIP (CALcul en Midi-Pyrénées, projects 1247, 09115 and 1325) and GENCI (Grand Equipement National de Calcul Intensif, project 010569)

References

- Andreas, E. L.: Approximation formulas for the microphysical properties of saline droplets, *Atmospheric Research*, 75, 323–345, doi:10.1016/j.atmosres.2005.02.001, <https://doi.org/10.1016/j.atmosres.2005.02.001>, 2005.
- Andreas, E. L. and Wang, S.: Predicting significant wave height off the northeast coast of the United States, *Ocean Engineering*, 34, 1328–1335, doi:10.1016/j.oceaneng.2006.08.004, wOS:000247148700023, 2007.
- Andreas, E. L., Edson, J. B., Monahan, E. C., Rouault, M. P., and Smith, S. D.: The spray contribution to net evaporation from the sea: A review of recent progress, *Boundary-Layer Meteorology*, 72, 3–52, doi:10.1007/BF00712389, <https://link.springer.com/article/10.1007/BF00712389>, 1995.
- Andreas, E. L., Mahrt, L., and Vickers, D.: An improved bulk air–sea surface flux algorithm, including spray-mediated transfer, *Quarterly Journal of the Royal Meteorological Society*, 141, 642–654, doi:10.1002/qj.2424, <http://onlinelibrary.wiley.com/doi/10.1002/qj.2424/abstract>, 2015.
- Bechtold, P., Bazile, E., Guichard, F., Mascart, P., and Richard, E.: A mass-flux convection scheme for regional and global models, *Quarterly Journal of the Royal Meteorological Society*, 127, 869–886, <http://onlinelibrary.wiley.com/doi/10.1002/qj.49712757309/full>, 2001.
- Bentsen, M., Evensen, G., Drange, H., and Jenkins, A. D.: Coordinate Transformation on a Sphere Using Conformal Mapping, *Monthly Weather Review*, 127, 2733–2740, doi:10.1175/1520-0493(1999)127<2733:CTOASU>2.0.CO;2, [http://journals.ametsoc.org/doi/abs/10.1175/1520-0493\(1999\)127%3C2733:CTOASU%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0493(1999)127%3C2733:CTOASU%3E2.0.CO%3B2), 1999.
- Buongiorno Nardelli, B., Tronconi, C., Pisano, A., and Santoleri, R.: High and Ultra-High resolution processing of satellite Sea Surface Temperature data over Southern European Seas in the framework of MyOcean project, *Remote Sensing of Environment*, 129, 1–16, doi:10.1016/j.rse.2012.10.012, <http://www.sciencedirect.com/science/article/pii/S003442571200394X>, 2013.
- Caniaux, G., Redelsperger, J.-L., and Lafore, J.-P.: A Numerical Study of the Stratiform Region of a Fast-Moving Squall Line. Part I: General Description and Water and Heat Budgets, *Journal of the Atmospheric Sciences*, 51, 2046–2074, doi:10.1175/1520-0469(1994)051<2046:ANSOTS>2.0.CO;2, [http://dx.doi.org/10.1175/1520-0469\(1994\)051<2046:ANSOTS>2.0.CO;2](http://dx.doi.org/10.1175/1520-0469(1994)051<2046:ANSOTS>2.0.CO;2), 1994.
- Charnock, H.: Wind stress on a water surface, *Quarterly Journal of the Royal Meteorological Society*, 81, 639–640, doi:10.1002/qj.49708135027, <https://doi.org/10.1002/qj.49708135027>, 1955.
- Cuxart, J., Bougeault, P., and Redelsperger, J.-L.: A turbulence scheme allowing for mesoscale and large-eddy simulations, *Quarterly Journal of the Royal Meteorological Society*, 126, 1–30, doi:10.1002/qj.49712656202, <http://dx.doi.org/10.1002/qj.49712656202>, 2000.
- Deardorff, J. W., Willis, G. E., and Lilly, D. K.: Laboratory investigation of non-steady penetrative convection, *Journal of Fluid Mechanics*, 35, 7–31, http://journals.cambridge.org/article_S0022112069000942, 1969.
- Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) system, *Remote Sensing of Environment*, 116, 140–158, doi:10.1016/j.rse.2010.10.017, <http://www.sciencedirect.com/science/article/pii/S0034425711002197>, 2012.
- Drobinski, P., Ducrocq, V., Alpert, P., Anagnostou, E., Béranger, K., Borga, M., Braud, I., Chanzy, A., Davolio, S., Delrieu, G., Estournel, C., Boubrahmi, N. F., Font, J., Grubišić, V., Gualdi, S., Homar, V., Ivančan-Picek, B., Kottmeier, C., Kotroni, V., Lagouvardos, K., Lionello, P., Llasat, M. C., Ludwig, W., Lutoff, C., Mariotti, A., Richard, E., Romero, R., Rotunno, R., Roussot, O., Ruin, I., Somot, S., Taupier-Letage, I., Tintore, J., Uijlenhoet, R., and Wernli, H.: HyMeX: A 10-Year Multidisciplinary Program on the Mediterranean Water Cycle, *Bulletin of the American Meteorological Society*, 95, 1063–1082, doi:10.1175/BAMS-D-12-00242.1, <http://journals.ametsoc.org/doi/abs/10.1175/BAMS-D-12-00242.1>, 2013.

- Ducrocq, V., Braud, I., Davolio, S., Ferretti, R., Flamant, C., Jansa, A., Kalthoff, N., Richard, E., Taupier-Letage, I., Aral, P.-A., Belamari, S., Berne, A., Borga, M., Boudevillain, B., Bock, O., Boichard, J.-L., Bouin, M.-N., Bousquet, O., Bouvier, C., Chiggiato, J., Cimini, D., Corsmeier, U., Coppola, L., Cocquerez, P., Defer, E., Delanoë, J., Di Girolamo, P., Doerenbecher, A., Drobinski, P., Dufournet, Y., Fourrié, N., Gourley, J. J., Labatut, L., Lambert, D., Le Coz, J., Marzano, F. S., Molinié, G., Montani, A., Nord, G., Nuret, M., Ramage, K., Rison, W., Roussot, O., Said, F., Schwarzenboeck, A., Testor, P., Van Baelen, J., Vincendon, B., Aran, M., and Tamayo, J.: HyMeX-SOP1: The Field Campaign Dedicated to Heavy Precipitation and Flash Flooding in the Northwestern Mediterranean, *Bulletin of the American Meteorological Society*, 95, 1083–1100, doi:10.1175/BAMS-D-12-00244.1, <http://journals.ametsoc.org/doi/abs/10.1175/BAMS-D-12-00244.1>, 2013.
- Estournel, C., Auclair, F., Lux, M., Nguyen, C., and Marsaleix, P.: "Scale oriented" embedded modeling of the North-Western Mediterranean in the frame of MFSTEP, *Ocean Sci.*, 5, 73–90, doi:10.5194/os-5-73-2009, <http://www.ocean-sci.net/5/73/2009/>, 2009.
- Estournel, C., Testor, P., Damien, P., D'Ortenzio, F., Marsaleix, P., Conan, P., Kessouri, F., Durrieu de Madron, X., Coppola, L., Lellouche, J.-M., Belamari, S., Mortier, L., Ulses, C., Bouin, M.-N., and Prieur, L.: High resolution modeling of dense water formation in the north-western Mediterranean during winter 2012–2013: Processes and budget, *Journal of Geophysical Research: Oceans*, 121, 5367–5392, doi:10.1002/2016JC011935, <http://onlinelibrary.wiley.com/doi/10.1002/2016JC011935/abstract>, 2016a.
- Estournel, C., Testor, P., Taupier-Letage, I., Bouin, M.-N., Coppola, L., Durand, P., Conan, P., Bosse, A., Brilouet, P.-E., Beguery, L., and others: HyMeX-SOP2 The Field Campaign Dedicated to Dense Water Formation in the Northwestern Mediterranean, *Oceanography*, 29, 196–206, <https://hal.archives-ouvertes.fr/hal-01443236/>, 2016b.
- Fairall, C. W., Bradley, E. F., Rogers, D. P., Edson, J. B., and Young, G. S.: Bulk parameterization of air-sea fluxes for tropical ocean-global atmosphere coupled-ocean atmosphere response experiment, *Journal of Geophysical Research: Oceans*, 101, 3747–3764, <http://onlinelibrary.wiley.com/doi/10.1029/95JC03205/full>, 1996.
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B.: Bulk Parameterization of Air–Sea Fluxes: Updates and Verification for the COARE Algorithm, *Journal of Climate*, 16, 571–591, doi:10.1175/1520-0442(2003)016<0571:BPOASF>2.0.CO;2, [http://journals.ametsoc.org/doi/abs/10.1175/1520-0442\(2003\)016%3C0571:BPOASF%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0442(2003)016%3C0571:BPOASF%3E2.0.CO%3B2), 2003.
- Flamant, C.: Alpine lee cyclogenesis influence on air-sea heat exchanges and marine atmospheric boundary layer thermodynamics over the western Mediterranean during a Tramontane/Mistral event, *Journal of Geophysical Research: Oceans*, 108, 8057, doi:10.1029/2001JC001040, <http://onlinelibrary.wiley.com/doi/10.1029/2001JC001040/abstract>, 2003.
- Foken, T.: 50 Years of the Monin–Obukhov Similarity Theory, *Boundary-Layer Meteorology*, 119, 431–447, doi:10.1007/s10546-006-9048-6, <http://link.springer.com/article/10.1007/s10546-006-9048-6>, 2006.
- Fouquart, Y. and Bonnel, B.: Computations of solar heating of the earth's atmosphere- A new parameterization, *Beitraege zur Physik der Atmosphaere*, 53, 35–62, 1980.
- Gaspar, P., Grégoris, Y., and Lefevre, J.-M.: A simple eddy kinetic energy model for simulations of the oceanic vertical mixing: Tests at station Papa and long-term upper ocean study site, *Journal of Geophysical Research: Oceans*, 95, 16 179–16 193, doi:10.1029/JC095iC09p16179, <http://dx.doi.org/10.1029/JC095iC09p16179>, 1990.
- Hauser, D., Branger, H., Bouffies-Cloché, S., Despiau, S., Drennan, W. M., Dupuis, H., Durand, P., Durrieu de Madron, X., Estournel, C., Eymard, L., Flamant, C., Graber, H. C., Guérin, C., Kahma, K., Lachaud, G., Lefèvre, J.-M., Pelon, J., Pettersson, H., Pigué, B., Queffeuilou, P., Tailliez, D., Tournadre, J., and Weill, A.: The FETCH experiment: An overview, *Journal of Geophysical Research: Oceans*, 108, 8053, doi:10.1029/2001JC001202, <http://onlinelibrary.wiley.com/doi/10.1029/2001JC001202/abstract>, 2003.

- Herrmann, M., Somot, S., Sevault, F., Estournel, C., and Déqué, M.: Modeling the deep convection in the northwestern Mediterranean Sea using an eddy-permitting and an eddy-resolving model: Case study of winter 1986–1987, *Journal of Geophysical Research: Oceans*, 113, C04 011, doi:10.1029/2006JC003991, <http://onlinelibrary.wiley.com/doi/10.1029/2006JC003991/abstract>, 2008.
- Herrmann, M., Sevault, F., Beuvier, J., and Somot, S.: What induced the exceptional 2005 convection event in the northwestern Mediterranean basin? Answers from a modeling study, *Journal of Geophysical Research: Oceans*, 115, C12 051, doi:10.1029/2010JC006162, <http://onlinelibrary.wiley.com/doi/10.1029/2010JC006162/abstract>, 2010.
- Herrmann, M. J. and Somot, S.: Relevance of ERA40 dynamical downscaling for modeling deep convection in the Mediterranean Sea, *Geophysical Research Letters*, 35, L04 607, doi:10.1029/2007GL032442, <http://onlinelibrary.wiley.com/doi/10.1029/2007GL032442/abstract>, 2008.
- 10 Houpert, L., Durrieu de Madron, X., Testor, P., Bosse, A., D’Ortenzio, F., Bouin, M., Dausse, D., Le Goff, H., Kunesch, S., Labaste, M., Coppola, L., Mortier, L., and Raimbault, P.: Observations of open-ocean deep convection in the northwestern Mediterranean Sea: Seasonal and interannual variability of mixing and deep water masses for the 2007–2013 period, *Journal of Geophysical Research: Oceans*, pp. n/a–n/a, doi:10.1002/2016JC011857, <http://onlinelibrary.wiley.com/doi/10.1002/2016JC011857/abstract>, 2016.
- Kurihara, Y., Tuleya, R. E., and Bender, M. A.: The GFDL hurricane prediction system and its performance in the 1995 hurricane season, *Monthly weather review*, 126, 1306–1322, [http://journals.ametsoc.org/doi/abs/10.1175/1520-0493\(1998\)126%3C1306:TGHPSA%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0493(1998)126%3C1306:TGHPSA%3E2.0.CO%3B2), 1998.
- 15 Lafore, J. P., Stein, J., Asencio, N., Bougeault, P., Ducrocq, V., Duron, J., Fischer, C., Héreil, P., Mascart, P., Masson, V., Pinty, J. P., Redelsperger, J. L., Richard, E., and Arellano, J. V.-G. d.: The Meso-NH Atmospheric Simulation System. Part I: adiabatic formulation and control simulations, *Annales Geophysicae*, 16, 90–109, doi:10.1007/s00585-997-0090-6, <http://link.springer.com/article/10.1007/s00585-997-0090-6>, 1997.
- 20 Lebeauin Brossier, C. and Drobinski, P.: Numerical high-resolution air-sea coupling over the Gulf of Lions during two tramontane/mistral events, *Journal of Geophysical Research: Atmospheres*, 114, n/a–n/a, doi:10.1029/2008JD011601, <http://dx.doi.org/10.1029/2008JD011601>, d10110, 2009.
- Lebeauin-Brossier, C., Arsouze, T., Béranger, K., Bouin, M.-N., Bresson, E., Ducrocq, V., Giordani, H., Nuret, M., Rainaud, R., and Taupier-Letage, I.: Ocean Mixed Layer responses to intense meteorological events during HyMeX-SOP1 from a high-resolution ocean simulation, *Ocean Modelling*, 84, 84–103, doi:10.1016/j.ocemod.2014.09.009, <http://www.sciencedirect.com/science/article/pii/S1463500314001401>, 2014.
- 25 Lellouche, J.-M., Le Galloudec, O., Drévillon, M., Régnier, C., Greiner, E., Garric, G., Ferry, N., Desportes, C., Testut, C.-E., Bricaud, C., Bourdallé-Badie, R., Tranchant, B., Benkiran, M., Drillet, Y., Daudin, A., and De Nicola, C.: Evaluation of global monitoring and forecasting systems at Mercator Océan, *Ocean Sci.*, 9, 57–81, doi:10.5194/os-9-57-2013, <http://www.ocean-sci.net/9/57/2013/>, 2013.
- 30 Léger, F., Lebeauin Brossier, C., Giordani, H., Arsouze, T., Beuvier, J., Bouin, M.-N., Bresson, E., Ducrocq, V., Fourrié, N., and Nuret, M.: Dense water formation in the north-western Mediterranean area during HyMeX-SOP2 in 1/36° ocean simulations: Sensitivity to initial conditions, *Journal of Geophysical Research: Oceans*, 121, 5549–5569, doi:10.1002/2015JC011542, <http://onlinelibrary.wiley.com/doi/10.1002/2015JC011542/abstract>, 2016.
- 35 Maraldi, C., Chanut, J., Levier, B., Ayoub, N., De Mey, P., Reffray, G., Lyard, F., Cailleau, S., Drévillon, M., Fanjul, E. A., Sotillo, M. G., Marsaleix, P., and Team, M.: NEMO on the shelf: assessment of the Iberia-Biscay-Ireland configuration., *Ocean Science*, 9, 745–771, doi:10.5194/os-9-745-2013, <https://hal.archives-ouvertes.fr/hal-00766544>, 2013.

- Marsaleix, P., Auclair, F., Floor, J. W., Herrmann, M. J., Estournel, C., Pairaud, I., and Ulses, C.: Energy conservation issues in sigma-coordinate free-surface ocean models, *Ocean Modelling*, 20, 61–89, doi:10.1016/j.ocemod.2007.07.005, <http://www.sciencedirect.com/science/article/pii/S1463500307000984>, 2008.
- Marsaleix, P., Auclair, F., and Estournel, C.: Low-order pressure gradient schemes in sigma coordinate models: The seamount test revisited, *Ocean Modelling*, 30, 169–177, doi:10.1016/j.ocemod.2009.06.011, <http://www.sciencedirect.com/science/article/pii/S1463500309001358>, 2009.
- Marsaleix, P., Auclair, F., Duhaut, T., Estournel, C., Nguyen, C., and Ulses, C.: Alternatives to the Robert–Asselin filter, *Ocean Modelling*, 41, 53–66, doi:10.1016/j.ocemod.2011.11.002, <http://www.sciencedirect.com/science/article/pii/S1463500311001788>, 2012.
- Marshall, J. and Schott, F.: Open-ocean convection: Observations, theory, and models, *Reviews of Geophysics*, 37, 1–64, doi:10.1029/98RG02739, <http://onlinelibrary.wiley.com/doi/10.1029/98RG02739/abstract>, 1999.
- Marsland, S. J., Haak, H., Jungclaus, J. H., Latif, M., and Röske, F.: The Max-Planck-Institute global ocean/sea ice model with orthogonal curvilinear coordinates, *Ocean modelling*, 5, 91–127, 2003.
- Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouysse, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essaouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of Earth surface variables and fluxes, *Geoscientific Model Development Discussions*, 5, 3771–3851, doi:10.5194/gmdd-5-3771-2012, <http://adsabs.harvard.edu/abs/2012GMDD....5.3771M>, 2012.
- Michaud, H., Marsaleix, P., Leredde, Y., Estournel, C., Bourrin, F., Lyard, F., Mayet, C., and Ardhuin, F.: Three-dimensional modelling of wave-induced current from the surf zone to the inner shelf, *Ocean Science*, 8, 657, 2012.
- Miglietta, M. M., Laviola, S., Malvaldi, A., Conte, D., Levizzani, V., and Price, C.: Analysis of tropical-like cyclones over the Mediterranean Sea through a combined modeling and satellite approach, *Geophysical Research Letters*, 40, 2400–2405, doi:10.1002/grl.50432, <http://onlinelibrary.wiley.com/doi/10.1002/grl.50432/abstract>, 2013.
- Millot, C.: Circulation in the western Mediterranean Sea, *Journal of Marine Systems*, 20, 423–442, <http://www.sciencedirect.com/science/article/pii/S0924796398000785>, 1999.
- Millot, C. and Taupier-Letage, I.: Circulation in the Mediterranean sea, in: *The Mediterranean Sea*, pp. 29–66, Springer, <http://link.springer.com/chapter/10.1007/b107143>, 2005.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *Journal of Geophysical Research: Atmospheres*, 102, 16 663–16 682, doi:10.1029/97JD00237, <http://onlinelibrary.wiley.com/doi/10.1029/97JD00237/abstract>, 1997.
- Moon, I.-J., Ginis, I., Hara, T., and Thomas, B.: A Physics-Based Parameterization of Air–Sea Momentum Flux at High Wind Speeds and Its Impact on Hurricane Intensity Predictions, *Monthly Weather Review*, 135, 2869–2878, doi:10.1175/MWR3432.1, <http://journals.ametsoc.org/doi/abs/10.1175/MWR3432.1>, 2007.
- Pergaud, J., Masson, V., Malardel, S., and Couvreur, F.: A parameterization of dry thermals and shallow cumuli for mesoscale numerical weather prediction, *Boundary-layer meteorology*, 132, 83–106, <http://www.springerlink.com/index/K5P218150R2J424P.pdf>, 2009.
- Pinty, J. and Jabouille, P.: A mixed-phase cloud parameterization for use in mesoscale non-hydrostatic model: simulations of a squall line and of orographic precipitations, in: *Conf. on Cloud Physics*, pp. 217–220, Amer. Meteor. Soc Everett, WA, 1998.

- Rainaud, R., Brossier, C. L., Ducrocq, V., Giordani, H., Nuret, M., Fourrié, N., Bouin, M.-N., Taupier-Letage, I., and Legain, D.: Characterization of air-sea exchanges over the Western Mediterranean Sea during HyMeX SOP1 using the AROME-WMED model, *Quarterly Journal of the Royal Meteorological Society*, 142, 173–187, <https://hal.archives-ouvertes.fr/hal-01443601/>, 2016.
- Renault, L., Chiggiato, J., Warner, J. C., Gomez, M., Vizoso, G., and Tintoré, J.: Coupled atmosphere-ocean-wave simulations of a storm event over the Gulf of Lion and Balearic Sea, *Journal of Geophysical Research: Oceans*, 117, n/a–n/a, doi:10.1029/2012JC007924, <http://dx.doi.org/10.1029/2012JC007924>, c09019, 2012.
- Schott, F. and Leaman, K. D.: Observations with Moored Acoustic Doppler Current Profilers in the Convection Regime in the Golfe du Lion, *Journal of Physical Oceanography*, 21, 558–574, doi:10.1175/1520-0485(1991)021<0558:OWMADC>2.0.CO;2, [http://journals.ametsoc.org/doi/abs/10.1175/1520-0485\(1991\)021%3C0558:OWMADC%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0485(1991)021%3C0558:OWMADC%3E2.0.CO%3B2), 1991.
- Schott, F., Visbeck, M., Send, U., Fischer, J., Stramma, L., and Desaubies, Y.: Observations of deep convection in the Gulf of Lions, Northern Mediterranean, during the winter of 1991/92, *Journal of Physical Oceanography*, 26, 505–524, <http://oceanrep.geomar.de/3810/>, 1996.
- Sempéré, R., Madron, X. D. d., and Guieu, C.: The MERMeX Program for the Mediterranean Sea, pp. 389–392, doi:10.1007/978-90-481-8630-3_70, http://link.springer.com/chapter/10.1007/978-90-481-8630-3_70, 2010.
- Small, R., deSzoeko, S., Xie, S., O'Neill, L., Seo, H., Song, Q., Cornillon, P., Spall, M., and Minobe, S.: Air–sea interaction over ocean fronts and eddies, *Dynamics of Atmospheres and Oceans*, 45, 274 – 319, doi:<http://dx.doi.org/10.1016/j.dynatmoce.2008.01.001>, <http://www.sciencedirect.com/science/article/pii/S0377026508000341>, {OCEANIC} {FRONTS}, 2008.
- Small, R. J., Carniel, S., Campbell, T., Teixeira, J., and Allard, R.: The response of the Ligurian and Tyrrhenian Seas to a summer Mistral event: A coupled atmosphere–ocean approach, *Ocean Modelling*, 48, 30–44, doi:10.1016/j.ocemod.2012.02.003, <http://www.sciencedirect.com/science/article/pii/S1463500312000339>, 2012.
- Testor, P. and Gascard, J. C.: Post-convection spreading phase in the Northwestern Mediterranean Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 53, 869–893, doi:10.1016/j.dsr.2006.02.004, <http://www.sciencedirect.com/science/article/pii/S0967063706000550>, 2006.
- Ulses, C., Estournel, C., Bonnin, J., Durrieu de Madron, X., and Marsaleix, P.: Impact of storms and dense water cascading on shelf-slope exchanges in the Gulf of Lion (NW Mediterranean), *Journal of Geophysical Research: Oceans*, 113, n/a–n/a, doi:10.1029/2006JC003795, <http://dx.doi.org/10.1029/2006JC003795>, c02010, 2008.
- Valcke, S., Craig, T., and Coquart, L.: OASIS3-MCT User Guide, OASIS3-MCT 3.0, Technical Report TR/CMGC/15/38, Cerfacs, France, 1875, http://www.cerfacs.fr/oa4web/oasis3-mct_3.0/oasis3mct_UserGuide.pdf, 2015.
- Voldoire, A., Decharme, B., Pianezze, J., Lebeaupin Brossier, C., Sevault, F., Seyfried, L., Garnier, V., Bielli, S., Valcke, S., Alias, A., Accensi, M., Arduin, F., Bouin, M.-N., Ducrocq, V., Faroux, S., Giordani, H., Léger, F., Marsaleix, P., Raynaud, R., Redelsperger, J.-L., Richard, E., and Riette, S.: The seamless and multi-model coupling between atmosphere, land, hydrology, ocean, waves and sea-ice models based on SURFEX surface model using OASIS3-MCT, *Geosci. Model Dev. Discuss.*, 2017, 1–39, doi:10.5194/gmd-2017-91, <http://www.geosci-model-dev-discuss.net/gmd-2017-91/>, 2017.
- Waldman, R., Somot, S., Herrmann, M., Bosse, A., Caniaux, G., Estournel, C., Houpert, L., Prieur, L., Sevault, F., and Testor, P.: Modeling the intense 2012–2013 dense water formation event in the northwestern Mediterranean Sea: Evaluation with an ensemble simulation approach, *Journal of Geophysical Research: Oceans*, <http://onlinelibrary.wiley.com/doi/10.1002/2016JC012437/full>, 2016a.
- Waldman, R., Somot, S., Herrmann, M., Testor, P., Estournel, C., Sevault, F., Prieur, L., Mortier, L., Coppola, L., Taillandier, V., Conan, P., and Dausse, D.: Estimating dense water volume and its evolution for the year 2012–2013 in the Northwestern Mediterranean Sea: An observing

system simulation experiment approach, *Journal of Geophysical Research: Oceans*, 121, 6696–6716, doi:10.1002/2016JC011694, <http://onlinelibrary.wiley.com/doi/10.1002/2016JC011694/abstract>, 2016b.

Webster, P. J. and Lukas, R.: TOGA COARE: The coupled ocean—atmosphere response experiment, *Bulletin of the American Meteorological Society*, 73, 1377–1416, 1992.




Figure 1. Full frame: spatial extension of the atmospheric model. Topography (meters, horizontal colour bar, positive above sea level) and bathymetry (metres, vertical colour bar, negative below sea level) of the ocean model. The black dotted rectangle indicates the area studied. The red and magenta dotted line represent oceanic surface circulation with the Northern Current (NC) and North Balearic Front (NBF), respectively. The black dotted circle indicates the Dense Water Formation (DWF) zone




Figure 2. Turbulent fluxes of momentum (a), sensible heat (b) and latent heat (c) as functions of the surface wind speed for each bulk parameterization (COARE 3.0 in blue, COARE-2.6 ANDREAS no-spray in cyan, ANDREAS in green and MOON in red). The computation was performed with SURFEX offline, with the air surface temperature fixed at 10 °C, the sea surface temperature at 15 °C , the relative humidity at 70 %, and the surface atmospheric pressure at 1013 hPa.




Figure 3. Stratification Index at 1000 m derived from CTD (squares) and ARGO (dots). Observations during (a) summer 2012 (MOOSE), (b) autumn 2012 (DOWEX), (c) winter 2013 (DEWEX Leg 1) and (d) spring 2013 (DEWEX leg 2). (e) Time series of Stratification Index at 1000 m. The red contour corresponds to the convective area defined in section 3.




Figure 4. Time series of (a) the 10 m wind speed (b) the 2 m air temperature (c) the 2 m air humidity and (d) the sea surface temperature measured at the Lion meteorological buoy (in black) and computed by each simulation (COARE in blue, ANDREAS in green and MOON in red). The grey bars correspond to the strong wind periods (hourly wind speed > 15 m s⁻¹)




Figure 5. Scatter plots of simulations against observations at Lion buoy for (a) the 10 m wind speed, (b) 2 m air temperature and (c) humidity, and (d) sea surface temperature.




Figure 6. Time series of computed (a) wind stress, (b) sensible heat flux and (c) latent heat flux at the location of Lion buoy for each simulation (COARE in blue, ANDREAS in green and MOON in red). The grey bars correspond to the strong wind periods (hourly wind speed $> 15 \text{ m s}^{-1}$)




Figure 7. Time series of (a) instantaneous buoyancy mass fluxes, (b) cumulated buoyancy mass fluxes (Eq. 5), and (c) stratification index (Eq. 4) relative to 2000 m at Lion buoy computed for each simulation (COARE in blue, ANDREAS in green and MOON in red). The grey bars correspond to the strong wind periods (hourly wind speed > 15 m s⁻¹)




Figure 8. Buoyancy mass fluxes (in kg m^{-2}): (a-c) during preconditioning and convective period (16 August 2012 to 21 March), (d-f) only during preconditioning period (16 August 2012 to 15 January 2013), (g-i) only during convective period (15 January 2013 to 21 March 2013) computed for each simulation: (left) COARE, (middle) ANDREAS, and (right) MOON. Red line: Convective area defined in section 3. Black star: location of Lion meteorological buoy.




Figure 9. Bias of the stratification index relative to 1000 m calculated for each simulation (left) COARE, (middle) ANDREAS, (right) MOON, during (a-c) DEWEX leg 1 cruise and (d-f) DEWEX leg 2 cruise. Red line: Convective area defined in section 3.




Figure 10. Coloured areas: extend of mixed patch zone computed for each simulation: (left) COARE, (middle) ANDREAS, (right) MOON. The mixed patch is defined as the area where the stratification index, (a-c) at 1000 m and (d-f) at 2000 m, reaches 0 during simulation. Black squares correspond to the position of the CTD and ARGO mixed profiles: (a-c) $SI(1000) = 0$. and (d-f) $SI(2000) = 0$.. Cyan points show the position of the CTD and ARGO stratified profiles: (a-c) $SI(1000) > 0$. and (d-f) $SI(2000) > 0$. Red line: Convective area defined in section 3.




Figure 11. Time series of 0–2000 m sea water density (a) observed at LION mooring and computed for each simulation (b) COARE, (c) ANDREAS, (d) MOON at LION mooring location. The black line correspond to mixed layer depth (computed with a density criterion of 0.01 kg m⁻³)




Figure 12. Time series of dense water volume (m^3) formed from 8/1/12 for sea water density up to 29.11 kg m^{-3} computed for each simulation (COARE in blue, ANDREAS in green and MOON in red) in the convection area (as defined in section 3). The black dot and error bars correspond to the dense water formation volume estimated from the observations by Waldman et al. (2016b) for the same density thresholds. The grey bars correspond to the strong wind periods (hourly wind speed $> 15 \text{ m s}^{-1}$)

Table 1. Main characteristics of the atmospheric and oceanic model configurations

Physical processes	Schemes	References
Atmospheric model		
Horizontal grid	Arakava-C grid, 10 km resolution	
Vertical levels	Sigma-z <u>Sigma-z</u> coordinates, 52 vertical levels from 15m to 15000 m	
Turbulence	TKE, 1D vertical	(Cuxart et al., 2000)
Convection	mass-flux	(Bechtold et al., 2001), (Pergaud et al., 2009)
Radiation	longwave : RRTM shortwave : ECMWF	(Mlawer et al., 1997), (Fouquart and Bonnel, 1980)
Microphysics	ICE3	(Caniaux et al., 1994), (Pinty and Jabouille, 1998)
Initial and boundary condition	ECMWF analyses	
Oceanic model		
Horizontal grid	Curvilinear Arakava-C grid, 1km resolution,	(Bentsen et al., 1999)
Vertical levels	40 generalized sigma vertical levels	(Ulses et al., 2008)
Sea surface conditions	Craig & Banner TKE boundary conditions	(Estournel et al., 2009)
Mixing	Eddy Kinetic Energy	(Gaspar et al., 1990)
Convection	penetrative convection	(Deardorff et al., 1969), (Estournel et al., 2016a)
River input	Lateral condition (15 river inputs)	(Estournel et al., 2009)
Initial and boundary condition	MERCATOR-OCEAN MOOSE 2012-2013	(Lellouche et al., 2013), (Estournel et al., 2016a)

Table 2. Statistics of surface atmospheric and oceanic parameters at Lion meteorological buoy.

EXP	COARE			ANDREAS			MOON		
STAT	BIAS	RMS	R	BIAS	RMS	R	BIAS	RMS	R
U10M	-0.08	2.19	0.90	-0.16	2.21	0.90	0.11	2.17	0.90
T2M	-0.22	01.07 <u>1.07</u>	0.98	-0.18	01.02 <u>1.02</u>	0.98	-0.08	0.96	0.98
HU2M	3.24	07.02 <u>7.02</u>	0.85	3.45	7.23	0.85	1.15	6.17	0.85
SST	-0.50	1.11	0.98	-0.15	0.72	0.99	0.05	0.61	0.99

Table 3. Number of CTD-ARGO observed profiles and observations averaged, simulations bias and rmse averaged, for Stratification Index (kg m^{-2}) at ~~1000m~~1000 m, ~~1500m~~1500 m and ~~2000m~~2000 m for DEWEX oceanographic cruises.

Number of profiles CTD-ARGO		COARE	Observations SI averaged	ANDREAS	<u>COARE</u> bias, rmse	MOON	<u>ANDREAS</u> bias, rmse	<u>MOON</u> bias, rmse
DEWEX leg 1								
1000 m	62-230		<u>13.8</u>		-4.9, <u>19.5</u>		-3.8, <u>18.4</u>	1.8, <u>16.6</u>
1500 m	56-202		<u>12.6</u>		-7.4, <u>19.0</u>		-5.7, <u>17.5</u>	-0.4, <u>14.9</u>
2000 m	48-29		<u>22.0</u>		-9.2, <u>24.0</u>		-6.0, <u>22.5</u>	0.9, <u>22.8</u>
DEWEX leg 2								
1000 m	78-103		<u>27.0</u>		4.2, <u>19.4</u>		6.6, <u>19.0</u>	9.1, <u>20.0</u>
1500 m	72-88		<u>31.2</u>		2.9, <u>19.5</u>		6.0, <u>19.5</u>	7.4, <u>19.2</u>
2000 m	60-7		<u>42.1</u>		3.6, <u>26.7</u>		4.5, <u>26.5</u>	7.4, <u>23.5</u>