

Orbit related sea level errors for TOPEX altimetry at seasonal to decadal time scales

Saskia Esselborn¹, Sergei Rudenko^{1,2}, Tilo Schöne¹

¹GFZ German Research Centre for Geosciences, Department 1: Geodesy, Potsdam, 14473, Germany

5 ²Deutsches Geodätisches Forschungsinstitut (DGFI-TUM), Technische Universität München, Munich, 80333, Germany (since August 2016, until that – at GFZ)

Correspondence to: Saskia Esselborn (Saskia.Esselborn@gfz-potsdam.de)

Abstract. Interannual to decadal sea level trends are indicators of climate variability and change. A major source of global and regional sea level data is satellite radar altimetry, which relies on precise knowledge of the satellite's orbit. Here, we assess the error budget of the radial orbit component for the TOPEX/Poseidon mission for the period 1993 to 2004 from a set of different orbit solutions. Upper bound errors for seasonal, interannual (5 years), and decadal periods are estimated on global and regional scales based on radial orbit differences from three state-of-the-art orbit solutions provided by different research teams (GFZ, GSFC, and GRGS). The global mean sea level error related to the orbit is of the order of 7 mm (more than 10 % of the sea level variability) with negligible contributions on the annual and decadal time scale. In contrast, the orbit related error of the interannual trend is 0.1 mm/year (18 % of the corresponding sea level variability) and might hamper the estimation of an acceleration of the global mean sea level rise. For regional scales, the gridded orbit related error is up to 11 mm and for about half the ocean the orbit error accounts for at least 10 % of the observed sea level variability. The seasonal orbit error amounts to 10 % of the observed seasonal sea level signal in the Southern Ocean. At interannual and decadal time scales, the orbit related trend uncertainties reach regionally more than 1 mm/year. The interannual trend errors account for 10 % of the observed sea level signal in the Tropical Atlantic and the south-eastern Pacific. For decadal scales, the orbit related trend errors are prominent in a couple of regions including: South Atlantic, western North Atlantic, central Pacific, South Australian Basin, and Mediterranean Sea. Based on a set of test orbits calculated at GFZ, the sources of the observed orbit related errors are further investigated. Main contributors on all time scales are uncertainties in Earth's time variable gravity field models and on annual to interannual time scales discrepancies of the tracking station sub-networks, i.e., SLR and DORIS.

30 **1 Introduction**

Sea level is an important indicator of climate variability and change. Based on tide gauge data using different techniques, the global mean sea level rise for the last century is estimated to be 1.2-1.9 mm/year (Douglas, 1997; Church and White, 2011; Jevrejeva et al., 2008, 2014; Hay et al., 2015). Based on satellite altimetry data since 1993, the current rate of global mean sea level has been estimated to be more than 3 mm/year (Cazenave et al., 2014; Ablain et al., 2016, Quartly et al., 2017). The main sources of the current rise are thermal expansion of the sea water and melting of glaciers and ice sheets. At interannual time scales, changes of terrestrial water storage imprint additionally on the global mean sea level (Llovel et al., 2011). Recent work (Watson et al., 2015; Fasullo et al., 2016) has focussed on the detectability of accelerations in global mean sea level trends during the last decades. Regionally, sea level rates during the last 24 years show higher variability, they range from -1 mm/year to more than 10 mm/year. They are mainly linked to regional changes in the oceans density field, which might be induced by internal ocean variability, atmosphere-ocean interaction, or influx of freshwater. Satellite altimeters are a unique source of global and regional sea level data and are available continuously since the beginning of the 1990s. Precise orbits of altimetry satellites are a precondition for global and regional mean sea level investigations (Rudenko et al., 2012; Rudenko et al., 2014) and errors related to precise orbit determination (POD) are demonstrably one of the major error sources for global and regional sea level products (Ablain et al., 2015). A detailed description of the main factors contributing to the radial orbit errors is given by Fu and Haines (2013). The orbit errors have typically long wavelengths and may contain systematic contributions at seasonal to decadal timescales.

Couhert et al. (2015) investigated the main contributions to the radial orbit error budget for the Jason-1 and Jason-2 series based on Geophysical Data Records (GDR)-D at seasonal to decadal time scales for the second altimetry decade (2002-2013). According to their analysis, the orbit related uncertainty of the global mean interannual and decadal trends is less than 0.1 mm/year. As main factors for regional errors they identified contributions from tracking data and from reference frame (up to 8 mm) at seasonal time scales, contributions from tracking data (up to 3 mm/year) and Earth's time variable gravity field (up to 2 mm/year) at interannual time scales, and contributions from tracking data (up to 2 mm/year) and Earth's time variable gravity field (up to 1.5 mm/year) at decadal time scales. A correspondent assessment for the first altimetry decade (1992-2001) has still been missing and is the rationale of this paper.

We assess the error budget of the radial orbit component for the TOPEX/Poseidon mission for the period 1993 to 2004 from a set of different orbit models. We have chosen TOPEX/Poseidon, since it is the reference altimetry mission used in the European Space Agency's (ESA) Climate Change Initiative (CCI) Sea Level project over this time span (Ablain et al., 2016). We assess the upper bound estimates of the radial orbit error budget at regional and global scales at seasonal, interannual, and decadal time scales by the analysis of three state-of-the-art orbit solutions derived and provided by different research teams from the German Research Centre for Geosciences (GFZ), the Groupe de Recherche de Geodesie Spatiale (GRGS), and the Goddard Space Flight Centre (GSFC). Note, that our assessment necessarily excludes contributions from errors common to these three orbits. In our further analyses, we use test orbits calculated at GFZ to investigate the impact of

uncertainties of the tracking station sub-networks, of the reference frame, and of the Earth's time variable gravity field models on the radial orbit component, and hence the derived sea level.

65 A detailed description and assessment of the analysed orbits as well as specifications of the altimeter data processing are given in Sect. 2. Sect. 3 describes the methods implemented to assess the upper bound orbit errors for the different time scales and the corresponding results for global and regional scales. The main findings are summarized and discussed in Sect. 4.

2 Orbit and altimetry data

70 2.1 Description of the analysed orbit solutions

Our aim is to assess the range and the characteristics of radial orbit errors on regional and global scales. Therefore, the differences between three independent state-of-the-art orbit solutions available for the TOPEX/Poseidon mission are analysed. All orbit solutions are derived in the International Terrestrial Reference Frame (ITRF) 2008 reference frame (Altamimi et al., 2011) and use Satellite Laser Ranging (SLR) and Doppler Orbitography and Radiopositioning Integrated by
75 Satellite (DORIS) tracking data, but are based on different software and on distinct models. The actual multi-mission GFZ orbit solution VER11 (Rudenko et al., 2017) is used as a reference in this paper and is called REF hereafter. The GSFC std1504 orbit (Lemoine et al., 2010; Beckley et al., 2015) has been chosen by the ESA CCI Sea Level Phase 2 project and differs in many aspects from the GFZ orbit, regarding software as well as the suite of implemented models including another Earth's gravity field model. As the third model, we have chosen the GRGS orbit solution (Soudarin et al., 2016), which is
80 derived using models similar to those of the GFZ solution, but employing another software package. The main models used for GFZ REF, GRGS, and GSFC std1504 orbits are described in Table 1. The main differences in these three orbit solutions are related to the choice of the Earth's time variable gravity (TVG) field models, ocean tide model, modelling of non-tidal atmospheric and oceanic gravity, and the treatment of geocenter variations in station displacements. Proper modelling of the Earth's gravity field, in particular of its time-variable part, is crucial for the computation of precise orbits of altimetry
85 satellites and has been shown to contribute to errors in regional sea level trends and seasonal signals (Rudenko et al., 2014; Esselborn et al., 2015). For the pre-GRACE period the TVG field is poorly constrained. The weekly TVG solutions used for the GSFC orbit were derived up to degree and order 5 from the analysis of SLR and DORIS observations to 20 geodetic satellites starting from 1993 (Lemoine et al., 2016). The TVG part used for the GFZ REF (GRGS) orbits consists of the combination of yearly coefficients, drift terms and annual and semi-annual variations for degree and order 1 to 80 (2 to 50)
90 derived from GRACE data and SLR measurements to LAGEOS-1/2. The annual and semi-annual coefficients used for the GFZ REF orbit are fitted yearly starting from August 2002. For the pre-GRACE period before August 2002 (January 2003) only the degree 2 terms exhibit yearly values and drift terms, however, the annual and semi-annual variations, which were derived for the GRACE-period, are applied for degree and order 1-80 (2-50) (Rudenko et al., 2014, Förste et al., 2016).

The approach adopted for the estimation of the radial orbit errors implicates that errors common to all three orbits can not be detected. In particular, all three orbits rely on the ITRF2008 reference frame and basically the same set of tracking stations. To further estimate the orbit related radial orbit error budget due to the most significant factors, we have derived five test orbits based on the GFZ REF orbit. The errors related to inconsistencies of the tracking data networks are tested by using only one tracking network instead of two. Since the GRGS orbit was derived without estimation of the DORIS system time bias, we have studied the impact of this bias on the radial orbit differences with special focus on systematic differences between ascending and descending passes. The effect of errors in the realization of the terrestrial reference frame is tested by the implementation of the most recent ITRF2014 version. The effects of uncertainties in Earth's TVG field models are tested by the implementation of the EIGEN-6S2 model which is the predecessor of the EIGEN-6S4 model. For each case, the same background models and estimated parameters were used as for the REF orbit, except for those that represent the changes for the specific test case. The five test orbits and the differences with respect to the GFZ REF orbit are:

- SLR orbit: derived by using SLR tracking observations only,
- DORIS orbit: computed by using DORIS tracking observations only,
- TBias orbit: calculated without estimation of the DORIS system time bias,
- ITRF14 orbit: calculated by using the information on station positions and velocities from ITRF2014 (Altamimi et al., 2016) instead of ITRF2008,
- Geoid orbit: obtained by using EIGEN-6S2 (Rudenko et al., 2014) Earth's gravity field model instead of EIGEN-6S4 model (Förste et al., 2016). Note that the Geoid orbit is based on the same gravity field model as the GRGS orbit.

2.2 TOPEX altimeter data

In order to assess the orbit accuracy at crossover points and to relate the estimated errors to the total variability of the sea level data, along-track TOPEX Sea Level v1.1 ECV data (Ablain et al., 2015) released from the ESA CCI Sea Level project has been included in the analyses. The along-track data has been corrected for all instrumental and geophysical effects by the state-of-the-art models provided with the data. However, for some corrections updated models were applied. These include: the GSFC std1504 orbits, EOT11a ocean tides and loading tides (Savcenko and Bosch, 2012), solid earth tides following the IERS 2003 conventions, and updated GPD+ wet tropospheric corrections (Fernandes and Lazaro, 2016). The processing of the data, the crossover point and collinear analyses as well as the interpolation to a regular grid were performed using GFZ's Altimeter Database and Processing System (ADS) Central (Schöne et al., 2010).

2.3 Evaluation of the orbit solutions

In the following, the performance of the analysed orbits is evaluated. For the GFZ orbit solutions, the consistency with tracking data and at arc overlaps is assessed. Table 2 provides the main results of precise orbit determination of the GFZ reference and test orbits, namely, the average values of SLR and DORIS RMS fits, radial, cross-track, and along-track two-

day arc overlaps, illustrating the internal orbit consistency in these directions, and the number of the arcs used to compute these values for the reference and five test orbits. Smaller values of arc overlaps and observation fits, when using the same observation types and weighting between them, indicate improved orbit quality. Reduced radial arc overlaps characterise reduced radial orbit error. SLR observations were used at all 494 orbital arcs of five GFZ orbits, except for the DORIS orbit for which no SLR observations were used at all. Since DORIS data are available for TOPEX/Poseidon only until October 31, 2004, these data were used at 459 orbital arcs preceding this date, except for the SLR orbit for which no DORIS observations were used at all. All orbital arcs for GFZ orbits are manoeuvre-free. Thus, two-day arc overlaps were computed for 433 overlaps for the REF, TBias, ITRF14, and Geoid orbits. In case of the SLR and DORIS orbits, a few gaps in the observations caused radial arc overlap larger than 0.5 m. Those arc overlaps have been excluded from the statistics resulting in less arc overlaps shown for these orbits in Table 2.

Fig. 1 and 2 provide information on the quality of the reference and tests orbits. The four orbits derived using SLR and DORIS observations provide comparable levels of average SLR RMS fits (1.96 – 1.99 cm, Fig. 1), however, the largest value is obtained for the TBias orbit, when no DORIS system time bias is estimated. The smallest average SLR RMS fits (1.59 cm) are obtained with the orbit based on SLR observations only. This is related to the weighting of the observations used (3 cm for SLR and 0.05 cm/s for DORIS) and to the number of observations used. Among five orbits derived using DORIS observations, a slightly increased average value of DORIS RMS fits (0.04795 cm/s) is obtained for the DORIS orbit derived using only DORIS observations followed by the TBias orbit (0.04785 cm/s), while the other orbits derived using SLR and DORIS observations (REF, ITRF14, and Geoid) show comparable average values of DORIS RMS fits (0.04775 – 0.04778 cm/s). Fig. 2 shows two-day arc overlaps in the radial direction for the GFZ REF and test orbits. The smallest average value of the radial overlaps (0.83 cm) is obtained using the EIGEN-6S2 geopotential model. The radial arc overlaps of the TOPEX/Poseidon orbit derived using only SLR data are 1.95 times larger than those of the orbit derived using only DORIS data. Using the reference frame ITRF2014 instead of ITRF2008 eliminates many outliers in the radial arc overlaps (Fig. 2) and therefore reduces the average value of the radial overlaps from 0.90 to 0.84 cm.

The DORIS system time bias is regularly estimated and applied during GFZ's POD process to adjust the DORIS time system to the SLR time system. Zelensky et al. (2006) have shown, that there is a strong linear relationship between along-track orbit position and the DORIS time bias. The comparison of the fits and overlap values of the REF and the TBias orbit (Table 2) show, that the estimation of the DORIS time bias improves the orbit quality. The temporal behaviour of the DORIS system time bias derived for TOPEX/Poseidon REF, ITRF14 and Geoid test orbits is in close agreement (Fig. S1) and resembles the estimation given by Lemoine et al. (2016). For the GFZ VER11 (REF) orbit, it indicates variations between -22.4 microsecond (μs) and +4.4 μs from 1992.73 to 1994.18, followed by a period of a linear trend of 35.11 $\mu\text{s}/\text{year}$ from 1994.18 to 1995.00 that ends with a jump from -28.65 μs to +1.98 μs around 1995.00. Then the DORIS time bias shows two rather stable periods with a mean value of +3.70 μs with a standard deviation of 1.77 μs from 1995.0 to 1999.0 and a mean values -1.32 μs with a standard deviation 1.19 μs from 1999.0 to 2001.13, followed again by a period of a linear trend (-3.14

160 $\mu\text{s}/\text{year}$) from 2001.13 to 2004.83. The mean value of the DORIS system time bias is $0.04 \pm 0.36 \mu\text{s}$ for the DORIS test orbit, and it is equal to zero (not shown in the figure) for the TBias orbit.

For all orbit solutions, a crossover point analysis for the period April 1993 to September 2004 has been performed based on the altimeter data described in Sect. 2.2. Differences between the values of ascending and descending passes at crossover points are caused by oceanic variability and errors related to the measurements, the orbit, and the applied corrections. Since in our study errors related to the measurements and the applied corrections and oceanic variability are always identical, here, smaller absolute mean differences and decreased RMS values at crossover points are indicative for increased orbit quality. The median of the time series of global mean height differences and RMS values at the crossover points are provided in Table 3. The smallest ascending/descending differences (-1.6 mm) and as well the lowest RMS values (49.5 mm) at the crossover points are reached by the GSFC orbit solution. The mean global ascending/descending differences are -3.1 mm for the GFZ REF and -2.9 mm for the GRGS orbit solutions. However, while the RMS value of the GFZ REF solution (49.8 mm) is comparable to the one of the GSFC, the GRGS orbit solution shows degraded performance (51.3 mm RMS). Keeping the DORIS time bias fixed to zero deteriorates the mean differences between ascending and descending passes to -3.6 mm but does not change the RMS value. The median of the global mean ascending/descending differences is -2.7 mm for the SLR and -4.7 mm for the DORIS orbits. Both orbit solutions show degraded performance (51.1/50.7 mm RMS) with respect to the REF solution. This shows that using SLR and DORIS observations together improves the orbit quality considerably, even though the DORIS observations seem to aggravate the mean differences between ascending and descending tracks. Using ITRF2014 instead of ITRF2008 does not change the crossover point statistics. The Geoid orbit solution exhibits clearly improved ascending/descending differences (-2.1 mm) and as well a slight reduction of the RMS values. A further analysis of the temporal evolution of the ascending/descending differences reveals, that these improvements take place in the pre-GRACE period before August 2002.

180 **3 Estimation of the orbit related sea level error**

Sea level is varying on typical temporal and spatial scales, that are often connected to the driving processes. At the same time, orbit errors are not randomly distributed but exhibit also typical temporal and spatial pattern. Here, we apply statistical methods in order to assess upper bound errors related to the orbit solutions for global and regional sea level at seasonal to decadal time scales.

185 **3.1 Methods**

In order to estimate upper limits for orbit related errors in sea level height, the differences between the radial components of the GFZ REF orbit and the two independent orbit solutions (GSFC and GRGS) have been analysed. To assess the effect of uncertainties in the reference system, in the realisation of the tracking station networks, and in Earth's time variable gravity on the radial error budget, we have evaluated the differences of the radial orbit components between the GFZ's REF and

190 ITRF14, SLR, DORIS, and Geoid test orbits. Since the radial orbit components map directly to the derived sea level heights, we consider the differences presented here to represent estimates of the orbit related sea level error. However, since the orbit error analysis is based on orbit differences, any error common to all three orbits will be lacking in our assessment.

The differences of the radial orbit components at the time of the altimetry measurement (1 Hz, ~6.7 km on ground) are calculated and interpolated to a global $1^\circ \times 1^\circ$ grid for every cycle (9.92 days). In general, we merge both, ascending and descending, passes in our calculations. In addition, we analyse ascending and descending passes for some orbit combinations separately.

For the regional analyses, starting from the gridded radial orbit differences, the RMS values relative to the local temporal mean are calculated for each grid point over the entire time series. Decadal trends, annual and semi-annual signals, and the corresponding formal errors are estimated by a least-square fit. As a measure for errors at interannual time scales, we calculate the RMS of the five-year running trend series of the radial orbit differences at each grid point. Regional upper bound errors are guessed from the corresponding maximum RMS values over the ocean at the $1^\circ \times 1^\circ$ grid.

For the global analyses, the gridded radial orbit differences are averaged (with area weighting) over the ocean ($\pm 67^\circ$ latitude). Starting from these global mean height differences, global mean RMS, decadal trend, annual cycle, and the RMS of the five-year running trends series are derived, based on the same methods as used for the regional analyses. Global mean RMS values per cycle are calculated as the square root of the spatial mean of the radial orbit differences at each grid point over the ocean for the respective cycle.

In order to relate the estimated errors to the total variability of the sea level data, TOPEX altimeter data has been included as well. The data and the processing is described in Sect. 2.2. From the gridded sea level anomalies, seasonal, interannual and decadal trends were derived using the methods described above.

210 **3.2. Global mean errors**

In the following, we investigate the orbit related global sea level error, differentiating between the total error and its annual, interannual, and decadal components. The TBias orbit differences are not included in these analyses but will be further investigated for the study of changes between ascending and descending passes (Sect. 3.4). The time series of the global mean RMS of gridded radial orbit differences per cycle are shown in Fig. 3 for all orbit solutions relative to GFZ's REF orbit. The largest differences occur between the REF and the GRGS orbits, the smallest changes occur for the ITRF14 test orbit. Most orbit differences are dominated by sub-seasonal variability, only the Geoid and ITRF14 orbit differences are governed by seasonal and decadal periods. For the Geoid, GSFC, and GRGS orbit differences relative to the REF orbit, the RMS series exhibit a seasonal cycle, which is an indication for seasonal orbit differences on regional scales. The RMS of the *REF minus Geoid* orbit difference is decreased after August 2002 indicating that the main differences between the two orbits originate from the pre-GRACE period. In contrast, the differences between the REF and the ITRF14 orbits are slightly increasing from 2000 onwards. The orbit errors derived from the analysis of the global mean orbit difference series over the oceans are summarized in Table 4 for all orbit models together with the corresponding absolute sea level values. The global

mean RMS of the radial orbit differences between the REF and GRGS (GSFC) orbits amount to 7.0 (5.4) mm, which corresponds to more than 10 % of the global mean sea level variability of 52.5 mm. The restriction to one tracking station sub-network leads to large changes of the orbit, for the DORIS (SLR) orbit solution the RMS values of the radial differences with respect to the REF orbit amount to 5.1 mm (4.2 mm), which is almost of the size of the estimated upper bound orbit errors. This highlights the importance of manifold, precise, and consistent tracking data for accurate global mean sea level estimates. The substitution of the Earth's gravity field model (EIGEN-6S4 by EIGEN-6S2) and the ITRF realization (ITRF2008 by ITRF2014) accounts for 2.0 mm and 1.1 mm, respectively, of the mean orbit errors. A spectral analysis of the global mean radial differences (Fig. S2) exhibits peaks at ~60 days for all but the GSFC and TBias orbit differences and at ~90 and ~170 days for the SLR and DORIS orbit differences. A weak annual component can be observed for the GRGS and Geoid orbit differences. Since the annual amplitude is less than 1 mm only, it can be neglected and is not included in Table 4. The time series of the five-year running trends of the global mean radial orbit differences over the ocean are shown in Fig. 4 for the various orbit combinations. All curves range between ± 0.2 mm/year and show at least one zero-crossing and imply interannual changes of the estimated decadal sea level trends. The corresponding curve of the five-year running trends for the global mean sea level (not shown) range between 4 mm/year at the beginning of the time series and 2 mm/year at the end. Before 1998 the GSFC and GRGS solutions are close to each other and both suggest larger sea level trends for this period than the GFZ solution. After that, trends derived from GFZ and GSFC orbits show good agreement, while the GRGS solution exhibits smaller trends till 2001 which would result in smaller sea level trends for this period when using the GRGS orbit. Maximum interannual trend variability of 0.1 mm/year occurs between the REF and GRGS orbits (Table 4) which amounts to almost 20 % of the corresponding value derived for the global mean sea level curve (0.55 mm/year). Hence, the use of the GFZ instead of the GRGS orbit should result in a slightly stronger acceleration of the mean sea level curve during the TOPEX period. Since the exclusive use of DORIS tracking station leads to interannual trend variability of 0.1 mm/year, inconsistencies of the tracking stations sub-networks might explain large portions of the observed global mean interannual variability. The errors of the interannual trend variability are for all orbit combinations higher than for the decadal trends. The global mean decadal trends (calculated over the full mission time) are mostly significant but can be further neglected, since they are two orders of magnitude smaller than the observed sea level signal over this period (~3 mm/year).

3.3 Regional errors

The maximum regional errors derived from the analysis of the gridded orbit difference series over the oceans are summarized in Table 5. The TBias orbit differences are not included in these analyses but will be further investigated for the study of changes between ascending and descending passes (Sect. 3.4). Regionally, the maximum radial orbit differences on the $1^\circ \times 1^\circ$ grid between the REF and GRGS (GSFC) orbits amount to 10.7 (7.4) mm. The exclusive use of only one tracking station sub-network leads to distinct changes with RMS values of 9.3 mm (7.2) mm for the DORIS (SLR) sub-network. This suggests that for the weighting factors applied with GFZ's REF orbit especially inhomogeneity in the SLR station sub-network has the potential to produce notable regional orbit errors.

Annual difference signals with respect to the REF orbit are most prominent for the GSFC and GRGS solutions, while they are negligible for the SLR, DORIS, and ITRF14 orbits. The corresponding patterns of the annual amplitudes for the differences of REF versus GSFC, GRGS, and Geoid orbits and of GRGS versus GSFC orbits are shown in Fig. 5. The observed patterns for the GSFC and GRGS orbit differences consist of a dipole with centers in the southeastern Indian Ocean and the Caribbean. Since the two centres are phase shifted by half a year, the effect on the global mean differences is marginal. The pattern coincides with the patterns already shown to be related to the use of AOD1B products (Rudenko et al., 2016) and different Earth's time variable gravity fields for TOPEX/Poseidon POD (Esselborn et al., 2016). However, the annual differences between the REF and Geoid orbits can only explain part of the observed differences between the REF and GRGS orbits. In addition, the annual differences between GRGS and GSFC orbits are quite small and show no distinct pattern. Another plausible source of the relatively strong signal for the GSFC and GRGS orbit cases are the differences in the annual corrections for station coordinates by geocenter motion corrections and non-tidal atmospheric loading. The thorough reflection of the relevant models used for the POD of these three orbits suggests, that the observed differences originate in part from the non-tidal atmospheric loading effect on the stations which was applied for the GFZ but not the GRGS and GSFC orbits. There is evidence, that the annual signal from the EIGEN-6S2 gravity field model is closer to the solution applied for the GSFC orbits than to EIGEN-6S4 – at least in the pre-GRACE period.

The patterns of the interannual variability of the regional trends are shown in Fig. 6 for all orbit differences. The trend errors reach up to 1.2 (0.9) mm/year for the GSFC (GRGS) orbit differences (Table 5). The patterns of the trend variability from the GSFC and GRGS differences show coinciding maxima in the regions around South America and Australia. The differences for the Geoid orbit show similar features even though the absolute trend variability is smaller (up to 0.4 mm/year). For the SLR and DORIS orbit differences, the patterns of interannual variability (Fig. 6) are patchy and oriented along individual tracks. For the ITRF14 solution, the trend variability is slightly increased at high latitudes (up to 0.2 mm/year). The patterns of the interannual trend variability derived from the GFZ test orbits suggest, that differences in the TVG modelling and contributions from the tracking systems are the most plausible sources of the observed regional differences of trend variability between REF, GSFC, and GRGS orbits.

The strongest regional changes in the decadal trend (Fig. 7 and Table 5) are observed for the differences between the REF and GSFC orbits (up to 1.0 mm/year). For the GSFC orbit, high absolute decadal trend differences tend to coincide with maximum seasonal differences, but not with maximum interannual variability. The differences between the REF and GRGS orbit trends reach 0.7 mm/year at maximum and the patterns of maximum annual amplitudes, interannual and decadal trend differences coincide. The differences between the REF and Geoid orbit trends resemble these patterns, however, the trend values are smaller (up to 0.4 mm/year) and can explain only about half of the observed decadal trend differences. The source of decadal trends between EIGEN-6S2 and EIGEN-6S4 during the TOPEX period are presumably differences in the modelling of the TVG after August 2002, since before drift terms are only applied to degree 2 terms. The degree 2 terms, in turn, are defined by SLR data and show close agreement between the two TVG models for the pre-GRACE period. The ITRF14 orbit differences drift locally by a rate of up to 0.2 mm/year with positive values in the southern hemisphere and

290 negative values in the northern hemisphere, indicating a drift in the z-component between the reference system realisations. The observed values are in good agreement with the combined change of scale and rate of the z-component of the transformation between ITRF2008 and ITRF2014 (Altamimi et al., 2016). The regional decadal trends for the SLR and DORIS orbit differences are patchy and rather related to particular tracks without consistent long-wavelength behaviour. Higher trends of up to 0.4 mm/year emerge for the DORIS orbit. The patterns of the decadal trend differences derived from
295 the GFZ test orbits suggest, that differences in the TVG modelling are the most plausible source of the observed regional trend differences between REF, GSFC, and GRGS orbits.

3.4 Differences between ascending and descending passes

The crossover point analysis (Table 3) reveals considerable global mean differences between ascending and descending passes for most orbits. Fu and Haines (2013) have shown, that orbit errors might induce diverging drifts for sea level derived
300 from ascending and descending passes. In the following, we study whether there are systematic changes to the results obtained so far when ascending and descending passes are investigated separately. Therefore, for a subset of orbit solutions the same analyses were performed as before, but for data sets derived from ascending and descending passes only. Since the DORIS orbit reveals the most pronounced median ascending/descending differences we have chosen to study the *REF minus DORIS* and the *REF minus TBias* orbit differences further. During the POD of the GRGS orbit, the DORIS system time bias
305 is not estimated, therefore, we include the GRGS orbit in the analysis as well. However, in contrast to the previous analysis, we study the difference *Geoid minus GRGS* instead of *REF minus GRGS* in order to exclude the effects of different time variable gravity fields from the analysis.

The global mean radial orbit differences for ascending and descending passes are for all three cases in the range of ± 12 mm (Fig. S3). The ascending and descending radial orbit differences are significantly anti-correlated. The correlation coefficient
310 is almost -1 for the *REF minus TBias* case, almost -0.9 for the *Geoid minus GRGS* case, and still -0.5 for the *REF minus DORIS* case. The correlation is further increased for periods of more than one year. The *REF minus TBias* global mean time series resembles the DORIS system time bias applied for the REF orbit (Fig. S1). The global mean radial differences for the *Geoid minus GRGS* case reveal similar features as well. All three orbit differences exhibit diverging global mean radial differences for ascending and descending tracks after the year 2000. The interannual trend variability and decadal trends
315 derived from the analysis of the global mean radial orbit difference series over the oceans are summarized in Table 6 for the merged, ascending, and descending passes. If the ascending and descending passes are analysed separately, the interannual trend variability is increased by at least five times for the corresponding orbit differences. Ascending passes exhibit higher variability than the descending. The differences for the global mean decadal trends between ascending and descending passes are a multiple of the values for the merged data and reach up to 0.6 mm/year, where both data sets are drifting in
320 opposite directions.

The regional patterns of the decadal trend differences for ascending and descending passes are shown in Fig. 8. The DORIS orbit differences reveal a striking spread between the decadal trends of the ascending and descending passes. The trends are

opposite for ascending and descending passes for most areas of the global ocean and reach regionally absolute values of up to 0.8 mm/year. Trends for the *REF minus TBias* orbit differences are very similar but smaller than the *REF minus DORIS* orbit ones. The corresponding analysis for the *Geoid minus GRGS* orbit differences shows again very similar features as for the DORIS differences. This indicates that discrepancies in the reference systems of the tracking stations (distribution of tracking stations, observation sampling, etc.) might give rise to long-wavelength orbit errors being anti-correlated for ascending and descending passes. Relevant contributions are originating from uncertainties of the timing of the DORIS measurements. Increasing time biases are related to increasing along-track position errors and seem to be transferred to radial orbit errors. This mechanism is not fully understood but a further analysis is beyond the scope of this paper. The uncertainties are especially pronounced in tropical and subtropical regions. On regional scales, the interannual and decadal trend errors derived from ascending/descending passes separately can be many times higher than the values derived from the merged data. Even though such effects tend to cancel, whenever both components are merged, they might still introduce considerable errors in regional studies, that are based on along-track data, e.g. at calibration sites.

3.5 Regional orbit errors and sea level variability

Our analysis exhibits large scale patterns of the orbit related error. Errors for interannual to decadal sea level trends of more than 1 mm/year might hamper the interpretation of the observed sea level variability from altimetry, at least apart from the large oceanic currents. In order to define regions where the orbit related error should be considered when analysing sea level data from TOPEX, we have determined areas with orbit errors of at least 10 % of the corresponding sea level value. Fig. 9 shows the sea level variability, seasonal signal, interannual and decadal trends derived from ESA CCI TOPEX altimeter data for those regions where the orbit error amounts to at least 10 % of the corresponding sea level value. Taking into account the total orbit related error, about half the ocean is affected. This includes especially calm oceanic regions, whereas for energetic regions like the Circumpolar Current, Tropical Pacific, and the western boundary currents of the northern hemisphere the dynamic ocean signal is much larger than the orbit error. For the seasonal signal, mainly the Southern Ocean is concerned. Critical regions for the estimation of the interannual variability are the Tropical and Subtropical Atlantic and the south-eastern Pacific. For decadal scales, the orbit related trend errors are prominent in a couple of regions including: South Atlantic, western North Atlantic, central Pacific, and south-eastern Indian Ocean, but also several marginal seas including the Mediterranean, Red Sea, Yellow Sea and Sea of Japan.

4 Summary and Conclusions

We have investigated the radial orbit error budget associated with three state-of-the-art orbit solutions from GFZ, GSFC and GRGS over the first altimetry decade (1993-2004). It is crucial to know the accuracy of these early altimeter data in order to judge the reliability of long-term sea level trends and of estimations of the acceleration of global mean sea level rise. For this purpose, we have chosen the TOPEX/Poseidon mission, since it is the reference altimetry mission used in ESA's CCI Sea

Level project over this time span. We estimate the orbit errors from the radial orbit differences which implies that errors
355 common to all orbits can not be detected. A set of five test orbit solutions derived at GFZ is used to estimate the
contributions of the most significant factors to the error budget. We have focused on the impact of uncertainties of the
tracking station sub-networks (SLR and DORIS), of the reference frame, and of the Earth's time variable gravity field
models on the radial orbit component, and hence the derived sea level. The estimates of the upper bound radial orbit errors at
seasonal, interannual (5 years), and decadal time scales are given in Table 4 for the global mean sea level and in Table 5 for
360 the regional sea level.

According to our study, the global mean RMS radial orbit errors for the TOPEX period are of the order of 7 mm, which
corresponds to more than 10 % of the global mean sea level variability (52 mm). The global mean annual (seasonal)
component of the radial error is below 1 mm and can be neglected. The orbit related errors of the decadal trends are less than
0.05 mm/year and should not induce any significant artificial global mean sea level trends. However, on time scales of five
365 years the trend variability may reach up to 0.1 mm/year, which amounts to almost 20 % of the corresponding sea level
variability (0.55 mm/year), and could potentially hamper the detection of sea level acceleration from the altimeter data. The
major contributions to this error (0.04 – 0.10 mm/year) are, most probably, discrepancies of the station sub-networks
(DORIS or SLR) used. The contributions of Earth's time variable gravity field model and the ITRF realisation (ITRF2008
versus ITRF2014) to the global mean error are of only minor importance (0.02 mm/year). These values are in line with the
370 mean upper bound orbit errors given by Couhert et al. (2015) derived for Jason-1 and Jason-2 orbits for the second altimetry
decade (2002-2012).

For regional scales, the maximum RMS of the gridded radial orbit error is more than 10 mm. However, this error includes a
large fraction of sub-seasonal variability which is not subject of this study. The regional upper bound error of the seasonal
signal is 6 mm, of the interannual trend variability 1.2 mm/year, and of the decadal trend 1 mm/year. Errors for interannual
375 to decadal sea level trends of more than 1 mm/year might hamper the interpretation of the observed sea level variability from
altimetry. For about half of the ocean outside the energetic regions (e.g. Circumpolar Current, Tropical Pacific, Gulf Stream
and Kuroshio System) the orbit related errors reach at least 10% of the observed sea level variability. For the seasonal signal,
mainly the Southern Ocean is concerned. Critical regions for the estimation of the interannual variability are the Tropical and
Subtropical Atlantic and the south-eastern Pacific. For decadal scales, the orbit related trend errors are prominent in a couple
380 of regions including: South Atlantic, western North Atlantic, central Pacific, and south-eastern Indian Ocean, but also several
marginal seas including the Mediterranean, Red Sea, Yellow Sea and Sea of Japan.

When using ascending and descending passes separately, the interannual and decadal trend errors can reach multiples of the
values derived from the merged data. This is the case for global mean values as well as for regional values. The
corresponding large scale pattern is coherent for low and medium latitudes and is strongly anti-correlated for ascending and
385 descending passes. Even though such effects tend to cancel, whenever both components are merged, they might still
introduce considerable errors in regional studies, that are based on along-track data, e.g. at calibration sites.

Orbit errors related to discrepancies between the tracking station sub-networks (distribution of tracking stations, observation sampling, etc.) are studied based on GFZ's SLR, DORIS, and TBias orbit solutions. Using both SLR and DORIS observations for TOPEX POD together reduces (improves) the RMS of the altimetry single-satellite crossover differences considerably (2-3%), though the DORIS observations seem to aggravate the mean differences between ascending and descending passes. The proper estimation of the DORIS system time bias has proven to be a critical factor for the minimization of this effect. The most significant changes are observed for the DORIS orbit solution suggesting that uncertainties of the SLR station sub-network should have the most prominent effects on the orbit accuracy – at least for GFZ's orbit solutions. This fact is, most probably, related to the weighting factors applied to the observations within the GFZ orbit determination process. Using the latest reference frame ITRF2014 instead of the predecessor ITRF2008 slightly improves the accuracy of the TOPEX/Poseidon orbit solution. The contribution of the uncertainties in the ITRF realisation to the regional upper bound error is only marginal. Errors induced by uncertainties of the Earth's time variable gravity field model are studied on the base of GFZ's Geoid orbit solution. The orbit evaluations show that the Geoid orbit performs slightly better than the REF orbit in the pre-GRACE period due to differences in the time-invariant annual and semiannual variations applied to the TVG field models. Uncertainties of the gravity field model give rise to orbit errors at all analysed periods. We estimate regional upper bound errors of ~3 mm for the seasonal signal and of 0.4 mm/year for the interannual trend variability and the decadal trend. This accounts for about 60 % of the seasonal, about 30 % of the interannual, and about 40 % of the decadal orbit error which are related to differences between EIGEN-6S2 and EIGEN-6S4.

The regional upper bound radial orbit errors obtained from our study are by factor 2 to 5 smaller than the ones reported by Couhert et al. (2015) for the period 2002 to 2012. This might partly reflect recent improvements of the stability of reference frames which result in smaller changes from ITRF2008 to ITRF2014 than previously from ITRF2005 to ITRF2008. However, the accuracy of the Earth's time variable gravity model and the tracking observations for the 1990's should be inferior to more recent periods. The error related to the uncertainties of the tracking station sub-networks might be underrated in our study since all analysed orbits rely on basically the same set of tracking observations. The effect of uncertainties of the time variable gravity (TVG) field might be underestimated as well since EIGEN-6S4 and EIGEN-6S2 both model the TVG field in the pre-GRACE period by time-invariant annual and semi-annual variations derived from GRACE plus annual values and drift terms for degree two terms derived from SLR measurements. In contrast, the TVG field used for the GSFC orbit determination is weekly changing. Using SLR measurements to geodetic cannon-ball satellites (Sośnica et al., 2015, Bloßfeld et al., 2016) and in combination with DORIS measurements to altimetry and remote sensing satellites (Lemoine et al., 2016) allows to determine Earth's time variable gravity for the period 1993-2003, i.e. before GRACE, more precisely than just using SLR measurements to LAGEOS-1/2. Combined use of SLR and DORIS measurements to numerous geodetic satellites, especially for 1990-2003, with the GRACE measurements should further improve Earth's time variable gravity field models and hence further enhance orbit solutions for the ERS and the TOPEX/Poseidon altimetry missions.

420 **Acknowledgements**

We are grateful for the insightful comments of the reviewer Dr. Nikita Zelensky which helped to improve the manuscript substantially. We thank Goddard Space Flight Centre and Groupe de Recherche de Geodesie Spatiale for the provision of GSFC std1504 and GRGS orbit solutions, ESA CCI for along-track TOPEX Sea Level v1.1 ECV data, and Joana Fenandes for the provision of updated wet troposphere corrections (GPD+). This research was partly supported by the European Space Agency (ESA) within the Climate Change Initiative Sea Level (SLCCI) Phase II project and by the Deutsche Forschungsgemeinschaft (DFG) through grant (CoRSEA) as part of the Special Priority Program (SPP)-1889 "Regional Sea Level Change and Society" (SeaLevel) and within the projects "Consistent dynamic satellite reference frames and terrestrial geodetic datum parameters" and "Interactions of low-orbiting satellites with the surrounding ionosphere and thermosphere (INSIGHT)" and by the International Office of the BMBF under the grant 01DO17017 "Sea Level Change and its Hazardous Potential in the East China Sea and Adjacent Waters" (SEAHAP).

435

440

445

450 **References**

- Ablain, M., Cazenave, A., Valladeau, G. and Guinehut, S.: A new assessment of the error budget of global mean sea level rate estimated by satellite altimetry over 1993–2008, *Ocean Sci.*, 5(2), 193–201, doi:10.5194/os-5-193-2009, 2009.
- Ablain, M., Cazenave, A., Larnicol, G., Balmaseda, M., Cipollini, P., Faugère, Y., Fernandes, M. J., Henry, O., Johannessen, J. A., Knudsen, P., Andersen, O., Legeais, J., Meyssignac, B., Picot, N., Roca, M., Rudenko, S., Scharffenberg, M. G.,
455 Stammer, D., Timms, G. and Benveniste, J.: Improved sea level record over the satellite altimetry era (1993–2010) from the Climate Change Initiative project, *Ocean Sci.*, 11(1), 67–82, doi:10.5194/os-11-67-2015, 2015.
- Ablain, M., Legeais, J. F., Prandi, P., Marcos, M., Fenoglio-Marc, L., Dieng, H. B., Benveniste, J. and Cazenave, A.: Satellite Altimetry-Based Sea Level at Global and Regional Scales, *Surv. Geophys.*, 1–25, doi:10.1007/s10712-016-9389-8, 2016.
- 460 Altamimi, Z., Collilieux, X. and Métivier, L.: ITRF2008: an improved solution of the international terrestrial reference frame, *J. Geod.*, 85(8), 457–473, doi:10.1007/s00190-011-0444-4, 2011.
- Altamimi, Z., Rebischung, P., Métivier, L. and Collilieux, X.: ITRF2014: A new release of the International Terrestrial Reference Frame modeling nonlinear station motions, *J. Geophys. Res. Solid Earth*, 121(8), 2016JB013098, doi:10.1002/2016JB013098, 2016.
- 465 Beckley, B., Ray, R., Holmes, S., Zelensky, N., Lemoine, F., Yang, X., Brown, S., Desai, S., Mitchum, G. and Hausman, J.: Integrated Multi-Mission Ocean Altimeter Data for Climate Research TOPEX/Poseidon, Jason-1 and OSTM/Jason-2 User's Handbook Version 3.0, pp. 61, California Institute of Technology, ftp://podaac.jpl.nasa.gov/allData/merged_alt/L2/TP_J1_OSTM/docs/v121415.version3.0_multi_alt_handbook.pdf (Accessed May 2017), 2015.
- 470 Bloßfeld, M., Stefka, V., Müller, H., and Gerstl M.: Satellite laser ranging - a tool to realize GGOS? In: Rizos C., Willis P. (Eds.) IAG 150 Years, IAG Symposia 143, 540-547, doi:10.1007/1345_2015_202, 2016
- Boehm, J. and Schuh, H.: Vienna mapping functions in VLBI analyses, *Geophys. Res. Lett.*, 31(1), L01603, doi:10.1029/2003GL018984, 2004.
- Carrère, L., Lyard, F., Cancet, M., Roblou, L. and Guillot, A.: FES 2012: a new tidal model taking advantage of nearly 20
475 years of altimetry measurements, OSTST meeting, September 22-29, Venice-Lido, Italy, [online] Available from: http://www.avisio.altimetry.fr/fileadmin/documents/OSTST/2012/oral/01_thursday_27/03_tides/04_TID_Carrere2.pdf (Accessed May 2017), 2012.
- Cazenave, A., Dieng, H.-B., Meyssignac, B., von Schuckmann, K., Decharme, B. and Berthier, E.: The rate of sea-level rise, *Nature Clim. Change*, 4(5), 358–361, doi:10.1038/nclimate2159, 2014.
- 480 Cerri, L. and Ferrage, P.: DORIS satellites models implemented in POE processing, CNES, Paris, France, Tech. Rep. SALP-NTBORD-OP-16137-CN, Rev. 10, [online] Available from: <ftp://ftp.ids-doris.org/pub/ids/satellites/DORISSatelliteModels.pdf> (Accessed May 2017), 2016.
- Church, J. A. and White, N. J.: Sea-Level Rise from the Late 19th to the Early 21st Century, *Surv. Geophys.*, 32(4–5), 585–602, doi:10.1007/s10712-011-9119-1, 2011.

- 485 Couhert, A., Cerri, L., Legeais, J.-F., Ablain, M., Zelensky, N. P., Haines, B. J., Lemoine, F. G., Bertiger, W. I., Desai, S. D. and Otten, M.: Towards the 1 mm/y stability of the radial orbit error at regional scales, *Adv. Space Res.*, 55(1), 2–23, doi:10.1016/j.asr.2014.06.041, 2015.
- Dobslaw, H., Flechtner, F., Bergmann-Wolf, I., Dahle, C., Dill, R., Esselborn, S., Sasgen, I. and Thomas, M.: Simulating high-frequency atmosphere-ocean mass variability for dealiasing of satellite gravity observations: AOD1B RL05, *J. Geophys. Res. Oceans*, 118(7), 3704–3711, doi:10.1002/jgrc.20271, 2013.
- 490 Douglas, B. C.: Global Sea Rise: A Redetermination, *Surv. Geophys.*, 18(2–3), 279–292, doi:10.1023/A:1006544227856, 1997.
- Esselborn, S., Schöne, T. and Rudenko, S.: Impact of Time Variable Gravity on Annual Sea Level Variability from Altimetry, in *IAG 150 Years*, pp. 55–62, Springer, Cham., doi:10.1007/1345_2015_103, 2015.
- 495 Fasullo, J. T., Nerem, R. S. and Hamlington, B.: Is the detection of accelerated sea level rise imminent?, *Scientific Reports*, 6, 31245, doi:10.1038/srep31245, 2016.
- Fernandes, M. J. and Lázaro, C.: GPD+ Wet Tropospheric Corrections for CryoSat-2 and GFO Altimetry Missions, *Remote Sensing*, 8(10), 851, doi:10.3390/rs8100851, 2016.
- Förste, C., Bruinsma, S., Abrikosov, O., Rudenko, S., Lemoine, J.-M., Marty, J.-C., Neumayer, K. H. and Biancale, R.: EIGEN-6S4 A time-variable satellite-only gravity field model to d/o 300 based on LAGEOS, GRACE and GOCE data from the collaboration of GFZ Potsdam and GRGS Toulouse, [online] Available from: <https://doi.org/10.5880/icgem.2016.008> (Accessed May 2017), 2016.
- 500 Fu, L.-L. and Haines, B. J.: The challenges in long-term altimetry calibration for addressing the problem of global sea level change, *Adv. Space Res.*, 51(8), 1284–1300, doi:10.1016/j.asr.2012.06.005, 2013.
- 505 Goiginger, H., Hoeck, E., Rieser, D., Mayer-Guerr, T., Maier, A., Krauss, S., Pail, R., Fecher, T., Gruber, T. and Brockmann, J.: The combined satellite-only global gravity field model GOCO02S, *Geophysical Research Abstracts*, 13, EGU2011-10571, [online] Available from: <http://www.goco.eu/data/egu2011-10571-goco02s.pdf> (Accessed May 2017), 2011.
- Hedin, A. E.: MSIS-86 Thermospheric Model, *J. Geophys. Res.: Space Physics*, 92(A5), 4649–4662, doi:10.1029/JA092iA05p04649, 1987.
- 510 Hay, C. C., Morrow, E., Kopp, R. E. and Mitrovica, J. X.: Probabilistic reanalysis of twentieth-century sea-level rise, *Nature*, 517(7535), 481–484, doi:10.1038/nature14093, 2015.
- IERS Conventions (2003), McCarthy, D.D. and Petit, G. (Eds.): Bundesamt für Kartographie und Geodäsie, Frankfurt am Main. [online] Available from: <http://www.iers.org/TN32> (Accessed May 2017), 2004.
- 515 IERS Conventions (2010), Petit, G. and Luzum, B. (Eds.): Bundesamt für Kartographie und Geodäsie, Frankfurt am Main. [online] Available from: <https://www.iers.org/TN36> (Accessed May 2017), 2010.
- Jevrejeva, S., Moore, J. C., Grinsted, A. and Woodworth, P. L.: Recent global sea level acceleration started over 200 years ago?, *Geophys. Res. Lett.*, 35(8), L08715, doi:10.1029/2008GL033611, 2008.
- Jevrejeva, S., Moore, J. C., Grinsted, A., Matthews, A. P. and Spada, G.: Trends and acceleration in global and regional sea levels since 1807, *Global and Planetary Change*, 113, 11–22, doi:10.1016/j.gloplacha.2013.12.004, 2014.
- 520 Knocke, P., Ries, J. and Tapley, B.: Earth radiation pressure effects on satellites, in *AIAA Paper 88-4292*, pp. 577–587, Austin, TX, United States, 1988.

- Lemoine, F. G., Zelensky, N. P., Chinn, D. S., Pavlis, D. E., Rowlands, D. D., Beckley, B. D., Luthcke, S. B., Willis, P., Ziebart, M., Sibthorpe, A., Boy, J. P. and Luceri, V.: Towards development of a consistent orbit series for TOPEX, Jason-1, and Jason-2, *Adv. Space Res.*, 46(12), 1513–1540, doi:10.1016/j.asr.2010.05.007, 2010.
- Lemoine, F. G., Chinn, D. S., Zelensky, N. P., Beall, J. W. and Le Bail, K.: The development of the GSFC DORIS contribution to ITRF2014, *Adv. Space Res.*, 58(12), 2520–2542, doi:10.1016/j.asr.2015.12.043, 2016.
- Llovel, W., Becker, M., Cazenave, A., Jevrejeva, S., Alkama, R., Decharme, B., Douville, H., Ablain, M. and Beckley, B.: Terrestrial waters and sea level variations on interannual time scale, *Glob. Planet. Change*, 75(1–2), 76–82, doi: 10.1016/j.gloplacha.2010.10.008, 2011.
- Lyard, F., Lefevre, F., Letellier, T. and Francis, O.: Modelling the global ocean tides: modern insights from FES2004, *Ocean Dyn.*, 56(5–6), 394–415, doi:10.1007/s10236-006-0086-x, 2006.
- Masters, D., Nerem, R. S., Choe, C., Leuliette, E., Beckley, B., White, N. and Ablain, M.: Comparison of Global Mean Sea Level Time Series from TOPEX/Poseidon, Jason-1, and Jason-2, *Marine Geodesy*, 35(sup1), 20–41, doi:10.1080/01490419.2012.717862, 2012.
- Mendes, V. B. and Pavlis, E. C.: High-accuracy zenith delay prediction at optical wavelengths, *Geophys. Res. Lett.*, 31(14), L14602, doi:10.1029/2004GL020308, 2004.
- Pavlis, E. C., SLRF2008: The ILRS reference frame for SLR POD contributed to ITRF2008, presented at the Ocean Surf. Topogr. Sci. Team Meeting, Seattle, WA, USA, Jun. 2009. [online] Available from: http://www.avisioceanobs.com/fileadmin/documents/OSTST/2009/poster/Pavlis_2.pdf (Accessed May 2017), 2009.
- Quartly, G. D., Legeais, J.-F., Ablain, M., Zawadzki, L., Fernandes, M. J., Rudenko, S., Carrère, L., García, P. N., Cipollini, P., Andersen, O. B., Poisson, J.-C., Mbajon Njiche, S., Cazenave, A., Benveniste, J.: A new phase in the production of quality-controlled sea level data, *Earth Syst. Sci. Data*, 9(2), 557–572, doi:10.5194/essd-9-557-2017, 2017.
- Ray, R. D.: Precise comparisons of bottom-pressure and altimetric ocean tides, *J. Geophys. Res. Oceans*, 118(9), 4570–4584, doi:10.1002/jgrc.20336, 2013.
- Ries, J.: Annual geocenter motion from space geodesy and models, abstract G12A-08, AGU Fall Meeting, San Francisco, 9–13 December 2013, [online] Available from: <http://ids-doris.org/images/documents/report/publications/AGU2013-AnnualGeocenterMotion-Ries.pdf> (Accessed May 2017), 2013.
- Rudenko, S., Otten, M., Visser, P., Scharroo, R., Schöne, T. and Esselborn, S.: New improved orbit solutions for the ERS-1 and ERS-2 satellites, *Adv. Space Res.*, 49(8), 1229–1244, doi:10.1016/j.asr.2012.01.021, 2012.
- Rudenko, S., Dettmering, D., Esselborn, S., Schöne, T., Förste, C., Lemoine, J.-M., Ablain, M., Alexandre, D. and Neumayer, K.-H.: Influence of time variable geopotential models on precise orbits of altimetry satellites, global and regional mean sea level trends, *Adv. Space Res.*, 54(1), 92–118, doi:10.1016/j.asr.2014.03.010, 2014.
- Rudenko, S., Dettmering, D., Esselborn, S., Fagiolini, E. and Schöne, T.: Impact of Atmospheric and Oceanic De-aliasing Level-1B (AOD1B) products on precise orbits of altimetry satellites and altimetry results, *Geophys. J. Int.*, 204(3), 1695–1702, doi:10.1093/gji/ggv545, 2016.
- Rudenko, S., Neumayer, K.-H., Dettmering, D., Esselborn, S., Schöne, T. and Raimondo, J.-C.: Improvements in precise orbits of altimetry satellites and their impact on mean sea level monitoring, *IEEE Trans. Geosci. Remote Sens.*, 55(6), 3382–3395, doi:10.1109/TGRS.2017.2670061, 2017.

- 560 Savcenko, R. and Bosch, W.: EOT11a-empirical ocean tide model from multi-mission satellite altimetry, DGFI Report 89, [online] Available from: http://epic.awi.de/36001/1/DGFI_Report_89.pdf (Accessed May 2017), 2012.
- Schöne, T., Esselborn, S., Rudenko, S. and Raimondo, J.-C.: Radar Altimetry Derived Sea Level Anomalies – The Benefit of New Orbits and Harmonization, in *System Earth via Geodetic-Geophysical Space Techniques*, edited by F. M. Flechtner, T. Gruber, A. Güntner, M. Manda, M. Rothacher, T. Schöne, and J. Wickert, pp. 317–324, Springer Berlin Heidelberg, Berlin, Heidelberg. [online] Available from: <http://edoc.gfz-potsdam.de/gfz/16014> (Accessed May 2017), 2010.
- 565 Sośnica, K., Jäggi, A., Meyer, U., Thaller, D., Beutler, G., Arnold, D. and Dach, R.: Time variable Earth’s gravity field from SLR satellites, *J. Geod.*, 89, 945–960, doi:10.1007/s00190-015-0825-1, 2015.
- Soudarin, L., Capdeville, H. and Lemoine, J.-M.: Activity of the CNES/CLS Analysis Center for the IDS contribution to ITRF2014, *Adv. Space Res.*, 58(12), 2543–2560, doi:10.1016/j.asr.2016.08.006, 2016.
- 570 Watson, C. S., White, N. J., Church, J. A., King, M. A., Burgette, R. J. and Legresy, B.: Unabated global mean sea-level rise over the satellite altimeter era, *Nature Clim. Change*, 5(6), 565–568, doi:10.1038/nclimate2635, 2015.
- Willis, P., Zelensky, N. P., Ries, J., Soudarin, L., Cerri, L., Moreaux, G., Lemoine, F. G., Otten, M., Argus, D. F. and Heflin, M. B.: DPOD2008: A DORIS-Oriented Terrestrial Reference Frame for Precise Orbit Determination, in *IAG 150 Years*, pp. 175–181, Springer, Cham., doi: 10.1007/1345_2015_125, 2015.
- 575 Zelensky, N.P., Berthias, J.-P., Lemoine, F.G.: DORIS time bias estimated using Jason-1, TOPEX/Poseidon and ENVISAT orbits, *J Geod.*, 80, 497–506. doi:10.1007/s00190-006-0075-3, 2006.

580

585

590

595

Table 1. The main models used for calculation of GFZ VER11, GSFC std1504 and GRGS orbits

Parameter	GFZ REF (VER11) orbit	GSFC std1504 orbit	GRGS orbit
Terrestrial reference frame	ITRF2008 (Altamimi et al., 2011), SLRF2008 (Pavlis 2009), DPOD2008 (Willis et al., 2016)	ITRF2008, SLRF2008, DPOD2008	ITRF2008, SLRF2008, DPOD2008
Polar motion and UT1	IERS EOP 08 C04 (IAU2000A) series with IERS diurnal and semi-diurnal variations	IERS Bulletin A daily (consistent with ITRF2008), diurnal and semi-diurnal variations	IERS EOP 08 C04
Precession and nutation model	IERS Conventions (2010)	IAU2000	IERS 2010 using non-rotating origin
Station displacements due to annual geocenter variations	None	Ries (2013)	None
Non-tidal atmospheric loading effect on stations	Based on ECMWF ERA-Interim data	None	None
Ocean loading effect on stations	FES2004 (Lyard et al., 2006)	GOT4.10 (Ray, 2013)	FES2012 (Carrère et al., 2012)
Static Earth's gravity field model	EIGEN-6S4 (Förste et al., 2016) degree/order 81-90	GOCO2S (> d/o5) (Goiginger et al., 2011)	EIGEN-6S2 (Rudenko et al., 2014)
Time-variable Earth's gravity field model	EIGEN-6S4 degree 2: yearly value and drift term, time-invariant (semi-)annual variations for d/o 1-80 from 15.8.2002: yearly values, drift terms and (semi-)annual variations for d/o 1-80.	Updated harmonic piecewise fit weekly solutions (Lemoine et al., 2016) up to d/o 5	EIGEN-6S2 degree 2: yearly value and drift term, time in-variant (semi-)annual variations for d/o 2-50 from 1.1.2003: yearly values and drift terms for d/o 2-50
Solid Earth tide	IERS Conventions (2010)	IERS Conventions (2003)	IERS Conventions (2010)
Ocean tide model	EOT11a (Savchenko and Bosch, 2012) up to d/o 80	GOT4.10 up to d/o 50	FES2012 up to d/o 50
Non-tidal atmospheric and oceanic gravity	GFZ AOD1B RL05 up to d/o 100 (Dobslaw et al, 2013), including ECMWF 6-hourly fields and OMCT	ECMWF 6-hourly fields up to d/o 50	3-hourly ERA-interim and TUGO R12 up to d/o 50

Atmospheric density model	MSIS-86 (Hedin, 1987)	MSIS-86	DTM 94, with best available solar activity data
Earth radiation and albedo	Knocke et al. (1988)	Knocke et al. (1988)	Albedo and IR pressure values interpolated from ECMWF 6hr grids
Radiation pressure model	Tuned 8-panel (Cerri and Ferrage, 2016)	Tuned 8-panel	Thermo-optical coefficient from pre-launch box and wing model, with smoothed Earth shadow model
Tracking data	SLR, DORIS	SLR, DORIS	SLR, DORIS
SLR tropospheric correction model	Mendes and Pavlis (2004)	Mendes and Pavlis (2004)	Mendes and Pavlis (2004)
DORIS tropospheric correction model	Vienna Mapping Functions 1 (Boehm and Schuh, 2004)	Vienna Mapping Functions 1	GPT2/Vienna Mapping Functions 1
DORIS modelling	DORIS beacon frequency bias modelling	DORIS beacon phase center	DORIS beacon phase center
DORIS system time bias	Estimated once per arc	Estimated once per arc	None
SLR antenna reference	LRA model (note 1 below)	LRA model (note 1 below)	X: 1.2429, Y: -0.0012, Z: 0.8783 in [m]
DORIS antenna reference	pre-launch	pre-launch	pre-launch
SLR / DORIS observation weight	3 cm / 0.05 cm/s	10 cm / 0.2 cm/s	1 cm / 0.03 cm/s

Note 1: https://ilrs.cddis.eosdis.nasa.gov/missions/satellite_missions/past_missions/topx_com.html

600

605

Table 2. Average values of SLR and DORIS RMS fits, radial, cross-track and along-track two-day arc overlaps and the number of the arcs used to compute these values for the reference and five test orbits.

Orbit name	SLR RMS [cm]	DORIS RMS [cm/s]	Radial arc overlap [cm]	Cross-track arc overlap [cm]	Along-track arc overlap [cm]	Number of arcs used for SLR RMS	Number of arcs used for DORIS RMS	Number of arc overlaps used	Comment on the orbit
REF	1.96	0.04778	0.90	6.52	3.65	494	459	433	Reference
SLR	1.59		1.72	7.23	9.54	494	—	425	SLR only
DORIS	—	0.04795	0.88	6.84	2.96	—	459	392	DORIS only
TBias	1.99	0.04785	0.85	6.45	2.78	494	459	433	No DORIS system time bias estimated
ITRF14	1.97	0.04776	0.84	6.45	2.83	494	459	433	ITRF2014
Geoid	1.96	0.04775	0.83	6.43	2.80	494	459	433	EIGEN-6S2

Table 3: Median of time series of global mean height differences and RMS values at crossover points for maximum time lapses of 5 days for all orbit solutions during the period April 1993 — September 2004. The highest and lowest values of each quantity are marked bold.

	REF	GSFC	GRGS	SLR	DORIS	TBias	ITRF14	Geoid
Mean [mm]	-3.1	-1.6	-3.0	-2.7	-4.7	-3.6	-2.8	-2.1
RMS [mm]	49.8	49.5	51.3	51.2	50.7	49.8	49.8	49.7

Table 4: Global mean orbit related errors for the total signal, interannual trend variability, and decadal trend. Values were derived from the mean radial orbit differences over the oceans: REF minus SLR, REF minus DORIS, REF minus ITRF14, REF minus Geoid, REF minus GSFC, REF minus GRGS for the period April 1993 — June 2004. The corresponding values for the total sea level are tabulated under SLA.

	REF-SLR	REF-DORIS	REF-ITRF14	REF-Geoid	REF-GSFC	REF-GRGS	SLA
RMS [mm]	4.2	5.1	1.1	2.0	5.4	7.0	52.5
5-year trend [mm/year]	0.04	0.10	0.02	0.02	0.05	0.10	0.55
Decadal trend [mm/year]	0.01	0.05	0.00	0.00	0.04	0.02	2.89

625

Table 5: Regional orbit related errors for the total and seasonal signal, for interannual trend variability and decadal trend. Values were derived from the radial orbit differences: *REF minus SLR*, *REF minus DORIS*, *REF minus ITRF14*, *REF minus Geoid*, *REF minus GSFC*, *REF minus GRGS* for the periods April 1993 – June 2004.

	REF-SLR	REF-DORIS	REF-ITRF14	REF-Geoid	REF-GSFC	REF-GRGS
RMS [mm]	7.2	9.3	2.4	3.5	7.4	10.7
Annual amplitude [mm]	1.4	2.1	0.4	3.2	5.4	5.6
5-year trend [mm/year]	0.5	0.6	0.2	0.4	1.2	0.9
Decadal trend [mm/year]	0.2	0.4	0.2	0.4	1.0	0.7

630

Table 6: Global mean differences of interannual trend variability and decadal trend related to the orbit solution. Values were derived from the mean radial orbit differences over the oceans: *Geoid minus GRGS*, *REF minus DORIS*, and *REF minus TBias* for the period April 1993 – June 2004 from all passes and (in brackets) for ascending, descending passes separately.

	Geoid-GRGS	REF-DORIS	REF-TBias
Interannual trend [mm/year]	0.05 (0.36, 0.29)	0.10 (0.53, 0.37)	0.01(0.57,0.56)
Decadal trend [mm/year]	-0.01 (0.28, -0.34)	-0.05 (0.19, -0.27)	-0.01(0.08,-0.11)





Figure 1: SLR RMS fits of TOPEX/Poseidon REF, SLR, TBias, ITRF14, and Geoid orbits.

635

640



645 Figure 2. Radial arc overlaps of TOPEX/Poseidon REF, SLR, DORIS, TBias, ITRF14, and Geoid orbits.

650

655






Figure 3: Time series of the global mean RMS per cycle over the oceans of gridded radial orbit differences for *REF minus GSFC* (dark blue), and *REF minus GRGS* (red) on the top; for *REF minus DORIS* (dark blue), *REF minus SLR* (light blue), *REF minus Geoid* (green), and *REF minus ITRF14* (red) on the bottom.


660

665




670 **Figure 4: 5-year running trends for the global mean radial orbit differences over the oceans for *REF minus GSFC* (dark blue), *REF minus GRGS* (light blue), and *GRGS minus GSFC* (green) on the top; for *REF minus SLR* (dark blue), *REF minus DORIS* (light blue), *REF minus Geoid* (green), and *REF minus ITRF14* (red) on the bottom. Trend values are given for the central time of the corresponding running window.**

675



680 **Figure 5: Annual amplitude of the radial orbit differences for *REF minus GSFC*, *REF minus GRGS*, *GRGS minus GSFC*, and *REF minus Geoid*. The regions with formal errors larger than the fitted value are masked out (white). The maximum amplitude difference is given in Table 5.**


685



690 **Figure 6: RMS of 5-year running trend differences of the radial orbit components for *REF minus GSFC*, *REF minus GRGS*, *REF minus SLR*, *REF minus DORIS*, *REF minus ITRF14*, and *REF minus Geoid* for the period April 1993 – June 2004. The global mean RMS of the differences over the ocean is given in Table 4.**

695


700



705 **Figure 7:** Trend differences of radial orbit components for *REF minus GSFC*, *REF minus GRGS*, *REF minus SLR*, *REF minus DORIS*, *REF minus ITRF14*, and *REF minus Geoid* for the period April 1993–June 2004. Regions with formal errors larger than the fitted value are masked out (white). The global mean trend difference over the ocean is given in Table 4.

710

715



720 **Figure 8:** Trend differences of radial orbit components for ascending (left) and descending (right) passes for *Geoid minus GRGS*, *REF minus TBias*, and *REF minus DORIS* for the period April 1993 – June 2004. Regions with formal errors larger than the fitted value are masked out (white). The global mean trend difference over the ocean is given in Table 6.

725




Figure 9: RMS of sea level, annual amplitude, RMS of interannual (5 years) running trend, and decadal trends from TOPEX altimeter data for the period February 1993– October 2005. Colour coded are sea level values for which the local orbit errors (estimated from *GFZ minus GRGS*) reach more than 10 % of the local sea level values. All other regions are masked out (white).








Figure S1: DORIS system time bias of TOPEX/Poseidon REF, DORIS, ITRF14, and Geoid orbits.



735 **Figure S2: Power spectra of the global mean radial orbit differences for *REF minus GSFC*, *REF minus GRGS*, *GRGS minus GSFC*, *REF minus SLR*, *REF minus DORIS*, *REF minus ITRF14*, *REF minus Geoid*, and *REF minus TBias*. Vertical dashed lines mark periods of 59, 85, 170 days, 1 and 5 years.**



740 **Figure S3: Global mean radial orbit differences per cycle for *Geoid minus GRGS*, *REF minus DORIS*, and *REF minus Tbias* separately for ascending (blue, cyan) and descending (yellow, red) tracks and 1-year box-car filtered. The cross-correlation coefficient between the ascending and descending passes for the original and the filtered series is given at the lower part of each graph.**