



Consideration of various aspects in a drift study of MH370 debris

Oleksandr Nesterov¹

¹Independent Consultant (Physical Oceanography & Coastal Engineering)

Correspondence to: Oleksandr Nesterov (oleksandr.nesterov@gmail.com)

Abstract. On March 7, 2014, a Boeing 777-200ER aircraft operated by Malaysian Airlines on the route MH370 from Kuala Lumpur to Beijing abruptly ceased all communications and disappeared with 239 people aboard, leaving a mystery about its fate. The subsequent analysis of so-called satellite 'handshakes' supplemented by military radar tracking has suggested that the aircraft ended up in the southern Indian Ocean. Eventual recovery of a number of fragments washed ashore in several countries has confirmed its crash. A number of drift studies were undertaken to assist in locating the crash site, mostly focusing either on the spatial distribution of the washed ashore debris or efficacy of the aerial search operation. A recent biochemical analysis of the barnacles attached to the flaperon (the first fragment found in La Réunion) has indicated that their growth likely began in the water of 24°C, then the temperature dropped to 18°C, and then it rose up again to 25°C. An attempt was made in the present study to take into consideration all these aspects. The analysis was conducted by the means of numerical screening of 40 hypothetical locations of the crash site along the so-called 7th arc. Obtained results indicate the likelihood of the crash site to be located between 25.5°S and 30.5°S latitudes, with the segment from 28°S to 30°S being the most promising.

1 Introduction

On March 7, 2014, approximately 40 minutes after the takeoff, a Boeing-777 aircraft (registration 9M-MRO) operated by Malaysian Airlines (MAS) as MH370 on the route from Kuala Lumpur (Malaysia) to Beijing (China), abruptly ceased all data and voice communications, and disappeared with 239 people aboard, leaving investigators clueless about a possible cause. The subsequent analysis of so-called satellite 'handshakes' (Ashton et al., 2015) supplemented by the primary radar tracking has suggested that the plane turned back, crossed the Malay Peninsula along Malay-Thai border, then flew toward Nicobar Islands in the Strait of Malacca, where it finally turned into the Indian Ocean, as detailed by the Australian Transport Safety Bureau (ATSB, 2014a, b) and the Ministry of Transport Malaysia (MTM, 2017). Eventual recovery of a number of 9M-MRO fragments, which were washed ashore in several countries, has confirmed the crash of the aircraft in the Indian Ocean. A total of 27 suspected and confirmed fragments were found according to the report published by the Malaysian Safety Investigation Team for MH370 (MSIT, 2017) on March 27, 2017: 1 in La Réunion, 2 in Mauritius, 1 in Rodrigues, 6 in Mozambique, 5 in South Africa, 1 in Tanzania, and 11 in Madagascar, with the latest find in January 2017.

Shortly after the disappearance, the Australian Government, whose geographical responsibility for rescue and recovery covers respective region of the Indian Ocean, has established the Joint Agency Coordination Centre (JAAC, 2014) to assist in the search operation. The Australian Maritime Safety Authority (AMSA, 2014), JAAC and ATSB have conducted an extensive



aerial search operation, which lasted from March 18 to April 27, but failed to locate any debris related to MH370. Although some objects were spotted on March 18, subsequent attempts to recover them were unsuccessful. After expiration of the underwater locator beacon, a device emitting acoustic signal to facilitate underwater search, the Australian Government has commissioned an engineering company Fugro N.V., which specializes on marine and geotechnical surveys, to conduct a deep-
5 water high-resolution sonar survey of the seabed. The search domain assigned to Fugro N.V. was a band-shaped area in the South Indian Ocean along the so-called 7th arc from approximately 36°S to 39.5°S latitudes, defined by the ATSB (2015), and then later refined by the Australian Defense Science and Technology Group (Davey et al., 2016). The 7th arc is a geometric curve on the Earth surface, all points of which are equidistant from the satellite, through which the last 'handshake' was transmitted (Ashton et al., 2015). The actual 7th arc may slightly differ from the nominal arc due to the uncertainty in the
10 altitude of the aircraft, as well as truncation and measurement errors in the data (e.g., ATSB, 2015; Davey et al., 2016). Despite such an unprecedented effort, the underwater search was unsuccessful, and it was finally called off in January 2017 (MTM, 2017).

A number of drift studies were undertaken since then to assist in locating the crash site. Prior to the discovery of the flaperon in La Réunion on August 29, 2015, the studies focused on the analysis of the efficacy of the aerial search, such as García-
15 Garrido et al. (2015). Later, after the flaperon and other 9M-MRO fragments were found, the mainstream approach shifted to the analysis of the probabilities of debris to reach specific locations by known dates starting from various origins along the 7th arc: the series of numerical and experimental studies undertaken by Griffin et al. (2017) at the Commonwealth Scientific and Industrial Research Organisation (CSIRO); the screening of 25 hypothetical locations along the 7th arc by Pattiaratchi and Wijeratne (2016) at the University of Western Australia; the numerical modelling conducted by Ormond and Baart (2015) at
20 Deltares; the study conducted by Maximenko et al. (2015) at the International Pacific Research Center; the study by Jansen et al. (2016) at the Euro-Mediterranean Center on Climate Change. An alternative approach was based on the reverse drift modelling: the studies conducted by the French Government meteorological agency Météo France (Daniel, 2016) and GEOMAR Helmholtz Centre for Ocean Research Kiel (Durgadoo and Biastoch, 2015). The latter, however, did not account for wind forcing, what presumably explains the large difference in its conclusions compared to other studies.

25 There is an ongoing disagreement between conclusions of these studies with regard to the most likely origin of the debris at the 7th arc: the latest study conducted by CSIRO (Griffin et al., 2017) recommends a new search area at around 35°S, backed by the earlier IPRC study (34°S to 37°S). In contrast, Pattiaratchi and Wijeratne (2016) have suggested the crash site to be more likely located between 28.3°S and 33.2°S, narrowing down earlier Jansen et al. (2016) estimates (between 28° and 35°S), being also consistent with Ormond and Baart (2015). Assuming zero drift angle of the flaperon, Daniel (2016) favors the location
30 north of 25°S, but south of 35°S if the drift angle was set to 18° to the left with regard to wind at the leeway factor of 3.29% - both the parameters experimentally established by the Direction générale de l'Armement (DGA). The latest CSIRO study (Griffin et al., 2017) disagrees with these parameters, but confirms observed non-zero drift angles between 0° to 30°, explaining this effect by the longwise asymmetry of the flaperon.

An important feature of the fragment found in La Réunion are barnacles attached to it. Although it was not possible to
35 establish their age, according to De Deckker (2017), who conducted biochemical analysis of the barnacles at the Australian



National University, the start of their growth was in the water of approximately 24°C, and then for some time the temperature ranged between 20°C and 18°C, and then it went up again to around 25°C. This additional information has not been previously considered in the drift studies.

Consequently, this study comprises the three major elements to assess the most likely origin of the debris:

- 5 (1). Efficacy of the aerial search campaign.
- (2). Ambient water temperatures at the flaperon.
- (3). Spatial distribution of washed ashore debris.

2 Modelling

A total of 40 hypothetical locations of the debris origin along the 7th arc were screened against the three selection criterion
 10 by the means of forward particle tracking technique. The deterministic forcing of each particle in an ensemble was governed by the balance of water and air drag forces, magnitudes of which were assumed to be proportional to the squared relative (with respect to the particle) speeds of the ambient water and air respectively. Surface current velocities were sourced from the Hybrid Coordinate Ocean Model (HYCOM), wind data was sourced from the Global Data Assimilation System (GDAS), as detailed in Section 2.2.2. The stochastic component was modelled using the random walking technique (e.g., Al Rabeh et al.,
 15 2000; DHI, 2009; Jansen et al., 2016). Numerical integration was performed in the geocentric Cartesian coordinate system.

All the particles in an ensemble were 'released' from a single starting point at the 7th arc. Four models with regard to the leeway and drift angle properties of particles were considered. After individual particle tracks were obtained, they were supplemented by respective sea surface temperatures (SST) extracted from the publicly available archives. A subsequent analysis was undertaken in a statistical manner to: (1) estimate the maximum ensemble coverages during the aerial search; (2) assess
 20 percentage of particles in ensembles, which could reached La Réunion by August 29, 2015, and be subjected to the temperature variation matching the results of the barnacle analysis; and (3) compare spatial distribution of particles washed ashore with the locations, where MH370 debris were found.

2.1 Model description

2.1.1 Assumptions

25 In the frame of this model it was assumed that a particle was subjected to the drag forces induced by water and wind. Particles were assumed to be non-inertial. Impacts of the Coriolis force, Stokes' drift, waves, decay and sinking were neglected (e.g., Kraus, 1972; Al Rabeh et al., 2000). In the local coordinate system (x, y) , where the axes x and y correspond to the local west-to-east and south-to-north directions respectively (Figure 1), the location (x_i, y_i) of the i -th particle was described by Langevin's equation (DHI, 2009):

$$30 \quad \frac{dx_i}{dt} = u_i + D(t, x_i, y_i)\zeta, \quad \frac{dy_i}{dt} = v_i + D(t, x_i, y_i)\xi, \quad (1)$$




Figure 1. Local and geocentric Cartesian coordinate systems.

where (u_i, v_i) is the deterministic velocity of the i -th particle in the local coordinate system, D is the turbulent diffusion term, and ζ, ξ are the random numbers, as detailed in Sections 2.1.3 through 2.1.5.

2.1.2 Coordinate systems

Velocity of a particle and random walk are formulated in the local coordinate system, where the axis z is normal to the Earth surface. On the other hand, a large extent of the study domain dictates the necessity to properly take into consideration the Earth curvature. It is convenient to perform numerical integration of the governing equations in the geocentric Cartesian coordinate system (X, Y, Z) depicted in Figure 1, where the Earth surface is approximated by WGS'84 ellipsoid with the polar and equatorial axes radii $R_p = 6356752$ m and $R_e = 6378137$ m respectively. Transformation of coordinates and velocity vectors are required for this.

The Cartesian coordinates X , Y , and Z of a point on the surface of the ellipsoid described by the longitude ψ and geocentric latitude φ can be formulated as:

$$X = R \cos \varphi \cos \psi, \quad Y = R \cos \varphi \sin \psi, \quad Z = R \sin \varphi, \quad (2)$$

where $R = \frac{R_e R_p}{\sqrt{(R_p \cos \varphi)^2 + (R_e \sin \varphi)^2}}$ is the distance between this point and the center of the ellipsoid, what follows from the ellipse equation $\frac{(R \cos \varphi)^2}{R_e^2} + \frac{(R \sin \varphi)^2}{R_p^2} = 1$.

It should be noted that the backward transformation is required to extract and interpolate surface current and wind data at a location. Respective trigonometric transformations involve solving a 4th-degree polynomial equation.

The unit vectors \vec{L} , \vec{M} , and \vec{N} , which define directions of the axes x , y , and z of the local coordinate system (Figure 1) are introduced to obtain velocity components in the geocentric system. The outward unit vector $\vec{N} = \{N_X, N_Y, N_Z\}$ normal to the surface of the ellipsoid is formulated according to Korn and Korn (1968):



$$\vec{N} = \frac{\left\{ \frac{\partial X}{\partial \psi}, \frac{\partial Y}{\partial \psi}, \frac{\partial Z}{\partial \psi} \right\} \times \left\{ \frac{\partial X}{\partial \varphi}, \frac{\partial Y}{\partial \varphi}, \frac{\partial Z}{\partial \varphi} \right\}}{\left| \left\{ \frac{\partial X}{\partial \psi}, \frac{\partial Y}{\partial \psi}, \frac{\partial Z}{\partial \psi} \right\} \times \left\{ \frac{\partial X}{\partial \varphi}, \frac{\partial Y}{\partial \varphi}, \frac{\partial Z}{\partial \varphi} \right\} \right|} \Rightarrow$$

$$N_X = \frac{1}{\sqrt{1+\mu^2}} (\cos \varphi - \mu \sin \varphi) \cos \psi,$$

$$N_Y = \frac{1}{\sqrt{1+\mu^2}} (\cos \varphi - \mu \sin \varphi) \sin \psi,$$

$$N_Z = \frac{1}{\sqrt{1+\mu^2}} (\sin \varphi + \mu \cos \varphi), \quad (3)$$

$$5 \quad \text{where } \mu = R_e^2 \frac{R_e^2 - R_p^2}{R_e^2 R_p^2} \cos \varphi \sin \varphi.$$

The direction of the axis x is defined by the unit vector \vec{L} :

$$\vec{L} = \{L_X, L_Y, L_Z\} = \{-\sin \psi, \cos \psi, 0\}. \quad (4)$$

Eq. (3) and Eq. (4) allow for expressing the unit vector $\vec{M} = \{M_X, M_Y, M_Z\}$ collinear to the axis y in the form of the vector product $\vec{M} = \vec{N} \times \vec{L}$, so that its components are:

$$10 \quad M_X = -\frac{1}{\sqrt{1+\mu^2}} (\sin \varphi + \mu \cos \varphi) \cos \psi,$$

$$M_Y = -\frac{1}{\sqrt{1+\mu^2}} (\sin \varphi + \mu \cos \varphi) \sin \psi,$$

$$M_Z = \frac{1}{\sqrt{1+\mu^2}} (\cos \varphi - \mu \sin \varphi). \quad (5)$$

Therefore, a velocity vector, the components of which are $\{u, v, 0\}$ in the local system, has the following components in the geocentric Cartesian system:

$$15 \quad U = -u \sin \psi - v \frac{1}{\sqrt{1+\mu^2}} (\sin \varphi + \mu \cos \varphi) \cos \psi,$$

$$V = u \cos \psi - v \frac{1}{\sqrt{1+\mu^2}} (\sin \varphi + \mu \cos \varphi) \sin \psi,$$

$$W = v \frac{1}{\sqrt{1+\mu^2}} (\cos \varphi - \mu \sin \varphi). \quad (6)$$

Hence, Langevin's equation (1) can be formulated in the geocentric Cartesian system as follows:




Figure 2. Schematic representation of a thin floating object.

$$\begin{cases} \frac{dX_i}{dt} = U(u_i, v_i, \psi_i, \varphi_i) + D(L_X \zeta + M_X \xi), \\ \frac{dY_i}{dt} = V(u_i, v_i, \psi_i, \varphi_i) + D(L_Y \zeta + M_Y \xi), \\ \frac{dZ_i}{dt} = W(u_i, v_i, \psi_i, \varphi_i) + D(L_Z \zeta + M_Z \xi), \end{cases} \quad (7)$$

where the relations between the longitude ψ_i and geocentric latitude φ_i of particle's location and its geocentric Cartesian coordinates X_i, Y_i, Z_i is given by Eq. (2); the transformation of the velocity components is given by Eq. (6); and the local velocity components u_i , and v_i are defined from the balance of the deterministic forces.

5 2.1.3 Deterministic terms

Mathematical description of the dynamics of a floating object is not a trivial problem due to the variety of phenomenon and processes in a near-surface layer, such as surface waves, Stokes drift, flow-object interaction, variable water and air velocities, buoyancy, stratification, etc. (e.g., Kraus, 1972).

When relative (with respect to a floating object) velocities of the water and air induce drag forces collinear to these velocities, the steady-state balance can be formulated as follows:

$$\begin{aligned} \frac{1}{2} C_{Dw} S_w \rho_w |(\vec{u}_i - \vec{u}_w(x_i, y_i, t))| (\vec{u}_i - \vec{u}_w(x_i, y_i, t)) = \\ - \frac{1}{2} C_{Da} S_a \rho_a |(\vec{u}_i - \vec{u}_a(x_i, y_i, t))| (\vec{u}_i - \vec{u}_a(x_i, y_i, t)), \end{aligned} \quad (8)$$

where C_{Dw} , S_w are the water drag coefficient and corresponding area of the submerged part of the object, C_{Da} , S_a are the air drag coefficient and corresponding area of the part exposed to the air, ρ_w and ρ_a are the water and air densities respectively, $\vec{u}_w(x_i, y_i, t)$ and $\vec{u}_a(x_i, y_i, t)$ are the water and air velocities at the location of the i^{th} particle.

It is easy to see that the solution of Eq. (8) with respect to the velocity of the i -th particle can be written in the form:

$$\vec{u}_i = (1 - \alpha_i) \vec{u}_w(x_i, y_i, t) + \alpha_i \vec{u}_a(x_i, y_i, t), \quad (9)$$




Figure 3. Assumed distribution of leeway factors in an ensemble.

where the scalar α_i (leeway factor) is:

$$\alpha_i = \frac{\sqrt{C_{Da}S_a\rho_a}}{\sqrt{C_{Da}S_a\rho_a} + \sqrt{C_{Ds}S_w\rho_w}}. \quad (10)$$

For a thin horizontally-floating object, which has equal areas of the surfaces exposed to the seawater and air ($S_a = S_w$), and which is characterized by equal drag coefficients $C_{Da} = C_{Dw}$, the leeway factor is $\alpha \approx 3.33\%$. This theoretical value is in a good agreement with the experimental data for the flaperon (3.29%) estimated by the DGA in the hydrodynamic engineering test facility center in Toulouse (Daniel, 2016).

In those scenarios, where the force induced by relative wind was not collinear to its direction, a modified formulation was used:

$$\vec{u}_i = (1 - \alpha_i) \vec{u}_w + \alpha_i \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \vec{u}_a, \quad (11)$$

where θ is the drift angle, positive counter-clockwise.

2.1.4 Random leeway factor model

In contrast to earlier studies (e.g., Daniel, 2016; Pattiaratchi and Wijeratne, 2016; Griffin et al., 2017), an attempt was made in this work to take into consideration the variety of leeway factors, which describe random shapes and flotation characteristics of individual fragments generated by the crash. For this purpose, in one class of the carried out simulations particles were described by random leeway factors. It was assumed that particles in ensembles were representing thin objects of irregular shapes floating in tilted orientation, partially submerged (Figure 2). Such an object is subjected to the dynamic pressure and shear stress forces due to the action of the water and air; its steady-state orientation depends on its buoyancy characteristics and the moments of forces around its principal axes of inertia. According to the classic theory of the turbulent boundary layer (e.g., Gandin et al., 1955), a vertical velocity profile exhibits logarithmic dependence on height, being proportional to the wind speed at the height of the boundary layer (typically 10 m), whereas the turbulent shear stress remains constant within this layer,



Table 1. Longitudes and latitudes of the origins preselected for screening

No.	Lon.,°E	Lat.,°S	No.	Lon.,°E	Lat.,°S	No.	Lon.,°E	Lat.,°S	No.	Lon.,°E	Lat.,°S
1	85.72	39.17	11	92.29	35.29	21	98.07	30.03	31	102.64	23.69
2	86.43	38.83	12	92.91	34.82	22	98.59	29.44	32	103.02	23.00
3	87.13	38.48	13	93.53	34.34	23	99.09	28.84	33	103.39	22.31
4	87.82	38.11	14	94.14	33.84	24	99.58	28.23	34	103.75	21.62
5	88.51	37.73	15	94.73	33.33	25	100.05	27.61	35	104.18	20.74
6	89.01	37.44	16	95.32	32.81	26	100.52	26.98	36	104.50	20.03
7	89.69	37.03	17	95.89	32.28	27	100.97	26.34	37	104.82	19.31
8	90.35	36.62	18	96.46	31.74	28	101.40	25.69	38	105.20	18.41
9	91.00	36.19	19	97.01	31.18	29	101.83	25.03	39	105.48	17.68
10	91.65	35.74	20	97.55	30.61	30	102.24	24.36	40	105.75	16.94

being proportional to the squared wind speed at the same height. Hence, for the sake of simplification, it was assumed that the total force acting on a floating object was proportional to the squared relative velocities of the air and water, with the drag coefficients for the air and water being equal ($C_{Da} = C_{Ds}$). Then, according to Eq. (10), only the knowledge of the ratio of respective areas $k = S_a/S_w$ is required to estimate the leeway factor of this object:

$$\alpha_i = \frac{\sqrt{\rho_a/\rho_w} \sqrt{k_i}}{1 + \sqrt{\rho_a/\rho_w} \sqrt{k_i}}. \quad (12)$$

In the frame of this study it was assumed that dimensions of individual objects are log-normally distributed. Furthermore, it was assumed that the principal axis of inertia of the object splits it into two parts, areas of which are also log-normally distributed, so that $k_i = \frac{\gamma_i S_{i,1}}{S_{i,2} + (1.0 - \gamma_i) S_{i,1}}$, where $\{\ln S_{i,1}, \ln S_{i,2}\} \in \mathcal{N}(\mu, \sigma^2)$, and $\gamma \in [0, 1]$ is an independent random parameter to account for the draft of the object (0-fully submerged; 1 - the center of gravity is at the water surface). Hence the logarithm of the ratio $\ln(S_1/S_2) = \ln S_1 - \ln S_2 \in \mathcal{N}(0, 2\sigma^2)$ is also normally distributed, with the mean of zero. Here, the property was used that the sum of two independent normally distributed random variables is also normally distributed, with its mean being the sum of the means, and its variance being the sum of the two variances (Eisenberg and Sullivan, 2008). Hence, the ratio S_1/S_2 is log-normally distributed. The modelling was performed assuming $\ln(S_1/S_2) \in \mathcal{N}(0, 1)$. The resulting distribution of the leeway factors of particles in an ensemble in this class of simulations (hereafter referred as the "random leeway" model) is depicted in Figure 3.

2.1.5 Numerical realization and random walk

The whole integration interval was split into the time steps of $\Delta t = 15$ minutes. Similarly to DHI (2009) particle tracking model, integration of the system of equations (7) for each particle over the time step Δt was comprised of the two stages, namely deterministic and stochastic:



$$\begin{aligned}\vec{\mathbf{X}}_i(t + \Delta t) &= \vec{\mathbf{X}}_i(t) + \int_t^{t+\Delta t} \vec{\mathbf{U}}(u_i, v_i, \psi_i, \varphi_i) dt, \\ \vec{\mathbf{X}}_i(t + \Delta t) &= \vec{\mathbf{X}}_i(t + \Delta t) + \vec{\delta}_i(\dots),\end{aligned}$$

where $\vec{\mathbf{X}}_i = \{X_i, Y_i, Z_i\}$ is the location of the i -th particle; $\vec{\mathbf{X}}_i$ is the intermediate location prior to the superposition of the random displacement $\vec{\delta}_i(\dots)$; $\vec{\mathbf{U}}(u_i, v_i, \psi_i, \varphi_i) = \{U(\dots), V(\dots), W(\dots)\}$ is the velocity of the particle. Unlike DHI (2009) model, which uses the 1-st order discretization method to integrate deterministic terms, the fifth- and sixth-order Runge-Kutta method was used in this work utilizing FORTRAN libraries to ensure particles remain on ellipsoid's surface with sufficient accuracy. During the integration, input current and wind data were bi-linearly interpolated in space, and linearly in time.

The vector $\vec{\delta}_i(\dots)$ corresponds to a numerical solution for the diffusivity term of the Langevin equation (1). It was treated as a random displacement in the xy -plane locally tangential to ellipsoid's surface. Its components δ_x and δ_y in the local coordinate system are the random values from the trimmed two-dimensional Gaussian distribution \mathcal{N}_2 :

$$\delta_x = \sigma_L \zeta, \quad \delta_y = \sigma_L \xi, \quad (13)$$

where $\{\zeta, \xi\} \in \mathcal{N}_2(0, 1)$ are the random numbers, and the standard deviation of the turbulent dispersion $\sigma_L = \sqrt{2D\Delta t}$ is assumed to be a function of the horizontal eddy diffusivity coefficient $D = D(t, \psi_i, \varphi_i)$. Such a relation between D and σ_L was first established by A. Einstein in 1905, who studied diffusion associated with Brownian motion, and since then it was adopted in the variety of random walk models (e.g., DHI, 2009; Jansen et al., 2016). In this study trimming was imposed to discard values δ_x and δ_y that resulted in displacements, which exceeded 10 km distance over the time step Δt . If this criteria was violated, the next pair of random values δ_x and δ_y was computed. The distance of 10 km was selected as the representative resolution of the ocean circulation model HYCOM, used as a source of the surface current velocities (see Section 2.2.2).

In contrast to Jansen et al. (2016), who applied the constant eddy diffusivity coefficient $D = 2 \text{ m}^2/\text{s}$, in this work D was computed according to the well-known Smagorinsky (1963) parameterization, applied in various ocean circulation models (e.g., Blumberg and Mellor, 1987; DHI, 2009):

$$D = \kappa \Delta x \Delta y \sqrt{\left(\frac{\partial u}{\partial x}\right)^2 + \frac{1}{2} \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x}\right)^2 + \left(\frac{\partial v}{\partial y}\right)^2}, \quad (14)$$

where $\kappa = 0.1$ is the constant coefficient (a typical range is from 0.1 to 0.2 according to Blumberg and Mellor (1987)), Δx and Δy are the horizontal dimensions of the numerical grid cell, applied for the discretization of the velocity field. A discrete approximation of Eq. (14) was used to estimate the eddy diffusivity coefficient D , with subsequent bi-linear interpolation in space and linear interpolation in time at the actual locations of particles.

In the geocentric Cartesian coordinate system the random displacement translates into the 3D vector $\vec{\delta}_i = \{\delta X_i, \delta Y_i, \delta Z_i\}$, components of which are:



$$\begin{aligned}\delta X_i &= L_X \delta_x + M_X \delta_y, \\ \delta Y_i &= L_Y \delta_x + M_Y \delta_y, \\ \delta Z_i &= L_Z \delta_x + M_Z \delta_y.\end{aligned}\tag{15}$$

Such a displacement in the tangential plane causes a particle to move away from the surface of the ellipsoid. However, due to the imposed limitation on the distance, the elevation of the particle does not exceed 8 m after superposition of the random walk. Thus, particle elevations were forced to zeros after applying random walk procedure at each integration time step, while preserving longitudes and latitudes.

Box-Muller (1958) transform was used to obtain a pair of pseudo-random numbers $\{\zeta, \xi\} \in \mathcal{N}_2(0, 1)$:

$$\zeta = \sqrt{-2 \ln \tau} \cos(2\pi\omega), \quad \xi = \sqrt{-2 \ln \tau} \sin(2\pi\omega),\tag{16}$$

where τ and ω are the two pseudo-random numbers from the interval $(0, 1]$. To obtain τ and ω , the use was made of the generator developed by Marsaglia and Zaman (1987), claimed to have the period of 2^{144} .

In the proximity to a shore, flow velocity was linearly interpolated between zero and velocity in the adjacent cell of the numerical grid of HYCOM. If a particle moved onshore as a result of wind action or random walk, all the subsequent forcing acting on such a particle was nullified, so that it remained at the location where it beached. No specific properties of the shores (e.g. type, slope), or contributing factors such as waves, tides, or storm surges, were considered.

It is worth of noting that the transformations between the longitude, latitude and geocentric Cartesian coordinates were performed using extended precision accuracy (80 bits). Conducted tests have shown that the maximum errors in the elevation of a particle arising in a result of a single forward-backward conversion of the coordinates were of order 0.5 mm using the extended precision compared to 0.5 m using the double precision arithmetics.

2.2 Model setup

Four scenarios with respect to the leeway factor and drift angle of particles in an ensemble were considered:

- (1). Leeway factor of 3.29%; zero drift angle;
- (2). Leeway factor of 3.29%; drift angle of 18° to the left;
- (3). Leeway factor of 2.76%; drift angle of 32° to the left;
- (4). Random leeway factor; zero drift angle.

All the ensembles released at various locations were identical with respect to the properties of particles in these ensembles. Each ensemble comprised 50,000 particles, same as in Pattiaratchi and Wijeratne (2016) study. The second and third models focused specifically on the flaperon path: respective settings corresponded to the flotation characteristics established by the




Figure 4. Locations of the selected hypothetical origins on the 7th arc, and snapshots of the sea surface temperatures.

DGA (Daniel, 2016). The fourth model aimed to achieve more realistic representation of the flotation characteristics of the debris generated by the crash by assigning random leeway factors to the particles of an ensemble.

2.2.1 Screened locations

The extent of the study domain covered zone from 20°E to 140°E, 55°S to 15°N. Integration was performed from March 8, 2014 to December 31, 2016 inclusive. The locations of the 40 hypothetical debris origins are summarized in Table 1 and indicated in Figure 4. The coordinates of these locations were estimated from the burst time offset (ATSB, 2015) of the last 'handshake' 00:19, assuming that the aircraft was at the altitude of 10 km.

2.2.2 Model forcing and SST data

The use of the following datasets was made in this study for the model forcing and temperature analysis:

- **Surface currents** were extracted from the Hybrid Coordinate Ocean Model, a data-assimilative isopycnal-sigma-pressure coordinate ocean circulation model (Chassignet et al., 2007). Spatial resolution: 0.08°x 0.08°; temporal resolution: daily. The HYCOM consortium is a multi-institutional effort sponsored by the National Ocean Partnership Program (<https://hycom.org/>); data are available at: ftp://ftp.hycom.org/datasets/global/GLBa0.08_rect/data/.




Figure 5. Snapshots of the particle ensembles originated from the locations [1](#), [4](#), [11](#), [21](#), [23](#), [27](#), [28](#), [36](#), [40](#), and progressing search maps (by Mar 28 and Apr 05). Sources of maps and photograph: JAAC (2014), AMSA (2014).

- **Wind velocities** were extracted from the Global Data Assimilation System, provided by the Air Resources Laboratory (ARL) of the U.S. National Oceanic and Atmospheric Administration (NOAA). Spatial resolution: $1^\circ \times 1^\circ$; temporal resolution: 3 hours. Further details are available at <http://www.ready.noaa.gov/gdas1.php>; archived data in a proprietary format are available at: <ftp://arlftp.arlhq.noaa.gov/pub/archives/gdas1>.

- 5
- **SST data** were sourced from a Group for High Resolution Sea Surface Temperature (GHR SST) Level 4 MUR Global Foundation Sea Surface Temperature Analysis, a product developed by the Jet Propulsion Laboratory under NASA



MEaSURES program (JPL, 2015). Spatial resolution: $0.011^\circ \times 0.011^\circ$; temporal resolution: daily. Detailed information on this data is available at: <https://mur.jpl.nasa.gov/> and <https://podaac.jpl.nasa.gov/dataset/JPL-L4UHfnd-GLOB-MUR>.

3 Results

3.1 Efficacy of the aerial search

5 An extensive aerial search for MH370 debris, which lasted from March 18 to April 27, 2014, (e.g., JAAC, 2014; AMSA, 2014), failed to find any debris relevant to MH370. Although some suspected objects were observed on March 28 (AMSA, 2014), such as a rectangular object photographed from the Royal New Zealand Airforce (RNAZF) Orion P3K survey aircraft, attempts to recover them were unsuccessful.

To understand reasons contributing to this failure, the efficacy of the aerial search was analyzed in terms of the coverages
10 of the ensembles of particles on each day of the aerial search. Respective areas were digitized from the progressing maps published at AMSA (2014) and JAAC (2014) portals.

Snapshots of the modelled particle locations in the ensembles originated from the nine selected locations along the 7th arc on March 28 and April 5 are depicted in Figure 5 for the random and constant 3.29% leeway factor models. Snapshots on several other days can be found in Figure S1 in the Supplement. As seen, the leeway factors had significant influence on the dispersion,
15 which was notably more intense for the ensembles that comprised particles of various leeway factors. Furthermore, differences in the leeway factors could presumably explain the failed attempts to track suspected objects with the help of deployed buoys. It is worth of noting that the aerial survey would have been rather inefficient for the objects characterized by the leeway factor of 3.29% (such as the flaperon), which originated from the locations around 30°S or from the segment from 25.5°S to 28°S of the 7th arc.

20 As the value of a single-day peak coverage can be downplayed by other factors affecting detection capability, such as poor weather conditions, the five largest estimated daily coverages of particle ensembles are shown in Figure 6 for the random and constant 3.29% leeway factor models versus origin's latitudes along the 7th arc. Respective dates of the maximum coverage and cumulative coverages over the entire duration of the search campaign are summarized in Table 2.

As seen, if the crash site was located between 30.5°S and 34.5°S , or between 20°S and 25°S , debris would have been
25 relatively well covered, thus maximizing chances of their detection. Interestingly, the coverage of MH370 debris would have been notably lower if the crash site was located between 25.5°S and 27.5°S . In particular, for the origins No. 27 and 28, the maximum coverages of approximately 10% occurred more than 6 weeks after the crash, which is a rather low figure bearing in mind that decay and sinking processes were not modelled. A relatively poor coverage is also noted for the origins No. 21 and 23.

30 Interestingly, a rectangular object photographed from RNZAF Orion P3K on March 28 was in the area, where some debris originated from the location No. 21 (98.07°E , 30.03°S) could be expected according to the modelling results (Figure 5). In addition, the timing of the peak coverage (March 28 - 31; see Table 2) for the debris originated from this and nearby locations is consistent with the observation date.




Figure 6. Efficacy of the aerial search 18 March - 27 April: the five largest estimated daily coverages of the debris field for each origin.

Table 2. Computed maximum daily coverage (%), respective date (2014), and cumulative coverage (%) for the random leeway model.

No.	Max. daily	Cumu-lative	Date	No.	Max. daily	Cumu-lative	Date	No.	Max. daily	Cumu-lative	Date	No.	Max. daily	Cumu-lative	Date
1	90.7	91.9	Mar 19	11	24.9	25.1	Apr 01	21	30.4	114.1	Mar 31	31	25.8	132.9	Apr 04
2	19.8	37.2	Mar 19	12	10.7	16.9	Apr 01	22	45.1	279.0	Mar 28	32	66.7	419.3	Apr 04
3	18.6	18.6	Mar 18	13	66.0	107.0	Apr 01	23	28.6	116.1	Apr 02	33	53.4	137.4	Apr 05
4	4.3	4.3	Mar 19	14	36.8	62.6	Apr 01	24	43.1	192.3	Mar 28	34	63.1	254.8	Apr 05
5	13.6	13.6	Mar 19	15	71.1	142.0	Mar 31	25	34.9	199.1	Mar 28	35	62.4	234.6	Apr 05
6	0.4	0.4	Mar 19	16	67.3	160.7	Mar 31	26	19.5	136.2	Mar 29	36	42.3	138.9	Apr 03
7	<0.1	<0.1	Mar 19	17	65.3	244.3	Apr 01	27	10.5	68.8	Apr 20	37	0.3	0.6	Apr 05
8	0	0	-	18	61.7	295.1	Mar 29	28	12.5	46.3	Apr 18	38	0.2	0.6	Apr 05
9	<0.1	<0.1	Apr 01	19	50.0	256.8	Mar 29	29	46.4	126.3	Apr 13	39	0	0	-
10	0.8	0.8	Apr 01	20	68.3	245.1	Mar 29	30	19.0	100.8	Apr 04	40	0	0	-

3.2 SST along the path of the flaperon

One of the goals of this study is to address a question whether the ambient water temperatures along modelled particle tracks could match the temperature variation derived from the biochemical analysis of the barnacles attached to the flaperon (De Deckker, 2017), and if yes, whether this information could help to further refine the search area. Therefore, those particles

5 were of interest, which satisfied the two following conditions:

1. A particle must have arrived to the Sain-Andre beach, a place where the flaperon was found (assumed coordinates 55.67°E, 20.93°S), by July 29, 2015;
2. The ambient SST at particle's location should have first exceeded 24°C, then dropped down below 18°C, and then risen up again to 25°C or higher.




Figure 7. Percentages of particles satisfying distance and temperature criterion with the score $S > 0.01$.




Figure 8. Sample tracks and temperatures for the 6 particles of the ensemble released at 99.09°E , 28.84°S (leeway 3.29%; drift angle 18°).

Due to the inherent uncertainty in the temperature estimations based on the barnacle analysis, a score-based function S was introduced to identify those particles, which approximately satisfied the two aforementioned conditions:



$$S_i = S_{i,d} S_{i,\theta}. \quad (17)$$

Here the term $S_{i,d} = \exp(-0.5 d_i^2 / d_{ref}^2)$ is responsible for the first condition above; $d_i = d_i(\psi_i, \varphi_i)$ is the ground distance between the i -th particle location on August 29, 2015 and the Sain-Andre beach; $d_{ref} = 50$ km is the reference distance, which was chosen to be the approximate linear dimension of the Réunion Island.

5 The term $S_{i,\theta}$ responsible for the second condition, was formulated in the following way:

$$S_{i,\theta} = \begin{cases} \hat{S}_{i,\theta}, & \text{if } \exists \{t_1, t_2, t_3\} \in [T_s, T_e]: \max_{0 \leq t \leq t_1} \theta_i(t) \geq 23^\circ\text{C}, \\ & \min_{t_1 \leq t \leq t_2} \theta_i(t) \leq 19^\circ\text{C}, \max_{t_2 \leq t \leq T} \theta_i(t) \geq 24^\circ\text{C}, \\ 0 & \text{otherwise,} \end{cases}$$

where $\theta_i(t)$ is the SST at the i -th particle location at the time t ($T_s \leq t \leq T_e$, where T_s and T_e correspond to March 8, 2014 and July 29, 2015 respectively), and

$\hat{S}_{i,\theta} = (S_{i,\theta_1} + S_{i,\theta_2} + S_{i,\theta_3})/3$, where:

$$\begin{aligned} 10 \quad S_{i,\theta_1} &= \min(\max(\max_{0 \leq t \leq t_1} \theta_i(t) - 23, 0), 1), \\ S_{i,\theta_2} &= \min(\max(19 - \min_{t_1 \leq t \leq t_2} \theta_i(t), 0), 1), \\ S_{i,\theta_3} &= \min(\max(\max_{t_2 \leq t \leq T} \theta_i(t) - 24, 0), 1). \end{aligned} \quad (18)$$

The purpose of such a formulation is to relax the selection criteria and avoid discontinuities by assigning a positive score to a particle even if it did not arrive precisely to the Sain-Andre beach, or if it did not strictly satisfy the temperature condition
 15 (the tolerance allowing for a positive score was set to 1°C). As a result, the maximum score a particle could receive was one. If a particle arrived to La Réunion before August 29, it was still assigned a positive score. If a particle was located at the distance greater than 152 km from the Sain-Andre beach on August 29, it could not receive a score higher than 0.01. If the SST at particle's location never reached 23°C , or never dropped below 19°C after that, such a particle received zero score.

The percentages of particles in ensembles, which received scores $S > 0.01$ are shown in Figure 7 vs. origin's latitudes along
 20 the 7th arc for all the four model setups, together with the normal distribution fitting. As seen, for the leeway factors and non-zero drift angles of the flaperon determined by the DGA (Daniel, 2016), the segment centered at $28.2 \pm 3^\circ$ appears to be the most likely area, where the flaperon began its journey. A fraction of particles in the two other models (characterized by zero drift angle), which satisfied the two conditions, could reach a surprisingly high value of 1%, however, both the random leeway factor and constant 3.29% leeway factor models predicted the peak fitted probabilities at 30.8°S and 31.9°S respectively. The
 25 screened origins south of 36.5°S or north of 20°S are deemed to be unlikely starting locations of the flaperon, as none of the four models has predicted notable percentage of particles meeting the two requirements above for the ensembles released at these locations.

Examples of the first six particle tracks, which received the highest scores S , are depicted in Figure 8 for the scenario corresponding to the leeway factor of 3.29% and drift angle 18° to the left, starting from the origin No. 23 (99.09°E , 28.84°S).




Figure 9. Snapshot of particle locations on Dec 31, 2016 (random leeway model; origin No. 23), and percentages of particles washed ashore.

All these particles received the scores $S_i > 0.83$, and they arrived to La Réunion between June 17 (black track) and July 17 (yellow track), 2015. As seen, the two main reasons for the drop in the ambient water temperature from 23–25°C to as low as 16°C are:

1. Seasonal cooling of the water surface (see comparison of the SSTs on March 8 and June 8, 2014 in Figure 4);
2. Entrapment in counter-clockwise eddies, which could first carry the flaperon northwestward up to 22–25°S latitudes and then southward to 30–33°S (Figure 8).

3.3 Beached debris distribution

A total of 27 possibly relevant and confirmed fragments of 9M-MRO were recovered in La Réunion, Mauritius, Rodrigues, Madagascar, Mozambique, South Africa and Tanzania as on April 2017 according to the MSIT (2017); none was ever found in Australia, although a suspected object, the unopened towelette bearing MAS logotype, was discovered at the Thirsty Point. Distribution of the found debris offers a useful insight into the possible location of the crash site (e.g., Jansen et al., 2016; Pattiaratchi and Wijeratne, 2016; Griffin et al., 2017), although no consensus was reached up to date with regard to the origin.




Figure 10. Expected number of objects to be found in several countries by Dec 31, 2016 vs. origin's latitudes.




Figure 11. Expected along-shore concentration of beached MH370 objects to be found by the end of 2016, origin No. 23.

Similarly to the aforementioned studies, an effort was made in this study to analyze spatial distribution of the washed ashore fragments.




Figure 12. Arrival times of particles to the proximity of the Thirsty Point (Australia) during 2014 and Mossel Bay (South Africa).

A sample snapshot of the computed particle locations on December 31, 2016 is depicted in Figure 9 for the random leeway model, origin No. 23. Corresponding animation S2 is included in the Supplement. Considerable beaching of particles of this ensemble was modelled in Africa, Madagascar, La Réunion and Mauritius, but negligible in Australia. The total fractions of particles landed by the ends of 2015 and 2016 are shown in the right top corner for all the screened origins, both the random and constant 3.29% leeway factor models. This result is, in part, due to the West Australian Current, which entrains large percentages of particles from the northern and southern origins (see Animations S3 and S4 in the Supplement), while most of particles from the middle of the screened segment of the 7th arc remain trapped in the Indian Ocean Gyre.

An interesting result can be obtained by comparing the modelled ratios of the fractions of particles washed ashore in several countries against the ratios of the number of fragments actually recovered in these countries. Such a comparison provides an indication of a number of fragments expected to be found by certain date assuming the same reporting factors. Figure 10 shows computed ratios for the random leeway model, using South Africa as the reference (a total of 5 objects were found there). As seen, more than 5 objects could be expected in Australia, at least 2 objects in Sri Lanka, 9 objects in Tanzania and 3 objects in Kenya, should the origin be south of 35°S. At least one fragment could be expected in Sri Lanka, and several in Kenya and Tanzania for the origins north of 23°S. The best matching segment of the 7th arc is located approximately between 26.5°S and 31°S. In particular, the ratio of the percentages of particles beached in Mozambique to those in South Africa was in the range 1.4 to 1.7, being in a good agreement with the actual ratio of 6:5.

To estimate along-shore concentration of MH370 objects expected to be found by certain time, the two-dimensional Gaussian smoothing filter was first applied to obtain smoothed concentration of beached particles:

$$P(\psi, \varphi) = \frac{1}{2\pi d_{ref}^2} \sum_{i=1}^M \exp\left(-\frac{d_i^2(\psi, \varphi, \psi_i, \varphi_i)}{2 d_{ref}^2}\right),$$

where d_i is the ground distance between the i -th beached particle and the location of interest (ψ, φ) , $d_{ref} = 5$ km is the size of the smoothing filter, and M is the number of beached particles. To compute along-shore density distribution of objects expected to be found, this function was numerically integrated over a relatively narrow band-shaped areas Ω_i long-wise centered at the



shorelines, divided by the respective lengths of the shoreline segments Δs_i , and prorated by the ratio of the number of fragments found in South Africa $N_{SA} = 5$ to the number of particles landed in South Africa M_{SA} :

$$\frac{dN}{ds} \approx \frac{N_{SA}}{M_{SA}} \frac{1}{\Delta s_i} \iint_{\Omega_i} P(\psi, \varphi) d\Omega.$$

An example of the estimated in this way concentration of objects is shown in Figure 11 for the ensemble released from the origin No. 23 (random leeway model); concentrations for all the other studied origins are presented in Figure S5 in the Supplement. As seen, locations of the elevated concentrations are in a fairly good agreement with the locations, where the fragments were found, except the Rodrigues Island, which was not properly resolved by HYCOM. More fragments could be expected in Tanzania at 7°S and 8.2°S.

Screening of the origin No. 23 has revealed that the elevated concentration in the proximities to the Cape Leeuwin in Australia is due to the beaching, which mainly occurred in 2016. During 2014-2015 notable arrival of particles from this origin to Australia took place only around the Windy Harbor and Thirsty Point, where the unopened MAS towelette was found on July 2 (assumed coordinates 115.3°E, 31°S). A detailed analysis has shown that 8 particles of the respective ensemble landed in the proximity to the Thirsty Point during July 8-11, and 10 more particles arrived before July 28, 2014. All the first 8 particles were characterized by relatively low leeway factors ranging from 0.1% to 0.4%. This suggests a systematic feature rather than a random occurrence. Systematic arrival of particles from the origins north of 27°S or south of 31°S was not predicted earlier than in the last week of August, as seen in Figure 12.

Figure 12 also shows the arrival times of particles to the Mossel Bay in South Africa, where the engine cowling fragment was found. Being the fourth found object, it has covered long distance over a relatively short time interval. The modelling has shown possibility of the sporadic arrival of high-windage particles to the Mossel Bay as early as in February 2015, but the systematic arrival was predicted only in March-May 2016, being consistent with the discovery date.

4 Conclusions

The drift study of MH370 debris was conducted by the means of numerical modelling using forward particle tracking technique. A total of 40 hypothetical locations of the crash site along the 7th arc were screened. The three major aspects were considered: (1) efficacy of the aerial search; (2) ambient water temperatures along the path of the flaperon to La Réunion; (3) the spatial distribution of the washed ashore debris.

The governing equations were numerically integrated in the geocentric Cartesian coordinate system, where the Earth surface was approximated by the WGS'84 ellipsoid. Four models with regard to the leeway factors and drift angles were considered, including a proposed model of random distribution of the leeway factors of particles in an ensemble.

The obtained results indicate significance of the leeway factor in all the three aspects considered. In addition to the uncertainties in the model forcing, assumptions and simplification, judgment about the most likely location of the crash site depends on weights assigned to the aerial search, accuracy of the barnacle biochemical analysis, and probability of fragments not only



to be washed ashore, but also recovered and reported. While it does not appear to be possible to confidently point out location of the crash site based on the drift study alone, a few observations can be made with regard to various segments of the 7th arc:

- **South of 36°S:** Considerable beaching, where no fragments were found, particularly in Australia and Sri Lanka; incompatibility with the water temperatures suggested by the barnacle biochemical analysis.
- 5 • **34.5 to 36°S:** While corresponding areas were poorly surveyed during the aerial search, considerable beaching could be expected in several countries, where no fragments were found, particularly in Australia.
- **30.5 to 34.5°S:** Excellent aerial coverage of the debris cloud originated from this segment makes the crash site unlikely to be located within it.
- **25.5 to 30.5°S:** Consistency with the barnacle temperature analysis; elevated concentration of beached particles, where
 10 the fragments of 9M-MRO were found; several 'gaps' in the aerial search; floating objects detected on March 28-31; possible consistency with the early arrival of the MAS towelette to the Thirsty Point.
- **North of 25.5°S:** Inconsistency with the distribution of the washed ashore debris; incompatibility with the barnacle temperature analysis; good aerial coverage of the areas corresponding to the origins from 20°S to 25°S.

Summarizing all the above, the most likely area of the crash site appears to be between 25.5°S and 30.5°S, with the segment
 15 from 28°S to 30°S being the most promising. This area is consistent with the original definition of a high-priority search zone by the ATSB in June 2014.

Competing interests. This study was conducted solely at author's expense. The author declares that he has no conflict of interest.

Acknowledgements. The use was made of current data from HYCOM, provided by the National Ocean Partnership Program and the Office of Naval Research; the wind data was sourced from NOAA ARL GDAS; the SST data was obtained from NASA EOSDIS PODAAC at the
 20 JPL. The author is thankful to the anonymous reviewers, who helped to improve clarity and content of this paper.



References

- Australian Maritime Safety Authority (AMSA). MH370 Search - Media kit, 2014. Available at: <http://www.amsa.gov.au/media/incidents/mh370-search.asp>.
- Australian Transport Safety Bureau: MH370 - Definition of Underwater Search Areas, 26 June 2014. Available at: <http://www.atsb.gov.au>.
- 5 Australian Transport Safety Bureau: MH370 - Flight Path Analysis Update, 30 July 2014. Available at: <http://www.atsb.gov.au>
- Australian Transport Safety Bureau: MH370 - Definition of Underwater Search Areas, 3 December 2015. Available at: <http://www.atsb.gov.au>.
- Ashton C., Bruce A.S., Colledge G., and Dickinson M. The search for MH370, *The Journal of Navigation*, 68, pp. 1-22, 2015.
- Blumberg A.F., Mellor G.L. A description of a three-dimensional coastal ocean circulation model. *Three-Dimensional Coastal Ocean Models*,
10 N. Heaps (Ed.), Washington D.C., Am. Geoph. Union, pp. 1-16, 1987.
- Box G. E. P. and Muller M.E. A Note on the Generation of Random Normal Deviates. *Ann. Math. Statist.*, 29(2), pp. 610-611, 1958.
- Chassignet, E.P., Hurlburt, H.E., Smedstad, O.M., Halliwell, G.R., Hogan, P.J., Wallcraft, A.J., Baraille, R., and Bleck, R.: The HYCOM (HYbrid Coordinate Ocean Model) data assimilative system, *J. Mar. Syst.*, 65, pp. 60-83, 2007.
- Daniel P. Rapport d'étude. Dérive à rebours de flaperon. Météo France, 8 février 2016. Direction des Opérations pour la Prévision, Département
15 Prévision marine et Océanographique, 32 p., in French.
- Davey S., Gordon N., Holland I., Rutten M., and Williams J. Bayesian Methods in the Search for MH370. *Springer Briefs in Electrical and Computer Engineering*, 2016, 114 p.
- De Deckker P. Chemical investigations on barnacles found attached to debris from the MH370 aircraft found in the Indian Ocean. Appendix F in ATSB report: "The Operational Search for MH370", AE-2014-054, October 3, 2017. Available at: <http://www.atsb.gov.au>.
- 20 DHI Water and Environment. MIKE21 and MIKE3 Flow Model FM. Particle Tracking Module. Scientific Documentation, 28 pp., Hørsholm, Denmark, 2009. Available at: www.dhigroup.com.
- Durgadoo J. and Biastoch A. Where is MH370? GEOMAR Helmholtz Centre for Ocean Research Kiel, 28 August 2015. Available at: <http://www.geomar.de>.
- Eisenberg B. and Sullivan R., Why is the Sum of Independent Normal Random Variables Normal, *Math. Magazine*, 81, pp. 362-366, 2008.
- 25 Gandin A.S., Lughtman D.L., Matveyev L.T., and Yudin M.I., *Fundamentals of dynamic meteorology*, Gidrometizdat, Leningrad (in Russian), 647 pp., 1955.
- García-Garrido V.J., Mancho A.M., Wiggins S., and Mendoza C. A dynamical systems approach to the surface search for debris associated with the disappearance of flight MH370. *Nonlin. Processes Geophys.*, 22, pp. 701-712, 2015. Available at: <http://www.nonlin-processes-geophys.net/22/701/2015/npg-22-701-2015.pdf>.
- 30 Griffin D., Oke P. and Jones E. The search for MH370 and ocean surface drift - Part II. Report No. EP172633, prepared for the Australian Transport Safety Bureau, 13 April 2017. Available at: <http://https://www.atsb.gov.au>.
- The Joint Agency Coordination Centre (JACC). Media Releases - 2014. Available at: <http://jacc.gov.au/media/releases/2014/april>.
- Jansen E., Giovanni Coppini G., and Pinardi N. Drift simulation of MH370 debris using superensemble techniques. *Natural Hazards and Earth System Sciences*, pp. 1623 - 1628, 2016.
- 35 JPL MUR MEaSUREs Project. 2015. GHRSSST Level 4 MUR Global Foundation Sea Surface Temperature Analysis (v4.1). Ver. 4.1. PO.DAAC, CA, USA. Available at: <https://podaac.jpl.nasa.gov/dataset/MUR-JPL-L4-GLOB-v4.1>.



- Korn G.A. and Korn T.M. Mathematical Handbook for Scientists and Engineers. Definitions, Theorems and Formulas for Reference and Reviews. Second Enlarged and Revised Edition. McGraw-Hill Book Company, 832 pp., 1968.
- Kraus E.B. Atmosphere-Ocean Interaction. Ed. Shepard P.A. Clarendon Press - Oxford, 1972.
- Malaysian ICAO Annex 13 Safety Investigation Team for MH370, Ministry of Transport, Malaysia. Summary of Possible MH370 Debris Recovered, April 2017. Available at: <http://www.mh370.gov.my>.
- Marsaglia G. and Zaman A. Toward a Universal Random Number Generator. Florida State University Report: FSU-SCRI-87-50, 1987.
- Maximenko N., Hafner J., Speidel J., and Wang K.L. IPRC Ocean Drift Model Simulates MH370 Crash Site and Flow Paths. International Pacific Research Center, School of Ocean and Earth Science and Technology at the University of Hawaii, August 4, 2015. Available at: http://iprc.soest.hawaii.edu/news/MH370_debris/IPRC_MH370_News.php.
- 10 Ministry of Transport Malaysia. Official Portal for MH370 Air Incident Investigation: <http://www.mot.gov.my/en/aviation/air-incident-investigation/MH370>.
- van Ormondt M., Baart F. Aircraft debris MH370 makes Northern part of the search area more likely. Deltares News, 31 July 2015. <https://www.deltares.nl/en/news/aircraft-debris-mh370-makes-northern-part-of-the-search-area-more-likely>.
- Pattiaratchi C., Wijeratne S. Ocean currents suggest where we should be looking for missing flight MH370. The Conversation, July 28, 2016: <https://theconversation.com/ocean-currents-suggest-where-we-should-be-looking-for-missing-flight-mh370-63100>.
- 15 Al-Rabeh A.H., Lardner R.W., Gunay N. Gulfspill Version 2.0: a software package for oil spills in the Arabian Gulf. Environmental Modelling & Software, 15 (2000), pp. 425-442.
- Smagorinsky J. General circulation experiments with primitive equations. The basic experiment. Monthly Weather Rev., 91, pp. 99-164, 1963.