

Evaluation of extreme wave probability on the basis of long-term data analysis.

5 Kirill Bulgakov^{1,2}, Vadim Kuzmin², Dmitry Shilov²

¹Shirshov Institute of Oceanography, Russian Academy of Sciences, 36, Nahimovsky prospect, Moscow, 117997, Russia,

²Russian State Hydrometeorological University, 98, Maloochtinsky Pr., Saint-Petersburg, 195196, Russia

10 Correspondence to Kirill Bulgakov (bulgakov.kirill@gmail.com)

Abstract. A method of calculation of wind wave height probability based on the significant wave height probability is described (Chalikov and Bulgakov, 2017). The method can be also used for estimation of height of extreme waves of any given cumulative probability. The application of the method on the basis of long-term model data is presented. Examples of averaged annual and seasonal fields of extreme wave heights obtained by the above method are given. Areas where extreme wave can appear are shown.

Introduction

Highest risks of economic and environmental damage for the sea-based human activities, i.e., cargo shipments, fishery, oil production etc., are mostly connected with extreme weather conditions on sea surface among which strong storms are the foremost. It is especially difficult to predict emergency situations caused by extreme waves for those cases of the sea-based activities which require people's long stay at sea or prolonged use of the equipment in the ocean.

One of the methods to minimize such possible risks is use of climatic data based on long-term series of observations. At present there are archives consisting of the reanalysis data on surface waves based on wave forecast corrected by different methods, i.e., direct measurements using accelerometers and GPS-buoys; remote measurements by satellite-borne altimetry and various types of radars. The main characteristic of wave field included in the archive is significant wave height H_s defined as a mean value (trough to crest) of one third of the highest of all the waves (Ochi 2005). The value of H_s is calculated in the following way:

$$H_s = 4 \left(\int_0^\infty \int_0^\infty S(k_x, k_y) dk_x dk_y \right)^{1/2}, \quad (1)$$

where k_x, k_y are wave numbers, while $S(k_x, k_y)$ is wave spectrum.

It is evident that significant wave height is not enough to evaluate real wave height for a given wave field. Extreme waves of the same height can appear with different probability for different values of H_s . For example, a wave 10 meters high can appear both in a wave field with $H_s=10$ m. and in a wave field with $H_s=5$ m. Or there can be waves with height 15 meters and 17 meter in wave field characterized by H_s 10 meters. Thus, H_s data does not give enough information about the probability of real wave heights.

The nature of freak waves was investigated analytically (Onorato et al., 2009) and numerically (Chalikov, 2009). Recently it was found that the statistical properties of trough-to-crest wave height are quite different from those of the wave height above mean level. Paper (Chalikov and Babanin, 2016; Chalikov, 2017) shows that linear and nonlinear statistics of extreme waves (defined as trough-to-crest waves) are identical not only for broad spectrum but for one-dimensional wave field too. It means that generation of a trough-to-crest extreme wave is the result of simple superposition of linear modes, no matter how broad the spectrum is. This

property is not found for the wave height above mean level. Thus, the statistical properties of trough-to-crest wave height can be investigated with linear modeling, but just by generation of large ensembles of superposition of linear modes with random phases and the spectrum prescribed. It is obvious that the problem of the trough-to-crest statistics becomes quite straightforward. From the practical point of view the data on the probability of wave height above mean level are important for a fixed construction type of offshore platforms. For floating objects the data on full height (trough-to-crest) of wave are more important. Contrary to such approach, investigation of the statistics of wave height above mean level remains a subject of nonlinear wave theory.

The theoretical probability distribution for wave crest height (or wave height above mean level) was suggested by Weibull (1951). Later it was studied on a basis of observational data in nature and wave channels (see review by Kharif et al., 2009). Extended data for estimation of probability of wave height can be obtained with integration of nonlinear modes based on full potential equations (Touboul and Kharif, 2010; Chalikov et al, 2009). Methods of probability calculations were considered in many papers (see, for example Bitner-Gregersen and Toffoli, 2012; Dyachenko et al, 2016).

The most popular method of trough-to-crest wave height detection is based on zero-crossing technique. Direct method is based on use of moving windows, which is applicable both for 1-D and 2-D cases.

Estimation of extreme waves nowadays mostly is made by analysis of data of significant wave height. (Jiangxia Li 2018) analyzed long-term data considered than extreme wave is wave exceeding two significant wave heights. In (Larsen et al, 2015) It was suggested that there is impact of unresolved part of spectrum in wave forecast model (frequencies more than 2.5 10⁻⁵ Hz) on calculation extreme waves. The spectral correction method was developed to fill in the missing variability of the modeled variable at high frequencies. In (Lanli Guo and Jinyu Sheng, 2015) the peaks-over-threshold method was used to estimate the extreme significant wave heights from 30-year wave simulations. In (Samayam S. et al 2017) estimation of extreme wave was made by using of the methods of extreme value theory. The main advantage of the method (Chalikov and Bulgakov, 2017) to compare with methods mentioned above and others that method is based on results of direct modeling of wave fields.

This paper is devoted to investigation of the statistics and geographical distribution of wave height above mean sea level.

1. Description of the method

In paper (Chalikov and Bulgakov, 2017) an algorithm for estimation of cumulative probability of waves exceeding a specific value of wave height above mean level ($P(h)$ and h below) was developed using long-term data on H_s . The description of the method is given below.

The probability of wave exceeding specific height h , if significant wave height is in a small range dH_s around H_s , equals $\tilde{P}(\tilde{H})$ for specific $\tilde{H} = h/H_s$ multiplied by probability of H_s in this range ($\tilde{P}(\tilde{H}) \cdot P(H_s)$), by the standard definition of conditional probability. Consequently, $P(h)$ can be determined as integral of $\tilde{P}(\tilde{H}) \cdot P(H_s)$: over all possible value of H_s :

$$P(h) = \int_0^{H_{smax}} \tilde{P}(\tilde{H}) P(H_s) dH_s, \quad (2)$$

where $P(H_s)$ is probability distribution of H_s for a specific point, while H_{smax} is the maximum value of H_s in the dataset for a specific point.

The model H_s data used for $P(H_s)$ were calculated with the latest version of WAVEWATCH III model (Tolman 2014) and GFS-2 wind analysis 2 (Sasha et al. 2014). The hindcasts cover the period from August 1999 to July 2015. The spatial resolution of the dataset fields is 0.5×0.5 degree. Calibration of the model and its validation are carried out using a great number of wave buoys. The data and results of its validation are described in (Chawla et al., 2013).

The approximation of $\tilde{P}(\tilde{H})$ was based on results of 3-D model of potential fluid (the curl of the velocity field is zero). The model used spectral definitions of fields, finite differences for vertical derivatives calculation, fourth-order Runge–Kutta scheme for time integration. Fourier resolution is 256×64 wave number, resolution in physical space is 1024×256 (more detail in (Chalikov et al., 2014)). The calculations were done for 350 units of nondimensional time, i.e., for 70,000 time steps. The initial conditions were generated on basic JONSWAP spectrum. Model runs were calculated under condition when input energy from wind to waves equals wave energy dissipation. This condition corresponds fully developed wind waves. Totally 50 experiments were made (more detail in (Chalikov and Bulgakov, 2017)). The results of the series of experiments were processed in the following way: each wave field of surface height above mean level (η) reproduced by numerical model was normalized by the value of significant wave height corresponding to this field. ($\tilde{H} = \eta/H_s$). (Note that η is variable of 3-D model of potential waves. It should be distinguished from h despites the fact that both (η and h) have the same physical sense.)

Then, a nondimensional wave field was used for calculation of cumulative probability of nondimensional wave height $\tilde{P}(\tilde{H})$. The distribution obtained was approximated by the following function:

$$\tilde{P}(\tilde{H}) = \exp(-3.97\tilde{H} - 4.02\tilde{H}^2) \quad (3)$$

Note, that $\tilde{P}(\tilde{H})$ is cumulative probability of the height of free surface above mean level. This probability for $\tilde{H} = 1$ (the height of free surface equals significant wave height) is quite small (0.0003).

The above expression can be used for the interval $0 \leq \tilde{H} \leq 1.85$. The probability of wave higher than 1.85 (it's maximal value of $\tilde{H}$ in data) can be considered as extremely low and therefore - neglected. It should be noted that approximation (3) was obtained with use of the precise 3-D model based on full nonlinear equations. The volume of data used for approximation (3) includes more than 4.5 billion values of η (number point in single field multiplied by number of record in experiment multiplied by number of experiments). Currently, this approximation is considered as universal for wind wave fields where cases of freak waves are most likely. Waves of other types of spectrum (swells) have a small steepness and don't influence on extreme wave generation except rare cases when long-wave currents can steepen shorter waves.

The examples of the calculations using the method (Chalikov and Bulgakov, 2017) where space distribution of the extreme wave probability was investigated.

In these paper results of another approach of this method are considered. The global fields of wave height with cumulative probability 10^{-7} were calculated with using data Chawla et al., 2013)..

5

2. Results




Fig.1 — Wave heights (m) with cumulative probability 10^{-7} , annual average

10

Figure 1 shows an average annual field of wave heights with cumulative probability 10^{-7} . It can be seen that waves with the height up to 20 m can appear with such probability, some of the extreme waves (16 m and more) being found in the areas of active navigation (eastern part of the Atlantic Ocean, East China Sea, Philippine Sea, Yellow Sea, south-western part of the Pacific Ocean).

15




Figure 2 – Average annual significant wave height (m).

Distribution of **annual average** significant wave height **provided by model** is shown in Fig. 2. As
 5 seen, the maximum value of field of **annual average** significant wave height does not exceed 5 m
 (**southern area of Indian and Pacific oceans**), while the height of real extreme wave can reach **16**
m there. The data in Fig 1 have a more complicated structure, due, for example, to the periods
 with strong wind along trajectories of tropical storms. Consequently, the calculations of
 distribution of real wave height should be done for shorter periods, i.e., for seasonal or monthly
 10 averaged data on significant wave heights.




Figure 3. – Wave height (m) with cumulative probability 10^{-7} for December-February.

In Fig. 3 the field of wave height with probability 10^{-7} averaged for December-February, is shown.

When comparing Figs 3 and 1 it is seen that in mid-latitudes of the Northern Hemisphere wave heights become higher. In some areas appearance of extreme wave heights exceeding 16 m is possible. At the same time there are actually no extreme waves in the eastern part of the Arctic Ocean, which is connected with seasonal ice formation in the area. In equatorial and tropical areas of the World Ocean wave heights are less in winter (Northern Hemisphere), as compared with the average annual wave heights. It should be noted that in the western part of the Atlantic Ocean tracking trajectories of hurricanes disappeared while the number of such trajectories increased in the Indian Ocean.




Figure 4. – Wave height (m) with cumulative probability 10^{-7} averaged for March-May.

An increase of wave heights over March-May can be registered (Fig.4) in the Southern Hemisphere. Actually all the area of mid-latitudes from the latitude of 40 degrees S. to the latitude of 60 degrees S. is characterized by probability $>10^{-7}$ of wave heights over 14 m. In mid-latitudes of the Northern Hemisphere in spring wave height values are more than the average annual values, though less than the winter values, while in some areas (Atlantic Ocean near Iceland, Pacific Ocean near Bering Sea) appearance of waves exceeding 14 m in height is quite possible.




Fig.5 –Wave heights (m) with the cumulative probability of 10^{-7} , for June-August

Summer months (Fig.5) are characterized by general decrease of extreme wave probability. It is especially noticeable in the northern areas of the Atlantic and Pacific Oceans. Also, wave heights slightly decreased in the Southern Hemisphere. It should be noted that storm tracks appear off the eastern coast of North America and disappear in the southern part of the Pacific Ocean. Besides, quite distinct trajectories of storms appeared in the eastern part of the Pacific Ocean. Small wave heights can be observed in the Arctic Ocean, in the area free from ice.




Figure 6. – Wave heights (m) with cumulative probability 10^{-7} , averaged for September - November

During autumn months (Fig. 6) an increase of wave heights is observed in the Arctic Ocean, the extreme wave height values sometimes reaching 20 m. Among other peculiarities is an increase of wave-free area in polar latitudes of the Southern Hemisphere, which is obviously connected with seasonal ice formation.

It is quite evident that the average monthly fields of cumulative wave-height probability will allow us to obtain more exact information on the areas of extreme wave probability.

The above approach can be used with different values of cumulative probability. It is not expedient to use the values less than 10^{-9} as this value is outside the range of validity for equation (2) ($\tilde{P}(1.85)$ is approximately 10^{-9}). Hence, on the whole, the method considered is suitable for estimation of extreme values of wave heights with minor probability.

3. Conclusions

The paper describes a method of calculation of extreme wave probability, based on long-term wave hindcast data on significant wave height developed in (Chalikov and Bulgakov 2017). Such method can be used for estimation of probability of extreme waves, which is important for designing of engineering constructions. Another approach of the methods is presented in the paper; algorithm (Chalikov and Bulgakov 2017) was used to evaluate height of waves of any given cumulative probability. The maps of global distribution of wave heights of certain probability for main seasons illustrate the approach of the method.

Authors are thankful to Professor D. Chalikov for his useful consultations.

The investigation was fulfilled with financial support of Russian Scientific Fund (Project No 16-17-00124).

References

- Bitner-Gregersen E. M. and Toffoli A.: On the probability of occurrence of rogue waves, Nat. Hazards Earth Syst. Sci., 2012, 12, 751–762, doi:10.5194/nhess-12-751-2012
- Chalikov D.: Freak waves: their occurrence and probability, Phys. of Fluid, 2009, 21, 076602; doi:10.1063/1.3175713
- Chalikov D., Babanin A., and Sanina E.: Modeling of Three-Dimensional Fully Nonlinear Potential Periodic Waves, Ocean Dynamics, 2014, 64, 10, 1469–1486. doi: 10.1007/s10236-014-0755-0
- Chalikov D. and Babanin A.V.: Comparison of linear and nonlinear extreme wave statistics, Acta Oceanologica Sinica, 2016, 35, 5, 99–105, doi:10.1007/s13131-016-0862-5
- Chalikov D.: Numerical modeling of sea waves, 2016, Springer, ISBN 978-3-319-32914-7 ISBN 978-3-319-32916-1 (eBook), doi 10.1007/978-3-319-32916-1, 330 pp
- Chalikov D.V.: Linear and nonlinear statistics of extreme waves, Russian Journal of Numerical Analysis and Mathematical Modelling, 2017, 32, 2, 91–99.
- Chalikov D. and Bulgakov K.: Estimation of wave height probability based on the statistics of significant wave height, J. Ocean Eng. Mar. Energy, 2017. <https://doi.org/10.1007/s40722-017-0093-7>
- Chawla A., Spindler D.M., and Tolman H.L.: Validation of a thirty year wave hindcast using the climate forecast system reanalysis winds, Ocean Modelling, 2013, 70, 189–206.
- Ochi M.K. Ocean waves: the stochastic approach, 2005, Cambridge university press, ISBN-13 978-0-521-56378-9? ISBN-10 0-521-01767-X, 332 pp.

- Dyachenko A.I., Kachulin D.I., and Zakharov V.E.: Probability distribution function of freak waves: nonlinear versus linear model, *Studies in applied mathematics*, 2016, 132, 2, 189-198
- 5 Guo L. and Sheng J.: Statistical estimation of extreme ocean waves over the eastern Canadian shelf from 30-year numerical wave simulation, 2015, 65, 11, 1489-1507
- Jiangxia L., Shunqi P. Yongping C., Yang-Ming Fan and Yi P.: Numerical estimation of extreme waves and surges over the northwest Pacific Ocean, *Ocean Engineering*, 2018, 153, 225-241
- 10 Kharif C., Pelinovsky E., and Slunyaev A.: Rogue Waves in the Ocean, *Advances in geophysical and environmental Mechanics and Mathematics*, 2009, DOI 1007/978/-3-540-88419-4
- Larsen X.G., Kalogeri C., Galanis G. and Kallos G.: A statistical methodology for the estimation of extreme wave conditions for offshore renewable applications, *Renewable Energy*, 2015, 205-218.
- 15 Onorato M., T. Waseda T., T, Toffoli A, Cavaleri L, Gramstad O, Janssen PAEM, Kinoshita T, Monbaliu J, Mori N, Osborne AR, Serio M, Stansberg CT, Tamura H, and K. Trulsen.: K Statistical Properties of Directional Ocean Waves: The Role of the Modulational Instability in the Formation of Extreme Events, *Phys. Rev. Lett.*, 2009, 102, 114502
- 20 Samayam S., Laface V., Annamalaisamy S.S., Areana F., Valaam S. and Ponikov V.G.: Assessment of reliability of extreme wave height prediction models, *Natural Hazards and Earth System Science*, 2017, 17, 409-421.
- Sasha S., Moorthi S., Pan H. and Goldberg M., The NCEP climate forecast reanalysis version 2, *Journal of Climate*, 2014, 27, 6, 2185-2208.
- 25 Tolman H.L.: 2014. User manual and system documentation of WAVEWATCH III version 4.18. NOAA / NWS / NCEP / MMAB Technical Note 316, 194 pp.+ Appendices (pdf)
- Touboul J. and Kharif C.: Two-dimensional direct numerical simulations of the dynamics of rogue waves under wind action, *Advances in Numerical Simulation of Nonlinear Water Waves*, 2010, 11, 43-74.
- 30 Weibull W.: A statistical distribution function of wide applicability, *Journal of applied mechanics*, 1951, 18, 293-297.