



Evaluation of extreme wave probability on the basis of long-term data analysis.

5 Kirill Bulgakov^{1,2}, Vadim Kuzmin², Dmitry Shilov²

¹Shirshov Institute of Oceanography, Russian Academy of Sciences, 36, Nahimovsky prospect, Moscow, 117997, Russia,

²Russian State Hydrometeorological University, 98, Maloochtinsky Pr., Saint-Petersburg, 195196, Russia

10 Correspondence to Kirill Bulgakov (bulgakov.kirill@gmail.com)

Abstract. A method of calculation of wave height probability based on the significant wave height probability is described. An application of the method on the basis of long-term data analysis is presented. Examples of averaged annual and seasonal fields of extreme wave heights obtained by the above method are given.

1. Introduction

Highest risks of economic and environmental damage for the sea-based human activities, i.e., cargo shipments, fishery, oil production etc., are mostly connected with extreme weather conditions on sea surface among which strong storms are the foremost. It is especially difficult to predict emergency situations caused by extreme waves for those cases of the sea-based activities which require people's long stay at sea or prolonged use of the equipment in the ocean.

One of the methods to minimize such possible risks is use of climatic data based on long-term series of observations. At present there are archives consisting of the reanalysis data on surface waves based on wave forecast corrected by different methods, i.e., direct measurements using accelerometers and GPS-buoys; remote measurements by satellite-borne altimetry and various types of radars. The main characteristic of wave field included in the archive is significant wave height H_s defined as a mean value (trough to crest) of one third of the highest of all the waves (Ochi 2005). The value of H_s is calculated in the following way:

$$H_s = 4 \left(\int_0^\infty \int_0^\infty S(k_x, k_y) dk_x dk_y \right)^{1/2}, \quad (1)$$

where k_x, k_y are wave numbers, while $S(k_x, k_y)$ is wave spectrum.

It is evident that significant wave height does not provide any information on real wave height for a given wave field. Extreme waves of the same height can appear with different probability for different values of H_s . For example, a wave 10 meters high can appear both in a wave field with $H_s=10$ m and in a wave field with $H_s=5$ m. Thus, there are not enough data on H_s to evaluate the probability of real wave heights exceeding certain values and to calculate the probability of real wave heights for specific values H_s .

Up to the present some vagueness of definition of extreme ('freak') wave has existed. The standard definition suggests that any wave exceeding two significant wave heights H_s can be considered as "freak" wave (Chalikov 2016). Firstly, such definition is not universal, since it does not define real dimensional wave height. This definition makes sense for theoretical investigations only, as the adiabatic equations of wave dynamics are self-similar, i.e., when transformed to the nondimensional form, they do not contain any nondimensional parameters, which makes the whole approach quite convenient due to its universal character. The non-adiabatic factors (for example, input energy) violate self-similarity; anyway, since the time scale of wave energy transformation is by many orders of magnitude larger than the wave periods, a nondimensional approach in theoretical investigations is still acceptable. However, practical



conclusions of the theoretical investigation should be formulated in a dimensional form. Secondly, the value $2H_s$ is large, which means that the definition refers to the full height of wave, i.e., trough-to-crest height. This definition is straightforward for linear monochromatic waves only when the trough-to crest height is simply equal to the doubled wave height above mean level. For a complicated multi-mode wave field the trough-to crest height should be defined as a vertical distance between crest and the closest wave trough. The question is what the term ‘closest’ means. The only way recommended in (Chalikov, 2009) (see also (Chalikov, 2016)) is based on the consideration of moving window with horizontal size L_w . Since extreme wave actually occurs at peak frequency, it is reasonable to assume $L_w = 1.5L_p$ (where L_p is the wavelength in peak of spectrum). Coefficient 1.5 is introduced to take into account the variability of length of ‘wave’ (which in fact is a transient local superposition of several modes).

The nature of freak waves was investigated analytically (Onorato et al., 2009) and numerically (Chalikov, 2009). Recently it was found that the statistical properties of trough-to-crest wave height are quite different from those of the wave height above mean level. Paper (Chalikov and Babanin, 2016; Chalikov, 2017) shows that linear and nonlinear statistics of extreme waves (defined as trough-to-crest waves) are identical not only for broad spectrum but for one-dimensional wave field too. It means that generation of a trough-to-crest extreme wave is the result of simple superposition of linear modes, no matter how broad the spectrum is. This property is not found for the wave height above mean level. Thus, the statistical properties of trough-to-crest wave height can be investigated with no nonlinear modeling, but just by generation of large ensembles of superposition of linear modes with random phases and the spectrum prescribed. It is obvious that the problem of the trough-to-crest statistics becomes quite straightforward. Contrary to such approach, investigation of the statistics of wave height above mean level remains a subject of nonlinear wave theory.

The paper is devoted to investigation of the statistics and geographical distribution of wave height above mean sea level.

2. Description of the method

In paper (Chalikov and Bulgakov, 2017) an algorithm for estimation of cumulative probability of waves exceeding a specific value h ($P(h)$ below) was developed using long-term data on H_s . The results of the series of experiments carried out with use of 3-D model of potential waves (described in detail in (Chalikov et al., 2014)), were processed in the following way: for each wave field reproduced by numerical model the significant wave height H_s was calculated. This value was used for scaling of wave fields ($\tilde{H} = \eta/H_s$). Then, a nondimensional wave field was used for calculation of cumulative probability of nondimensional wave height \tilde{H} . The distribution obtained was approximated by the following function:

$$\tilde{P}(\tilde{H}) = \exp(-3.97\tilde{H} - 4.02\tilde{H}^2) \quad (2)$$

Note, that $\tilde{P}(\tilde{H})$ is cumulative probability of the height of free surface above mean level. This probability for $\tilde{H} = 1$ (the height of free surface equals significant wave height) is quite small (0.0003).

The above expression can be used for the interval $0 \leq \tilde{H} \leq 1.85$. The probability of wave higher than 1.85 can be considered as extremely low and therefore - neglected. It should be noted that approximation (2) was obtained with use of the precise 3-D model based on full nonlinear equations. The volume of data used for approximation (2) includes more than 3 million values.



Currently, this approximation is considered as universal. Calculation of $P(h)$ can be done by the following formula:

$$P(h) = \int_0^{H_{smax}} \tilde{P}\left(\frac{h}{H_s}\right) P(H_s) dH_s, \quad (3)$$

5

where $P(H_s)$ is distribution of cumulative probability H_s for a specific point, while H_{smax} is the maximum value of H_s in the initial data.

The data (Chawla et al., 2013) used were calculated with WAVEWATCH III (Tolman, 2014). Calibration of the model and its validation are carried out using a great number of wave buoys.

10 The examples of the calculations using the method (1-3) are given in (Chalikov and Bulgakov, 2017) where space distribution of the extreme wave probability was investigated. The method 1-3 can be also used for estimation of height of extreme waves of any given cumulative probability.

15 3. Results




Fig.1 — Wave heights (m) with cumulative probability 10^{-7} , annual average

20 Figure 1 shows an average annual field of wave heights with cumulative probability 10^{-7} . It can be seen that waves with the height up to 20 m can appear with such probability, some of the extreme waves (16 m and more) being found in the area of active navigation (eastern part of the Atlantic Ocean, East China Sea, Philippine Sea, Yellow Sea, south-western part of the Pacific Ocean).

The probability of extreme wave can be given in terms of expectance time $T_e = T_p / P(h)$,
 25 where T_p is a period of wave with height h .




Figure 2 – Average annual significant wave height (m).

Distribution of mean annual significant wave height calculated by data (Chawla et al., 2013) is shown in Fig. 2. As seen, the maximum value of significant wave height does not exceed 5 m, while the height of real extreme wave can reach 20 m. The data in Fig 1 have a more complicated structure, due, for example, to the periods with strong wind along trajectories of tropical storms. Consequently, the calculations of distribution of real wave height should be done for shorter periods, i.e., for seasonal or monthly averaged data on significant wave heights.

10




Figure 3. – Wave height (m) with cumulative probability 10^{-7} for December-February.

In Fig. 3 the field of wave height with probability 10^{-7} calculated by the data (Chawla et al., 2013)], averaged for December-February, is shown. When comparing Figs 3 and 1 it is seen

15



that in mid-latitudes of the Northern Hemisphere wave heights become higher. In some areas appearance of extreme wave heights exceeding 16 m is possible. At the same time there are actually no extreme waves in the eastern part of the Arctic Ocean, which is connected with seasonal ice formation in the area. In equatorial and tropical areas of the World Ocean wave heights are less in winter (Northern Hemisphere), as compared with the average annual wave heights. It should be noted that in the western part of the Atlantic Ocean tracking trajectories of hurricanes disappeared while the number of such trajectories increased in the Indian Ocean.




Figure 4. – Wave height (m) with cumulative probability 10^{-7} averaged for March-May.

An increase of wave heights over March-May can be registered (Fig.4) in the Southern Hemisphere. Actually all the area of mid-latitudes from the latitude of 40 degrees S. to the latitude of 60 degrees S. is characterized by probability of wave heights over 14 m. In mid-latitudes of the Northern Hemisphere in spring wave height values are more than the average annual values, though less than the winter values, while in some areas (Atlantic Ocean near Iceland, Pacific Ocean near Bering Sea) appearance of waves exceeding 14 m in height is quite possible.




Fig.5 –Wave heights (m) with the cumulative probability of 10^{-7} , for June-August

Summer months (Fig.5) are characterized by general decrease of extreme wave probability. It is especially noticeable in the northern areas of the Atlantic and Pacific Oceans. Also, wave heights slightly decreased in the Southern Hemisphere. It should be noted that storm tracks appear off the eastern coast of North America and disappear in the southern part of the Pacific Ocean. Besides, quite distinct trajectories of storms appeared in the eastern part of the Pacific Ocean. Small wave heights can be observed in the Arctic Ocean, in the area free from ice.




Figure 6. – Wave heights (m) with cumulative probability 10^{-7} , averaged for September - November



During autumn months (Fig. 6) an increase of wave heights is observed in the Arctic Ocean, the extreme wave height values sometimes reaching 20 m. Among other peculiarities is an increase of wave-free area in polar latitudes of the Southern Hemisphere, which is obviously connected with seasonal ice formation.

5 It is quite evident that the average monthly wave fields with the cumulative probability will allow us to obtain more exact information on the areas of extreme wave probability.

The above approach can be used with different initial values of cumulative probability. It is not expedient to use the values less than 10^{-8} as this value is outside the approximation area (2). Mapping of wave heights with the cumulative probability exceeding 10^{-7} may have a certain
10 practical importance. Hence, on the whole, the method considered is more suitable for estimation of extreme values of wave heights with minor probability.

4. Conclusions

15 The paper describes a method of calculation of extreme wave probability, based on long-term statistical data on significant wave height. Such method can be used for estimation of probability of extreme waves, which is important for designing of engineering constructions, as well as for other purposes. The maps of global distribution of wave heights of certain probability for main seasons illustrate the method.

20

Authors are thankful to Professor D. Chalikov for his useful consultations.

The investigation was fulfilled with financial support of Russian Scientific Fund (Project No 16-17-00124).

25

References

- Chalikov D.: Freak waves: their occurrence and probability. *Phys of Fluid*, 2009 ,21, 076602; doi:10.1063/1.3175713
- Chalikov D., Babanin A., and Sanina E.: Modeling of Three-Dimensional Fully
30 Nonlinear Potential Periodic Waves, *Ocean Dynamics*, 2014, 64, 10,1469–1486. doi: 10.1007/s10236-014-0755-0.
- Chalikov D. and Babanin AV.: Comparison of linear and nonlinear extreme wave statistics, *Acta Oceanologica Cinica*, 2016, 35, 5, 99-105, doi:10.1007/313131-016-0862-5
- 35 Chalikov D.: Numerical modeling of sea waves. 2016, Springer, ISBN 978-3-319-32914-7 ISBN 978-3-319-32916-1 (eBook), doi 10.1007/978-3-319-32916-1, 330pp
- Chalikov D.V.: Linear and nonlinear statistics of extreme waves, *Russian Journal of Numerical Analysis and Mathematical Modelling*, 2017, 32, 2, 91–99.
- Chalikov D. and Bulgakov K.: Estimation of wave height probability based on the
40 statistics of significant wave height, *J. Ocean Eng. Mar. Energy*, 2017. <https://doi.org/10.1007/s40722-017-0093-7>
- Chawla A., Spindler D.M., and Tolman H.L.: Validation of a thirty year wave hindcast using the climate forecast system reanalysis winds, *Ocean Modelling*, 2013, 70, 189–206.
- Ochi M.K. *Ocean waves: the stochastic approach*, 2005, Cambridge university press,
45 ISBN-13 978-0-521-56378-9? ISBN-10 0-521-01767-X, 332 pp.
- Onorato M, T. Waseda T., T. Toffoli A, Cavaleri L, Gramstad O, Janssen PAEM, Kinoshita T, Monbaliu J, Mori N, Osborne AR, Serio M, Stansberg CT, Tamura H, and K. Trulsen.: K Statistical Properties of Directional Ocean Waves: The Role of the Modulational Instability in the Formation of Extreme Events, *Phys. Rev. Lett.*, 2009,
50 102, 114502



Tolman H.L., 2014. User manual and system documentation of WAVEWATCH III version 4.18. NOAA / NWS / NCEP / MMAB Technical Note 316, 194 pp.+ Appendices (pdf)