

Water Masses in the Atlantic Ocean: Characteristics and Distributions

Mian Liu^{1, 2}

Toste Tanhua^{2, *}

¹ *College of Ocean and Earth Sciences,
Xiamen University, Xiamen, 361005, China*

² *GEOMAR Helmholtz Centre for Ocean Research Kiel,
Marine Biogeochemistry, Chemical Oceanography
Düsternbrooker Weg 20, 24105 Kiel, Germany*

Correspondence to: T. Tanhua (ttanhua@geomar.de)

Abstract: A large number of water masses are presented in the Atlantic Ocean and knowledge of their distributions and properties are important for understanding and monitoring of a range of oceanographic phenomena. The characteristics and distributions of water masses in biogeochemical space are useful for, in particular, chemical and biological oceanography to understand the origin and mixing history of water samples. Here we define the characteristics of the major water masses in the Atlantic Ocean as Source Water Types (SWTs) from their formation areas, and map out their distributions. The SWTs are described by six properties taken from the biased adjusted data product GLODAPv2, including both conservative (Conservative Temperature and Absolute Salinity) and non-conservative (oxygen, silicate, phosphate and nitrate) properties. The distributions of these water masses are investigated with the use of the Optimal Multi-Parameter (OMP) method and mapped out. The Atlantic Ocean is divided into four vertical layers by distinct neutral densities and four zonal layers to guide the identification and characterization. The water masses in the upper layer originate from winter-time subduction and are defined as Central Waters. Below the upper layer, the intermediate layer consists of three main water masses; Antarctic Intermediate Water (AAIW), Subarctic Intermediate Water (SAIW) and Mediterranean Water (MW). The North Atlantic Deep Water (NADW, divided into its upper and lower components) is the dominating water mass in the deep and overflow layer. The origin of both the upper and lower NADW is the Labrador Sea Water (LSW), the Iceland-Scotland Overflow Water (ISOW) and the Denmark Strait Overflow Water (DSOW). The Antarctic Bottom Water (AABW) is the only natural water mass in the bottom layer and this water mass is redefined as North East Atlantic Bottom Water (NEABW) in the north of equator due to the change of key properties, especial silicate. Similar with NADW, two additional water masses, Circumpolar Deep Water (CDW) and Weddell Sea Bottom Water (WSBW), are defined in the Weddell Sea region in order to understand the origin of AABW.

Key Words: Atlantic Ocean, Water Mass, Source Water Types, GLODAP, Optimal-Multi-Parameter Analysis

1 Introduction

The ocean is composed a large number of water masses without clear boundaries but gradual transformations between each other (e.g. Castro et al., 1998). Properties of the water in the ocean are not uniformly distributed and the characteristics vary with regions and depths (or densities). The water masses, which are defined as bodies of water with similar properties and common formation history, are referred to as a body of water with a measurable extent both in the vertical and horizontal, and thus a quantifiable volume (e.g. Helland-Hansen, 1916; Montgomery, 1958). Mixing occurs inevitably between water masses, both along and across density surfaces, and result in mixtures with different properties away from their formation areas. Understanding of the distributions and variations of water masses have significance to several disciplines of oceanography, for instance while investigating the thermohaline circulation of the world ocean or predicting climate change (e.g. Haine and Hall, 2002; Tomczak and Godfrey, 2013; Morrison et al., 2015).

The concept of water masses is also important for biogeochemical and biological applications, where the transformations of properties over time can be successfully viewed in the water masses frame-work. For instance, the formation of Denmark Strait Overflow Water in the Denmark Strait was described using mixing of a large number of water masses from the Arctic Ocean and the Nordic Seas (Tanhua et al., 2005). A number of investigations show the significance of knowledge about water masses to the biogeochemical oceanography, for instance the investigation of mineralization of biogenic materials (Alvarez et al., 2014), or the change of ventilation in the Oxygen Minimum Zone (Karstensen et al., 2008). In a more recent work, Garcia-Ibanez et al. (2015) considered 14 water masses combined with velocity fields to estimate transport of water masses, and thus chemical constituents, in the north Atlantic. Similarly, Jullion et al. (2017) used water mass analysis in the Mediterranean Sea to better understand the dynamics of dissolved Barium. However, the lack of a unified definition of overview water masses on an basin or global scale leads to additional and repetitive amount of work by redefining water masses in specific regions. The goal of this study is to facilitate water mass analysis in the Atlantic Ocean and in particular, we aim at supporting biogeochemical and biological oceanographic work in a broad sense.

Understanding the formation, transformation, and circulation of water masses has been a research topic in oceanography since the 1920s (e.g. Jacobsen, 1927; Defant, 1929; Wüst and Defant, 1936; Sverdrup, 1942 etc.). The early studies were mainly based on (potential) temperature and (practical) salinity as summarized by Emery and Meincke (1986). The limitation of the analysis based on T—S relationship is obvious; distributions of more (than three) water masses cannot be analyzed at the same time with only these two parameters, so physical and chemical oceanographers has worked to add more parameters to the characterization of water masses (e.g. Tomczak and Large, 1989; Tomczak, 1981; 1999). The Optimum Multi-parameter (OMP) method extends the analysis so that more water masses can be considered by adding parameters/water properties (such as phosphate and silicate) and solving the equations of linear mixing without assumptions. The OMP analysis has been successfully applied in a

range of studies, for instance for the analysis of mixing in the thermocline in Eastern Indian Ocean (Poole and Tomczak, 1999).

An accurate definition and characterization is the prerequisite for the analysis of water masses. In this study, the concepts and definitions of water masses given by Tomczak (1999) are used and we seek to define the key properties of the main water masses in the Atlantic Ocean and to describe their distributions. In order to facilitate the analysis, the data product GLODAPv2 is used to identify and define the characteristics of the most prominent water masses based on 6 commonly measured physical and biogeochemical properties (Figure 1). The water masses are defined in a static sense, i.e. they are assumed to be steady and not change over time, and subtle differences between closely related water masses are not considered in this basin-scale focused study. The so defined water masses are in a subsequent step used to estimate their distributions in the Atlantic Ocean, again based on the GLODAPv2 data product. Detailed investigations on temporal variability of water masses, or their detailed formation processes, for instance, may find this study useful but will certainly want to use a more granular approach to water mass analysis in their particular areas.

2 Data and Methods

2.1 The GLODAPv2 data product

Oceanographic surveys conducted by different countries have been actively organized and coordinated since late 1950s. WOCE (the World Ocean Circulation Experiment), JGOFS (Joint Global Ocean Flux Study) and OACES (Ocean Atmosphere Carbon Exchange Study) are three typical representatives of international coordination in the 1990s. The GLODAP (Global Ocean Data Analysis Project) data product was devised and implemented in this context with the aim to create a global dataset suitable to describe the distribution and interior ocean inorganic carbon variables (Key et al., 2004; 2010). The first edition (GLODAPv1.1) contains data up to 1999 whereas the updated and expanded versions GLODAPv2 (Key et al., 2015; Olsen et al., 2016) was published in 2016 and the GLODAP team is striving for annual updates (Olsen et al., 2019; 2020). Since GLODAPv2 is a comprehensive and, more importantly, biased adjusted data product, this is used to quantify the characteristics of water masses. The data in the GLODAPv2 product has passed both a primary quality control (QC), aiming at precision of the data and unify the units, and a secondary quality control, aiming at the accuracy of the data (Tanhua et al., 2010). The GLODAPv2 data product is adjusted to correct for any biases in data through these QC routines and is unique in its internal consistency, and is thus an ideal product to use for this work. Armed with the internally consistent data in GLODAPv2, we utilize previously published studies on water masses and their formation areas to define areas and depth/density ranges that can be considered to be representative samples of a water masses.

The variables of Absolute Salinity (SA in g kg^{-1}), Conservative Temperature (CT in $^{\circ}\text{C}$) and neutral density (γ in kg m^{-3}), which consider the thermodynamic properties such as entropy, enthalpy and

chemical potential (Jackett et al., 2006; Groeskamp et al., 2016), are used in this study because they systematically reflects the spatial variation of seawater composition in the ocean, as well as the impact from dissolved neutral species on the density and provides a more conservative, actual and accurate description of seawater properties (Millero et al., 2008; Pawlowicz et al., 2011; Nycander et al., 2015).

2.2 Water Masses (WMs) and Source Water Types (SWTs)

In practice, defining properties of water masses (WMs) is often a difficult and time-consuming part, particularly when analyzing water masses in a region distant from their formation areas. Tomczak (1999) defined a water mass as “a body of water with a common formation history, having its origin in a particular region of the ocean” whereas Source Water Types (SWTs) describe “the original properties of water masses in their formation areas”. The distinction between the WMs and SWTs is that WMs define physical extents, i.e. a volume, while SWTs are only mathematical definitions, i.e. SWTs are defined values of properties without physical extents. Knowledge of the SWTs, on the other hand, is essential in labeling WMs, tracking their spreading or mixing progresses, since the values from SWTs describe their initial characteristics and can be considered as the fingerprints of WMs. The SWT of a WM is defined by the values of key properties, while some of them, like Central Waters, require more than one SWT to be defined (Tomczak, 1999). In this study, the terminology “water mass” is used in the discussions, realizing that the properties of the WMs used for the further analysis actually refer to SWTs.

2.3 OMP Analysis

2.3.1 Principle of OMP Analysis

For the analysis, six key properties are used to define SWTs, including two conservative (Conservative Temperature and Absolute Salinity) and four non-conservative (oxygen, silicate, phosphate and nitrate) properties. In order to determine the distributions of WMs, the OMP analysis is invoked as objective mathematical formulations of the influence of mixing (Karstensen and Tomczak, 1997; 1998). The starting point is the 6 key properties (Figure 1) from observations (such as CT_{obs} is the observed Conservative Temperature). The OMP model determines the contributions from predefined SWTs (such as CT_i that describes the Conservative Temperature in each SWT), which represent the values of the “unmixed” WMs in the formation areas, through a linear set of mixing equations, assuming that all key properties of water masses are affected similarly by the same mixing processes. The fractions (x_i) in each sampling point are obtained by finding the best linear mixing combination in parameter space defined by 6 key properties and minimizing the residuals (R , such as R_{CT} is the residual of Conservative Temperature) in a nonnegative least squares sense (Lawson and Hanson, 1974) as shown in the following equations:

$$x_1CT_1 + x_2CT_2 + \cdots + x_nCT_n = CT_{obs} + R_{CT}$$

$$x_1SA_1 + x_2SA_2 + \dots + x_nSA_n = SA_{obs} + R_{SA}$$

$$x_1O_1 + x_2O_2 + \dots + x_nO_n = O_{obs} + R_O$$

$$x_1Si_1 + x_2Si_2 + \dots + x_nSi_n = Si_{obs} + R_{Si}$$

$$x_1Ph_1 + x_2Ph_2 + \dots + x_nPh_n = Ph_{obs} + R_{Ph}$$

$$x_1N_1 + x_2N_2 + \dots + x_nN_n = N_{obs} + R_N$$

$$x_1 + x_2 + \dots + x_n = 1 + R$$

Where the CT_{obs} , SA_{obs} , O_{obs} , Si_{obs} , Ph_{obs} and N_{obs} are the observed values of properties, the CT_i , SA_i , O_i , Si_i , Ph_i and N_i ($i = 1, 2 \dots, n$) represent the predetermined (known) values in each SWT for each property. The last row expresses the condition of mass conservation.

OMP analysis represent an inversion of an overdetermined system in each sampling point, so that the sampling points are required to be located “downstream” from the formation areas, i.e. on the spreading pathway. The total number of WMs which can be analyzed simultaneously within one OMP run is limited by the number of variables/key properties, because mathematically, 6 variables ($x_1 - x_6$) can be solved with 6 equations. In our analysis, one OMP run can solve up to 6 WMs. The above system of equations can be written in matrix notation as:

$$\mathbf{G} \cdot \mathbf{x} - \mathbf{d} = \mathbf{R};$$

Where $\mathbf{G}$ is a parameter matrix of defined SWTs with 6 key properties, $\mathbf{x}$ is a vector containing the relative contributions from the “unmixed” water masses to the sample (i.e. solution vector of the SWT fractions), $\mathbf{d}$ is a data vector of water samples (observational data from GLODAPv2 in this study) and $\mathbf{R}$ is a vector of residual. The solution is to find the minimum of the residual ($\mathbf{R}$) with a linear fit of parameters (key properties) for each data point with a non-negative values. In this study, the mixed layer is not considered as its properties tend to be strongly variable on seasonal time-scales so that water mass analysis is inapplicable. The solution is dependent on, and sensitive to, the prior assumptions of the properties of the SWTs. Here we have not explicitly explored this sensitivity, but note that a common difficulty in OMP analysis is to properly define the SWT properties, and that this study provide a generally applicable set of SWT properties for the major water masses in the Atlantic Ocean.

2.3.2 Extended OMP Analysis

The prerequisite (or restriction) for using (basic) OMP analysis is that the water masses are formed close enough to the water samples with short transport times within a limited ocean region, for instance an oceanic front or intertidal belt, so that the mixing can be assumed not influenced by biogeochemical processes (i.e. assume all the parameters to be quasi-conservative). However, biogeochemical processes

cannot be ignored in a basin-scale analysis (Karstensen and Tomczak, 1998). Obviously, this prerequisite does not apply to our investigation for the entire Atlantic, so the “extended” OMP analysis is required. In this concept, non-conservative parameters (phosphate and nitrate) are converted into conservative parameters by introducing the “preformed” nutrients PO and NO, where PO and NO denotes the concentrations of phosphate and nitrate in seawater by considering the consumption of dissolved oxygen by respiration (in other words, the alteration due to respiration is eliminated) (Broecker, 1974; Karstensen and Tomczak, 1998). In addition, a new column should be added to the equations for non-conservative properties ($a\Delta O_2$, $a\Delta Si$, $a\Delta Ph$ and $a\Delta N$) to express the changes in SWTs due to biogeochemical impacts, namely, the change of oxygen concentration with the remineralization of nutrients:

$$x_1CT_1 + x_2CT_2 + \dots + x_nCT_n = CT_{obs} + R_{CT}$$

$$x_1SA_1 + x_2SA_2 + \dots + x_nSA_n = SA_{obs} + R_{SA}$$

$$x_1O_1 + x_2O_2 + \dots + x_nO_n - a\Delta O_2 = O_{obs} + R_O$$

$$x_1Si_1 + x_2Si_2 + \dots + x_nSi_n + a\Delta Si = Si_{obs} + R_{Si}$$

$$x_1Ph_1 + x_2Ph_2 + \dots + x_nPh_n + a\Delta Ph = Ph_{obs} + R_{Ph}$$

$$x_1N_1 + x_2N_2 + \dots + x_nN_n + a\Delta N = N_{obs} + R_N$$

$$x_1 + x_2 + \dots + x_n = 1 + R$$

As a result, the number of water masses should be further reduced in one OMP run if the biogeochemical processes are considered and extended OMP analysis is used. In this study, a total number of 5 water masses are included in each OMP run.

2.3.3 Presence of mass residual

The fractions of WMs in each sample are obtained by finding the best linear mixing combination in parameter space defined by 6 key properties which minimizes the residuals (R) in a non-negative least squares sense. Ideally, a value of 100% is expected when the fractions of all the water masses are added together. However, mass residuals where the sum of water masses for a sample differ from 100%, is inevitable during the analysis and is due to sample properties outside the input SWTs to the OMP formulation. There are two different cases. The first is that one single water mass is larger than 100% and other water masses are all 0%. This mostly happens in the Central Waters ($\gamma < 27.10 \text{ kg m}^{-3}$, Figure 2). The reason is that key properties, for instance CT, of Central Waters are variable. When the CT increases beyond the range of this water mass, the OMP analysis considers the fraction is over 100%. In this case, all such samples are set to 100% after confirming the absence of any other water mass. The second case is that none of each single water mass is more than 100%, but the total fraction is more than

100% when added together. In this study, the total fractions are generally less than 105% ($\gamma > 27.10 \text{ kg m}^{-3}$, Figure 2).

In order to map the distributions of water masses, all GLODAPv2 data in the Atlantic Ocean (below the mixed layer) are analyzed with the OMP method by using 6 key properties. In order to solve the contradiction between the limitation of water masses in one OMP run and the total number of 16 water masses (Figure 3), the Atlantic Ocean is divided into 17 regions (Table 1) and each with its own OMP formulation, by only including water masses that are likely to appear in the area. In the vertical, neutral density intervals are used to separate boxes. In the horizontal direction, the division lines are 40°N , the equator and 50°S where the area south of 50°S is one region, independent of density, and additional divisions are set between equator and 40°N (γ at 26.70 and 27.30 kg m^{-3} , latitude of 30°N , Table 1). In this way, we end up with a set of 17 different OMP formulations that are used for estimating the fraction(s) of water masses in each water sample. The neutral density and the latitude of the water sample are thus used to determine which OMP should be applied (Table 1). Note that all water masses are present in more than one OMP so that reasonable (i.e. smooth) transitions between the different areas can be realized.

3 Overview of the water masses in the Atlantic Ocean and the Criteria of Selection

In line with the results from Emery and Meincke (1986) and from our interpretation of the observational data from GLODAPv2, the water masses in the Atlantic Ocean are considered to be distributed in four main isopycnal (vertical) layers separated by surfaces of equal (neutral) density (Figure 4). The upper (shallowest) layer with lowest neutral density is located within upper $\sim 500\text{--}1000 \text{ m}$ of the water column (below the mixed layer and $\gamma < 27.10 \text{ kg m}^{-3}$). The intermediate layer is located between ~ 1000 and 2000 m (γ between 27.10 and 27.90 kg m^{-3}). The deep and overflow layer occupies the layer between $\sim 2000\text{--}4000 \text{ m}$ (γ between 27.90 and 28.10 kg m^{-3}) whereas the bottom layer is the deepest layer and mostly located below $\sim 4000 \text{ m}$ ($\gamma > 28.10 \text{ kg m}^{-3}$).

To define the main water masses in the Atlantic Ocean, the determination of their formation areas is the first step (Figure 5) and then the selection criteria are listed to define SWTs based on the CT—SA distribution, pressure (P) or neutral density (γ) (Table 2). For some SWTs, additional properties such oxygen or silicate are also required for the definition. With these criteria, which are taken from the literature and also based on data from GLODAPv2 product, the SWTs of all the main water masses can be defined for further estimating their distributions in the Atlantic Ocean by using OMP analysis.

For the water masses in the upper layer, i.e. the Central Waters, properties cover a “wide” range instead of a “narrow” point value due to their variations, especially in CT and SA space i.e. The Central Waters are labeled by two SWTs to identify the upper and lower boundaries of properties (Karstenson and Tomczak, 1997; 1998). In order to determine these two SWTs, one property is taken as a benchmark

(neutral density in this investigation) and the relationships to the others are plotted to make a linear fit and the two endpoints are selected as SWTs to label Central Waters (Figure 6).

During the determination of each SWT, two figures are displayed to characterize them, including a) depth profiles of the 6 key properties under consideration (same color coding), and b) bar plots from the distributions of the samples within the criteria (the blue dots in Figure 6 and 7) for a SWT with a Gaussian curve to show the statistics (Figure 7). The plots of properties vs. pressure provides an intuitive understanding of each SWT compared to other WMs in the region. The distributions of properties with the Gaussian curves are the basis to visually determine and confirm the SWT property values and associated standard deviations.

Most water masses maintain their original characteristics away from their formation areas. However, some are worthy to be mentioned as products from mixing of several original water masses (for instance, North Atlantic Deep Water is the product from Labrador Sea Water, Iceland-Scotland Overflow Water and Denmark Strait Overflow Water). Also, characteristics of some water masses changes sharply during their pathways (namely, the sharp drop silicate concentration of Antarctic Bottom Water after passing the equator). As a result, it is advantageous to redefine their SWTs. In order to distinguish such water masses from the other original ones, their defined specific areas are mentioned as “redefining” areas instead of formation areas, because, strictly speaking, they are not “formed” in these areas.

4 The Upper Layer, Central Waters

The upper layer is occupied by four Central Waters known formed by winter subduction with upper and lower boundaries of properties. All values between these boundaries are used to calculate the means and standard deviations (Figure 7 and Figure 1—4 in Supplement), and occupies two SWTs in one OMP run.

Central Waters can be easily recognized by their linear CT—SA relationships (Pollard et al., 1996; Stramma and England, 1999). In this study, the upper layer is defined to be located above the neutral density isoline of 27.10 kg m^{-3} (below the mixed layer). The formations and transports of the Central Waters are influenced by the currents in the upper layer and finally form relative distinct bodies of water in both the horizontal and vertical directions (Figure 8). The concept of Mode Water is referred to as the sub-regions of Central Water, which describes the particularly uniform properties of seawater within the upper layer and more refers to the physical properties (such as: CT—SA relationship and potential vorticity). In this study, the unified name “Central Water”, which more refers to the biogeochemical properties (Cianca et al., 2009; Alvarez et al., 2014), is used to avoid possible confusions.

4.1 Eastern North Atlantic Central Water (ENACW)

The main Central Water in the region east of the Mid-Atlantic-Ridge (MAR) is the East North Atlantic Central Water (ENACW, Harvey, 1982). This water mass is formed in the inter-gyre region during the winter subduction (Pollard and Pu, 1985). One component of the Subpolar Mode Water (SPMW) is

carried south and contributes to the properties of ENACW (McCartney and Talley, 1982). The intergyre region limited by latitudes between 39 and 48 °N and longitudes between 20 and 35 °W (Pollard et al., 1996) is considered as the formation area of ENACW (Figure 5). Neutral densities of 26.50 and 27.30 kg m⁻³ are selected as the upper and lower boundaries to define the SWT of ENACW (Cianca et al., 2009; Prieto et al., 2015), which is in contrast to Garcia-Ibanez et al. (2015) that used potential temperature (θ) as the upper limit. The core of ENACW is located within the upper 500 m of the water column (Figure 7, a) with the iconic linear T-S relationship (Figure 6, b) consistent with Pollard et al. (1996). The main character of ENACW is the large ranges of temperature and salinity and low nutrient concentrations, especially silicate (Figure 7, b).

4.2 Western North Atlantic Central Water (WNACW)

Western North Atlantic Central Water (WNACW) is another water mass formed through winter subduction (Worthington, 1959; McCartney and Talley, 1982) with the formation area at the southern flank of the Gulf Stream (Klein and Hogg, 1996). In some studies, this water mass is referred to as 18 °C water since a temperature of around 18 °C is one symbolic feature (e.g. Talley and Raymer, 1982; Klein and Hogg, 1996). In general, seawater in the Northeast Atlantic has higher salinity than in the Northwest Atlantic due to the stronger winter convection (Pollard and Pu, 1985) and input of MW (Pollard et al., 1996; Prieto et al., 2015). However, for the Central Waters, the situation is the opposite. WNACW has a significantly higher salinity (SA) by ~0.9 g kg⁻¹ than ENACW (Table 4). In this study, the work from McCartney and Talley (1982) is followed and the region 24—37°N, 50—70°W shallower than 500 m is considered as the formation area (Figure 5). By defining the SWT of WNACW, the neutral density between 26.20 and 26.70 kg m⁻³ is selected due to the discrete CT—SA distribution outside this range (Table 2). Besides the linear CT—SA relationship, another property of this water mass is, as the alternative name suggests, a temperature of around 18 °C, which is the warmest in the four Central Waters due to the low latitude of the formation area and the impact from the warm Gulf Stream (Cianca et al., 2009; Prieto et al., 2015). In addition, low nutrient is also a significant property compared to other Central Waters (Figure 2 in Supplement).

4.3 Eastern South Atlantic Central Water (ESACW)

The formation area of ESACW is located in area southwest of South Africa and south of the Benguela Current (Peterson and Stramma, 1991). In this region the Agulhas Current brings water from the Indian Ocean (Deruijter, 1982; Lutjeharms and van Ballegooyen, 1988) that mixes with the South Atlantic Current from the west (Stramma and Peterson, 1990; Gordon et al., 1992). The origin of ESACW can partly be tracked back to the WSACW, but defined as a new SWT since seawater from Indian Ocean is added by the Agulhas Current. The mixing region of Agulhas Current and South Atlantic Current (30—40 °S, 0—20 °E) is selected as the formation area of ESACW (Figure 5). To investigate the properties of ESACW, results from Stramma and England (1999) is followed and consider 200—700m as the core of this water mass. For the properties, neutral density (γ) between 26.00 and 27.00 kg m⁻³ and oxygen

concentration higher than $230 \mu\text{mol kg}^{-1}$ are used to define ESACW (Table 2). Similar as ENACW, ESACW also exhibits relative large CT and SA ranges and low nutrient concentrations (especially low in silicate) compared to the AAIW below. The properties in ESACW are similar to that of WSACW, although with higher nutrient concentrations due to input from the Agulhas current (Figure 3 in Supplement).

4.4 Western South Atlantic Central Water (WSACW)

The WSACW is formed in the region near the South American coast between 30 and 45 °S, where surface South Atlantic Current brings Central Water to the east (Kuhlbrodt et al., 2007). The WSACW is formed with little directly influence from other Central Water mass (Sprintall and Tomczak, 1993; Stramma and England, 1999), while the origin of other Central Waters (e.g. ESACW or ENACW) can be traced back, to some extent at least, to WSACW (Peterson and Stramma, 1991). This water mass is a product of three Mode Waters mixed together: the Brazil current brings Salinity Maximum Water (SMW) and Subtropical Mode Water (STMW) from the north, while the Falkland Current brings Subantarctic Mode Water (SAMW) from the south (Alvarez et al., 2014). Here we follow the work of Stramma and England (1999) and Alvarez et al. (2014) that choose the meeting region of these two currents (25–60 °W, 30–45 °S) as the formation area of WSACW (Figure 5). Neutral density (γ) between 26.0 and 27.0 kg m^{-3} is selected to define the SWT of WSACW and the requirement of silicate concentrations lower than $5 \mu\text{mol kg}^{-1}$ and oxygen concentrations lower than $230 \mu\text{mol kg}^{-1}$ is also added (Table 2). WSACW shows the similar hydrochemical properties to other Central Waters such as linear T-S relationship with large T and S ranges and low concentration of nutrients, especially silicate (Figure 4 in Supplement).

4.5 Atlantic Distribution of Central Waters

Based on the OMP analysis of the GLODAPv2 data product, the physical extent of the Central Waters can be described over the Atlantic Ocean. The horizontal distributions of four Central Waters in the upper layer are shown on the maps in Figure 8 and the vertical distributions along selected GO-SHIP sections are found in Figure 9. Note that the Central Waters are found at different densities, the eastern variations being denser, so that there is significant overlap in the horizontal distribution. The vertical extent of the Central Waters is clearly seen in Figure 9.

The ENACW is mainly found in the northeast part of North Atlantic, near the formation area in the intergyre region (Figure 8). High fractions of ENACW is also found in a band across the Atlantic at around 40 °N, where the core of this water mass is found at close to 1000 m depth in the western part of the basin (Figure 9).

The WNACW is predominantly found in the western basin of the North Atlantic in a zonal band between ~10 °N and 40 °N (Figure 8). The vertical extent of WNACW is significantly higher in the western basin with an extent of about 500 meter in the west, tapering off towards the east (Figure 9).

The ESACW is found over most of the South Atlantic, as well as in the tropical and subtropical north Atlantic (Figure 8). The extent of ESACW do reach particular far north in the eastern part of the basin where it is an important component over the Eastern Tropical North Atlantic Oxygen Minimum Zone, roughly south of the Cape Verde Islands. In the vertical direction, the ESACW is located below WSACW (Figure 9).

The horizontal distribution of the WSACW does reach into the northern hemisphere but is, obviously, concentrated in the western basin (Figure 8). In the vertical scale, the WSACW also tends to dominate the upper layer of the South Atlantic above the ESACW (Figure 9).

5 The Intermediate Layer

The intermediate water masses have an origin in the upper 500m of the ocean and subduct into the intermediate depth (1000—1500m) during their formation process. Similar to the Central Waters, the distributions of the Intermediate Waters are significantly influenced by the major currents (Figure 10, left). The neutral density (γ) of the Intermediate Waters is in general between 27.10 and 27.90 kg m⁻³ and selected as the definition of Intermediate Layer.

In the Atlantic Ocean, two main intermediate water masses, Subarctic Intermediate Water (SAIW) and Antarctic Intermediate Water (AAIW), are formed in the surface of sub-polar regions in north and south hemisphere, respectively. In addition to AAIW and SAIW, Mediterranean Water (MW) is also considered as an intermediate water mass due to the similarity in density ranges, although the formation history is different (Figure 10).

5.1 Antarctic Intermediate Water (AAIW)

The Antarctic Intermediate Water (AAIW) is the main Intermediate Water in the South Atlantic Ocean. This water mass originates from the surface region north of the Antarctic Circumpolar Current (ACC) in all three sectors of the Southern Ocean, in particular in the area east of the Drake Passage in the Atlantic sector (McCartney, 1982; Alvarez et al., 2014), then subducts and spreads northward along the continental slope of South America (Piola and Gordon, 1989).

Based on the work by Stramma and England (1999) and Saenko and Weaver (2001), the region between 55 and 40 °S (east of the Drake Passage) at depths below 100 m is selected as the formation area of AAIW as well as the primary stage during the subduction and transformation (Figure 5). Previous work is considered to distinguish AAIW from surrounding water masses, including SACW in the north and NADW in the deep. Piola and Georgi (1982) and Talley (1996) define AAIW as potential densities (σ_θ) between 27.00/27.10 and 27.40 kg m⁻³ and Stramma and England (1999) define the boundary between AAIW and SACW at $\sigma_\theta = 27.00$ kg m⁻³ and the boundary between AAIW and NADW at $\sigma_1 = 32.15$ kg m⁻³. The following criteria are used as selection criteria to define AAIW: neutral density between 26.95 and 27.50 kg m⁻³ and depth between 100 and 300 m. In addition, high oxygen (> 260 $\mu\text{mol kg}^{-1}$) and low temperature (CT < 3.5 °C) are used to distinguish AAIW from Central Waters (WSACW and

ESACW), while the relative low silicate concentration ($< 30 \mu\text{mol kg}^{-1}$) of AAIW is an additional boundary to differentiate AAIW from AABW (Table 2). The AAIW is distributed across most of the Atlantic Ocean up to $\sim 30^\circ \text{N}$ and the water mass fraction shows a decreasing trend towards the north (Kirchner et al., 2009). AAIW is found at depths between 500 and 1200 m (Talley, 1996) with the two significant characteristic features of low salinity and high oxygen concentration (Figure 5 in Supplement, Stramma and England, 1999).

5.2 Subarctic Intermediate Water (SAIW)

The Subarctic Intermediate Water (SAIW) originates from the surface layer in the western boundary of the North Atlantic Subpolar Gyre, along the Labrador Current (Lazier and Wright, 1993; Pickart et al., 1997). This water mass subducts and spreads southeast in the region north of the NAC, advects across the Mid-Atlantic-Ridge and finally interacts with MW (Arhan, 1990; Arhan and King, 1995). The formation of SAIW is a mixture of two surface sources: Water with high temperature and salinity carried by the NAC and cold and fresh water from the Labrador Current (Read, 2000; Garcia-Ibanez et al., 2015). In Garcia-Ibanez et al. (2015), there are two definitions of SAIW, SAIW₆, which is biased to the warmer and saltier NAC, and SAIW₄, which is closer to the cooler and fresher Labrador Current. Here, only the combination of these two end-members is considered on the whole-Atlantic Ocean scale.

For defining the spatial boundaries we followed Arhan (1990) and selected the region between 35 and 55°W and 50 and 60°N , i.e. the region along the Labrador Current and north of the NAC as the formation area of SAIW (Figure 5). Within this area, neutral densities higher than 27.65 kg m^{-3} and CT higher than 4.5°C is selected to define SAIW by following Read, (2000). Samples in the depth range from the MLD to 500 m are considered as the core layer of SAIW, which included the formation and subduction of SAIW (Table 2).

5.3 Mediterranean Water (MW)

The predecessor of the Mediterranean Water (MW) is the Mediterranean Overflow Water (MOW) flowing out through the Strait of Gibraltar, whose main component is the modified Levantine Intermediate Water. This water mass is recognized by high salinity and temperature and intermediate neutral density in the Northeast Atlantic Ocean (Carracedo et al., 2016). After passing the Strait of Gibraltar, the MOW mixes rapidly with the overlying ENACW and forms the MW, leading to a sharp decrease of salinity (Baringer and Price, 1997). In Gulf of Cadiz, the outflow of MW turns into two branches: One branch continues to the west, descending the continental slope, mixing with surrounding water masses in the intermediate depth and influence the water mass composition as far west as the MAR (Price et al., 1993). The other branch spreads northwards along the coast of Iberian Peninsula and along the European coast and its influence can be observed as far north as the Norwegian Sea (Reid, 1978; 1979). The impact from MW is significant in almost the entire Northeast Atlantic in the Intermediate Layer (east of the MAR, Figure 7 in Supplement), with high temperature and Salinity but low nutrients compared to other water masses.

Here we followed Baringer and Price (1997) and define the SWT of MW by the high salinity (SA between 36.5 and 37.00 g kg⁻¹, Table 2) samples in the formation area west of the Strait of Gibraltar (Figure 5).

5.4 Atlantic Distributions of Intermediate Waters

A schematic of the main currents in the intermediate layer (γ between 27.10 and 27.90 kg m⁻³) is shown in Figure 10 (left panel).

The SAIW is mainly formed north of 30 °N in the western basin by mixing of two main sources, the warmer and saltier NAC and the colder and fresher Labrador Current and characterized with relative low CT (< 4.5 °C), SA (< 35.1 g kg⁻¹) and silicate (< 11 μ mol kg⁻¹). The SAIW and MW can be easily distinguished by the OMP analysis due to significantly different properties. The meridional distributions of three Intermediate Waters along the A16 section are shown in Figure 10 (upper panel) as well as the zonal distributions of SAIW and MOW along the A03 section. A “blob” of MW centered around 35°N can be seen to separate the AAIW from the SAIW in the eastern North Atlantic. The fractions of SAIW in the western basin are definitely higher (Figure 10, right).

The MW enters the Atlantic from Strait of Gibraltar and spreads in two branches to the north and the west. MW is mainly formed close to its entry point to the Atlantic, near the Gulf of Cadiz, with low fractions in the western North Atlantic. The distribution of MW can be seen as roughly following the two intermediate pathways following two branches (Figure 10, left): One spreads to the north into the West European Basin until ~50°N, while the other branch spreads in a westward direction past the MAR (Figure 11), mainly at latitudes between 30 and 40 °N. The density of MW is higher than SAIW, and the distributions of the two water masses are complementary in the North Atlantic (Figure 10, right).

The AAIW has a southern origin and is found at slightly lighter densities (core neutral density ~27.20 kg m⁻³, Figure 10, right) compared to SAIW and MW. The AAIW is formed in the region south of 40 °S where it sinks and spreads to the north at depth between ~1000 and 2000 m with neutral densities between 27.10 and 27.90 kg m⁻³. The AAIW is the dominate Intermediate Water in the South Atlantic and it is clear that the AAIW represents a reduction of fractions during the pathway to the north with only a diluted part to be found the equator and 30 °N (Figure 10 and Figure 11).

6 The Deep and Overflow Layer

The Deep and Overflow Waters are found roughly between 2000 to 4000 m with neutral densities between 27.90 and 28.10 kg m⁻³. These water masses play an indispensable role in the Atlantic Meridional Overturning Circulation (AMOC). The source region of these waters is confined to the North Atlantic with their formation region either south of the Greenland-Scotland ridge, or in the Labrador Sea (Figure 5 and Figure 12, left). The Denmark Strait Overflow Water (DSOW) and the Iceland-Scotland Overflow water (ISOW) originate from Arctic Ocean and the Nordic Seas and enter the North Atlantic

through either the Denmark Strait of the Faroe Bank Channel (Figure 12, left). In the North Atlantic, these two water masses descend, mainly following the topography meet and mix in the Irminger Basin (Stramma et al., 2004; Tanhua et al., 2005) and form the bulk of the lower North Atlantic Deep Water (LNADW) (Read, 2000; Rhein et al., 2011). The Labrador Sea Water (LSW) is formed through winter deep convection in the Labrador and Irminger Seas, and makes up the bulk of the upper North Atlantic Deep Water (uNADW). Due to intense mixing processes the LSW, DSOW and ISOW are defined as the water masses in north of 40 °N whereas south of this latitude the presence of the two variations of NADW are considered (Figure 12, right).

In south of 40 °N, both variations of the NADW spread south mainly with the Deep Western Boundary Current (DWBC, Figure 12, left) (Dengler et al., 2004) through the Atlantic until ~50 °S where they meet the Antarctic Circumpolar Current (ACC). During the southward transport, the NADW also spreads significantly in the zonal direction (Lozier, 2012), so that the distribution of NADW covers mostly the whole Atlantic basin in the Deep and Overflow Layer (Figure 12, right). The southward flow of NADW is also an indispensable component of Atlantic Meridional Overturning Circulation (AMOC) (Broecker and Denton, 1989; Elliot et al., 2002; Lynch-Stieglitz et al., 2007).

6.1 Labrador Sea Water (LSW)

As an important water mass that contributes to the formation of North Atlantic Deep Water (NADW), Labrador Sea Water (LSW) is predominant in mid-depth (between 1000m and 2500m depth) in the Labrador Sea region (Elliot et al., 2002). This water mass was firstly noted by (Wüst and Defant, 1936) due to its salinity minimum and later defined and named by Smith et al. (1937). The LSW is formed by deep convection during the winter and is typically found at depth with $\sigma_\theta = \sim 27.77 \text{ kg m}^{-3}$ (Clarke and Gascard, 1983). Since then the character has been identified as a contribution to the driving mechanism of northward heat transport in the Atlantic Meridional Overturning Circulation (AMOC) (Rhein et al., 2011). The LSW is characterized by relative low salinity (lower than 34.9) and high oxygen concentration ($\sim 290 \mu\text{mol kg}^{-1}$) (Talley & McCartney, 1982). Another important criterion of LSW is the potential density (σ_θ), that ranges from 27.68 to 27.88 kg m^{-3} (Clarke and Gascard, 1983; Gascard and Clarke, 1983; Stramma et al., 2004; Kieke et al., 2006). In the large spatial scale, LSW can be considered as one water mass (Dickson and Brown, 1994), however significant differences of different “vintages” of LSW exist (Stramma et al., 2004; Kieke et al., 2006). In some references, this water mass is also broadly divided into upper Labrador Sea Water (uLSW) and classic Labrador Sea Water (cLSW) with the boundary between them at potential density of 27.74 kg m^{-3} (Smethie and Fine, 2001, Kieke et al., 2006; 2007). The LSW is considered as the main origin of the upper NADW (Talley and McCartney, 1982; Elliot et al., 2002).

On the basis of the above work, the formation area of LSW is selected to include the Labrador Sea region between the Labrador Peninsular and Greenland and parts of the Irminger Basin (Figure 5). The neutral density (γ) between 27.70 to 28.10 kg m^{-3} as well as $\text{CT} < 4^\circ\text{C}$ are used to define SWT of LSW

(Table 2) by considering Clarke and Giscard (1983) and Stramma and England (1999) with the depth range of 500-2000m (Elliot et al., 2002). Trademark characteristics of LSW are relative low salinity and high oxygen. The relatively large spread in properties is indicative of the different “vintages” of LSW, in particular the bi-modal distribution of density, and partly for oxygen (Figure 8 in the Supplement).

6.2 Iceland-Scotland Overflow Water (ISOW)

The Iceland Scotland Overflow Water (ISOW) flows close to the bottom from the Iceland Sea to the North Atlantic in the region east of Iceland, mainly through the Faroe-Bank Channel (Swift, 1984; Lacan et al., 2004; Zou et al., 2020). ISOW turn into two main branches when passing the Charlie-Gibbs Fracture Zone (CGFZ), with the first one flowing through the Mid-Atlantic-Ridge, into the Irminger basin, where it meets and mixes with DSOW (Figure 12). The other branch is transported southward and mixes with Northeast Atlantic Bottom Water (NEABW) (Garcia-Ibanez et al., 2015). ISOW is characterized by high nutrient and low oxygen concentration and its pathway closely follows the Mid-Atlantic-Ridge in the Iceland Basin. The following criteria, Conservative Temperature between 2.2 and 3.3 °C and Absolute Salinity higher than 34.95 g kg⁻¹, are used to define the SWT of ISOW, and neutral density higher than 28.00 kg m⁻³ is added order to distinguish ISOW from LSW in the region west of MAR (Table 2 and Figure 9 in Supplement).

6.3 Denmark Strait Overflow Water (DSOW)

A number of water masses from the Arctic Ocean and the Nordic Seas flows through Denmark Strait west of Iceland. At the sill of the Denmark Strait and during the descent into the Irminger Sea, these water masses undergo intense mixing. This overflow water mass is considered as the coldest and densest component of the sea water in the Northwest Atlantic Ocean and constitute a significant part of the southward flowing NADW (Swift, 1980). Samples from the Irminger Sea (Figure 5) with neutral density higher than 28.15 kg m⁻³ (Table 2 and Figure 10 in Supplement) are used for the definition of DSOW (Rudels et al., 2002; Tanhua et al., 2005).

6.4. Upper North Atlantic Deep Water (uNADW)

The uNADW is mainly formed by mixing of ISOW and LSW and considered as a distinct water mass south of the Labrador Sea as this region is identified as the redefining area of upper and lower NADW (Dickson and Brown, 1994). The region between latitude 40 and 50 °N, west of the MAR is selected as the redefining area of NADW (Figure 5) and the criteria of neutral density between 27.85 and 28.05 kg m⁻³ and CT < 4.0 °C within the depth range from 1200 to 2000 m (Table 2 and Figure 11 in Supplement) are used to define the SWT of uNADW (Stramma et al., 2004). As a mixture from LSW and ISOW, the uNADW obviously inherits many properties from LSW, but is also significantly influenced by the ISOW. The relative high temperature (~3.3 °C) is a significant feature of the uNADW together with relatively low oxygen (~280 µmol kg⁻¹) and high nutrient concentrations, which is a universal symbol of deep water (Table 4).

6.5. Lower North Atlantic Deep Water (LNADW)

The same geographic region is selected as the formation area of LNADW (Figure 5). In this region, the ISOW and DSOW, influenced by LSW, mix with each other and form the lower portion of NADW (Stramma et al., 2004). Water samples between depths of 2000 and 3000 m with CT higher than $\sim 2.5^{\circ}\text{C}$ and neutral densities between 27.95 and 28.10 kg m^{-3} are selected to define the SWT of LNADW (Table 2 and Figure 12 in Supplement).

6.6. Atlantic Distributions of Deep and Overflow Waters

The water masses dominate the neutral density interval 27.90 – 28.10 kg m^{-3} in the Atlantic Ocean north of 40°N are Labrador Sea Water (LSW), Iceland-Scotland Overflow Water (ISOW) and Denmark Strait Overflow Water (DSOW). In the region south of 40°N the upper and lower NADW, considered as products from these three original overflow water masses, dominate the deep and overflow layer (Figure 12).

The LSW is commonly characterized as two variations, “upper” and “classic” although in this study we consider this as one water mass in the discussion above. Our analysis indicates that the LSW dominates the North West Atlantic Ocean in the characteristic density range. In Figure 12, we choose to display $\gamma = 27.95$ that corresponds to the main property of the LSW (Kieke et al., 2006; 2007). The LSW spreads east and southward in the North Atlantic Ocean, but is less dominant in the area west of the Iberian Peninsula where the presence of MW from the Gulf of Cadiz tends to dominate that density level. Note that although the LSW is slightly denser than the MW, their density ranges do overlap (Figure 12 and 13).

The ISOW is mainly found in the Northeast Atlantic north 40°N between Iceland and Iberian Peninsula with core at $\gamma = \sim 28.05 \text{ kg m}^{-3}$. The ISOW is also found west of the Reykjanes Ridge, in the Irminger and Labrador Seas between the DSOW and LSW (Figure 12 and 13).

The DSOW is mainly found in the Irminger and Labrador Seas as the densest layer close to the bottom (Figure 11). Our analysis indicates a weak contribution of DSOW also east of the MAR. South of the Grand Banks the DSOW is already significantly diluted and only low to moderate fractions are found (Figure 12 and 13).

After passing 40°N , the upper and lower NADW are considered as independent water masses and dominate the most of the Atlantic Ocean in this density layer. The map in Figure 12 shows that upper NADW covers the most area, while the lower NADW is found mainly found in the west region near the Deep Western Boundary Current (DWBC), especially in South Atlantic. In the vertical view based on sections (Figure 14), the southward transports of both upper and lower NADW can be seen until $\sim 50^{\circ}\text{S}$ where they meet AABW in the ACC region.

7. The Bottom Layer and the Southern water masses

The Bottom Waters are defined as the densest water masses that occupy the lowest layers of the water column, typically below 4000 m depth and with neutral densities higher than 28.10 kg m^{-3} . These water masses have an origin in the Southern Ocean (Figure 15, left) and are characterized by their high silicate concentrations. The Antarctic Bottom Water (AABW) is the main water mass in the Bottom Layer (Figure 15, right). This water mass is formed in the Weddell Sea region, south of Antarctic Circumpolar Current (ACC) through mixing of Circumpolar Deep Water (CDW) and Weddell Sea Bottom Water (WSBW) (van Heuven et al., 2011). After the formation, AABW spreads to the north across the equator and further northwards until $\sim 40^\circ \text{N}$ (Figure 16), where a new SWT is redefined as North East Atlantic Bottom Water (NEABW) due to the drastic change in properties (sharp decrease in silicate concentration). As the two main sources of AABW, CDW and WSBW are confined to the Southern Ocean (Figure 15, right), so they are referred as the southern water masses and discussed in this section together with bottom waters.

7.1. Antarctic Bottom Water (AABW)

Antarctic Bottom Water (AABW) is the symbolic Bottom Water in the whole Atlantic Ocean. As one of the important components in Atlantic Meridional Overturning Circulation (AMOC), AABW spreads northward below 4000m depth, mainly west of Mid-Atlantic-Ridge (MAR, Figure 15, right) and plays a significant role in the Thermohaline Circulation (Rhein et al., 1998; Andri  et al., 2003). The origin of AABW in Atlantic section can be traced back to the Weddell Sea as a product of mixing of Weddell Sea Bottom Water (WSBW) and Circumpolar Deep Water (CDW) (Foldvik and Gammelsrod, 1988; Alvarez et al., 2014).

The definition of AABW is all water samples formed south of the Antarctic Circumpolar Current (ACC), i.e. south of 63°S in the Weddell Sea (Figure 5), with neutral density (γ) larger than 28.27 kg m^{-3} (Weiss et al., 1979; Orsi et al., 1999). As an additional constraint, AABW is defined as water samples with silicate higher than $120 \mu\text{mol kg}^{-1}$ to distinguish from other water masses in this region as high silicate is a trade mark property of AABW (Table 2).

The formation process of AABW is a mixture of another two original water masses, CDW and WSBW, which are referred to as southern water masses, in the Weddell Sea region, consistent with Orsi et al. (1999) and van Heuven et al. (2011). The CDW, with relative warm temperature ($CT > 0.4^\circ \text{C}$), is advected with the ACC from the north, while the extremely cold Shelf Water ($CT < -0.7^\circ \text{C}$) comes as Weddell Sea Bottom Water (WSBW) from the south (Figure 17). AABW is found from 1000m to 5500m depth (Figure 16 and 17) with low temperature ($CT < 0^\circ \text{C}$), salinity ($SA < 34.68$) but high nutrient, especially silicate, concentrations (Figure 13 in Supplement).

7.2. Northeast Atlantic Bottom Water (NEABW)

Northeast Atlantic Bottom Water (NEABW), also called lower Northeast Atlantic Deep Water (lNEADW, Garcia-Ibanez et al., 2015), is mainly found below 4000m depth in the eastern basin of the

North Atlantic (Figure 5). This water mass is an extension of AABW during the way to the north, since the properties of AABW change significantly on the slow transport north. A new SWT is redefined for this water mass in north of the Equator, similar as the redefinition of NADW south of the Labrador Sea.

The region east of the MAR and between the equator and 30 °N, i.e. before NEABW enters the Iberian Basin, is selected as the redefining area of NEABW. The criteria of depth deeper than 4000 m and CT above 1.8 °C are also used (Table 2). In the CT—SA diagram in Figure 3, similar T—S distribution between NEABW and AABW can be seen but with higher CT and SA of ~1.95 °C and ~35.060 g kg⁻¹. Most NEABW samples have a neutral density higher than 28.10 kg m⁻³ and NEABW is characterized by low CT and SA, but high silicate concentration (Figure 14 in Supplement). This further suggests that NEABW originates from AABW, although most properties have been changed significantly from the origin in the South Atlantic.

7.3. Circumpolar Deep Water (CDW)

Circumpolar Deep Water (CDW), which has significance to the thermohaline circulation during the wind-driven upwelling in the Southern Ocean (Morrison et al., 2015), is the lighter of the two water masses contribute to AABW formation. The production of this water mass can be tracked to the southward flow of NADW and the large-scale mixing in the Antarctic Circumpolar Current (ACC) region (van Heuven et al., 2011). At about 50°S, NADW is deflected upward by AABW before reaching the ACC (Figure 14, upper panel), this part of NADW spreads further southward into the ACC region, where it contacts with other water masses, including AAIW above and AABW below. After passing the ACC region, CDW splits into two branches at ~60 °S. The upper branch is upwelled and partly joint into the AAIW, while the rest spreads towards the coastal region, mixes with the cold fresh shelf water, sinks to the bottom and finally forms the Weddell Sea Bottom Water (WSBW), which is another contribution to the AABW (Marshall and Speer, 2012; Abernathey et al., 2016). The lower branch sinks and mixes with the WSBW below and contributes to the formation of AABW.

In this study, the SWTs of CDW is defined by considering the lower branch and the region between 55 and 65 °S is selected as the formation area (Figure 5). To define SWT of CDW (lower branch), water samples are selected from depth between 200 and 1000m in this region and additional constraints are SA higher than 34.82 g kg⁻¹ and CT between -0.5 and 1.0 °C (Table 2). Similar to other bottom/southern SWTs, CDW is also defined by high nutrient (silicate, phosphate and nitrate) and low oxygen concentrations (Figure 15 in Supplement).

7.4 Weddell Sea Bottom Water (WSBW)

The Weddell Sea Bottom Water (WSBW) is the densest water mass in the bottom layer. As mentioned in the above section, part of CDW from the upper branch cools down rapidly by mixing with extremely cold shelf water and sinks down to the bottom along the continental slop (Gordon, 2001). WSBW is formed in the Weddell Sea basin below the depth of 3000m before it meets and mixes with CDW. The

low temperature of WSBW compared to CDW ($CT = \sim -0.8\text{ }^{\circ}\text{C}$) is a characteristic property (Figure 16 in Supplement, van Heuven et al., 2011).

Water samples in the latitudinal boundaries of $55 - 65\text{ }^{\circ}\text{S}$ in the Weddell Sea (Figure 5) with pressures larger than 3000 m and CT lower than $-0.7\text{ }^{\circ}\text{C}$ and silicate higher than $105\text{ }\mu\text{mol kg}^{-1}$ are selected to define the SWT of WSBW (Table 2), following Gordon (2001) and van Heuven et al. (2011).

7.5. Atlantic Distribution of the bottom waters and southern water masses

AABW and NEABW dominate the bottom layer ($\gamma > 28.10\text{ kg m}^{-3}$). From the horizontal distribution (Figure 15) it can be seen that AABW and NEABW cover the most bottom area of South and North Atlantic respectively. In fact, both water masses have the same origin but are distinguished by redefining a new SWT due to the sharp reduction of silicate after passing the equator (Figure 16). The AABW is formed in the Weddell Sea region south of the Antarctic Circumpolar Current (ACC). After leaving the formation area, AABW sinks to the bottom due to the high density during the way north. After passing the ACC, AABW suffers from water exchange with NADW between $50\text{ }^{\circ}\text{S}$ and the equator (van Heuven et al., 2011). Similar to AABW, NEABW also mainly contacts with lower NADW and its origin (ISOW) in the North Atlantic (Garcia-Ibanez et al., 2015).

In the Weddell Sea region, distributions of water masses mainly reflect the formation process of AABW as displayed based on SR04 sections (Figure 17). In the zonal section, AABW can be seen as the mixture of CDW and WSBW, where the core of CDW distributes in the upper 1000m and WSBW originates from the surface and sinks along the continental slope into the bottom below 4000m. Both original water masses meet each other at depth between ~ 2000 and 4000m, where AABW is formed, with main core located at ~ 3000 m. The meridional section shows the northward outflow of AABW into the Atlantic Ocean. AABW is located between 2000 and 4000m as a product from CDW and WSBW. After leaving Weddell Sea region, AABW is considered as an independent water mass and spreads further northward as the only bottom water mass until the equator.

8. Conclusions and Discussion

The characteristics of the main water masses in their formation areas are defined in a 6-dimensional hydro-chemical space in the Atlantic Ocean. The values of properties for these water masses form the fundamental basis to investigate their transport, distribution and mixing and referred to as SWTs. Table 4 and Figure 3 provides an overview of SWTs of all the 16 Atlantic Ocean main water masses considered in this study. The distribution of water masses are estimated by using OMP analysis based on the GLODAPv2 data product, and preliminarily divided into four vertical layers based on neutral densities.

The upper layer, which covers the most shallow layer (typically down to about 500 m depth) of the ocean below the mixed layer (the mixed layer is not considered in this analysis), is occupied by Central Waters. The intermediate layer is situated between the upper layer and the deep and overflow layer at roughly 1000 to 2000m depth. Of the three water masses in this layer, AAIW and SAIW are both characterized by relative low salinity and temperature, while the MW have high SA and CT. The SAIW and MW show a Northwest-Southeast distribution in the North Atlantic, while the AAIW dominates the intermediate layer of the region south of 30 °N. In the deep and overflow layer between roughly 2000 and 4000m, NADW, which contains upper and lower portions, is recognized as the dominant water mass with a relative complex origin from LSW, ISOW and DSOW. The bottom layer is occupied by AABW with a southern origin formed by CDW and WSBW. After passing the equator, this water mass is redefined as NEABW due to the changes in properties (silicate).

Besides the 16 main Atlantic Ocean water masses, additional water masses still exist and can be found in the Atlantic that cannot be explained by the mixing of any above listed original water masses. This tends to happen close to the coast by local oceanographic events, such water masses are not listed and considered as main water mass in this study, and no additional SWTs are defined. For instance, in the coastal region of Southern Benguela Upwelling System (15 – 20 °E, 30 – 34 °S), water samples are found with low temperature and oxygen (CT = ~8 °C, oxygen = ~150 $\mu\text{mol kg}^{-1}$). This cannot be explained by the mixing of ESACW and WSACW, which are the only two possible water masses in this region and depth, because the CT and oxygen of both water masses are higher than these values. One possible explanation is that low-oxygen water transported by upwelling (Flynn et al., 2020).

The here presented characteristics (property values and the standard deviations) of Atlantic Ocean water masses and their distributions are intended to guide water mass analysis of hydrographic data and expect to provide a basis for further biogeochemical research.

Acknowledgements

This work is based on the comprehensive and detailed data from GLODAP data set throughout the past few decades. In particular, we are grateful to the efforts from all the scientists and crews on cruises, who generated funding and dedicated time on committing the collection of data. We also would like to thank the working groups of GLODAP for their support and information of the collation, quality control and publishing of data. Their contributions and selfless sharing are prerequisites for the completion of this work. We are grateful to Johannes Karstensen for his support and advices in running OMP programs and to Marcus Dengler for the information and suggestions on physical oceanography during the writing process. We are very thankful for the ground breaking research by the late Matthias Tomczek that made this work possible, and for the constructive comments on a previous version of this manuscript. Thanks

694 goes to the China Scholarship Council (CSC) for providing the funding support to Mian Liu's PhD
695 study in GEOMAR Helmholtz Centre for Ocean Research Kiel. Thanks to Prof. Minggang Cai and all
696 the colleagues in the research group of Marine Organic Chemistry (MOC) for the help and support
697 during Mian Liu's post-doctoral work in College of Ocean and Earth Sciences, Xiamen University.

698

699 **References**

- 700 Abernathey, R. P., Cerovecki, I., Holland, P. R., Newsom E., Mazloff, M. and Talley, L. D.: Water-mass
701 transformation by sea ice in the upper branch of the southern ocean overturning, *Nat. Geosci.*, 9, 596–
702 601, doi:10.1038/ngeo2749, 2016.
- 703 Alvarez, M., Brea, S., Mercier, H., Alvarez-Salgado, X.A.: Mineralization of biogenic materials in the
704 water masses of the South Atlantic Ocean. I: Assessment and results of an optimum multiparameter
705 analysis. *Prog Oceanogr* 123, 1-23, 2014.
- 706 Andri , C., Gouriou, Y., Bourl s, B., Ternon, J.F., Braga, E.S., Morin, P., Oudot, C.: Variability of AABW
707 properties in the equatorial channel at 35 W. *Geophysical Research Letters* 30, n/a-n/a, 2003.
- 708 Arhan, M.: The North Atlantic current and subarctic intermediate water. *J Mar Res* 48, 109-144, 1990.
- 709 Arhan, M., King, B.: Lateral Mixing of the Mediterranean Water in the Eastern North-Atlantic. *J Mar*
710 *Res* 53, 865-895, 1995.
- 711 Baringer, M.O., Price, J.F.: Mixing and spreading of the Mediterranean outflow. *Journal of Physical*
712 *Oceanography* 27, 1654-1677, 1997.
- 713 Broecker, W.S.: No a Conservative Water-Mass Tracer. *Earth Planet Sc Lett* 23, 100-107, 1974.
- 714 Broecker, W.S., Denton, G.H.: The Role of Ocean-Atmosphere Reorganizations in Glacial Cycles.
715 *Geochimica Et Cosmochimica Acta* 53, 2465-2501, 1989.
- 716 Carracedo, L.I., Pardo, P.C., Flecha, S., and P rez, F.F.: On the Mediterranean Water Composition. *J*
717 *Phys Oceanogr.*, 46, 1339–1358, 2016.
- 718 Castro, C.G., Perez, F.F., Holley, S.E., Rios, A.F.: Chemical characterisation and modelling of water
719 masses in the Northeast Atlantic. *Prog Oceanogr* 41, 249-279, 1998.
- 720 Cianca, A., Santana, R., Marrero, J., Rueda, M., Llin s, O.: Modal composition of the central water in
721 the North Atlantic subtropical gyre. *Ocean Science Discussions* 6, 2487-2506, 2009.
- 722 Clarke, R.A., Gascard, J.-C.: The Formation of Labrador Sea Water. Part I: Large-Scale Processes. *Journal*
723 *of Physical Oceanography* 13, 1764-1778, 1983.
- 724 Defant, A.: *Dynamische Ozeanographie*. Springer, 1929.
- 725 Dengler, M., Schott, F.A., Eden, C., Brandt, P., Fischer, J., Zantopp, R.J.: Break-up of the Atlantic deep
726 western boundary current into eddies at 8  S. *Nature* 432, 1018, 2004.
- 727 Deruijter, W.: Asymptotic Analysis of the Agulhas and Brazil Current Systems. *Journal of Physical*
728 *Oceanography* 12, 361-373, 1982.
- 729 Dickson, R.R., Brown, J.: The Production of North-Atlantic Deep-Water - Sources, Rates, and Pathways.
730 *J Geophys Res-Oceans* 99, 12319-12341, 1994.
- 731 Elliot, M., Labeyrie, L., Duplessy, J.C.: Changes in North Atlantic deep-water formation associated with
732 the Dansgaard-Oeschger temperature oscillations (60-10 ka). *Quaternary Science Reviews* 21, 1153-
733 1165, 2002.

734 Emery, W.J., Meincke, J.: Global Water Masses - Summary and Review. *Oceanologica Acta* 9, 383-391,
735 1986.

736 Foldvik, A., Gammelsrod, T.: Notes on Southern-Ocean Hydrography, Sea-Ice and Bottom Water
737 Formation. *Palaeogeography Palaeoclimatology Palaeoecology* 67, 3-17, 1988.

738 Flynn R.F., Granger J., Veitch J.A. Siedlecki, S., Burger, J. M., Pillay, K., Fawcett, S. E.: On-Shelf Nutrient
739 Trapping Enhances the Fertility of the Southern Benguela Upwelling System. *Journal of Geophysical*
740 *Research-Oceans* 125, 2020.

741 Garcia-Ibanez, M.I., Pardo, P.C., Carracedo, L.I., Mercier, H., Lherminier, P., Rios, A.F., Perez, F.F.:
742 Structure, transports and transformations of the water masses in the Atlantic Subpolar Gyre. *Prog*
743 *Oceanogr* 135, 18-36, 2015.

744 Gascard, J.-C., Clarke, R.A.: The Formation of Labrador Sea Water. Part II. Mesoscale and Smaller-Scale
745 Processes. *Journal of Physical Oceanography* 13, 1779-1797, 1983.

746 Gordon, A.L., Weiss, R.F., Smethie, W.M., Warner, M.J.: Thermocline and Intermediate Water
747 Communication between the South-Atlantic and Indian Oceans. *J Geophys Res-Oceans* 97, 7223-7240,
748 1992.

749 Gordon, A.L.: Bottom Water Formation. In: Steele JH, ed. *Encyclopedia of Ocean Sciences*. Oxford:
750 Academic Press, 334-40, 2001.

751 Groeskamp, S., Abernathey, R.P., Klocker, A.: Water mass transformation by cabbeling and
752 thermobaricity. *Geophysical Research Letters* 43, 10835-45, 2016.

753 Haine, T.W.N., Hall, T.M.: A generalized transport theory: Water-mass composition and age. *Journal of*
754 *Physical Oceanography* 32, 1932-1946, 2002.

755 Helland-Hansen, B.r.: *Nogen hydrografiske metoder*. Scand. Naturforsker Mote. Kristiana. Oslo, 1916.

756 Harvey, J.: Theta-S Relationships and Water Masses in the Eastern North-Atlantic. *Deep-Sea Research*
757 *Part a-Oceanographic Research Papers* 29, 1021-1033, 1982.

758 Jackett D. R., McDougall T., Feistel R., Wright D., Griffies S.: Algorithms for Density, Potential
759 Temperature, Conservative Temperature, and the Freezing Temperature of Seawater. *Journal of*
760 *Atmospheric and Oceanic Technology* 23, 1706-28, 2006.

761 Jacobsen, J. P.: Line graphische Methode zur Bestimmung des Vermischungskoeffizienten im Meer.
762 *Gerlands Beitrage zur Geophysik*, 16, 404-412, 1927.

763 Jullion, L., Jacquet, S., Tanhua, T.: Untangling biogeochemical processes from the impact of ocean
764 circulation: First insight on the Mediterranean dissolved barium dynamics. *Global Biogeochemical*
765 *Cycles* 31, 1256-1270, 2017.

766 Karstensen, J., Tomczak, M.: Ventilation processes and water mass ages in the thermocline of the
767 southeast Indian Ocean. *Geophysical Research Letters* 24, 2777-2780, 1997.

768 Karstensen, J., Tomczak, M.: Age determination of mixed water masses using CFC and oxygen data. *J*
769 *Geophys Res-Oceans* 103, 18599-18609, 1998.

770 Karstensen J., Stramma L., Visbeck M.: Oxygen minimum zones in the eastern tropical Atlantic and
771 Pacific oceans. *Progress in Oceanography* 77, 331-50, 2008.

772 Key, R.M., Kozyr, A., Sabine, C.L., Lee, K., Wanninkhof, R., Bullister, J.L., Feely, R.A., Millero, F.J., Mordy,
773 C., Peng, T.H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP).
774 Global biogeochemical cycles 18, 2004.

775 Key, R.M., Tanhua, T., Olsen, A., Hoppema, M., Jutterström, S., Schirnick, C., van Heuven, S., Kozyr, A.,
776 Lin, X., Velo, A., Wallace, D.W.R., Mintrop, L.: The CARINA data synthesis project: introduction and
777 overview. *Earth Syst. Sci. Data* 2, 105-121, 2010.

778 Key, R.M., A. Olsen, S. van Heuven, S. K. Lauvset, A. Velo, X. Lin, C. Schirnick, A. Kozyr, T. Tanhua, M.
779 Hoppema, S. Jutterström, R. Steinfeldt, E. Jeansson, M. Ishi, F. F. Perez, and T. Suzuki: Global Ocean
780 Data Analysis Project, Version 2 (GLODAPv2), ORNL/CDIAC-162, ND-P093. Carbon Dioxide Information
781 Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee.
782 doi:10.3334/CDIAC/OTG.NDP093_GLODAPv2, 2015

783 Kieke, D., Rhein, M., Stramma, L., Smethie, W.M., Bullister, J.L., LeBel, D.A.: Changes in the pool of
784 Labrador Sea Water in the subpolar North Atlantic. *Geophysical Research Letters* 34, 2007.

785 Kieke, D., Rhein, M., Stramma, L., Smethie, W.M., LeBel, D.A., Zenk, W.: Changes in the CFC inventories
786 and formation rates of Upper Labrador Sea Water, 1997-2001. *Journal of Physical Oceanography* 36,
787 64-86, 2006.

788 Kirchner, K., Rhein, M., HuttI-Kabus, S., Boning, C.W.: On the spreading of South Atlantic Water into
789 the Northern Hemisphere. *J Geophys Res-Oceans* 114, 2009.

790 Klein, B., Hogg, N.: On the variability of 18 Degree Water formation as observed from moored
791 instruments at 55 degrees W. *Deep-Sea Research Part I-Oceanographic Research Papers* 43, 1777-&,
792 1996.

793 Kuhlbrodt, T., Griesel, A., Montoya, M., Levermann, A., Hofmann, M., Rahmstorf, S.: On the driving
794 processes of the Atlantic meridional overturning circulation. *Reviews of Geophysics* 45, 2007.

795 Lacan, F., Jeandel, C.: Neodymium isotopic composition and rare earth element concentrations in the
796 deep and intermediate Nordic Seas: Constraints on the Iceland Scotland Overflow Water signature.
797 *Geochemistry Geophysics Geosystems* 5, 2004.

798 Lawson, C.L. and Hanson, R.J.: *Solving Least Squares Problems*. Prentice-Hall, New York, 1974.

799 Lazier, J.R.N., Wright, D.G.: Annual Velocity Variations in the Labrador Current. *Journal of Physical*
800 *Oceanography* 23, 659-678, 1993.

801 Lozier, M.S.: Overturning in the North Atlantic. *Ann Rev Mar Sci* 4, 291-315, 2012.

802 Lutjeharms, J.R., van Ballegooyen, R.C.: Anomalous upstream retroflexion in the agulhas current.
803 *Science* 240, 1770, 1988.

804 Lynch-Stieglitz, J., Adkins, J.F., Curry, W.B., Dokken, T., Hall, I.R., Herguera, J.C., Hirschi, J.J., Ivanova,
805 E.V., Kissel, C., Marchal, O., Marchitto, T.M., McCave, I.N., McManus, J.F., Mulitza, S., Ninnemann, U.,
806 Peeters, F., Yu, E.F., Zahn, R.: Atlantic meridional overturning circulation during the Last Glacial
807 Maximum. *Science* 316, 66-69, 2007.

808 Marshall, J., Speer, K.: Closure of the meridional overturning circulation through Southern Ocean
809 upwelling. *Nat. Geosci.* 5: 171-180, 2012.

810 McCartney, M.S.: The subtropical recirculation of Mode Waters. *J Mar Res* 40, 427-464, 1982.

811 McCartney, M.S., Talley, L.D.: The subpolar mode water of the North Atlantic Ocean. *Journal of Physical*
812 *Oceanography* 12, 1169-1188, 1982.

813 Millero F.J., Feistel R., Wright D.G., McDougall T.J.: The composition of Standard Seawater and the
814 definition of the Reference-Composition Salinity Scale. *Deep-Sea Research Part I-Oceanographic*
815 *Research Papers* 55, 50-72, 2008.

816 Montgomery, R.B.: Water characteristics of Atlantic Ocean and of world ocean. *Deep Sea Research*
817 (1953) 5, 134-148, 1958.

818 Morrison, A.K., Frulich, T.L., Sarmiento, J.L.: Upwelling in the Southern Ocean. *Phys. Today*. 68: 27-
819 32, 2015.

820 Nycander, J., Hieronymus, M. and Roquet, F.: The nonlinear equation of state of sea water and the
821 global water mass distribution, *Geophys. Res. Lett.*, 42(18), 7714–7721, doi:10.1002/2015GL065525,
822 2015.

823 Olsen, A., Key, R.M., Heuven, S.V., Lauvset, S.K., Velo, A., Lin, X., Schirnack, C., Kozyr, A., Tanhua, T.,
824 Hoppema, M.: The Global Ocean Data Analysis Project version 2 (GLODAPv2) - an internally consistent
825 data product for the world ocean. *Earth System Science Data* 71, 300-308, 2016.

826 Olsen, A., Lange, N., Key, R. M., Tanhua, T., Álvarez, M., Becker, S., Bittig, H. C., Carter, B. R., Cotrim da
827 Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E., Jones, S. D., Jutterström,
828 S., Karlsen, M. K., Kozyr, A., Lauvset, S. K., Lo Monaco, C., Murata, A., Pérez, F. F., Pfeil, B., Schirnack, C.,
829 Steinfeldt, R., Suzuki, T., Telszewski, M., Tilbrook, B., Velo, A., and Wanninkhof, R.: Global Ocean Data
830 Analysis Project, version 2.2019 (GLODAPv2.2019), NOAA National Centers for Environmental
831 Information, <https://doi.org/10.25921/xnmewr20>, 2019.

832 Olsen, A., Lange, N., Key, R. M., Tanhua, T., Bittig, H. C., Kozyr, A., Álvarez, M., Azetsu-Scott, K., Becker,
833 S., Brown, P. J., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M.,
834 Jeansson, E., Jutterström, S., Landa, C. S., Lauvset, S. K., Michaelis, P., Murata, A., Pérez, F. F., Pfeil, B.,
835 Schirnack, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Velo, A., Wanninkhof, R., and Woosley, R. J.:
836 GLODAPv2.2020 – the second update of GLODAPv2, *Earth Syst. Sci. Data Discuss.*,
837 <https://doi.org/10.5194/essd-2020-165>, in review, 2020.

838 Orsi, A.H., Johnson, G.C., Bullister, J.L.: Circulation, mixing, and production of Antarctic Bottom Water.
839 *Prog Oceanogr* 43, 55-109, 1999.

840 Pawlowicz R., Wright D.G., Millero F.J.: The effects of biogeochemical processes on oceanic
841 conductivity/salinity/density relationships and the characterization of real seawater. *Ocean Science* 7,
842 363-87, 2011.

843 Peterson, R.G., Stramma, L.: Upper-Level Circulation in the South-Atlantic Ocean. *Prog Oceanogr* 26,
844 1-73, 1991.

845 Pickart, R.S., Spall, M.A., Lazier, J.R.N.: Mid-depth ventilation in the western boundary current system
846 of the sub-polar gyre. *Deep-Sea Research Part I-Oceanographic Research Papers* 44, 1025-+, 1997.

847 Piola, A.R., Georgi, D.T.: Circumpolar properties of Antarctic intermediate water and Subantarctic
848 Mode Water. *Deep Sea Research Part A. Oceanographic Research Papers* 29, 687-711, 1982.

849 Piola, A.R., Gordon, A.L.: Intermediate Waters in the Southwest South-Atlantic. *Deep-Sea Research*
850 *Part a-Oceanographic Research Papers* 36, 1-16, 1989.

- 851 Pollard, R.T., Pu, S.: Structure and Circulation of the Upper Atlantic Ocean Northeast of the Azores.
852 Prog Oceanogr 14, 443-462, 1985.
- 853 Pollard, R.T., Griffths, M.J., Cunningham, S.A., Read, J.F., Pérez, F.F., Ríos, A.F.: Vivaldi 1991 – a study
854 of the formation, circulation and ventilation of Eastern North Atlantic Central Water. Progress in
855 Oceanography 37, 167–192, 1996.
- 856 Poole, R., Tomczak, M.: Optimum multiparameter analysis of the water mass structure in the Atlantic
857 Ocean thermocline. Deep-Sea Research Part I-Oceanographic Research Papers 46, 1895-1921, 1999.
- 858 Price, J.F., Baringer, M.O., Lueck, R.G., Johnson, G.C., Ambar, I., Parrilla, G., Cantos, A., Kennelly, M.A.,
859 Sanford, T.B.: Mediterranean outflow mixing and dynamics. Science 259, 1277-1282, 1993.
- 860 Prieto, E., Gonzalez-Pola, C., Lavin, A., Holliday, N.P.: Interannual variability of the northwestern Iberia
861 deep ocean: Response to large-scale North Atlantic forcing. J Geophys Res-Oceans 120, 832-847, 2015.
- 862 Read, J.: CONVEX-91: water masses and circulation of the Northeast Atlantic subpolar gyre. Prog
863 Oceanogr 48, 461-510, 2000.
- 864 Reid, J.L.: On the middepth circulation and salinity field in the North Atlantic Ocean. Journal of
865 Geophysical Research: Oceans 83, 5063-5067, 1978.
- 866 Reid, J.L.: On the contribution of the Mediterranean Sea outflow to the Norwegian-Greenland Sea.
867 Deep Sea Research Part A. Oceanographic Research Papers 26, 1199-1223, 1979.
- 868 Rhein, M., Kieke, D., Huttel-Kabus, S., Roessler, A., Mertens, C., Meissner, R., Klein, B., Boning, C.W.,
869 Yashayaev, I.: Deep water formation, the subpolar gyre, and the meridional overturning circulation in
870 the subpolar North Atlantic. Deep-Sea Research Part II-Topical Studies in Oceanography 58, 1819-1832,
871 2011.
- 872 Rhein, M., Stramma, L., Krahnemann, G.: The spreading of Antarctic bottom water in the tropical Atlantic.
873 Deep-Sea Research Part I-Oceanographic Research Papers 45, 507-527, 1998.
- 874 Rudels, B., Fahrbach, E., Meincke, J., Budéus, G., Eriksson, P.: The East Greenland Current and its
875 contribution to the Denmark Strait overflow. ICES Journal of Marine Science 59, 1133-1154, 2002.
- 876 Saenko, O. A., and Weaver, A. J.: Importance of wind-driven sea ice motion for the formation of
877 antarctic intermediate water in a global climate model, Geophys. Res. Lett., 28(21), 4147–4150,
878 doi:10.1029/2001GL013632, 2001.
- 879 Smethie, W.M., Fine, R.A.: Rates of North Atlantic Deep Water formation calculated from
880 chlorofluorocarbon inventories. Deep-Sea Research Part I-Oceanographic Research Papers 48, 189-215,
881 2001.
- 882 Smith, E.H., Soule, F.M., Mosby, O.: The Marion and General Greene Expeditions to Davis Strait and
883 Labrador Sea, Under Direction of the United States Coast Guard: 1928-1931-1933-1934-1935:
884 Scientific Results, Part 2: Physical Oceanography. US Government Printing Office, 1937.
- 885 Sprintall, J., Tomczak, M.: On the formation of Central Water and thermocline ventilation in the
886 southern hemisphere. Deep Sea Research Part I: Oceanographic Research Papers 40, 827-848, 1993.
- 887 Stramma, L., England, M.H.: On the water masses and mean circulation of the South Atlantic Ocean. J
888 Geophys Res-Oceans 104, 20863-20883, 1999.

889 Stramma, L., Kieke, D., Rhein, M., Schott, F., Yashayaev, I., Koltermann, K.P.: Deep water changes at
890 the western boundary of the subpolar North Atlantic during 1996 to 2001. *Deep Sea Research Part I:*
891 *Oceanographic Research Papers* 51, 1033-1056, 2004.

892 Stramma, L., Peterson, R.G.: The South-Atlantic Current. *Journal of Physical Oceanography* 20, 846-859,
893 1990.

894 Sverdrup: *The Oceans: Their Physics, Chemistry and General Biology*, 1942.

895 Swift, J.H.: The Circulation of the Denmark Strait and Iceland Scotland Overflow Waters in the North-
896 Atlantic. *Deep-Sea Research Part a-Oceanographic Research Papers* 31, 1339-1355, 1984.

897 Swift, S.M.: Activity patterns of pipistrelle bats (*Pipistrellus pipistrellus*) in north - east Scotland.
898 *Journal of Zoology* 190, 285-295, 1980.

899 Talley, L.: Antarctic intermediate water in the South Atlantic, *The South Atlantic*. Springer, pp. 219-238,
900 1996.

901 Talley, L., Raymer, M.: Eighteen degree water variability. *J. Mar. Res* 40, 757-775, 1982.

902 Talley, L.D., McCartney, M.S.: Distribution and Circulation of Labrador Sea-Water. *Journal of Physical*
903 *Oceanography* 12, 1189-1205, 1982.

904 Tanhua, T., Olsson, K.A., Jeansson, E.: Formation of Denmark Strait overflow water and its hydro-
905 chemical composition. *Journal of Marine Systems* 57, 264-288, 2005.

906 Tanhua, T., van Heuven S., Key R. M., Velo A., Olsen A., and Schirnack C.: Quality control procedures
907 and methods of the CARINA database, *Earth Syst. Sci. Data*, 2(1), 35-49, 2010.

908 Tomczak, M.: A multi-parameter extension of temperature/salinity diagram techniques for the analysis
909 of non-isopycnal mixing. *Prog Oceanogr* 10, 147-171, 1981.

910 Tomczak, M.: Some historical, theoretical and applied aspects of quantitative water mass analysis. *J*
911 *Mar Res* 57, 275-303, 1999.

912 Tomczak, M., Godfrey, J.S.: *Regional oceanography: an introduction*. Elsevier, 2013.

913 Tomczak, M., Large, D.G.: Optimum multiparameter analysis of mixing in the thermocline of the
914 eastern Indian Ocean. *Journal of Geophysical Research: Oceans* 94, 16141-16149, 1989.

915 van Heuven, S.M.A.C., Hoppema, M., Huhn, O., Slagter, H.A., de Baar, H.J.W.: Direct observation of
916 increasing CO₂ in the Weddell Gyre along the Prime Meridian during 1973–2008. *Deep Sea Research*
917 *Part II: Topical Studies in Oceanography* 58, 2613-2635, 2011.

918 Weiss, R.F., Ostlund, H.G., Craig, H.: *Geochemical Studies of the Weddell Sea*. *Deep-Sea Research Part*
919 *a-Oceanographic Research Papers* 26, 1093-1120, 1979.

920 Worthington, L.: The 18 water in the Sargasso Sea. *Deep Sea Research* (1953) 5, 297-305, 1959.

921 Wüst, G., Defant, A.: *Atlas zur Schichtung und Zirkulation des Atlantischen Ozeans: Schnitte und Karten*
922 *von Temperatur, Salzgehalt und Dichte*. W. de Gruyter, 1936.

923 Zou S.J., Bower A., Furey H., Lozier M.S., Xu X.B.: Redrawing the Iceland-Scotland Overflow Water
924 pathways in the North Atlantic. *Nature Communications* 11, 2020.




Figure 1: The Atlantic distribution of key properties required by the OMP analysis along the A16 section as occupied in 2013 (Expocode: 33RO20130803 in North Atlantic & 33RO20131223 in South Atlantic)
The dashed lines show the neutral densities at 27.10, 27.90 and 28.10 kg m⁻³.




Figure 2: An example of a mass conservation residual in OMP analysis for the A03 section. This figure indicate that in density layers outside of the water masses included in the analysis, we find a high residual, i.e. the OMP analysis should only be used for a certain density interval.

926




Figure 3: T - S diagram of all Atlantic data from the GLODAPv2 data product (gray dots) indicating the 16 main SWTs in the Atlantic Ocean discussed in this study. The colored dots with letters A--D show the upper and lower boundaries of Central Waters, and E--P show the mean values of other SWTs.

927




Figure 4: Left panel: Distributions of water masses in the Atlantic Ocean based on the A16 section in 2013, see Figure 1. The background color shows the Absolute Salinity (g kg^{-1}). The dashed lines show the boundary of the four vertical layers divided by Neutral Density.

Right panel: 5 selected WOCE/GO-SHIP sections that are selected in this work to represent the vertical distribution of the main water masses.




Figure 5: Formation/Redefining areas of the 16 main water masses in the Atlantic Ocean.

The red dots show stations in formation area and the blue dots stations that where the SWT was found, and the grey dots show all the stations from GLODAPv2 dataset.




Figure 6: Example of a selection of water samples to define a water mass (here ENACW):

Panel a) the formation area, **b)** the T-S diagram

The red dots show all the data in formation area, the blue dots show the selected data as ENACW and the grey dots show all the data from GLODAPv2 dataset.

Panel c) Six key Properties vs Neutral Density (γ) as independent variable. Blue dots show the selected data as ENACW from Panel a) and b) and the red line shows the linear fit of blue dots. The start and end points of the red line are the upper and lower boundaries of ENACW.





Figure 7: Example of the definition of a SWTs (here ENACW):

Panel a) the distribution of key properties vs. pressure, **b)** bar plots of the data distribution of samples used to define the SWTs. Conservative Temperature ($^{\circ}\text{C}$), Absolute Salinity (g kg^{-1}), Neutral Density (kg m^{-3}), Oxygen and Nutrients ($\mu\text{mol kg}^{-1}$)
The red Gaussian fit shows mean value and standard deviation of selected data.




Figure 8: Currents (left) and Water Masses (right) in the Upper Layer.

Left: The warm (red) and cold (blue) currents (arrows) and the formation areas (rectangular shadows) of water masses in the Upper Layer.

Right: The colored dots show fractions from 20% to 100% of water masses in each station around core neutral densities (kg m^{-3}). Stations with fractions less than 20% are marked by black dots while gray dots show the all the GLODAPv2 stations.




Figure 9: Distribution of Central Water Masses based on A16 (upper), A03 (middle), A10 (lower) sections for the top 3000 m depth.

The contour lines show fractions of 20% 50% and 80%, blue lines show cross section of other cruises, yellow dashed lines show the boundaries of vertical water columns layers (neutral density at 27.10, 27.90 and 28.10 kg m⁻³)




Figure 10: Currents (left) and Water Masses (right) in the Intermediate Layer

Left: The currents (arrows) and the formation areas (rectangular shadows) of water masses in the Intermediate Layer.

Right: The colored dots show fractions from 20% to 100% of water masses in each station around core neutral densities (kg m^{-3}).

Stations with fractions less than 20% are marked by black dots while gray dots show all the GLODAPv2 stations.




Figure 11: Distribution of Water Masses in the Intermediate Layer based on A16 (upper) and A03 (lower) sections. Contour lines show fractions of 20% 50% and 80%, blue lines show cross section of other cruises, yellow dashed lines show the boundaries of vertical water columns layers (neutral density at 27.10, 27.90 and 28.10 kg m⁻³)

937




Figure 12: Currents (left) and Water Masses (right) in the Deep and Overflow Layer

Left: The currents (arrows) and the formation areas (rectangular shadows) of water masses in the Deep and Overflow Layer.

Right: The colored dots show fractions (from 20% to 100%) of water masses in each station around core neutral density (kg m^{-3}).

Stations with fractions less than 20% are marked by black dots while gray dots show all the GLODAPv2 stations.




Figure 13: Distribution of SAIW (upper left), LSW (upper right), ISOW (lower left) and DSOW (lower right) based on the A25 section.

Contour lines show fractions of 20% 50% and 80% and yellow dashed lines show the boundaries of vertical water columns layers (neutral density at 27.10, 27.90 and 28.10 kg m^{-3})

939

940




Figure 14: Distribution of upper and lower NADW based on A16 (upper), A03 (middle) and A10 (lower) sections. Contour lines show fractions of 20% 50% and 80%, blue lines show cross section of other sections, yellow dashed lines show the boundaries of vertical water columns layers (neutral density at 27.10, 27.90 and 28.10 kg m^{-3})




Figure 15: Currents (upper) and Water Masses (lower) in the Bottom Layer (AABW and NEABW) and the Southern Area (CDW and WSBW).

Left: The arrows show the main currents in the Southern Area.

Right: The colored dots show fractions (from 20% to 100%) of water masses in each station around core neutral density (kg m^{-3}). Stations with fractions less than 20% are marked by black dots while gray dots show all the GLODAPv2 stations.




Figure 16: Distribution of AABW and NEABW based on A16 (upper), A10 (lower left) and A03 (lower right) sections. Contour lines show fractions of 20% 50% and 80%, blue lines show cross sections from other sections, yellow dashed lines show the boundaries of vertical water columns layers (neutral density at 27.10, 27.90 and 28.10 kg m⁻³)

944

945




Figure 17: Distribution of Southern Water Masses (CDW, AABW and WSBW) based on SR04 sections
 Left figures show the west (zonal) part and right figures show the east (meridional) part
 Contour lines show fractions of 20% 50% and 80%.

946

947

948

949 **Table 1:** Schematic of the selection criteria for the OMP analysis (runs) in this study.

950

951	50 °S	Equator	40 °N
#17 AAIW AABW CDW WSBW ($\gamma = 27.10 \text{ kg m}^{-3}$)	#13 WSACW ESACW AAIW ($\gamma = 26.70 \text{ kg m}^{-3}$)	#5 WSACW WNACW ESACW(upper) ENACW(upper) 30°N	#1 WNACW ENACW SAIW MOW ($\gamma = 26.70 \text{ kg m}^{-3}$)
		#7 ESACW(lower) ENACW(lower) AAIW MOW uNADW	#6 ESACW(lower) ENACW(lower) MOW SAIW uNADW
	#14 ESACW AAIW uNADW ($\gamma = 27.30 \text{ kg m}^{-3}$)	#9 ENACW(lower) ESACW(lower) AAIW MOW uNADW	#8 ENACW(lower) ESACW(lower) MOW SAIW uNADW
		30°N #10 AAIW MOW uNADW	#2 ENACW SAIW MOW LSW (uNADW) ($\gamma = 27.30 \text{ kg m}^{-3}$)
#15 AAIW uNADW INADW CDW AABW ($\gamma = 28.10 \text{ kg m}^{-3}$)		#11 AAIW MOW uNADW INADW NEABW	#3 SAIW LSW ISOW DSOW (uNADW INADW) NEABW
	#16 INADW AABW	#12 INADW (ISOW DSOW) NEABW	#4 ISOW DSOW (INADW) NEABW
952	50 °S	Equator	40 °N

953
954

Table 2: Summary of the criteria used to select the water samples considered to represent the Source Water Types discerned in this study.
For convenience, they are grouped into four depth layers.

Layer	SWT	Longitude	Latitude	Pressure (dbar)	Conservative Temperature (°C)	Absolute Salinity (g kg ⁻¹)	Neutral Density (kg m ⁻³)	Oxygen (μmol kg ⁻¹)	Silicate (μmol kg ⁻¹)
Upper Layer	ENACW	20°W—35°W	39°N—48°N	100 — 500	---	---	26.50—27.30	---	---
	WNACW	50°W—70°W	24°N—37°N	100 —500	---	---	26.20—26.70	---	< 2
	ESACW	0—15°E	30°S—40°S	200 — 700	---	---	26.00—27.50	200—230	< 8
	WSACW	25°W—60°W	30°S—45°S	100 —1000	---	---	26.00—27.00	< 230	< 5
Intermediate Layer	AAIW	25°W—55°W	45°S—60°S	100 — 300	< 3.5	< 34.40	26.95—27.50	> 260	< 30
	SAIW	35°W—55°W	50°N—60°N	100 — 500	> 4.5	---	> 27.70	---	---
	MW	6°W—24°W	33°N—48°N	> 300	---	36.50—37.00	---	---	---
Deep and Overflow Layer	uNADW	32°W—50°W	40°N—50°N	1200—2000	< 4.0	---	27.85—28.05	---	---
	INADW	32°W—50°W	40°N—50°N	2000—3000	> 2.5	---	27.90—28.10	---	---
	LSW	24°W—60°W	48°N—66°N	500 —2000	< 4.0	---	27.70—28.10	---	---
	ISOW	0—45°W	50°N—66°N	1500—3000	2.2—3.3	> 34.95	> 28.00	---	< 18
	DSOW	19°W—46°W	55°N—66°N	>1500	< 2.0	---	> 28.15	---	---
Bottom Layer	AABW	---	> 63°S	---	---	---	> 28.20	> 220	> 120
	CDW	< 60°W	55°S—65°S	200—1000	-0.5—1	> 34.82	> 28.10	---	---
	WSBW	---	55°S—65°S	3000---6000	< -0.7	---	---	---	---
	NEABW	10°W—45°W	0—30°N	> 4000	> 1.8	---	---	---	---

955

956 **Table 3:** *The full name of the water masses discussed in this study, and the abbreviation.*

Full name of Water Mass	Abbreviation
East North Atlantic Central Water	ENACW
West North Atlantic Central Water	WNACW
West South Atlantic Central Water	WSACW
East South Atlantic Central Water	ESACW
Antarctic Intermediate Water	AAIW
Subarctic Intermediate Water	SAIW
Mediterranean Overflow Water	MOW
Upper North Atlantic Deep Water	uNADW
Lower North Atlantic Deep Water	lNADW
Labrador Sea Water	LSW
Iceland-Scotland Overflow Water	ISOW
Denmark Strait Overflow Water	DSOW
Antarctic Bottom Water	AABW
Circumpolar Deep Water	CDW
Weddell Sea Bottom Water	WSBW
Northeast Atlantic Bottom Water	NEABW

957

958

959 **Table 4:** *Table of the mean value and the standard deviation of all variables for all the water masses*
960 *(i.e. Source Water Types) in this study*

Layer	SWTs	Conservative Temperature (°C)	Absolute Salinity	Neutral Density (kg m ⁻³)	Oxygen (μmol kg ⁻¹)	Silicate (μmol kg ⁻¹)	Phosphate (μmol kg ⁻¹)	Nitrate (μmol kg ⁻¹)
Upper Layer	ENACW (upper)	13.72	36.021	26.887	243.1	2.49	0.41	7.03
	ENACW (lower)	11.36	35.689	27.121	216.3	5.33	0.75	12.14
	WNACW (upper)	18.79	36.816	26.344	213.3	0.72	0.08	2.00
	WNACW (lower)	17.51	36.634	26.554	193.9	1.60	0.24	4.88
	ESACW (upper)	13.60	35.398	26.500	217.1	3.68	0.65	8.26
	ESACW (lower)	9.44	34.900	26.928	214.2	6.60	1.19	16.42
	WSACW (upper)	16.30	35.936	26.295	222.2	1.60	0.32	3.15
	WSACW (lower)	12.30	34.294	26.707	209.8	3.58	0.80	10.43
Intermediate Layer	AAIW	1.78±1.02	34.206±0.083	27.409±0.111	300.7±16.2	21.09±4.66	1.95±0.11	27.33±1.92
	SAIW	3.62±0.43	34.994±0.057	27.831±0.049	294.6±9.7	8.53±0.85	1.04±0.07	15.55±1.06
	MW	12.21±0.77	36.682±0.081	27.734±0.150	186.2±10.7	7.17±1.75	0.74±0.11	12.71±1.96
Deep and Overflow Layer	Upper NADW	3.33±0.31	35.071±0.027	27.942±0.027	279.4±8.0	11.35±0.78	1.11±0.04	16.99±0.49
	Lower NADW	2.96±0.21	35.083±0.019	28.000±0.029	278.0±4.6	13.16±1.42	1.10±0.05	16.80±0.48
	LSW	3.24±0.32	35.044±0.031	27.931±0.042	287.4±8.5	9.79±0.85	1.08±0.06	16.30±0.58
	ISOW	3.02±0.26	35.098±0.028	28.001±0.044	277.2±3.3	12.21±1.18	1.10±0.05	16.58±0.48
	DSOW	1.27±0.29	35.052±0.016	28.194±0.028	300.3±3.6	8.66±0.77	0.95±0.05	13.93±0.44
Bottom Layer	AABW	-0.46±0.24	34.830±0.009	28.357±0.048	239.0±9.3	124.87±2.36	2.27±0.03	32.82±0.45
	CDW	0.41±0.19	34.850±0.011	28.188±0.037	203.8±8.5	115.53±7.72	2.31±0.06	33.46±0.91
	WSBW	-0.79±0.05	34.818±0.005	28.421±0.010	251.8±3.7	119.93±3.26	2.24±0.03	32.50±0.36
	NEABW	1.95±0.06	35.061±0.008	28.117±0.005	245.9±3.7	47.06±2.32	1.49±0.04	22.27±0.53

961