

# DUACS DT2018: 25 years of reprocessed sea level altimetry products

Guillaume Taburet<sup>1</sup>, Antonio Sanchez-Roman<sup>2</sup>, Maxime Ballarotta<sup>1</sup>, Marie-Isabelle Pujol<sup>1</sup>, Jean-François Legeais<sup>1</sup>, Florent Fournier<sup>1</sup>, Yannice Faugere<sup>1</sup> and Gerald Dibarboure<sup>3</sup>

5 <sup>1</sup>Collecte Localisation Satellites, Parc Technologique du Canal, 8-10 rue Hermès, 31520 Ramonville-Saint-Agne, France

<sup>2</sup>Instituto Mediterráneo de Estudios Avanzados, C/Miquel Marqués, 21 – 07190 Esporles – Illes Balears, Spain

<sup>3</sup>Centre National d’Etudes Spatiales, 18 avenue Edouard Belin, 31400 Toulouse, France

*Correspondence to:* Guillaume Taburet (gtaburet@groupcls.com)

**Abstract.** For more than 20 years, the multi-satellite Data Unification and Altimeter Combination System (DUACS) has  
10 been providing Near Real Time (NRT) and Delayed Time (DT) altimetry products. DUACS datasets range from along-track measurements to multi-mission Sea Level Anomaly (SLA) and Absolute Dynamic Topography (ADT) maps. The DUACS DT2018 ensemble of products is the most recent and major release. For this, twenty five years of altimeter data have been reprocessed and are available through the Copernicus Marine Environment Monitoring Service (CMEMS) and the Copernicus Climate Change Service (C3S).

15 Several changes were implemented in DT2018 processing in order to improve the product quality. New altimetry standards and geophysical corrections were used, data selection was refined and Optimal Interpolation (OI) parameters were reviewed for global and regional map generation.

This paper describes the extensive assessment of DT2018 reprocessing. The error budget associated with DT2018 products at global and regional scales was defined and improvements on the previous version were quantified (DT2014; Pujol et al.,  
20 2016). DT2018 mesoscale errors were estimated using independent and in-situ measurements. They have been reduced by nearly 3 to 4 % for global and regional products compared to DT2014. This reduction is even greater in coastal areas (up to 10%) where it is directly linked to the geophysical corrections applied to DT2018 processing. The conclusions are very similar concerning geostrophic currents, where error was globally reduced by around 5% and as much as 10% in coastal areas.

## 25 1 Introduction

Since 1992, high precision sea level measurements have been provided by satellite altimetry. They have largely contributed to better understand both the ocean circulation and the response of the Earth’s system to climate change. Following Topex-Poseidon in 1992, the constellation has grown from one to six satellites flying simultaneously (see Figure 1). The combination of these missions permits to resolve the ocean circulation both on a mesoscale and global scale and on different  
30 time scales (annual and inter-annual signals and decadal trends). This has been made possible thanks to the DUACS

altimeter multi-mission processing system, initially developed in 1997. Ever since, it has been producing altimetry products for the scientific community in either Near Real Time (NRT), with a delay ranging from a few hours to one day, or Delayed Time (DT), with a delay of a few months. The processing unit has been redesigned and regularly upgraded as knowledge of altimetry processing has been refined (Le Traon et al., 1998; Ducet et al., 2000; Dibarboure et al., 2011; Pujol et al., 2016).

5 Every few years, a complete reprocessing is performed through DUACS that includes all altimetry missions and that uses up-to-date improvements and recommendations from the international altimetry community.

This paper presents the latest reprocessing of DUACS DT reanalysis (written hereafter DT2018) and focuses on improvements that have been implemented since the version DT2014 (Pujol et al., 2016). Previously reprocessed products (including DT2014) were distributed by Aviso from 2003 to 2017. Since May 2015, the European Copernicus Program  
10 (<http://www.copernicus.eu/>) has taken responsibility for all the processing, along with the operational production and distribution of along-track (level 3) and gridded (level 4) altimetry sea level products.

The daily DT2018 products time series starts from January 1, 1993 and temporal extensions of the sea level record are regularly updated with a delay of nearly six-months. Multi-mission products are based on all the altimetry satellites representing a total of 76 mission-years as shown in Figure 1. The DT2018 reprocessing is characterized by major changes in  
15 terms of standards and data processing compared to the DT2014 version. These changes are highlighted in section 2 and have a significant impact on sea level products quality. Two types of gridded altimetry sea level products are available in DT2018. The first is dedicated to retrieving mesoscale signals in the context of ocean modeling and analysis of ocean circulation on a global or regional scale. This type of dataset is produced and distributed by the Copernicus Marine Service (CMEMS). The second is dedicated to monitoring long-term evolution of sea level, for use in both climate applications and  
20 the analysis of ocean/climate indicators (such as the evolution of the global and regional Mean Sea Level (MSL)). This second type of dataset is produced and distributed by the Copernicus Climate Change Service (C3S). More details on the differences between the products distributed by these two Copernicus Services can be found in section 2.4.

The paper is organized as follows: section 2 considers DUACS processing , from the level 2 altimeter standards to the inter-mission calibration (level 3) and the mapping procedure (level 4). Sections 3 and 4 focus respectively on the quality of  
25 global and regional products at spatial (coastal, mesoscale) and time (climate scales) scales. Finally, section 5 discusses the key results and future prospects.

## **2 Data processing**

### **2.1 Altimeter constellation**

The 25 year period [1993-2017] involves 76 mission-years, and 12 different altimeters. The evolution of the altimeter  
30 constellation is shown in Figure 1. The most notable change in the constellation compared to DT2014 concerns availability of data from Sentinel-3A and Hayaing-2A altimetry missions. For Sentinel-3, an additional six months of data (from June

2016 to December 2016) have been incorporated into the system. For Hayaing-2A, data from March 2016 to February 2017 have also been added.

## 2.2 Altimetry standards

DUACS takes Level 2P (L2P) altimetry products as its input data. These data are disseminated by CNES and EUMETSAT.

5 L2P products are supplied by L2 products that are distributed by different agencies: NASA, NSOAS, ISRO, ESA, CNES, EUMETSAT. They include the altimetry standard, that is algorithms and parameters used to retrieve the sea level anomalies from the altimeter measurements, (i.e. instrumental, geophysical and environmental corrections together with Mean Sea Surface (MSS)), as well as a validity flag that is used to remove spurious measurements .

Indeed, the altimeter measurement is affected by various disturbances (atmospheric, instrumental...) that must be estimated  
10 to correct it. Specific corrections are also applied to remove high frequency signal that cannot be taken into account in the DUACS processing (Escudier et al., 2017). The Dynamic Atmospheric Correction (DAC) and ocean tide correction are the two main examples. The DUACS DT2018 global reprocessing was an opportunity to take into account new recommendations and new corrections from the altimetry community (Ocean Surface Topography Science Team, OSTST).

The altimetry standards were carefully selected in order to be as consistent and homogeneous as possible between the  
15 various missions, whatever their purpose (in particular the retrieval of mesoscale signals or climate applications). This selection was made possible between 2014 and 2017 in the framework of phase II of ESA's Sea Level Climate Change Initiative (SL\_cci) project. Part of the project activities included selecting a restricted number of altimetry standards (Quarty et al., 2017; Legeais et al., 2018a). Table 1 presents the altimetry standards used in DT2018 and the changes compared with the previous version (written in bold). The orbit standards from Jason-1, Jason-2, Cryosat-2, AltiKa, Jason-3 and Sentinel-  
20 3A altimeter missions were upgraded from Precise Orbit Estimation (POE)-D to a new POE-E. The new POE-E standards are of a very high quality (Ollivier et al., 2015; AVISO, 2017b). In this version, the main developments concerns the evolutions of gravity field model that has a positive impact on regional MSL error and greatly reduce geographically-correlated errors.

Various corrections were updated, of which the new MSS CNES-CLS-15 and ocean tide model (FES2014) have led to the  
25 greatest improvements in product quality. Valuable enhancements were made in the MSS to improve performance at short wavelengths (Pujol et al., 2018a). Furthermore, the sea level in coastal areas and the Arctic region is determined more accurately in the updated version, and errors were greatly reduced globally. Concerning the ocean tide correction, FES2014 is the latest version of the FES (Finite Element Solution) tide model developed between 2014 and 2016. This new release gives improved results in the deep ocean, at high latitudes and in shallow/coastal regions (Carrère et al., 2016).

## 30 2.3 Developments in DUACS processing

DUACS processing involves an initial preprocessing step during which data from the various altimeters are acquired and homogenized. Next, along-track products (L3) and multi-missions gridded products (L4) can be estimated. Finally, the

derived products are computed and disseminated to users. This section is not intended to describe the entire data processing system in detail, but rather to expose the major changes made for this DT2018 version. For a detailed description of DUACS processing, readers are advised to consult Pujol et al., 2016.

### 2.3.1 Acquisition and preprocessing

- 5 The DUACS processing sequence can be divided into multiple steps: acquisition, homogenization, input data quality control, multi-mission cross calibration, along-track SLA generation, multi-mission mapping and final quality control.

The acquisition stage consists in retrieving altimeter and ancillary data and applying to those data the most recent corrections, models and references recommended by expert (as described in section 2.1 and 2.2). This up-to-date selection is available in Table 1.

- 10 Input Data Quality Control is a process related to the calibration/validation activities carried out for CNES, ESA and EUMETSAT. It is composed of several editing processes designed to detect and fix spurious measurements and to ensure the long-term stability of L2P products. The up-to-date editing process is described in annual Cal/Val reports for each mission (AVISO, 2017c). Since 2014, and learning from experts' experience, great efforts have been made to refine this global process and notably to tailor some parts to specific regions such as high-latitude and coastal areas. At high latitudes the idea  
15 is to filter an altimeter parameter which has a specific signature for ice, compared to the ocean, and then to flag associated data as ice. But such a filtering solution affects all data, with the risk that potentially compromised data outside icy areas can be inaccurately flagged as ice. The updated development consists in using a mask so that the chosen filtering solution always provides relevant results (Ollivier et al., 2014). The mask is based on the sea ice concentration product from the EUMETSAT Ocean and Sea Ice Satellite Application Facility (OSI SAF, [www.osi-saf.org](http://www.osi-saf.org)) and gives a maximum estimation  
20 of ice extent. In coastal areas, along-track SLA measurements for non-repetitive missions were rejected for L2P DT2014 products, mainly due to the lower quality of MSS for area less than 20 km from the coast (Pujol et al., 2016). DT2018 benefits from a solution for improved MSS quality (Pujol et al., 2018), so efforts were made to retain as many valid measurements as possible close to the coast. The data selection strategy is based on a median filter applied in a 30km wide strip off the coastline (Ollivier et al., 2014). As a result, substantially more valid data can be used in DUACS, especially for  
25 geodetic measurements.

- Finally, the cross-calibration step ensures that all data from all satellites provide consistent and accurate information (Pujol et al., 2016). Mean sea level continuity between altimeter missions is ensured by reducing global and regional biases for each transition between reference missions (TP-J1, J1-J2 and J2-J3). In order to minimize geographically-correlated errors, two algorithms using empirical process methods are then applied, namely: Orbit Error Reduction (OER) and Long Wavelength  
30 Error Reduction (LWER).

### 2.3.2 Along-track product generation

The along-track generation for repetitive altimeter missions is based on the use of a mean profile (MP) (Table 2, Pujol et al., 2016; Dibarboure et al., in review). These MPs are necessary in order to co-locate the sea surface heights of the repetitive tracks and to retrieve a precise mean reference in order to compute sea level anomalies. The methodology used to compute the DT2018 MP was the same as for DT2014. The differences arise from the upstream measurements, as new altimetry standards were used in DT2018 (described in section 2.2), along with new data selection (section 2.3.1) and reviewed temporal periods for the different altimeters considered. Table 2 presents the altimeter missions and time periods used to compute the four different MPs available along the following tracks: TopexPoseidon/Jason1/OSTM-Jason2/Jason3, TopexPoseidon Interleaved Phase/Jason1 Interleaved/Jason2 Interleaved, ERS-1/ERS-2/Envisat/Saral-AltiKa and Geosat Follow On.

Following the previous MPs version, additional measurements collected by OSTM/Jason-2 and SARAL/AltiKa between 2012 and 2015 were exploited for DT2018. Since March 2015, however, AltiKa has been considered a non-repetitive mission for Delayed-Time products. As a result, no measurements after that date were taken into account when computing the ERS-1/ERS-2/EN/AL MP. To limit the ionospheric correction error in this MP, no ERS-2 data collected between January 2000 and October 2002 were used to compute the MP because the ionospheric activity was much more intense during this period than between 1995 and 2000.

New DT2018 MPs were defined as close to the coast as possible as illustrated in Figure 2. This improvement is associated with the use of the new MSS (Pujol et al., 2018a) and ocean tide correction and the refined selection of valid data (sections 2.2 and 2.3.1). It has a direct and positive impact on along-track product generation that provides extended coastal coverage.

Globally, comparisons at crossovers provides good results in this new version. Compared to the DT2014 version, we observe a decrease in the mean of the difference at crossovers by around 0.3cm globally and up to 1cm locally (data not shown here). It should be noted that for the Sentinel-3A, it was impossible to estimate a precise MP for this reprocessing, due to the short time period (i.e. a few months) available to compute it. Consequently, data from the Sentinel-3A mission were only interpolated into theoretical positions (Dibarboure et al., 2011), then the gridded MSS (Pujol et al., 2018) was removed. Since the reprocessing, an MP has been calculated (Dibarboure et al., in rev.; Pujol et al., 2018b) and the Sentinel-3A dataset has been reprocessed in a CMEMS version in 2019.

For non-repetitive missions (ERS-1 during its geodetic phase, Cryosat-2, Hayaing-2A, both Jason-1 and Jason-2 in their geodetic phase, and SARAL-AltiKa in its geodetic phase), no MP can be estimated. The SLA is in this case derived along the real altimeter tracks using the gridded MSS (Pujol et al., 2016; Dibarboure et al., in rev).

The final step of along-track processing consists of noise reduction using low-pass Lanczos filtering and subsampling. This process remains unchanged from the DT2014 version (Pujol et al., 2016).

DT2018 reprocessing was also the opportunity to propose new products. New along-track products were tailored for assimilation purposes to provide users with the specific geophysical corrections, used to compute the sea level anomaly in

the DUACS processing: DAC, ocean tide and LWER. As explained in section 2.2, these geophysical effects are taken into account in DUACS because their temporal variability is too high to be resolved by altimeter measurements and to be mapped using the OI method.

### **2.3.3 Gridded product generation: multi-mission mapping**

- 5 The multi-mission mapping procedure in DUACS is based on an optimal interpolation (OI) technique derived from LeTraon et al., 1998; Ducet et al., 2000 and LeTraon et al., 2003. This method is designed to generate regularly gridded products for Sea Level Anomalies by combining measurements from different altimeters. The main objective in the DT2018 reprocessing framework was to improve gridded altimetry products in the tropics, in coastal areas and at mesoscale. To do so, OI parameters were adjusted. The sea level spatial and temporal variability were more accurately defined based on the 25 years of observations available. Particular attention was paid to coastal areas, where spurious peaks of high variability were able to be reduced. An optimized selection of along-track data was incorporated into OI processing by changing the size of the suboptimal interpolation window, decreasing it by one third in regions of high variability and in the equatorial belt. OI observation errors were increased in the equatorial belt, as the impact of filtering and subsampling had been previously underestimated in this area where they generate noise at small scales in gridded products. Errors generated when using the gridded MSS were updated with the use of the new MSS version (Pujol et al., 2018a). Correlation scales were only reviewed for regional Mediterranean products. While set to constant values (100 km and 10 days) in the DT2014 version, precise covariance and propagation models were computed for DT2018 regional mapping. Spatial scales now range from 75 km to 200 km while temporal scales remain at 10 days. These changes have contributed to improving the retrieval of mesoscale signals' in Mediterranean regional products (section 4).
- 20 For Black Sea processing, OI parameters are now similar to parameters used for the global ocean processing, except for the correlation scales which are still set to 100km and 10days.

### **2.4 Different products for different applications**

- Two different types of sea level gridded altimetry products are available in DT2018 version. The first type, produced and distributed within the Copernicus Marine Service (CMEMS), is dedicated to mesoscale observation. The other type, produced and distributed within the Copernicus Climate Change Service (C3S), is dedicated to monitoring the long-term evolution of the sea level for use in climate applications and for analyzing ocean/climate indicators (such as global and regional MSL evolution). Two types of altimeter processing configurations are exploited to build these two products. The first difference of configuration is related to the number of altimeters used in the satellite constellation. Mesoscale observation requires the most accurate sea level estimation at each time step, along with the best spatial sampling of the ocean. All available altimeters are thus included in CMEMS products, and the sampling can vary with time depending on the constellation status. In contrast, the temporal stability of surface sampling is more important when monitoring the long-term sea level evolution. A steady number of altimeters (two) are thus used in C3S products. This corresponds to the
- 30

minimum number of satellites required to retrieve mesoscale signals in delayed time conditions (Pascual et al., 2006; Dibarboure et al., 2011). Within the production process, long-term stability and large-scale changes are established on the basis of records from the reference missions (TOPEX-Poseidon, Jason-1, Jason-2 and Jason-3) used in both CMEMS and C3S products. Any additional missions (e.g. as many as five additional missions in 2017) are then homogenized with respect to the reference missions and help to improve mesoscale process sampling, providing high-latitude coverage and increasing product accuracy. However, the total number of satellites has greatly varied over the altimetry era and biases may develop when a new satellite on a drifting orbit is introduced. Each addition, may affect the stability of the global and regional MSL by several millimeters (data not shown here). Although spatial sampling is reduced when there are fewer satellites, the risk of introducing such anomalies is thus also reduced in C3S products, resulting in improved stability. In CMEMS products, stability is ensured by the calibration with the reference missions and mesoscale errors are reduced due to the improved ocean surface sampling made possible by using all the satellites available in the constellation.

As a second difference of configuration, the reference used to compute Sea Level Anomalies for C3S products was a MSS for all missions whereas for CMEMS products, a MP was used along the theoretical track of satellites following a repetitive orbit (see section 2.3.2). Considering the regional mean sea level temporal evolution, the combined use of MSS and MP for successive missions in the merged product give rise to regional centimetric bias (data not shown here). Consequently, the systematic use of MSS for all missions has been privileged in the C3S products to ensure MSL stability and the use of MP for repetitive missions has been selected in the CMEMS products to increase their accuracy.

The differences between CMEMS and C3S product quality are discussed on a climate scale in section 3.4.

### **3 DT2018 Global product quality**

This section focuses on the quality of gridded (L4) products. Sea surface height and derived current products were analyzed at different spatial scales (open ocean, coastal areas), distinguishing different temporal scales (from mesoscale to climatic scales). DT2018 L4 products were compared with those of DT2014 over the 1993-2017 time period. Except when explicitly mentioned otherwise, the results presented in this section are valid for all DUACS DT2018 products distributed via both Copernicus services.

#### **3.1 Mesoscale signals in along-track and gridded products**

Optimizing the mapping process (section 2.3.3) and incorporating the new altimetry corrections (section 2.2) had a direct impact on the observation of ocean sea level and surface circulation dynamics in the gridded products. To characterize this impact, the difference between DT2014 and DT2018 temporal variability is shown in Figure 3. An additional variance of between 2% and 5% is observed for high variability regions in DT2018 products. This increase is due to having changed the OI spatial and temporal scales used in the mapping process and decreased the suboptimal interpolation window size. The OI selection window is more focused on close observations (both spatial and temporal). In coastal areas, a substantial reduction

in SLA variance is observed due to both the FES2014 tidal correction and, to a more limited extent, the new MSS. For the tidal correction, Carrere et al., (2016) have shown a reduction in SLA variance at nearshore crossovers. Pujol et al., (2018a) have emphasized that the new gridded MSS shows less SLA degradation near the coast. These improved standards contribute to a valuable local reduction in SLA variance (up to 50% alongshore). At high latitudes, the difference of variance is significant ( $\pm 100\text{cm}^2$  to  $\pm 200\text{cm}^2$ ) and is due to the new MSS correction. Indeed, Pujol et al., (2018a) have shown that the CNES\_CLS 2015 MSS improves both coverage in the Arctic and resolution of the shortest wavelengths at high latitudes.

Compared to DT2014, the new version reveals more intense western boundary currents (geostrophic part). This has a direct impact on the Eddy Kinetic Energy (EKE) derived from these products. Figure 4 presents the spatial difference in the mean EKE over global ocean between DT2018 and DT2014 products, along with their temporal evolution. As observed before for the differences of SLA variance, a higher energy is evident in high variability areas. This corresponds to a 2% increase in EKE in DT2018. However, in the equatorial belt ( $\pm 20^\circ\text{N}$ ), the EKE in DT2018 is lower (-17%). This is a direct consequence of the noise measurement that is taken into consideration in the mapping process for all satellites: observation errors prescribed during OI in the tropical belt have been increased, so the SLA signal is smoother and less energy is observed in this region. In coastal areas, DT2018 version presents fewer spurious peaks of high EKE (Figure 4 b). As already stated, this is related to the improved altimetry correction and lower SLA variance. Considering the mean EKE time series, a global reduction of  $26\text{ cm}^2$  (17%) is observed for dataset DT2018. This is directly due to the lower tropical EKE. Another important point to note is that the standard deviation of EKE in these products is lower than in DT2014. This illustrates that EKE variations are less important, there are fewer isolated anomalies (and these are mostly coastal) in the new DT2018 products.

The gridded SLA accuracy was estimated by comparison with independent along-track measurements. Maps produced by merging only two altimeters were compared with SLAs measured along-track from the tracks of another mission that was kept independent from the mapping process (see Pujol et al., 2016 for full methodology). Topex-Poseidon interleaved was compared with gridded products that merged Jason-1 and Envisat over 2003-2004. It must be pointed out that these results are much more representative of gridded products combining two altimetry missions. Products combining all available missions can usually benefit from improved sampling when three to six altimeters are used. The errors described here should thus be considered the upper limit. Table 3 summarizes the results of comparisons over different areas. Figure 5 shows the percentage of the difference in variance between gridded products and TP independent along-track measurements for DT2018 and DT2014 products. The gridded product error for mesoscale wavelengths ranges between  $1.4\text{ cm}^2$  (for a low variability area) and  $37.7\text{ cm}^2$  (for a high variability region). The improvements in DT2018 compared with DT2014 affect all areas. Offshore, the improvement is fairly low (around 3%) and is associated with the enhanced version of the OI mapping parameter. In coastal areas, the improvements are more significant (around 10%) and caused by the new tidal correction (FES2014) and, to a lesser extent, to the MSS and MPs. In the tropical belt, improvements are also significant (around 9%) and related to the observation errors that were increased in this area for the OI processing.



### 3.2 Geostrophic current quality

Absolute geostrophic currents for DT2018 were assessed using drifter data for the 1993-2017 time period. The AOML (Atlantic Oceanographic & Meteorological Laboratory) database was used for the comparison (Lumpkin et al. 2013). These in-situ data were corrected for Ekman drift (Rio et al., 2011) and wind if a drifter's drogue had been lost (Rio et al., 2012) so as to be comparable with the altimetry absolute geostrophic currents. Drifters positions and velocities were interpolated using a 3-day low-pass filter in order to remove high-frequency motions (Rio et al., 2011). The absolute geostrophic currents derived from altimetry products were then interpolated onto drifter positions for comparison.

The distribution of the current's intensity shows an overall underestimation of magnitude in altimetry products compared to drifter observations (data not shown). Figure 6 shows the RMS difference between the DT2018 geostrophic current and that of drifters. The mean RMS is nearly 10 cm/s and the main errors are located nearshore and in high variability region with peaks higher than 20 cm/s. Taylor skill scores (Taylor, 2001) were computed for the zonal and meridional components of the current in DT2018. This assessment took into consideration both the signal's correlation and its standard deviation. Results are quite robust: 0.89 for the zonal and 0.87 for the meridional component.

Table 4 summarizes the mean RMS of the differences between geostrophic current maps and drifter measurements over different areas for versions DT2018 and DT2014. DT2018 products are more consistent with drifter measurements than DT2014 version products. The improvement is clearly visible in the intra-tropical belt. The variance of the differences with drifters is reduced around 20 to 40% in this area. Additional noise-like signals present in the DT2014 version had reduced consistency with drifter measurement (Pujol et al., 2016). This degradation was corrected for by the change in mapping parameters used for this updated version (section 2.3.3). A significant improvement can also be observed in coastal areas, where the variance of differences with drifter measurements is reduced by nearly 15% (Table 4). Elsewhere, this reduction in the variance of difference ranges from 4 to 7%.

### 3.3 Coastal areas

As described in sections 2.3.1 and 2.3.2 the new DUACS DT2018 processing has a key impact on coastal areas and overall, all missions have more measurements available in DT2018 compared to DT2014.

The assessment of gridded products in coastal areas included a comparison with tide-gauge (TG) measurements. We used mean monthly TG measurements from the PSMSL network (Permanent Service for Mean Sea Level, PSMSL, 2016) from 1993 to 2017. We used only long-term monitoring stations with a lifetime of more than two years. Sea surface height measured by TG was compared with gridded SLA by considering the maximum correlation with the nearest neighboring pixel (Valladeau et al., 2012; AVISO, 2017a). In Figure 7 the variance of the difference between DT2018 altimetry products and TG measurements is compared with that obtained from the differences using DT2014 altimetry products. The results show a global reduction in the variance (0.6%) when DT2018 data are used. There is a clear improvement along the Indian coast, Oceania and northern Europe. Local degradation can be observed along the coast of Spain and along the United

States' West coast. These degradations, which are not observed in other diagnoses such as independent along track measurements still need to be further investigated.

### 3.4 Climate scales

The global mean sea level (GMSL) is a key indicator of climate change since it reflects both the amount of heat added in the ocean and the land ice melt coming mainly from Antarctic and Greenland ice sheets and glaciers. Three different altimeter products can be used to compute three GMSL estimates: the time series of the box-averaged along-track measurements of the reference missions only (Ablain et al., 2017), L4 merged gridded sea level products from CMEMS and C3S (e.g. Figure 8, left). For the same product versions and computation periods, these three GMSL estimates are considered to be equivalent since almost the same altimetry standards are used to compute sea level anomalies and, the long-term stability for all products is ensured by using the same reference missions. The remaining GMSL differences observed ( $\sim 0.17$  mm/year) are not significant given the uncertainty on different scales (uncertainty in the GMSL trend is 0.4 mm/year at the 90% confidence level given by Ablain et al. 2019). Note that as aforementioned (section 2.4), differences can be found between the two different Copernicus gridded products (CMEMS/C3S) when computing regionally-averaged MSL.

When computing area-averaged MSL time series, users are advised that DUACS products are not corrected for the effect of Glacial Isostatic Adjustment (GIA) due to post glacial rebound. A GIA model should be used to estimate the associated sea level trends.

In addition, between 1993 and 1998, GMSL is known to have been affected by instrumental drift in the TOPEX-A measurement, as quantified by several studies (Watson et al., 2015; Beckley et al., 2017; Dieng et al., 2017). The sea level altimetry community agrees that it is necessary to correct the TOPEX-A record for instrumental drift to improve accuracy and reduce uncertainty in the total sea level record. However, there is not consensus so far on the best approach to estimate drift correction at global and regional scales. DUACS sea level altimetry products are not corrected for TOPEX-A drift, pending on-going TOPEX reprocessing by CNES and NASA/JPL but users can apply their own correction. Adjusting for this TOPEX-A anomaly create a GMSL acceleration of  $0.10 \text{ mm yr}^{-2}$  for the 1993–2017 time span that does not otherwise appear (WCRP 2018).

Figure 8 (left) shows the GMSL's temporal evolution and associated trend computed with the new DT2018 and former DT2014 versions of DUACS C3S products. In the latest version, the global mean sea level trend is of 3.3 mm/year (including GIA correction of -0.3 mm/year). The origin of the associated uncertainty is discussed by Legeais et al., 2018b. The map of the differences for the local MSL trend derived from the latest and previous product versions (Figure 8, right) displays a pattern predominantly associated with the different orbit standards used in the two product versions (GDR-E versus GDR-D, see Table 1). Such a result is confirmed by comparing altimetry products with the independent dynamic height measurements derived from in-situ Argo profiles (Valladeau et al., 2012; Legeais et al., 2016).

## 4 DT2018 regional product quality

### 4.1 SLA field quality

As previously discussed for global ocean products, the quality of regional gridded SLA products is estimated through comparison with independent altimeter along-track and tide gauge measurements.

- 5 Figure 9 shows the spatial distribution of the RMS of the differences between regional DT2018 SLA gridded products and independent along-track measurements (Topex/Poseidon interleaved along-track measurements over the 2003-2004 period). The main statistics for these comparisons, as well as a comparison with the previous DT2014 version, are also given in Table 5. In contrast with the global products assessment, the evaluation of regional products cannot include the mesoscale signal analysis: the short length of the tracks segments available over the regional seas does not allow accurate filtering of the
- 10 signal in order to focus specifically on mesoscale. The results obtained show that for the DT2018 Mediterranean product, the main errors are located in coastal areas and in the Adriatic and Aegean Seas, with RMS values ranging from 6 to 9 cm. The Black Sea products also show higher errors in coastal areas (results not shown here). The mean Variance of the differences between gridded products and along-track measurements is nearly 17 cm<sup>2</sup> and 23 cm<sup>2</sup> over the Mediterranean Sea and the Black Sea. This value is higher than the mean error observed over low variability areas in the global ocean (Table 3), mainly
- 15 due to the different wavelengths addressed in these comparisons. Compared to the previous regional DT2014 version, the error is reduced by 4.2% for the Mediterranean Sea and 3.5% for the Black Sea. It is important to note that these results are representative of gridded product quality when only two altimeters are available. These products can be considered to be degraded products for mesoscale mapping since they use minimal altimeter sampling.
- Compared to the previous version, consistency with monthly TG measurements (Figure 10) is improved locally in the
- 20 regional DT2018 Mediterranean gridded product in the western part of the Mediterranean basin. Degradation is observed, however, in some other coastal areas, especially in the center of the basin and along the Turkish coast. For the Black Sea gridded product, only nine Tide Gauges were available for the comparison. With the exception of a tide gauge at the eastern end of the Black Sea, on the Georgian coast, these DT2018 regional products are improved of the order of 1 %.

### 4.2 Geostrophic current quality in the Mediterranean Sea

- 25 DT2018 regional absolute geostrophic current in the Mediterranean basin was assessed using drifter data for the 1993-2017 period. The data were collected from drifters released in the Mediterranean Sea as part of AlborEx (Pascual et al., 2017) and MEDESS-GIB (EU MED Program; <http://www.medess4ms.eu/> ; Sotillo et al., 2016) multi-platform experiments as well as other experiments incorporated into CMEMS In Situ Thematic Centre (INS TAC) products. These data are processed similarly to the global product (section 3.2).
- 30 Table 6 summarizes the main statistical results for the whole basin. The DT2018 regional product presents a correlation coefficient with drifter data 4% greater than that obtained when using the DT2014 regional product. Moreover, the errors in the later version are slightly lower 1%, whilst its improvement in explained variance is as high as 14%.

The analysis was then repeated for the different dynamical sub-regions of the basin (see Figure 11.a) reported by Manca et al. (2004). This differentiation is based on the typical permanent features in the upper 200 m of the water column. Overall, comparisons between geostrophic velocities derived from the DT2018 regional product and absolute surface velocities retrieved by the drifters (Figure 11. b – e) reveal a correlation coefficient greater than 0.40 in most of the boxes. Correlations greater than 0.50 are mainly located in the southernmost part of the basin where stronger mesoscale activity occurs; namely the Alboran Sea (DS1), the Algerian Basin (DS3 and DS4), the Sardinian Channel (DI1), the Strait of Sicily (DI3), the Ionian Sea (boxes DJ7, DJ8 and DJ5), and the Cretan passage (DH3). The overall RMS difference between both datasets ranges between 8 and 11 cm/s, although it reaches 20 cm/s in DS1 due this area's strong dynamics. Slightly larger errors are obtained when comparing the DT2014 product with drifter observations (not shown here). Furthermore, drifter data collected in boxes DS1, DS3, and DS4 have the largest variability due to the aforementioned mesoscale activity. This fact is also reflected in the two altimetry products, which have the largest variance values in the Mediterranean basin.

Overall, the correlation coefficient between the DT2018 regional product and in-situ drifter data is improved by 5-10% with respect to that obtained when using the DT2014 product (Figure 11.g). Here, positive values denote an improvement in DT2018 over DT2014. This fact is mainly observed in areas of strong mesoscale activity. Moreover, the errors (Figure 11.f) are reduced by 2% in the northernmost part of the western Mediterranean basin and Adriatic Sea. However, negative values lower than 2% (slightly larger errors when using DT2018) are observed in the Algerian Basin and most of the eastern part of the Mediterranean basin. The main improvement in DT2018 with respect to DT2014 lies in the variance explained (Figure 11.h), which presents values nearly 20% higher in the later product in some areas of the western part of the basin and nearly 10% higher in the eastern part. This is due to better capturing mesoscale activity. This improvement is not observed in the northernmost part of the basin, where less mesoscale activity occurs.

## 5 Discussions and Conclusions

More than 25 years of Level-3 and Level-4 altimetry products were reprocessed and delivered as version DT2018. This reprocessing takes into account the most up-to-date altimetry corrections and also includes changes in the mapping processing parameters. These changes impact SLA signals at multiple temporal and spatial scales.

A notable change concerns the gridded sea level altimetry products that are available in version DT2018. They are produced and distributed through two different Copernicus Services that correspond to different applications. CMEMS distributes maps that include all the available altimeter missions. These maps provide the most accurate sea level estimation with the best spatial and temporal sampling of the ocean at all times. Through C3S, maps that include only two satellites are used to compute the most homogeneous and stable sea level record over time and space. Sea level C3S products are dedicated to monitoring long-term sea level evolution for climate applications and analyzing ocean/climate indicators (such as global and regional MSL evolution).

Other changes were implemented in DT2018 processing: the altimetry standards and geophysical corrections were brought up-to-date with expert recommendations, and mapping parameters, including spatial and temporal correlation scale and measurement errors were refined. We also focused on improving coastal editing to obtain many relevant sea level data, mainly from drifting altimeters. Additional sea level data were incorporated into DT2018, in particular Sentinel-3A measurements taken over a 6-month extension period.

Discussing these key changes, we then focused on describing their impact on gridded sea level products. SLA variability has increased in energetic areas (from 5 to 10%) and decreased locally along coasts (up to 50%). A 10% EKE decrease in the equatorial belt has also been observed and related to the reduced measurement errors prescribed for OI in this area.

To achieve independent comparisons, geostrophic currents were examined to in-situ observations. Compared to the version DT2014, offshore improvements (+4-5%) particularly in the tropics (+5-10%) and coastal improvements (+10%) have been demonstrated using independent drifter data. An independent along-track sea level comparison and tide gauge comparisons have strengthened these conclusions.

Regional products are also enhanced with DT2018, taking advantage of new standards and processing. The SLA gridded product errors in the regional products have decreased by 3% to 4% when estimated using independent along-track measurements.

The limitations exposed by Pujol et al. (2016) are still valid and the errors observed in retrieving mesoscale features also highlight the L4 product's spatial resolution capability. To estimate the spatial resolution of gridded products, an evaluation was done based on a spectral coherence approach. A full description of this approach can be found in Ballarotta et al. (2019). Many applications are derived from these global and regional gridded products and greatly benefit from the products quality: the Lagrangian products (FSLE d'Ovidio et al. 2015) and eddy tracking application (Delepoulle et al., 2018) are prominent examples.

Medium-term developments concern new Level-3 products that will be dedicated to data assimilation and the CMEMS Monitoring Forecasting Centre. The Mean Dynamic Topography will also be updated, and the Black Sea area will be integrated. Finally, a new regional European product will substitute the current Mediterranean and Black Sea products.

In the coming years, DUACS will face major challenges with the arrival of new altimeter missions. SWOT, for example, will observe fine-scale dynamics with swath SSH observations (Morrow et al., 2018) that will need to be integrated into DUACS. The next step, therefore, will consist in moving towards a higher resolution for along-track and gridded products. New mapping techniques should also be taken into consideration and are currently being studied such as dynamical advection (Rogé al., 2017, Ubelmann et al., 2016).

## 6 Data availability

Datasets are available from the CMEMS web-portal (<http://marine.copernicus.eu/services-portfolio/access-to-products/>) and the C3S data store (<https://cds.climate.copernicus.eu> Level 2P (L2P) altimetry products are disseminated by CNES and

EUMETSAT. L2P products are supplied by L2 products that are distributed by different agencies: NASA, NSOAS, ISRO, ESA, CNES, EUMETSAT.

The L3 products for Sentinel-3's altimetry mission are processed at CLS on behalf of EUMETSAT, funded by the European Union.

- 5 The MEDESS-GIB dataset is available through the PANGAEA (Data Publisher for Earth and Environmental Science) repository, with the following DOI:10.1594/PANGAEA.853701. The AlborEx dataset is available at the SOCIB web page (<http://www.socib.eu>).

## **7 Acknowledgements**

- 10 The DT2018 reprocessing exercise was supported by the CNES/SALP project and the CMEMS and C3S services funded by the European Union. Global L3 Sentinel-3 production is coordinated by EUMETSAT and funded by the European Union.

## References

- Ablain, M., Legeais, J.F., Prandi, P., Fenoglio-Marc L., Marcos M., Benveniste J., Cazenave A., Satellite Altimetry-Based Sea Level at Global and Regional Scales, *Surv Geophys* 38, 9-33. doi:10.1007/s10712-016-9389-8, 2017.
- Ablain, M., Meyssignac, B., Zawadzki, L., Jugier, R., Ribes, A., Cazenave, A., and Picot, N.: Uncertainty in Satellite  
5 estimate of Global Mean Sea Level changes, trend and acceleration, *Earth Syst. Sci. Data Discuss.*,  
<https://doi.org/10.5194/essd-2019-10>, in review, 2019.
- AVISO. Annual report (2017) of Mean Sea Level activities, edition 1.0, available at:  
[https://www.aviso.altimetry.fr/fileadmin/documents/calval/validation\\_report/SALP-RP-MA-EA-23189-](https://www.aviso.altimetry.fr/fileadmin/documents/calval/validation_report/SALP-RP-MA-EA-23189-CLS_AnnualReport_2017_MSL.pdf)  
CLS\_AnnualReport\_2017\_MSL.pdf (last access: 4 October 2018), 2017a.
- 10 AVISO. Annual report (2017) of the assessment of Orbit Quality through the Sea Surface Height calculation, available at:  
[https://www.aviso.altimetry.fr/fileadmin/documents/calval/validation\\_report/SALP-RP-MA-EA-](https://www.aviso.altimetry.fr/fileadmin/documents/calval/validation_report/SALP-RP-MA-EA-CLS_1_0_YearlyReportOrbito_2017.pdf)  
[CLS\\_1\\_0\\_YearlyReportOrbito\\_2017.pdf](https://www.aviso.altimetry.fr/fileadmin/documents/calval/validation_report/SALP-RP-MA-EA-CLS_1_0_YearlyReportOrbito_2017.pdf) (last access: 4 October 2018), 2017b.
- AVISO. Cal/Val and cross calibration annual reports. <https://www.aviso.altimetry.fr/en/data/calval/systematic-calval.html>  
(last access: 12 December 2018), 2017c.
- 15 AVISO, Mesoscale Eddy Trajectory Atlas Product Handbook, SALP-MU-P-EA-23126-CLS, issue 2.0  
[https://www.aviso.altimetry.fr/fileadmin/documents/data/tools/hdbk\\_eddytrajectory\\_2.0exp.pdf](https://www.aviso.altimetry.fr/fileadmin/documents/data/tools/hdbk_eddytrajectory_2.0exp.pdf) (last access: 27 May 2019),  
2018.
- Ballarotta, M., Ubelmann, C., Pujol, M.-I., Taburet, G., Fournier, F., Legeais, J.-F., Faugere, Y., Delepoulle, A., Chelton, D.,  
Dibarboure, G., and Picot, N.: On the resolutions of ocean altimetry maps, *Ocean Sci. Discuss.*, <https://doi.org/10.5194/os->  
20 2018-156, in review, 2019.
- Dieng, H. B., Cazenave, A., Meyssignac, B., & Ablain, M. New estimate of the current rate of sea level rise from a sea level  
budget approach. *Geophysical Research Letters*, 2017.
- Carrere, Loren & Lyard, F. Modeling the barotropic response of the global ocean to atmospheric wind and pressure forcing  
Comparisons with observations. *Geophysical Research Letters*. 30. 10.1029/2002GL016473, 2003.
- 25 Carrere L., F. Lyard, M. Cancet, A. Guillot, N. Picot: *FES 2014, a new tidal model - Validation results and perspectives for*  
*improvements*, presentation to ESA Living Planet Conference, Prague, 2016.
- Cartwright, D. E. and Tayler, R. J.: New computations of the tide generating potential, *Geophys. J. R. Astr. Soc.*, 23, 45–74,  
1971.
- Cartwright, D. E. and Edden, A. C.: Corrected tables of tidal harmonics, *Geophys. J. R. Astr. Soc.*, 33, 253–264, 1973.
- 30 Delepoulle A., Faugere Y., Chelton D., Dibarboure G.: A new 25 year mesoscale eddy trajectory atlas on AVISO, poster  
presentation at OSTST 2018, available at:  
[https://meetings.aviso.altimetry.fr/fileadmin/user\\_upload/tx\\_ausycslseminar/files/Poster\\_OSTST2018\\_DUACS\\_eddy\\_aviso.](https://meetings.aviso.altimetry.fr/fileadmin/user_upload/tx_ausycslseminar/files/Poster_OSTST2018_DUACS_eddy_aviso.pdf)  
pdf (last access: 17 April 2019), 2018.

- Desai S., Wahr J., Beckley B.: Revisiting the pole tide for and from satellite altimetry, *J Geod* 89:12333. <https://doi.org/10.1007/s00190-015-0848-7>, 2015.
- Dibarboure G., M-I. Pujol, F. Briol, P.-Y. Le Traon, G. Larnicol, N. Picot, F. Mertz, P. Escudier, M. Ablain, and C. Dufau: Jason-2 in DUACS: first tandem results and impact on processing and products, *Mar. Geod.*, OSTM Jason-2 Calibration/Validation Special Edition – Part 2, (34), 214-241, doi:10.1080/01490419.2011.584826, 2011
- 5 Dibarboure G., Pujol M-I., “Improving the quality of Sentinel-3A with a hybrid mean sea surface model, and implications for Sentinel-3B and SWOT”, in review.
- Dieng, H. B., Cazenave, A., Meyssignac, B., & Ablain, M. New estimate of the current rate of sea level rise from a sea level budget approach. *Geophysical Research Letters*, 2017.
- 10 d’Ovidio, F., Della Penna, A., Trull, T. W., Nencioli, F., Pujol, M.I., Rio, M.-H., Park, Y.-H., Cotté, C., Zhou, M., and Blain, S.: The biogeochemical structuring role of horizontal stirring: Lagrangian perspectives on iron delivery downstream of the Kerguelen Plateau, *Biogeosciences*, 12, 5567–5581, doi:10.5194/bg12-5567-2015, 2015.
- Ducet, N., Le Traon, P.-Y., and Reverdun, G.: Global highresolution mapping of ocean circulation from TOPEX/Poseidon and ERS-1 and -2, *J. Geophys. Res.*, 105, 19477–19498, 2000.
- 15 Escudier, P., Couhert, A., Mercier, F., Mallet, A., Thibaut, P., Tran, N., Amarouche, L., Picard, B., Carrère, L., Dibarboure, G., Ablain, M., Richard, J., Steunou, N., Dubois, P., Rio, M. H., and Dorandeu, J.: Satellite radar altimetry: principle, geophysical correction and orbit, accuracy and precision, in: *Satellite Altimetry Over Oceans and Land Surfaces*, edited by: Stammer, D. and Cazenave, A., CRC Press, Taylor & Francis, Boca Raton, 2017.
- Fernandes MJ, La´zaro C, Ablain M, Pires N: Improved wet path delays for all ESA and reference altimetric missions.
- 20 Remote Sens Environ 169(2015):50–74. doi:10.1016/j.rse.2015.07.023, 2015.
- Gaspar P., Ogor F., Le Traon P.Y., Zanife O.Z.: Estimating the sea state bias of the TOPEX and POSEIDON altimeters from crossover differences - *Journal of Geophysical Research: Oceans*, 1994.
- Guibbaud, M., A. Ollivier and M. Ablain, A new approach for dual-frequency ionospheric correction filtering, ENVISAT Altimetry Quality Working Group (QWG), available in the Section 8.5 of the 2012 Envisat annual activity report:
- 25 [http://www.aviso.altimetry.fr/fileadmin/documents/calval/validation\\_report/EN/annual\\_report\\_e\\_n\\_2012.pdf](http://www.aviso.altimetry.fr/fileadmin/documents/calval/validation_report/EN/annual_report_e_n_2012.pdf), 2015
- Ijima, B. A., Harris, I. L., Ho, C. M., Lindqwiste, U. J., Mannucci, A. J., Pi, X., Reyes, M. J., Sparks, L. C., and Wilson, B. D.: Automated daily process for global ionospheric total electron content maps and satellite ocean altimeter ionospheric calibration based on Global Positioning System data, *J. Atmos. Sol.-Terr. Phys.*, 61, 16, 1205–1218, 1999.
- Keihm S. J., “TOPEX/Poseidon Microwave Radiometer (TMR): II. Antenna Pattern Correction and Brightness Temperature
- 30 Algorithm,” *IEEE Trans. Geosci. Remote Sens.*, vol. 33, no. 1, pp. 138–146, 1995.
- Legeais, J.-F., Prandi, P., and Guinehut, S.: Analyses of altimetry errors using Argo and GRACE data, *Ocean Sci.*, 12, 647-662, doi:10.5194/os-12-647-2016, 2016.
- Legeais, J.-F., Ablain, M., Zawadzki, L., Zuo, H., Johannessen, J. A., Scharffenberg, M. G., Fenoglio-Marc, L., Fernandes, M. J., Andersen, O. B., Rudenko, S., Cipollini, P., Quartly, G. D., Passaro, M., Cazenave, A., and Benveniste, J.: An



- improved and homogeneous altimeter sea level record from the ESA Climate Change Initiative, *Earth Syst. Sci. Data*, 10, 281–301, <https://doi.org/10.5194/essd-10-281-2018>, 2018a.
- Legouis J.-F., K. von Schuckmann, A. Melet, A. Storto and B. Meyssignac: Sea Level. In von Schuckmann et al., 2018. The Copernicus Marine Environment Monitoring Service Ocean State Report, *J Oper Oceanogr*, 11:sup1, s1–s142, DOI:10.1080/1755876X.2018.1489208, 2018b.
- Le Traon, P.-Y. and Ogor, F.: ERS-1/2 orbit improvement using TOPEX/POSEIDON: The 2 cm challenge, *J. Geophys. Res.*, 103, 8045–8057, 1998.
- LeTraon, P.-Y, Faugere, Y., Hernamdez, F., Dorandeu, J., Mertz, F., and Abalin, M.: Can We Merge GEOSAT Follow-On with TOPEX/Poseidon and ERS-2 for an Improved Description of the Ocean Circulation? *J. Atmos. Ocean. Technol.*, 20, 889–895, 2003.
- Lumpkin, R., Grodsky, S., Rio, M.-H., Centurioni, L., Carton, J., and Lee, D.: Removing spurious low-frequency variability in surface drifter velocities, *J. Atmos. Ocean. Technol.*, 30, 353–360, doi:10.1175/JTECH-D-12-00139.1, 2013.
- Manca, B., Burca, M., Giorgetti, A., Coatanoean, C., Garcia, M.-J., Ion, A.: Physical and biochemical averaged vertical profiles in the Mediterranean regions: an important tool to trace the climatology of water masses and to validate incoming data from operational oceanography. *J. Mar. Syst.* 48, 83–116. <http://dx.doi.org/10.1016/j.jmarsys.2003.11.025>, 2004.
- Mertz F., Mercier F., Labroue S., Tran N., Dorandeu J.: ERS-2 OPR data quality assessment long-term monitoring – Particular investigation. CLS.DOS.NT-06-001 [https://www.aviso.altimetry.fr/fileadmin/documents/calval/validation\\_report/E2/annual\\_report\\_e2\\_2005.pdf](https://www.aviso.altimetry.fr/fileadmin/documents/calval/validation_report/E2/annual_report_e2_2005.pdf), 2005.
- Morrow, R., Blumstein, D., & Dibarboure, G.: Fine-scale Altimetry and the Future SWOT Mission. *New Frontiers In Operational Oceanography*. Retrieved from [http://purl.flvc.org/fsu/fd/FSU\\_libsubv1\\_scholarship\\_submission\\_1536170512\\_b3d57dea](http://purl.flvc.org/fsu/fd/FSU_libsubv1_scholarship_submission_1536170512_b3d57dea), 2018.
- Obligis, Estelle & Rahmani, Abdel & Eymard, Laurence & Labroue, Sylvie & Bronner, Emilie. An Improved Retrieval Algorithm for Water Vapor Retrieval: Application to the Envisat Microwave Radiometer. *Geoscience and Remote Sensing, IEEE Transactions on*. 47. 3057 - 3064. 10.1109/TGRS.2009.2020433, 2009.
- Ollivier A., Guibbaud M., Faugere Y., Labroue S., Picot N., Boy F. Cryosat-2 altimeter performance assessment over ocean. 2014 Ocean Surface Topography Science Team Meeting oral presentation. Available at: [https://meetings.aviso.altimetry.fr/fileadmin/user\\_upload/tx\\_ausyclsseminar/files/29Ball1615-7\\_Pres\\_PerfoC2\\_Ollivier.pdf](https://meetings.aviso.altimetry.fr/fileadmin/user_upload/tx_ausyclsseminar/files/29Ball1615-7_Pres_PerfoC2_Ollivier.pdf) (last access: 4 October 2018), 2014.
- Ollivier A., Philipps S., Couhert A., Picot N. Assessment of Orbit Quality through the SSH calculation: POE-E orbit standards. Ocean Surface Topography Science Team Meeting 2015 presentation Available at: [https://meetings.aviso.altimetry.fr/fileadmin/user\\_upload/tx\\_ausyclsseminar/files/Poster\\_OSTST15\\_Orbit.pdf](https://meetings.aviso.altimetry.fr/fileadmin/user_upload/tx_ausyclsseminar/files/Poster_OSTST15_Orbit.pdf) (last access: 4 October 2018), 2015.
- Pascual A., Y. Faugere, G. Larnicol, P.-Y. Le Traon: Improved description of the ocean mesoscale variability by combining four satellite altimeters, *Geophys. Res. Lett.*, 33(2), doi:10.1029/2005GL024633, 2006.

- Pascual A, Ruiz S, Olita A, Troupin C, Claret M, Casas B, Mourre B, Poulain P-M, Tovar-Sanchez A, Capet A, Mason E, Allen JT, Mahadevan A and Tintoré J.: A Multiplatform Experiment to Unravel Meso- and Submesoscale Processes in an Intense Front (AlborEx). *Front. Mar. Sci.* 4:39. doi: 10.3389/fmars.2017.00039, 2017.
- Permanent Service for Mean Sea Level (PSMSL): “Tide Gauge Data”, available at: <http://www.psmsl.org/data/obtaining/>,  
5 (last access 1 June 2014), 2016.
- Picard B., Frery M. L., Obligis E., Eymard L., Steunou N., and Picot N., “SARAL/AltiKa Wet Tropospheric Correction: In-Flight Calibration, Retrieval Strategies and Performances,” *Mar. Geod.*, vol. 38, no. May, pp. 277–296, 2015.
- Pujol, M.-I., Faugère, Y., Taburet, G., Dupuy, S., Pelloquin, C., Ablain, M., and Picot, N.: DUACS DT2014: the new multi-mission altimeter data set reprocessed over 20 years, *Ocean Sci.*, 12, 1067-1090, doi:10.5194/os-12-1067-2016, 2016.
- 10 Pujol I, Schaeffer P, Faugere Y, Raynal M, Dibarboure G, Picot N.: Gauging the Improvement of Recent Mean Sea Surface Models: A New Approach for Identifying and Quantifying Their Errors. *Journal of Geophysical Research: Oceans*. 10.1029/2017JC013503, 2018a.
- Pujol M-I., Schaeffer P., Faugère Y., Davanne F-X., Dibarboure G., Picot N.: Improvements and limitations of recent mean sea surface models: importance for Sentinel-3 and SWOT, Poster at 2018 Ocean Surface Topography Science Team  
15 Meeting,  
[https://meetings.aviso.altimetry.fr/fileadmin/user\\_upload/tx\\_ausycslseminar/files/OSTST2018\\_PMS3A\\_Poster\\_Pujol.pdf](https://meetings.aviso.altimetry.fr/fileadmin/user_upload/tx_ausycslseminar/files/OSTST2018_PMS3A_Poster_Pujol.pdf)  
(last access December 12, 2018), 2018b.
- Quartly, G. D., Legeais, J.-F., Ablain, M., Zawadzki, L., Fernandes, M. J., Rudenko, S., Carrère, L., García, P. N., Cipollini, P., Andersen, O. B., Poisson, J.-C., Mbajon Njiche, S., Cazenave, A., and Benveniste, J.: A new phase in the production of  
20 quality-controlled sea level data, *Earth Syst. Sci. Data*, 9, 557-572, <https://doi.org/10.5194/essd-9-557-2017>, 2017.
- Rio, M. H., Guinehut, S., and Larnicol, G.: New CNES-CLS09 global mean dynamic topography computed from the combination of GRACE data, altimetry, and in-situ measurements, *J. Geophys. Res.*, 116, C07018, doi:10.1029/2010JC006505, 2011.
- Rio, M.-H.: Use of altimeter and wind data to detect the anomalous loss of SVP-type drifter’s drogue, *J. Atmos. Ocean. Technol.*, 1663–1674, doi:10.1175/JTECH-D-12-00008.1, 2012.  
25
- Rogé, M., Morrow, R., Ubelmann, C., Dibarboure, G.: Using a dynamical advection to reconstruct a part of the SSH evolution in the context of SWOT, application to the Mediterranean Sea. *Ocean Dynamics*. 1-20. 10.1007/s10236-017-1073-0, 2017.
- Rudenko, S., Otten, M., Visser, P., Scharroo, R., Schöne, T., and Esselborn, S.: New improved orbit solutions for the ERS-1  
30 and ERS-2 satellites, *Adv. Space Res.*, 49, 1229–1244, 2012.
- Scharroo, R. and Smith, W. H. F.: A global positioning system based climatology for the total electron content in the ionosphere, *J. Geophys. Res.*, 115, A10318, doi:10.1029/2009JA014719, 2010.
- Sotillo, M. G., Garcia-Ladona, E., Orfila, A., Rodríguez-Rubio, P., Maraver, J. C., Conti, D., Padorno, E., Jiménez, J. A., Capó, E., Pérez, F., Sayol, J. M., de los Santos, F. J., Amo, A., Rietz, A., Troupin, C., Tintore, J., and Álvarez-Fanjul, E.:

The MEDESS-GIB database: tracking the Atlantic water inflow, *Earth Syst. Sci. Data*, 8, 141-149, doi:10.5194/essd-8-141-2016, 2016.

Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *J. Geophys. Res.*, 106, 7183–7192, doi:10.1029/2000JD900719, 2001.

- 5 Tran, N., Labroue, S., Philipps, S., Bronner, E., and Picot, N.: Overview and Update of the Sea State Bias Corrections for the Jason-2, Jason-1 and TOPEX Missions, *Mar. Geod.*, 33, 348– 362, 2010.

Tran N., Philipps, S., Poisson, J.-C., Urien, S., Bronner, E., and Picot, N.: Impact of GDR\_D standards on SSB corrections, Presentation OSTST2012 in Venice, [http://www.aviso.altimetry.fr/fileadmin/documents/OSTST/2012/oral/02\\_friday\\_28/01\\_instr\\_processing\\_I/01\\_IP1\\_Tran.pdf](http://www.aviso.altimetry.fr/fileadmin/documents/OSTST/2012/oral/02_friday_28/01_instr_processing_I/01_IP1_Tran.pdf) (last access: 31 August 2016), 2012.

- 10 Ubelmann, C., Cornuelle, B., Fu L.-L. : Dynamic Mapping of Along-Track Ocean Altimetry: Method and Performance from Observing System Simulation Experiments. *Journal of Atmospheric and Oceanic Technology*. 33. 10.1175/JTECH-D-15-0163.1, 2016.

Valladeau, G., Legeais, J.-F., Ablain, M., Guinehut, S., and Picot, N.: Comparing Altimetry with Tide Gauges and Argo Profiling Floats for Data Quality Assessment and Mean Sea Level Studies, *Mar. Geod.*, OSTM Jason-2 Applications Special Edition – Part 3, 35, 42–60, doi:10.1080/01490419.2012.718226, 2012.

- 15 Watson, C. S., White, N. J., Church, J. A., King, M. A., Burgette, R. J., and Legresy, B.: Unabated global mean sea level over the satellite altimeter era, *Nat. Clim. Change*, 5, 565–568, DOI: 10.1038/NCLIMATE2635, 2015.

WCRP Global Sea Level Budget Group: Global sea-level budget 1993–present, *Earth Syst. Sci. Data*, 10, 1551-1590, <https://doi.org/10.5194/essd-10-1551-2018>, 2018.

20

25

**Table 1: Altimeter standards used in DT2018. Changes with the DT2014 solution are underlined in bold format.**

	J3	J2	J1	TP	ERS-1	ERS-2	EN	GFO	C2	AL	H2A	S3A
Orbit	POE-E	POE-E		GFSC STD15 until cycle 365, STD12 afterwards	Reaper [Rudenko et al., 2012]		POE-D	GSFC	POE-E		POE-D	POE-E
Sea State Bias	Non-Parametric SSB [Tran et al., 2012]		SSB issued from GDR-E	Non- parametric SSB [Tran et al., 2010]	BM3 [Gaspar et al., 1994]	Non- parametric [Mertz et al., 2005]	Non-Parametric SSB [Tran et al., 2012]	Non- Parametric SSB [Tran et al., 2010]	Non- Parametric SSB from J1 with unbiased sigma0.	Non- Parametric SSB [Tran et al., 2012]	Non- Parametric SSB from J1	Non- Parametric SSB [Tran et al., 2012]
Ionospheric	Filtered dual- frequency altimeter range measurements [Guibbaud et al., 2015]	Filtered dual- frequency altimeter range measurements [Guibbaud et al., 2015]	Filtered dual-frequency altimeter range measurements [Guibbaud et al., 2015]		Reaper NIC09 model [Scharroo et al., 2010]	Cycle<37 Reaper NIC09 model [Scharroo et al., 2010] Cycle>36 GIM [Ijima et al., 1999]	Dual-frequency altimeter range measurement [Guibbaud et al., 2015] (6≤cycles≤64)/GIM [Ijima et al., 1999] Corrected for 8mm bias (c≥65)	GIM [Ijima et al., 1999]		GIM [Ijima et al., 1999]		Filtered dual- frequency altimeter range measurements [Guibbaud et al., 2015]
Wet troposphere	From J3-AMR radiometer	Neural Network correction [Keihm et al. 1995]	JMR issued from GDR-E	GNSS derived Path Delay [Fernandes et al., 2015]			Neural Network correction (5 entries) [Obligis et al., 2009 and Picard et al., 2015]	From GFO radiometer	From ECMWF model	Neural Network correction (5 entries) [Obligis et al., 2009 and Picard et al., 2015]	From ECMWF model	From S3A- AMR radiometer
Dry troposphere	Model based on ECMWF Gaussian grids		Model based on ECMWF rectangular grids	Model based on ERA-INTERIM			Model based on ECMWF Gaussian grids	Model based on ECMWF rectangular grids	Model based on ECMWF Gaussian grids	Model based on ECMWF Gaussian grids		Model based on ECMWF Gaussian grids
Dynamic Atmospheric Correction	MOG2D High frequencies forced with analysed ECMWF pressure and wind field [Carrere et al., 2003; operational version used, current version is 3.2.0] + inverse barometer Low frequencies			MOG2D High frequencies forced with analysed ERA-INTERIM pressure and wind field + inverse barometer Low frequencies			MOG2D High frequencies forced with analysed ECMWF pressure and wind field [Carrere et al., 2003; operational version used, current version is 3.2.0] + inverse barometer Low frequencies			MOG2D High frequencies forced with analysed ECMWF pressure and wind field [Carrere et al., 2003; operational version used, current version is 3.2.0] + inverse barometer Low frequencies		MOG2D High frequencies forced with analysed ECMWF pressure and wind field [Carrere et al., 2003;

					operational version used, current version is 3.2.0] + inverse barometer Low frequencies
Ocean tide	FES2014 [Carrere et al., 2016]				
Pole tide	[Desai et al., 2015]				
Solid earth tide	Elastic response to tidal potential [Cartwright and Tayler, 1971], [Cartwright and Edden, 1973]				
Mean Sea Surface	CNES-CLS-2015 [Pujol et al., 2018a]				

**Table 2: Time periods and cycles used to compute Mean Profile in the DT2018 version.**

	<b>Satellite used in Mean Profile computation</b>	<b>Periods used in Mean Profile computation</b>	<b>Cycles</b>
<b>Topex/Poseidon – Jason-1 – Jason-2 – Jason-3</b>	Topex/Poseidon	January 1993 – April 2002 (9 years)	11 – 353
	Jason-1	April 2002 – October 2008 (6 years)	10 – 249
	OSTM/Jason-2	October 2008 – December 2015 (7 years)	10 – 273
<b>Ers-1 – Ers-2 – Envisat - AltiKa</b>	Ers-2	Mai 1995 – January 2000 (5 years)	1 – 49
	Envisat	October 2002 – October 2010 (8 years)	10 – 94
	AltiKa	March 2013 – March 2015 (2 years)	1 – 22
<b>Topex/Poseidon Interleaved orbit – Jason-1</b>	Topex/Poseidon Interleaved orbit	September 2002 – October 2005 (3 years)	368 – 481
<b>Interleaved orbit – Jason-2 Interleaved orbit</b>	Jason-1 Interleaved orbit	February 2009 – March 2012 (3 years)	262 – 374
<b>Geosat Follow On</b>	Geosat Follow On	January 2000 – September 2008 (8 years)	37 – 222

**Table 3: Variance of the differences between gridded (L4) DT2018 two-sat-merged products and independent TP interleaved along-track measurements for different geographic selections (unit = cm<sup>2</sup>). In parenthesis: variance reduction (in %) compared with the results obtained with the DT2014 products. Statistics are presented for wavelengths ranging between 65-500 km and after latitude selection (|LAT|<60°).**

	<b>TP [2003-2004]</b>
<b>Reference area*</b>	1.4 (-0.3%)
<b>Low variability (&lt;200 cm<sup>2</sup>) &amp; offshore (distance coast &gt;200 km) areas</b>	5.0 (-3.0%)
<b>High variability (&gt;200 cm<sup>2</sup>) &amp; offshore (distance coast &gt;200 km) areas</b>	37.7 (-3.1%)
<b>Coastal areas (distance coast &lt; 200km)</b>	8.2 (-10.1%)
<b>Intertropical belt (±20°N)</b>	4.8 (-9.1%)

*\*The reference area is defined by [330, 360°E]; [-22, -8°N] and corresponds to a very low-variability area (between 0 and 7 cm<sup>2</sup>) in the South Atlantic subtropical gyre where the observed errors are small.*

**Table 4: Variance of the differences between gridded geostrophic current (L4) DT2018 products and independent drifter measurements (unit = cm2/s2). In parenthesis: variance reduction (in %) compared with the results obtained with the DT2014 products. Statistics are presented for latitude selection (5°N<|LAT| < 60°N).**

	Zonal	Meridional
Reference area*	44.3 (-1.8%)	33.4 (-0.9%)
Low variability (<200 cm <sup>2</sup> ) & offshore (distance coast >200 km) areas	91.6 (-6.1%)	88.6 (-6.7%)
High variability (>200 cm <sup>2</sup> ) & offshore (distance coast >200 km) areas	229.6 (-4.3%)	260.5 (-4.5%)
Coastal areas (distance coast < 200km)	189.7 (-14.7%)	195.3 (-15.5%)
Intertropical belt (±20°N)	170.5 (-18.8%)	176.2 (-37.9%)

*\*The reference area is defined by [330, 360°E]; [-22, -8°N] and corresponds to a very low-variability area (between 0 and 7 cm<sup>2</sup>) in the South Atlantic subtropical gyre where the observed errors are small*

**Table 5: Variance of the differences between gridded (L4) DT2018 two-sat-merged regional Mediterranean (first line) and Black sea (second line) products and independent TP interleaved along-track measurements without filtering over the time period 2003-2004 (unit = cm<sup>2</sup>). In parenthesis: variance reduction (in %) compared with the results obtained with the DT2014 products.**

	TP [2003-2004] unfiltered
Mediterranean Sea product	16.7 (-4.2%)
Black Sea product	23.2 (-3.5%)

**Table 6: RMSE (m/s) and correlation coefficient between the absolute geostrophic velocities derived from DT-2018 regional products for the Mediterranean Sea; and absolute surface velocities as obtained from drifters collected in the basin. The variance of the datasets (m<sup>2</sup>/s<sup>2</sup>) and the data used to conduct the comparison are also displayed.**

	DT-2018 regional	DUACS-DT2018 improvements
R	0.49	4 %
RMS diff (m/s)	0.12	1 %
variance drifter (m <sup>2</sup> /s <sup>2</sup> )	0.017	-
variance altimetry (m <sup>2</sup> /s <sup>2</sup> )	0.008	14 %





Figure 1: Timeline of the altimeter missions used in the multi-mission DUACS DT2018 system.

5


10






**Figure 2: Gain of measurements in the Topex/Poseidon-Jason1-OSTM/Jason-2 Mean Profile used in DT2018 compared to DT2014. Gain of points in DT2018 is in red, loss of points is in blue.**

5




**Figure 3: Difference between SLA variance observed with DT2018 gridded products and SLA variance observed with DT2014 gridded products over the 1993-2017 period.**

10




**Figure 4: Map of the difference between mean EKE for DT2018 and DT2014 gridded products (left frame) and evolution of the mean EKE over global ocean, computed from DT2014 (blue line) and DT2018 (red line) SLA gridded products over the 1993-2017 period. The  $\pm 5^\circ\text{N}$  equatorial belt has been removed.**

5



**Figure 5: Difference of the RMS of the difference between gridded SLA products and independent Topex/Poseidon interleaved along-track SLA measurements, using successively DT2018 and DT2014 version. Negative values represent reduced differences between DT2018 altimetry products and independent along-track measurements.**

10



5 **Figure 6: Upper panels: Zonal (left) and meridional (right) RMS of the difference between DUACS DT2018 absolute geostrophic current and drifter measurements over the 1993-2017 period. Lower panels: Zonal (left) and meridional (right) difference of the RMS of the altimeter geostrophic currents minus drifters measurements, using successively DT2018 and DT2014 gridded products. Negative values represent reduced differences between DT2018 altimetry products and drifters. The statistic is expressed as a percentage of RMS of drifter measurements. Statistics have been computed in boxes of 5°x5°. Boxes with less than 1000 points have been masked.**




Figure 7: Difference of the variance between gridded SLA products and TG, using successively DT2018 and DT2014 gridded products. We used mean monthly TG measurements from the PSMSL network. Negative values represent reduced differences between DT2018 altimetry gridded SLA and TG. The statistic is expressed as a percentage of RMS of TG measurements.

5






Figure 8: Left panel: temporal evolution of the GMSL estimated from DT2018 (red line) and DT2014 (blue line) gridded SLA products. The annual and semi-annual signals were adjusted and no GIA correction was applied. Right panel: map of the differences of the local MSL trend estimated from the DT2018 and DT2014 gridded SLA products. MSL was estimated over the 1993-2017 period.



**Figure 9:** Difference of the RMS of the difference between gridded regional Mediterranean Sea (left frame) and regional Black Sea (right frame) SLA products and independent Topex/Poseidon interleaved along-track SLA measurements, using successively DT2018 and DT2014 version. Negative values represent reduced differences between DT2018 altimetry products and independent along-track measurements. The statistic is expressed as a percentage of RMS of the independent along-track product.

5



**Figure 10:** Difference of the variance between regional Mediterranean gridded products (upper frame) and regional Black Sea products (lower frame) SLA products and TG, using successively DT2018 and DT2014 gridded products. We used mean monthly TG measurements from the PSMSL network. Negative values represent reduced differences between DT2018 altimetry gridded SLA and TG. The statistic is expressed as a percentage of RMS of TG measurements. The statistic is expressed as a percentage of RMS of the independent along-track product.

10






Figure 11: (a) Map of the Mediterranean Sea showing the geographical limits and the nomenclatures of the regions (blue boxes) as defined in Manca et al. (2004) where drifter data is available in the western sub-basin: Alboran Sea (DS1), Balearic Sea (DS2), western and eastern Algerian (DS3 and DS4), Algero-Provençal (DF1), Liguro-Provençal (DF3, DF4), Gulf of Lion (DF2), Tyrrhenian Sea (DT1), Sardinian channel (DI1), Tyrrhenian Sea (DT2, DT3) and Sicily Strait (DI3); and in the eastern sub-basin: Adriatic Sea (DJ1, DJ2, DJ3), Ionian Sea (DJ4, DJ5, DJ6, DJ7, DJ8), Aegean Sea (DH1, DH2), Cretan Passage (DH3) and Levantine basin (DL1, DL2, DL3, DL4). Left column: maps of the Mediterranean Sea showing the comparison between DT-2018 regional altimetry product with the drifter in-situ observations within the geographical limits and the nomenclatures of the regions

defined in (a). The statistical parameters showed are: (b) RMS difference; (c) correlation coefficient; (d) altimetry variance and (e) drifter variance. Right column: improvements (%) of the comparisons between the DT-2018 regional product and drifter in-situ observations with respect to the comparisons by using the DT-2014 product within the geographical limits and the nomenclatures of the regions defined in (a). The statistical parameters showed are: (f) RMS difference; (g) correlation coefficient and (h) altimetry variance. Positive values denote an improvement of DT-2018 regional product over DT-2014.