Reponse review #2 Tom Farrar

The authors have improved the manuscript dramatically in their revised version. I believe the new analysis method is sound, and the description of what was done is also much clearer. It seems like a substantially different paper than the previous version I reviewed, and I am happy to recommend publication after minor revisions. I would also like to thank the authors for their pleasant and constructive attitude in responding to my previous, critical review. (There were actually two versions of the previous review-- I made an updated one with more detail and gave it to the authors directly, but the OSD system would not allow me to upload that review because the online discussion period had ended. I am uploading that review now.)

My only substantial reservation about this version of the manuscript has to do with the approach and conclusion about the temporal resolution. Something does not add up, as described in the following paragraph. I suggest the authors investigate this a little more and either add this as a caveat and change the plot and conclusions accordingly or adjust the analysis to have finer frequency resolution.

Authors response (AR): We would like to thank Tom Farrar for the review and the suggestions he made on this new version of the manuscript. We provide below the answers to the two main concerns he raised.

Page 4, lines 26+: There is another important consequence of choosing to analyze 100-day segments. This sets the coarseness of the resulting evaluation of temporal resolution, by setting the frequency resolution of the spectral analysis. For 100-day segments, the frequency resolution is 1/100 cycles per day (cpd) (actually, the frequency resolution is effectively somewhat less because of the taper window, but I think it is the frequency grid that matters here). If the record chunks are actually 100 days long as stated (instead of 100 data points long), the frequency bands will be centered at frequencies corresponding to periods of 101, 50.5, 33.67, 25.25, 20.20, 16.83, and 14.43 days. This means that the analysis should only be able to coarsely determine the temporal resolution (e.g., 50.5 days versus 33.67 days). It seems to me that the color scale chosen for displaying the results is thus misleading, or else, there has been some kind of interpolation of the spectral results to arrive at values between 34 and 50 days. It is not clear to me that such interpolation would be a valid approach.

AR: We agree that the choice of 100-day long segment set the coarseness of the resulting evaluation. In our analysis, we set the segment length to 100 days long to get many realizations. It also corresponds to 100 continuous data points for both tide gauge and map datasets. In other words, segments that are shorter than 100 data points (100-days long) are excluded from the analysis. Consequently, analysis at some tide gauges are lost when choosing longer segment sizes. In total, we found that 2 sensors are lost when choosing 300 days long-segments, 3 sensors when choosing 1 year-long segment. When choosing 2-year long period, we lose 20 sensors. Since the number of lost sensors between 100-days to 365

days periods not significative (for a total of 190 tide gauges datasets), we propose to use the analysis performed over segment that are 365-days long to get finer spectral resolution. We modify the manuscript accordingly (key value and figures 3 and 5).

Additionally, we attach below the results to a sensitivity study we performed for 100 data points (100 days) long segment, 200 data points segments (200 days) and 365 data points segments (1 year). As for the estimation of the spatial scales, we linearly interpolate between successive NSR values to find the wavelength corresponding to the NSR of 0.5. This linear interpolation indeed explains the value found between frequency scales. We illustrate that the segment length has a weak impact on the estimation of the effective temporal resolution from 40° to the poles. Larger difference is found in the inter-tropical band (cf. Fig.4 below: zonally averaged time scale for various segment size). Figure 1 to 3 show the maps of estimated effective temporal resolution using 100 days long segment, 200 days long segment and 365 days long segment. The spatial distribution of the temporal resolution looks relatively similar between each test. To illustrate the differences, we plot the zonally averaged effective resolutions for each segment sizes in Figure 4. The largest difference take place in the inter-tropical band.

Test#1: segment length = 100 days corresponding to frequency (in days):



Fig. 1 : Effective temporal resolution computed from 100 days long segment

Test#2: segment length = 200 days corresponding to frequency (in days):

200.0, 100.0, 66.7, 50.0, 40.0, 33.3, 28.6, 25.0, 22.2, 20.0, 18.2, 16.7, 15.4, 14.3, 13.3, 12.5, 11.8, 11.1, 10.5, 10.0, 9.5, 9.1, 8.7, 8.3, 8.0 ...



Fig 2.: Effective temporal resolution computed from 200 days long segment

Test#3: segment length = 365 days corresponding to frequency (in days): 365.0 182.5 121.7 91.2 73.0 60.8 52.1 45.6 40.6 36.5 33.2 30.4 28.1 26.1 24.3 22.8 21.5 20.3 19.2 18.2 17.4 16.6 15.9 15.2 14.6 14.0 13.5 13.0 12.6 12.2 11.8 11.4 11.1 10.7 10.4...



Fig. 3: Effective temporal resolution computed from 365 days long segment



Fig.4: Zonally averaged effective temporal resolution for segments size of 100 data points, 200 data points and 365 data-points. Black line is two times the zonally averaged correlation scales used in the DUACS system.

Page 6, line 6: I do not think Figure 3 supports the claim that midlatitude temporal resolution is 14-28 days. I see lots of points in midlatitudes that are dark red (~45 days). Furthermore, as noted above, something seems inconsistent about the frequency resolution of the analysis and the higher frequency resolution at which the results are being stated.

AR: We updated the value in the manuscript: The zonally averaged resolution ranges from 14 days to 45 days from mid-to-high latitude, as seen on Figure 4 above.

Minor comment:

Figure B1c: What are the white areas?

AR: We updated the figures caption to inform what are the white area. They correspond to the regions where the SNR threshold criterion is not achieved.

On the resolutions of ocean altimetry maps

Maxime Ballarotta¹, Clément Ubelmann¹, Marie-Isabelle Pujol¹, Guillaume Taburet¹, Florent Fournier¹, Jean-François Legeais¹, Yannice Faugère¹, Antoine Delepoulle¹, Dudley Chelton², Gérald Dibarboure³, Nicolas Picot³

¹Collecte Localisation Satellite, Ramonville-Saint-Agne, 31520, France ²College of Earth, Ocean and Atmospheric Sciences, Oregon State University, Corvallis, OR ³Centre National d'Études Spatiales, Toulouse, 31400, France

Correspondence to: M. Ballarotta (mballarotta@groupcls.com)

Abstract. The DUACS system produces sea level global and regional maps that serve oceanographic applications, climate forecasting centers, geophysics and biology communities. These maps are generated using an optimal interpolation method applied to altimeter observations. They are provided on a global ¼°x ¼° (longitude x latitude) and daily grid resolution framework (1/8°x1/8° longitude x latitude grid for the regional products) through the Copernicus Marine Environment Monitoring Service (CMEMS). Yet, the dynamical content of these maps does not have a full ¼° spatial and 1-day resolution, due to the filtering properties of the optimal interpolation. In the present study, we estimate the "effective" spatial and temporal

- 15 resolutions of the newly reprocessed delayed-time DUACS maps (aka, DUACS-DT2018). Our approach is based on the ratio between the spectral content of the mapping error and the spectral content of independent true signals (along-track and tide gauge observations), also known as the Noise-to-Signal ratio. We found that the spatial resolution of the DUACS-DT2018 global maps based on sampling by three altimeters simultaneously ranges from ~100km-wavelength at high latitude to ~800km-wavelength in the Equatorial band and the mean temporal resolution is ~34, days period. The mean effective spatial resolution at mid-latitude is estimated to ~200km. The mean effective spatial resolution is ~130 km for the regional Mediterranean Sea and for the regional Black Sea products. An inter-comparison with previous DUACS reprocessing systems (aka, DUACS-DT2010 and DUACS-DT2014) highlights the progress of the system over the past 8 years, in particular a gain of resolution in highly turbulent regions. The same diagnostic applied to maps constructed with two altimeters and maps with three altimeters confirms a modest increase of resolving capabilities and accuracies in the DUACS maps with the number of 25 missione.
- 25 missions.

1 Introduction

The Data Unification and Altimeter Combination System (DUACS) generates, as part of the CNES/SALP project and the Copernicus Marine Environment and Monitoring Service (CMEMS), delayed-time (DT) multi-mission altimeter Sea Level Anomaly (SLA) Level 3 (along-track cross-calibrated) and Level 4 (multiple sensors merged as maps or time series) products.

30 A full reprocessing of these products is carried out approximately every 3 years and covers the period 1993 - now. The

Supprimé: 28

reprocessing benefits from improvements associated with optimized mapping parameters and new altimeter corrections which are based on standards recommended for altimeter products by the different agencies and expert groups (Ocean Surface Topography Science Team (OSTST), the ESA Quality Working groups and the ESA Sea Level Climate Change Initiative project members). The previous reprocessing was released in 2014 (DUACS-DT2014, see Pujol et al., 2016) and the new release, namely DUACS-DT2018, is available since April 2018 (Taburet et al., 2019).

- The Level 4 DUACS-DT global maps are constructed from optimal interpolation (Bretherton et al., 1976, Le Traon, 1998, Ducet et al., 2000) of Level 3 altimeter observations and are provided on a regular ¼°x ¼° longitude x latitude and daily grid resolution framework (1/8°x1/8° horizontal sampling for the regional Mediterranean and Black Sea products). However, the optimal interpolation used in DUACS does not allow the restitution of the full dynamical spectrum of the ocean, limiting the capability of retrieving small mesoscale in Level 4 products (Chelton et al, 2011 and 2014).
- The "effective" resolution corresponds to the spatio-temporal scales of the features that can be properly resolved in the maps. The spatio-temporal resolution of the previous Level 4 global SLA products was estimated by Chelton et al. (2003, 2011, 2014) based on estimates of the mapping errors in Sea Surface Height (SSH) fields constructed from altimeter data, or spectral ratio analysis between maps and along-track altimeter data. Their analysis suggested mid-latitude spatial resolution capability
- 15 of the observations ranging from ~2° to 6°, depending on the number of altimeters used in the merging and the sampling pattern of the ground track (~2° for tandem mission T/P-Jason 5 days offset between parallel tracks, 6° for T/P mono-mission merging). The temporal resolution capability of the observations for a tandem T/P-Jason mission was estimated ~20 days. In the present study, we further investigate the effective resolution of the DUACS-DT gridded products using a spectral approach. The objective of the paper is threefold: 1) to deliver the spatial distribution of the effective resolutions as key
- 20 information to the users about the quality and the limitations (in term of resolution) of the newly produced DUACS-DT2018 gridded products, 2) to access and compare the spatial and temporal resolution capabilities of the DUACS-DT2018, DUACS-DT2014 and DUACS-DT2010 maps (i.e., to identify the impact of system upgrades), and 3) to verify the impact of the varying satellite constellation on the effective resolutions of the maps. The paper is organized as follow: the data and method are introduced in section 2. In section 3, we present our results. Finally, a discussion and a conclusion are provided in section 4.
- 25 A sensitivity study on the choice spectral criterion to estimate the resolution and a comparison of various approaches to estimate the resolution is given in the Appendix.

2 Data and method

2.1 Input data

5

In the present study, we consider two kinds of data:

Independent dataset: we used two independent (i.e., not used in the mapping) datasets to evaluate the effective resolutions of the maps: 1) Level 3 CMEMS SLA from independent 1Hz along-track and 2) the SLA estimated from tide gauges. The along-track SLA are constructed using a procedure similar to Level 3 CMEMS products and are

used to estimate the effective spatial resolution. The SLA at tide gauge locations originate from the Global Sea Level Observing System and Climate and Ocean Variability, Predictability and Change (GLOSS-CLIVAR) network and is used to estimate the effective temporal resolution. GLOSS-CLIVAR data are worldwide and available with daily sampling.

5

10

15

• The maps of SLA are constructed using optimal interpolation, based on the a priori statistical knowledge of the field (e.g., variance, correlation scales, noise). The mapping procedure is based on merging of calibrated multi-satellite altimeter (Level 3) data and follows the same protocol as described by Pujol et al. (2016) for the DUACS DT2014. Taburet et al. (2019) give the full description and validation of the DUACS-DT2018 global and regional products. The main differences between the DUACS-DT2014 and the DUACS-DT2018 processing consist of an improved along-track processing (e.g., improved orbit correction, wet troposphere correction, ocean tide correction and a new mean sea surface) and updated a priori knowledge of the SLA variance and optimized selection of the data in the optimal interpolation. The maps tested here are computed specifically for this study in several constellation scenario, keeping at least 1 mission out to allow an independent assessment of the resolution. The DUACS-DT products, formerly known as AVISO products, are referenced in the CMEMS catalogue as "OCEAN GRIDDED L3/4 SEA SURFACE HEIGHTS AND DERIVED VARIABLES REPROCESSED" products.

2.2 Method

Our method to estimate the effective spatial resolution is based on the ratio between the spectral content of the mapping error and the spectral content of independent signal (along-track observations previously mentioned).

 $NSR(\lambda_s) = \frac{S_{diff}(\lambda_s)}{S_{obs}(\lambda_s)}$

Where, λ_s is the spatial wavelength, $S_{diff}(\lambda_s)$ is the power spectral density of the difference (SLA_{obs} - SLA_{map}), $S_{obs}(\lambda_s)$ is the spectral density of the independent observation.

(1)

The algorithm to compute the spatial effective resolution follows 4 main steps:

- A coastal editing is applied in a 100 km coastal band (only for the global products) to remove the increased errors in the coastal area.
- 25

30

Gridded data are interpolated to the locations of the independent along-track data.

- Along-track and interpolated data are divided into overlapping 1500km long segments every 300 km for the global products (500km long segments for the Mediterranean Sea products and 300km long segments for the Black sea products). Each segment is saved in a database and referenced by it median (longitude, latitude) coordinates.
- Finally, between latitudes 90°S-90°S and longitudes 0-360°E, we consider 10°x10° longitude x latitude boxes for the global products (5°x5° longitude x latitude boxes for the Mediterranean Sea product, and 3°x3° longitude x latitude boxes for the Black Sea product) every 1° incremental step. All available segments referenced within the 10°x10°

box are selected to compute the power spectral densities based on the Welch method (1967). Prior to spectral computation, signals are detrended and we applied a Hanning window. The effective resolution is then given by the wavelength λ_s where the $NSR(\lambda_s)$ is 0.5.

- 5 The method (applied to the altimetry product) is illustrated in Figure 1 with the data selection and interpolation step to the spectral analysis. The total number of averaged segments in each 1°x1° longitude x latitude box is shown in Figure A1a) for the global product, Figure A1b) for the Mediterranean Sea product and the Black Sea product. Due to the coastal editing, the number of computed segments in the global product analysis is less than 1000 near the coast and ~1500 in the open ocean. In the Mediterranean Sea the number of segments is ~400 and ~250 for the Black Sea. A limitation of the present spectral approach is the need to rely on coastal for estimation of the resolution in the two regional products. It is worth noting, that we
- probably underestimate the resolution capability of the maps since we are estimating the spatial effective resolution of degraded maps to keep an independent dataset aside.

A comparison of SLA maps with independent tide gauge dataset is carried out to estimate the effective temporal resolution.

15 The approach is like the estimate of the effective spatial resolution and based on the computation of the ratio between the spectral content of the mapping error and the spectral content of the true tide gauge signal (Eq. 2):

$$NSR(\lambda_t) = \frac{S_{diff}(\lambda_t)}{S_{obs}(\lambda_t)}$$
(2)

Where, λ_t is the temporal wavelength, $S_{diff}(\lambda_t)$ is the power spectral density of the difference (SLA_{obs} - SLA_{map}), $S_{obs}(\lambda_t)$ is 20 the spectral density of the independent observation.

We computed the effective temporal resolution from each tide gauge time series of the GLOSS-CLIVAR network. The temporal domain covers the period 19930101 - 20151231. The computation for each time series follows 3 main steps:

25

30

- At each tide gauge location, we extract the gridded SLA time series that is most highly correlated with tide gauge time series (note that the maximum distance separation of the grid point that is most highly correlated with each tide gauge is 100km on average and can be as large as 300km)
- Each highly correlated time series (based on correlation criterion > 0.8) is subsampled into <u>365</u>_rday segments to compute the spectral densities S_{diff} and S_{obs}. The length of each segment must be set when estimating power spectral density using Welch's method. By subsampling into <u>365</u>_rday segments, we limit the frequency range to only periods shorter than <u>365</u>₄days. Note that consideration of time series longer than <u>365</u> days reduced the number of realizations because of occasional gaps in some of the data records. This can have an impact on the local estimation of the effective temporal resolution, e.g. from <u>365</u>-day to <u>2-years</u> segments we lose some continuous time series that do not contain >=365_x day segment. In these cases, the spectral analysis cannot be performed.

Supprimé: 100
Supprimé: 100
Supprimé: 100
Supprimé: We performed a sensitivity study on longer segment length (200 days and 300 days periods) and found similar global averaged effective temporal resolution (26 days for 300 days long segment, 27 days for 200 days long segment and 28 days for 100 days long segment).
Supprimé: 100
Supprimé: 100
Supprimé: 300-day
Supprimé: 200

Supprimé:

The effective temporal resolution at each tide gauge location is given by the period λ_t where the ratio $NSR(\lambda_t)$ is equal to 0.5 (deduced by linear interpolation between successive NSR values).

Note that this estimation of the temporal resolution is subject to an important caveat: the estimation is based mainly on coastal locations which may be contaminated by altimetry errors. Additionally, it may be not be fully representative of the temporal resolution of the DUACS maps which combine various oceanic regimes (e.g., coastal, offshore high variability, offshore low variability regimes). Our results may therefore be crude but useful estimates of the temporal resolution.

We somewhat subjectively define the effective resolutions to be the wavelength above which the NSR exceeds 0.5. In other words, it corresponds to the threshold where the mapping error variance is two times smaller than the observed true signal

- 10 variance. The methodology used here is similar to that Chelton et al. (2018), except that Chelton et al. (2018) consider the NSR in the spatial domain, whereas we here consider the NSR in the wavenumber domain. To illustrate and discuss the impact of the choice of the NSR criterion on the resolution, a sensitivity study is provided in the Appendix B. We demonstrate that the resolution can be ~30% coarser with NSR = 0.25 (SNR = 4) and > 30 % coarser with a more conservative NSR criterion (e.g., SNR=10, as recommended by Chelton et al, 2018). It is worth mentioning that various approaches may exist to estimate
- 15 the resolution (e.g., spectral magnitude ratio, filter transfer function). All measures of resolution have their advantages and drawbacks. We discuss in Appendix A the impact of using these different approaches in the estimation of the effective resolution.

3 Results

5

3.1 Effective resolutions of the DUACS-DT2018 maps

- 20 The effective spatial resolution of the DUACS-DT2018 global maps is shown in Figure 2a. Resolution was computed for maps constructed with three altimeters (Cryosat-2, HY-2, Jason-2) over the period 20140412-20151231 and Saral/Altika data were used as an independent dataset. We believe that this assessment of the spatial resolution based on maps constructed with three altimeter missions may be considered as a reasonable averaged estimate since ~3 altimeter missions are used in the merging for the CMEMS products 70% of the time over the period 19930101-20170515. The resolution ranges from ~100 km
- 25 wavelength at high latitudes to ~800km wavelength near the Equator, with a mean resolution at mid-latitude near 200km. Considering that eddy radius characteristic can be estimated as 20-25% of the wavelength (Chelton et al., 2011; 2018), this means that ~25km radius structures are properly resolved in the maps at high latitudes, ~200km radius structures are resolved in the Equatorial band and ~50km radius structures are resolved at midlatitudes. The effective spatial resolution of the DUACS-DT2018 Mediterranean Sea maps ranges from 90 to 160 km wavelength (Figure 2b). The averaged resolution is ~130 km
- 30 wavelength over the basin. The effective spatial resolution of the DUACS-DT2018 Black Sea maps ranges from 100 to 150 km wavelength and the averaged resolution is ~130km wavelength over the basin (Figure 2b).

The effective temporal resolution of the DUACS-DT2018 maps ranges from 2 to 440 days period (Figure 3). The temporal resolution is heterogeneously distributed over the global ocean, particularly in the inter-tropical band where a wide range of scales are found, linked to the mixture of continental tide gauges and island tide gauges, with the latter being more representative of open-ocean conditions. At mid-to-high latitudes the zonally averaged temporal scales are between 14- and

5 $45_{\overline{c}}$ day periods, coherent with the temporal correlation scales applied in the mapping process. The globally averaged effective temporal resolution is estimated $\sim 34_{\overline{c}}$ days period.

The globally averaged resolutions of about 200 km by <u>34</u> days period are consistent with the resolutions reported by Chelton et al. (2011; 2014) and Pujol et al. (2016). Using the spectral ratio method (see Appendix B), they found spatial resolution slightly better than 200 km at mid-latitude in Pacific Ocean.

10 3.2 Evolution of the DUACS system

We here investigate the impact of the DUACS system upgrade from 2010 to 2018 to highlight the progress of the DUACS processing. Resolutions were computed for maps constructed with two altimeters (Topex-Poseidon and Jason1) over the period 20030101-20041231 and Geosat Follow On data were used as independent dataset. To identify the impact of the DUACS system upgrade, we computed the relative improvement/deterioration of the effective resolutions (expressed in percentage) for

- 15 the upgrade DT2010 to DT2014, and DT2014 to DT2018 (Figure 4). Negative (positive) value means finer (coarser) resolution with the upgrade. The comparison of the DT2010 and DT2014 processing shows finer resolution (improvement > 2%) in DT2014 than in DT2010 in the high variability regions, e.g. the Gulfstream system, the Kuroshio system and the Antarctic Circumpolar Current (ACC) (Figure 4a). These improvements are associated with updated processing such as improved instrumental and atmospheric correction, tide correction, inter-calibration method and smaller correlation scale in the mapping
- 20 process. Coarser resolutions in DT2014 than in DT2010 are found in the Equatorial band and are potentially linked to larger correlation scales applied in this region in the DT2014, as reported by Pujol et al. (2016). Although the DT2018 and DT2014 global maps have similar mean effective spatial resolution, regional investigation highlights ~2 to 10% improved resolution in DT2018 in highly turbulent regions (Figure 4b), such as the Equatorial region, the Gulfstream system, the Kuroshio system as well as some regions in the ACC. These improvements are linked to the new mapping standard (optimized selection of the
- 25 observations in the turbulent region and a priori knowledge of the SLA variance based on a longer period in DT2018). The loss of resolution in the South Equatorial Atlantic is not understood yet. Similar comparison is performed for the Mediterranean and Black Sea regional products focusing on the upgrade DT2014 to

DT2018. Resolutions were computed for regional DUACS maps constructed with three altimeters (Jason-2, Cryosat-2, HY-2) over the period 20140412-20151231 and Saral/Altika was used as an independent dataset. The resolution capability of the

30 Mediterranean Sea maps is slightly finer (~4%) in DT2018 than in DT2014 (Figure 4c). The largest improvements (>6%) are found the western Mediterranean basin. The resolution in DT2018 is slightly coarser in the closed seas (Adriatic Sea and Aegean Sea). In these regions, the limited number of along-track data restricts a reliable interpretation of the spectral signal (see Figure A1). The resolution capability of the Black Sea maps is on average slightly finer (~3%) in DT2018 than in DT2014,

Supprimé: 10 Supprimé: 49

-	Supprimé: temporal
-1	Supprimé: 28
-{	Supprimé: 28
-1	Supprimé: 28

although a deterioration is found in the central part of the basin, which is also linked to a reduced number of spectral computations (Figure 4c).

The DUACS-DT2018and the DUACS-DT2014, maps have mean effective temporal resolution of ~ 34, days period and 37 days period, respectively. The differences can be locally larger than 15%, near 30° along the Japanese coast (~10 days gain), as shown in Figure 5. In these regions, the temporal resolution in the DUACS-DT2018 is finer than in DUACS-DT2014. These regions also coincide with the largest increased correlation score in the DUACS-DT2018 between SLA time series from maps and from independent tide gauge sensors (Taburet et al., 2019). These coastal improvements are linked to the new altimeter standards in coastal regions in DT2018 (Taburet et al., 2019).

3.3 Impact of altimeter constellation on the effective spatial resolution

- Since the number of altimeter data processed by the DUACS system varies with time (according to the availability of satellites and the data quality), we investigated the impact of the constellation on the effective spatial resolution. Figure 6 illustrates the impact of the number of altimeters (2 or 3 missions) used in the mapping on the effective spatial resolution. We verify, with our diagnostic, modest increases of resolving capabilities in the DUACS maps with increasing number of altimeters and found a globally averaged gain of resolution of ~5% from maps constructed with three altimeters compared with two altimeters. The
- 15 resolution improvements may be considered as modest. The reason is that the same covariance parameters are used in the optimal interpolation (OI) procedure, regardless of how many altimeters are available. These OI parameters exert very strong constraints on the filter transfer function of the OI procedure.

Regional gains of resolution can be larger than 10%. Additionally, it is possible to identify the improved resolving capability when a new mission is introduced in the DUACS system: for example, Figure 6a illustrates the improved resolving capability

- 20 when mission HY-2 is introduced in the mapping, Figure 6b illustrates the improved resolving capability when mission Cryosat-2 is introduced in the mapping. It is shown that the major contribution of the HY-2 mission in the mapping is in the high variability regions (Gulfstream, Kuroshio, Agulhas systems) while Cryosat-2 contributes in the mid-to-high latitude regions. On the global scale, the distribution of the effective spatial resolution is shifted toward shorter scales when the number of missions used in the merging increases (Figure 7) or when recent altimeters are used in the interpolation (e.g., compare the
- 25 resolution maps from DT2018 constructed with historical Jason-1/Envisat versus the maps from DT2018 constructed with currently operational missions Jason2/HY-2 or Jason-2/Cryosat2).

4 Discussion and conclusions

30

The present study investigates the resolving capability of the DUACS delayed-time gridded products (Global, Mediterranean Sea and Black Sea) delivered through the CMEMS catalogue. The key results are summarized in Table 1. Our method is based on the Noise-to-Signal spectral ratio to estimate the resolution. While along-track altimeter data resolve wavelength scales in

the order of few tens of kilometers (Dussurget et al., 2011, Dufau et al. 2016), we found that the merging of these along-track 7

-{	Supprimé: and DUACS-DT2014
-{	Supprimé: similar
1	Supprimé: 28
1	Supprimé: (globally averaged difference < 4%).

data into continuous maps in time and space leads to properly resolved structures with a wavelength scale of 100 km (a feature radius scale of ~25 km) at high latitudes to 800 km (a feature radius scale of ~200 km) near the Equator in the global gridded product and with a temporal scale of about 24-day period. The same analysis applied to the regional Mediterranean Sea and Black Sea products showed resolving capability of structure with feature radius resolution of ~30 km, which corresponds to ~3 grid spacing.

5

These results are consistent with previous investigations. Based on a spectral ratio approach (cf. Appendix B), Chelton et al. (2011) estimated a wavelength resolution of ~200 km for DT2010 and Chelton et al. (2014) estimated a wavelength resolution of ~180 km for DT-2014 in the mid-latitude Pacific Ocean. Our analysis based on the DT2018 global maps, suggests a wavelength resolution of ~200 km at mid-latitudes. As illustrated in Figure 8, we verified our estimation of the zonally averaged feature radius resolution of the mesoscale structures that can be properly mapped is smaller than the eddy scales

- 10 averaged feature radius resolution of the mesoscale structures that can be properly mapped is smaller than the eddy scales computed by Chelton et al. (2011). The eddy length scales range from ~70 km at high latitudes to ~180km near the Equator. The effective resolution is ~1.6 smaller than the eddy length scale. Additionally, we confirm that the minimum 4 weeks lifetime criteria used by Chelton et al. (2011) to identify and follow eddies seems to be compatible with the 34-day resolution capability. Note that our time scale estimation is based mainly on coastal locations and might not be representative of all oceanic regime.
- 15 The comparison of the DUACS-DT2018 reprocessing with former DUACS reprocessing (DT2010 and DT2014) reveals that finer structures are mapped in the global and regional Mediterranean Sea DT2018 products. For the Black Sea product, the interpretation is more complex due to the small dimension of the basin and the limited amount of spectral computation. Globally, we found that the largest improvements reach 20% and are mainly in high variability regions, associated with the new mapping standard (e.g., optimized selection of the along-track data, new a priori knowledge of the signal variance based
- 20 on 25 years of altimetry data, updated correlation scales for the regional Mediterranean Sea product) and new altimeter standards (e.g., instrumental and atmospheric corrections, tide corrections, inter-calibration method). The improvement patterns between DT2014 and DT2010 global maps is similar to those found by Pujol et al. (2016) using statistical comparison between maps and independent along-track, and drifters' datasets. The improvement patterns between DT2018 and DT2014 global maps coincide with those found by Taburet el al. (2019) for the validation of the DT2018 products. Using statistical
- 25 comparison between maps and independent along-track altimeter data, Taburet et al. (2019) also found improvement (~3-4%) of the mapped mesoscale structures, in the high variability region and in the western Mediterranean Sea basin. Note that, at global scale, Taburet et al. (2019) diagnosed the largest improvements between DT2018 and DT2014 in the coastal regions which are partially edited (along 100km coastal band) in the processing for estimating the spatial scale in this study, but they are detected with the temporal scale analysis, showing shorter timescale in the DT2018 compared with DT2014.
- 30 Several studies showed that at least two altimeters are required to accurately map the SSH mesoscale structures (Le Traon and Dibarboure 1999; Ducet et al. 2000; Pujol and Larnicol 2005; Dibarboure et al. 2011; Chelton et al. 2007; 2011) and up to four altimeters are required for Near-Real-Time products (Pascual et al. 2006), because only past observation are available for the mapping. This reduced number of observations has an impact on the estimation of the sea surface height. The present study

Supprimé: 28

Supprimé: 28

reinforces these findings, showing that the resolution capability increased ~10-20% at regional scale from the merging of data from two to three altimeters.

It is worth noting, that we probably underestimate the resolution capability of the maps since we are estimating the spatial effective resolution of degraded maps to keep an independent dataset aside. The resolution might hence be somewhat finer in

- 5 the distributed CMEMS products. Although the satellite constellation ranges from 1 to 5 altimeter(s) between 19930101 and 20170715, we believe that our estimation of the spatial resolution based on maps constructed with three altimeter missions may be considered as a reasonable averaged estimate since ~3 altimeter missions are used in the merging for the CMEMS products 70% of the time over the period 19930101-20170515. We can expect > 5% finer resolution over period where more than 4 altimeters are available (i.e., recent period). Likewise, we can expect on average 5% coarser resolution when only 2
- 10 altimeters are available.

To conclude, the number and the quality of altimeter simultaneously operational, the along-track configuration and sampling pattern, the weight given to the altimeter data in the mapping procedure and the choice of threshold SNR are key factors controlling the resolution capability of the DUACS gridded products. One may expect that, in permitting to observe finer mesoscale/sub-mesoscale structures (Dufau et al., 2016, Pujol et al, 2012), future instrumental systems based on large-swath

15 altimeter (such as Surface Ocean and Water Topography (SWOT)) combined with new mapping technique based on dynamic interpolation (Ubelmann et al., 2016), will push the maps resolution toward new limit.

Code and data availability. The DUACS system source code is not publicly available. The code for the spectral analysis is released under GNU General Public License v3.0 and is available at https://github.com/mballaro/scuba. DUACS all satellites
 gridded data and along-track data are available through the CMEMS website: http://github.com/mballaro/scuba. DUACS all satellites
 gridded data and along-track data are available through the CMEMS website: http://marine.copernicus.eu/. Specific maps used in our study are based on merging of three or two satellites and are available on request by contacting M. Ballarotta (mballarotta@groupcls.com).

Acknowledgment

This work is a contribution to the CMEMS R&D activities, the CNES-SALP project and the BOOST-SWOT project funded by ANR (project number ANR-17-CE01-0009-01). We would like to thank Lee Lueng Fu for his suggestions on the manuscript, Tom Farrar for his detailed and inspiring review and one anonymous referee. All their comments significantly added value to this paper. Supprimé:

References

5

Bretherton, F.P., Davis, R.E., and C. Fandry: A technique for objective analysis and design of oceanographic instruments applied to MODE-73, *Deep-Sea Res.*, 23, 559–582,10 doi:10.1016/0011-7471(76)90001-2, 1976

Chelton, D. B. & Schlax, M. G. The accuracies of smoothed sea surface height fields constructed from tandem satellite altimeter datasets. J. Atmos. Oceanic Technol. 20, 1276–1302 (2003)

Chelton, D. B., Schlax, M. G., Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Prog. Oceanogr.*, 91, 167–216, doi:10.1016/j.pocean.2011.01.002, 2011.

Chelton D., G. Dibarboure , M.-I. Pujol, G. Taburet , M. G. Schlax: The Spatial Resolution of AVISO Gridded Sea Surface Height Fields, OSTST Lake Constance, Germany, October, 28-31 2014, available at

10 http://meetings.aviso.altimetry.fr/fileadmin/user_upload/tx_ausyclsseminar/files/29Red0900-1_OSTST_Chelton.pdf, 2014 Chelton, D., Schlax, M. G., Samelson, R. M., Farrar J.T., Molemaker M.J., McWilliams J.C., Gula J. : Prospects for Future Satellite Estimation of Small-Scale Variability of Ocean Surface Velocity and Vorticity, accepted for publication in Progress in Oceanography, 2018

Dibarboure, G., Pujol, M.-I., Briol, F., Le Traon, P. Y., Larnicol, G., Picot, N., Mertz F. & M. Ablain (2011) Jason-2 in

DUACS: Updated System Description, First Tandem Results and Impact on Processing and Products, Marine Geodesy, 34:3-4, 214-241, DOI: <u>10.1080/01490419.2011.584826</u>
 Ducet, N., Le Traon, P-Y, and Reverdin G.: Global high-resolution mapping of ocean circulation from TOPEX/Poseidon and ERS-1 and-2, *Journal Of Geophysical Research-oceans*, 105(C8), 19477-19498, doi:10.1029/2000JC900063, 2000
 Dufau,C., Orsztynowicz, M., Dibarboure, G., Morrow, R. and P.-Y. Le Traon: Mesoscale resolution capability of altimetry:

- 20 present and future, J. Geophys. Res., Oceans, 121 (7), pp. 4910-4927, doi: 10.1002/2015JC010904, 2016 Dussurget, R., Birol, F., Morrow, R. and P. De Mey: Fine Resolution Altimetry Data for a Regional Application in the Bay of Biscay, Marine Geodesy, 34:3-4, 447-476, doi: 10.1080/01490419.2011.584835, 2011 Le Traon, P.-Y., Faugère, Y., Hernandez, F., Dorandeu, J., Mertz, F. and M. Ablain: Can we merge GEOSAT Follow-On with TOPEX/POSEIDON and ERS-2 for an improved description of the ocean circulation?, J. Atmos. Oceanic Technol., 20,
- 25 889 895, doi: 10.1175/1520-0426(2003)020<0889:CWMGFW>2.0.CO;2, 2003 Le Traon, P. Y., Nadal, F., and Ducet, N.: An improved mapping method of multisatellite altimeter data, J. Atmos. Ocean. Tech., 15, 522–533, 1998.

Le Traon, P.Y.; Dibarboure, G. Mesoscale mapping capabilities from multiple altimeter missions. J. Atmos. Ocean. Technol., 16, 1208–1223, 1999

30 Le Traon, P-Y: A method for optimal analysis of fields with spatially-variable mean. Journal Of Geophysical Research-oceans, 95(C8), 13543-13547, doi: 10.1029/JC095iC08p13543, 1990

Pujol, M. I., & Larnicol, G. 2005. Mediterranean Sea eddy kinetic energy variability from 11 years of altimetric data. Journal of marine systems, 58 (3-4), 121–142

Pujol, M.-I., Dibarboure G., Le Traon P.-Y., Klin P.: Using high-resolution altimetry to observe mesoscale signals, *J Atmos Ocean Technol* ., 29(9):1409–141, doi: 10.1175/JTECH-D-12-00032.1, 2012

Taburet, G., Sanchez-Roman, A., Ballarotta, M., Pujol, M.-I., Legeais, J.-F., Fournier, F., Faugere, Y., and Dibarboure, G.: DUACS DT-2018: 25 years of reprocessed sea level altimeter products, Ocean Sci. Discuss., https://doi.org/10.5194/os-2018-150, in review, 2019.

Smith, W.H.F.: Resolution of Seamount Geoid Anomalies Achieved by the SARAL/AltiKa and Envisat RA2 Satellite Radar Altimeters, *Marine Geodesy*, 38,644-671, doi: 10.1080/01490419.2015.1014950, 2015

Stammer, D.: Global characteristics of ocean variability estimated from regional TOPEX/POSEIDON altimeter measurements, Journal of Physical Oceanography, 27, 1743–1769, 1997

- 10 Ubelmann, C., Cornuelle, B., and Fu, L.-L.: Dynamic Mapping of Along-Track Ocean Altimetry: Method and Performance from Observing System Simulation Experiments, Journal of Atmospheric and Oceanic Technology, 33, 1691–1699, doi:10.1175/JTECH-D-15-0163.1, <u>http://dx.doi.org/10.1175/JTECH-D-15-0163.1</u>, 2016 Yale, M.M., Sandwell, D.T., and W.H.F. Smith: Comparison of along-track resolution of stacked Geosat, ERS-1 and Topex satellite altimeters, *Journal of Geophysical Research*, 100 (B8):15117–15127, doi: 10.1029/95JB01308, 1995
- 15 Welch, P.: The use of the fast Fourier transform for the estimation of power spectra: A method based on time averaging over short, modified periodograms, *IEEE Trans. Audio Electroacoust.*, vol. 15, pp. 70-73, 1967.

20

5

Appendix A: other approach to estimate the resolution

1) Spectral magnitude ratio

Chelton et al. (2011, 2014) estimated the resolution of the DUACS DT2014 maps based on the calculation of the spectral magnitude ratio between the reference Stammer (1997) along-track spectrum and gridded SSH spectra. Similarly, we here

5 estimate the resolution based on the spectral ratio between independent along-track and gridded SSH signals (Eq. 3). It is defined as follows:

$$SR(\lambda_s) = \frac{S_{map}(\lambda_s)}{S_{obs}(\lambda_s)} \quad (3)$$

, where $S_{obs}(\lambda_s)$ denote the power spectral density of the independent SSH along-track, $S_{map}(\lambda_s)$ denotes the power spectral density of the SSH map interpolated onto the independent along-track segment, *SR* the spectral ratio and λ_s the wavelength.

- 10 The resolution is given by the wavelength λ_s where the spectral ratio is equal to 0.5 and is based on the conventional notion of a filter being characterized by its half-power filter cutoff wavelength (Chelton et al., 2011). To differentiate it from the effective resolution we named it "useful" resolution: "useful" for verifying the available and realistic amount of energy at a specific wavelength between two signals without considering their phase (e.g., useful for model sensitivity study).
- Chelton et al. (2011, 2014) estimated the resolution of the DUACS-DT2010 and DUACS-DT2014 as the wavenumber at which
 the power is a factor of 2 smaller than the Stammer (1997) spectrum. From their analysis, they estimated spatial resolution of ~2° for the DUACS-DT2010, ~1.7° for the DUACS-2014. Chelton et al. (2014) found essentially the same resolution between maps constructed with 2 satellites or 4 satellites.

The resolution estimated with the SR method is shown in Figure A2a and the difference between the effective and useful resolution is shown in Figure A2b. The useful resolution of the DUACS-DT2018 maps ranges from 100km at high latitude to

- 20 500km near the Equator. The ratio effective/useful resolution suggests somewhat finer resolution in the intertropical band using SR approach and somewhat finer resolution at high latitude with the NSR approach. In other words, the amplitude of the mapped SSH spectral content is better in the inter-tropical band than the phase, whereas it is the opposite at high latitude. This feature highlights the difficulty to properly map propagating equatorial waves in the DUACS system. The two methods are equivalent at mid-latitudes.
- 25

2) Transfer function

The transfer function (H) measures the filtering properties of a system (Eq. 4). It is defined as follows:

$$H(\lambda_s) = \frac{CS_{obs-map}(\lambda_s)}{S_{obs}(\lambda_s)} \quad (4)$$

where $CS_{obs-map}(\lambda_s)$ is the cross-spectral density between the along-track data and the map interpolated onto the along-track 30 segment, $S_{obs}(\lambda_s)$ is the power spectral density of the along-track. Note that in this case, the along-track is considered as nonindependent (i.e., it is used in the mapping system). The resolution is given by the wavelength λ_s where *H* is equal to 0.5. It is the same definition used by Chelton and Schlax (1993) to estimate the resolution capability of an arbitrarily sampled dataset. The resolution estimated with the transfer function method is shown in Figure A3a) and the difference between the effective resolution and the transfer function resolution is shown in Figure A3b). The transfer function resolution of the DUACS-DT2018 maps ranges from 100km at high latitude to 400km near the Equator. The difference between effective resolution versus transfer function resolution suggests somewhat finer resolution using the transfer function. This is directly linked to the

5 fact that the along-track is here non independent. The assessment is undertaken below the track that is used in the filtering system. This diagnostic gives the filtering property of the system but "suffers" from the non-independency of the along-track dataset. The resolution may be different off-track.

These methods share the same number of spectrum calculation and number of segments used in the calculation (see Fig. A1) and each method has advantages and drawbacks. The spectral magnitude ratio compares the amplitude of the signals and the transfer function estimates the filtering properties from assimilated along-track. The function $NSR(\lambda_s)$, $SR(\lambda_s)$, and $H(\lambda_s)$ are shown for location (330°N, 45°N) in Figure A5. At this location, each function has a transition between 100km and 200km wavelength, separating the high mapping error regime for wavelength < 100km to the low mapping error for wavelength > 200km (Fig. A5a), separating the high amplitude error regime for wavelength < 100km to the low amplitude error for the wavelength > 200km (Fig. A5b), and separating the filter regimes (Fig A5c).

15 wavelength > 200km (Fig. A50), and separating the filter regimes (Fig A5c)

Appendix B: Sensitivity to the Noise-to-Signal Ratio (SNR) criterion

We here investigate and discuss the impact of the NSR criterion on the estimation of the effective resolution. NSR criterion is used to define the resolution limit of the map. In the present study, we choose the NSR=0.5 criterion to define the resolution limit. This value may be considered too generous; therefore, we present below the effective resolution for three cases, motivated by the analysis performed in the spatial domain by Chelton et al. (2018):

- criterion of NSR=0.5
- criterion of NSR=0.25
- criterion of NSR=0.1
- 25

20

Figure B1a) represents the effective resolution using NSR=0.5 (SNR=2) criterion, Figure B1b) using NSR=0.25 (SNR=4) criterion and Figure B1c) NSR=0.1 (SNR=10). For each panel the resolution becomes finer poleward. The white areas correspond to the regions where the NSR threshold criterion is not achieved. These areas become larger in the inter-tropical region as well as at high latitudes when the NSR criterion decreases. For NSR=0.1, the resolution in the intertropical band cannot be computed with the method, since the NSR is above 0.1 for all scales. To further illustrate this, we show an example

of NSR at one specific point (lon= 346° E, lat= 16° S) in Figure B2. The analysis shows that the NSR is greater than 0.3 (i.e., SNR < 3) in this location. This large-scale low coherency between maps and along-track may be linked to the misrepresentation

³⁰

of the large scale and rapid equatorial waves (e.g., equatorial gravity waves) in the mapping process, which are filtered in the mapping process.

- Despite the areas of missing values in Figures B1b) and B1c), we quantify the difference of effective resolution between 5 criterion NSR=0.4 and NSR=0.25 (Figure B3) and NSR=0.5 and NSR=0.1. The difference between effective resolution computed with NSR=0.5 vs NSR=0.25 is < 30% (~60km) at mid-latitude and <=50% (400km) in the inter-tropical band. The difference between effective resolution computed with NSR=0.5 vs NSR=0.1 is < 50% (~100km) at mid-latitude and >50% in the inter-tropical band.
- 10 In conclusion, we here demonstrate that the choice of the NSR criterion has an impact on the estimation of the resolution. Setting more conservative criterion NSR=0.25 leads to ~30% coarser effective resolution. The strongly conservative criterion NSR=0.1 also reveals one of the major caveats in the DUACS maps processing: the poor representation of the large and rapid scale equatorial circulation. This issue should be addressed in the future version DUACS maps.

15

20



Figure 1: Schematic illustration of the methodology: a) input data selection, b) colocation SLA and gridded SLA and c) spectral analysis showing the ratio error spectrum to signal spectrum



Figure 2: Effective spatial resolution in km of the DUACS-DT2018 maps for a) the Global Ocean product, b) the Mediterranean Sea and the Black Sea products. Unit in km



Figure 3: Effective temporal resolution in days of the DUACS-DT2018 maps. Unit in days



Figure 4: Gain/loss of effective spatial resolution for a) the Global Ocean product between DT2014 and DT2010, b) the Global Ocean product between DT2018 and DT2014, c) the Mediterranean Sea product and the Black Sea products between DT2018 and DT2014. Negative value means that the resolution capability is finer. Note the different colorbar scale between global and regional products



Figure 5: Gain/loss of effective temporal resolution between DT2018 and DT2014. Negative value means that the resolution capability is better in DT2018 than DT2014



Figure 6: Impact of the satellite constellation on the effective resolution – Ratio of effective resolution of a) maps constructed with C2-H2-J2 vs C2-J2, and b) maps constructed with C2-H2-J2 vs H2-J2



Figure 7: Distribution of the effective spatial resolution for various altimeter merging configuration



Figure 8: Zonally averaged eddy scale Ls (as in Chelton et al., 2011; and computed from the DUACS-DT2018 two satellites maps) and feature radius resolution of the mesoscale structures that can be properly mapped in DUACS (i.e., derived as 0.25xeffective resolution). Units in km



Figure A1: Number of segments used in the spectral computation for a) the global product and b) the regional Mediterranean and Black Sea products



Gain(-) / Loss (+) of effective spatial resolution DT2018 NSR method vs SR method



Figure A2: a) DUACS-DT2018 Useful resolution derived from spectral ratio approach and b) ratio effective / useful resolution for the DT2018 maps. Blue means finer resolution with NSR $\,$



Figure A3: a) DUACS-DT2018 resolution derived from the transfer function approach and b) ratio effective / transfer function resolution for the DT2018 maps. Blue means finer resolution with NSR



Figure A5: Illustration of the various spectral function used to estimate the resolution a 330°E, 45°N: a) with NSR method, b) with SR method, and c) with the transfer function



Figure B1: Effective resolution computed for three different SNR criteria: a) SNR=2, b) SNR=4 and c) SNR=10. The white areas correspond to the regions where the SNR threshold criterion is not achieved.



Figure B2: Ratio NSR at longitude 346°E and latitude 16°S. We illustrate that the ratio at this location is always above >0.3 and so the resolution cannot be computed for SNR criterion >4 (i.e., NSR <0.25)



Figure B3: Ratio effective resolution computed with a) SNR=2 versus SNR=4 criterion and b) SNR=2 versus SNR=10 criterion. Blue means finer resolution with SNR=2. The white areas correspond to the regions where the SNR threshold criterion is not achieved.

	SPATIAL	FEATURE	TEMPORA	L FEATURE		
Product	Effective resolution	Grid spacing	Effective resolution	Grid spacing		
GLOBAL	100 to 800 km	4 to 30 km	~34 days	1 day		Supprimé: 28
MED-SEA	~130 km	~10km	(1)	1 day		
BLACK-SEA	~130 km	~10km	(1)	1 day		

Table 1: Summary of the DUACS products spatial and temporal resolutions. (1) Not estimated due to the limited amount of tide5gauges in the Mediterranean Sea and Black Sea