Dear Editor,

We thank the two reviewers for their critical and constructive comments on our research. Their comments have significantly improved our manuscript. The main modifications in the revision are las follows:

- Use the new version of the BEC SSS to replace the previously used in this study.
- Extend the SSS from 5 m to 8 m near the surface when extracted from insitu data, in order to involve the ITP observations.
- Rearrange the orders in Section 3 and Section 4 to highlight the evaluation against in-situ data, and which has been divided into two parts referred to dependent and independent in-situ dataset in the Sec.
 4.
- All figures have been updated due to the above two changes and two scatterplots showing the SSS against in-situ data from CORA5.1 are complemented in the revision.
- We have improved the English and expanded/shortened the text as recommended by the reviewers.

Below are the detailed responses to their comments: the reviewer comments are in black oud our response is in red.

Anonymous Referee #1

The paper shows an intercomparison among 6 arctic salinity products (2 based on SMOS acquisitions, 2 climatologies and 2 reanalysis products). All products are also compared with in-situ data CORA 5.1. In addition to the intercomparison by itself, the aim of the paper is to evaluate the best SMOS product to be assimilated by TOPAZ4 reanalysis product. -A: We thank the referee for the detailed evaluation of our manuscript and constructive suggestions. We appreciated this very much, all the comments are taken into account in the revised version.

General Comments

The paper needs a general improvement of the writing. In some cases the concepts are no clear and English should be improved.

-A: We thank the reviewer for pointing out this weakness in the manuscript. We have improved the English, in addition to extending, shortening and rearranging the text for improved clarity.

Other general comment refers to the version of the BEC SMOS product included in the comparison. The version of the SMOS BEC product is only clearly defined at the end of the conclusions (lines 543-546). This should be explained in section 2. According to the expressed in conclusions, the version of the BEC SMOS product used in this paper is version 1.0. This version is not accessible now because it has been superseded by version 2.0. Why authors have not included v2.0 instead v1.0 in this study?

-A: Thank you for this comment, we agree with the reviewer. Note that BEC product was released just before the submission of this manuscript. The SMOS BEC product used has been replaced by the version of 2.0, which is also defined in Section 2.1. We update all figures and results in the revised manuscript.

This reviewer knows the effort that implies to redo this validation using the new BEC product, but taking into account that v1.0 is not available, the inclusion of this product in the study is not interesting and v2.0 should be used. It is not necessary to proceed with all the period of the current v2.0, only the studied period (2011-2013) will be enough. Please, use v2.0 of 2011-2013 period instead v1.0. Change "BEC product" section accordingly. -A: Agreed. The revision will replace the previous BEC product with version 2.0 and the concerned figures and analysis are updated.

Specific Comments

Lines 142-145: The BEC Arctic product 1.0 is not created as is described here. Systematic bias of the retrieved salinity data is corrected computing the so-called SMOS climatology (the most probable value for a given lat-lon, incidence angle and across swath distance) and substituting this one by a reference. The used reference is the annual WOA13 (the same reference for all maps) and not Argo float extrapolated at 7.5km. The second correction (the temporal bias correction) was computed for version 1.0 of the Arctic product in the same way as in the global one: assuming that the quantity of salt is constant in the surface. This coarse approach has been refined in version 2.0 (the current one) using Argo to compute the mean value of salinity for each Arctic map.

-A: Thank you for this informative and constructive comment, the text is changed to:

Lines 135-143: "The BEC SSS product was generated from ESA L1B (v620) products, and accumulates salinity data over 9 days with a spatial grid resolution of 25 km. With respect to its previous version, a systematic bias in the retrieved salinity is corrected by computing the SMOS climatology (the most probable value for a given lat-lon, incidence angle and across-swath distance) which is substituted by a reference value from WOA13. In addition, a temporal bias correction has been refined in this version using near-surface Argo salinity to compute regional averages (see the details in Olmedo et al., 2018)."

Line 147: The anomaly is referenced to WOA13 (not WOA09) -A: Thank you. It is corrected.

Section 3: Many comparisons are made involving different regions and products. A table similar to table 1 but for intercomparisons would help to the reader.
-A: It is a good suggestion. A new table (Table 1) is added to clearly explain the product specifications. Moreover, the concerned sections are rearranged, and the evaluations against in-situ data are divide into two part according to dependent and independent observations in Section 4.

Line 466: Beware both SMOS products do not use different BT filtering flags. The main difference between both is that they are applying a completely different salinity retrieval method.

-A: Thanks for this point. It is deleted and replaced by other statement as Line 447: "... due to the different retrieval applied in these two datasets."

Lines 5-9: Suppress "respectively". This long sentence probably sounds better as "Recently, two independent gridded SSS products have been derived from mission: the developed by the Barcelona... and the one developed by Ocean...." Here a mention about the regional or global character of both products will help to the reader to know about the general characteristics of each product (one can expect that a product specifically developed for Arctic will provide better results)

-A: Thank you for this suggestion. Here is the revised text as

Lines of 2-6: "Recently two gridded Sea Surface Salinity (SSS) products that cover the Arctic Ocean have been derived from the European Space Agency's (ESA) Soil Moisture and Ocean Salinity (SMOS) mission: one developed by the Barcelona Expert Centre (BEC) and the other developed by the Ocean Salinity Expertise Center of the Centre Aval de Traitement des Données SMOS at IFREMER (CEC)."

Line 42: "northern North Atlantic". Authors are referring to the north of the North Sea (a relative mall region) or the authors are referring to the thermohaline circulation between Arctic Ocean and North Atlantic? (probably is this second option but "northern North Atlantic" sounds strange to me)

-A: Thank you for this comment. This sentence referred to the thermohaline circulation between Arctic Ocean and North Atlantic. And this sentence will be changed as

Lines of 41-43: "The SSS also affects the decadal variability of hydrography in the upper waters of the North Atlantic (Reverdin et al., 1997)."

Lines 47-50. This sentence is difficult to read. "a significant change in the global warming scenario" should be "or a significant"? Probably no, but I do not clearly understand what is the meaning of this sentence.

-A: This sentence will be corrected by the lines of 46-49:

"Additionally, the increased melting of glaciers and sea-ice in the Arctic (McPhee et al., 1998; Macdonald et al., 1999) leads to significant changes in the salinity distribution and fresh water pathways (Steele and Ermold, 2004; Morison et al., 2012)."

- Macdonald, R. W., Carmack, E. C., McLaughlin, F. A., Falkner, K. K., and Swift, J. H.: Connections among ice, runoff and atmospheric forcing in the Beaufort Gyre. Geophys. Res. Lett., 26, 2223–2226, 1999
- McPhee, M. G., Stanton, T. P., Morison, J. H. and Martinson, D. G.: Freshening of the upper ocean in the Arctic: is perennial sea ice disappearing? Geophys. Res. Lett. 25, 1729–1732, 1998.
- Morison, J., Kwok, R., Peralta-Ferriz, C., Alkire, M., Rigor, I., Andersen, R., and Steele, M.: Changing arctic ocean freshwater pathways. Nature, 481:66–70, 2012.

Steele, M. and W. Ermold (2004) Salinity Trends on the East Siberian Shelves, Geophysical Research Letters, Vol. 31, L24308, doi:10.1029/2004GL021302, 2004.

Line 112: There exist, at least, two different versions of WOA13 (1.0 and 2.0) with significant differences between them in the Arctic data. Please, indicate the one used in this study. -A: Thanks for this point. It is the WOA13 version 2.0, clearly stated in the revision.

Line 130: "non geophysical sources" should be better than "unphysical contaminations" -A: Thank you for this suggestion. It is corrected.

Line 131: ice-sea contamination should be mentioned because is an important source of biases in the Arctic.

-A: Thank you for this suggestion. The following statement has been added on 124-127:

"The SSS retrieval from SMOS is subject to biases originating from various nongeophysical sources such as the so-called land-sea contamination and the latitudinal biases, mainly caused by the thermal drift of the instrument. A particular challenge in the Arctic is the sea-ice edge because of ice-ocean contamination."

Line 193: Acronyms EnKF (Ensemble Kalman Filter) and DEnKF have not been defined in the text

-A: It is corrected.

Line 268: "Marches" should be "matches"? -A: It is corrected.

Lines 281-282: Have in mind that comparison of BEC product and WOA is not recommended because BEC product incorporates as reference WOA13.

-A: Thank you for pointing out this issue. In fact, even if the BEC product has incorporated WOA13as reference, the updated evaluation of the BEC version 2.0 still shows values far from the referred climatology as shown in Fig. A1. In the Barents

Sea, there is a clear salinity bias of BEC even if this product has been referred to WOA13.



Fig. A1 Monthly SSS (unit: psu) in March from satellite products of (a) BEC and (b) CEC, reanalyzes of (b) TP4 and (e) MOB, and climatology of (c) PHC and (f) WOA. The black shaded isoline represents the salinity of 35 psu near surface regarding to the product self.

The statement is changed as the Lines of 251-256:

"In comparison to the March situation, the BEC and CEC SSS in the Nordic Seas are both less saline, indicated by the 35 psu isoline. The sea ice masking of the two SMOS products differ considerably in the Canadian Basin and in the Arctic marginal seas. Although the SSS of TP4, MOB, PHC and WOA agree relatively well in the northern Atlantic Ocean, the discrepancies become dramatic in ice-covered areas."

Lines 285-286: For this reviewer is not clear what do you mean with "over the sea-ice conver" and "under the sea-ice cover"... Under sea-ice cover means "below the ice"? Probably the meanin is related with latitudes not covered by ice?

-A: Thank you for this point. It is corrected as the lines of 256-260:

"Below the ice or near the sea-ice edge (denoted by the brown thick line in Fig. 2 and 3), TP4 and PHC share common features, which can be explained by the model restoring to PHC. On the other hand, the MOB and WOA differ significantly in spite of WOA being used as input to the MOB."

Line 294: This sentence refers to figure 6? This figure is only referred in conclusions (line 487)

-A: No, it refers to Fig. 4 and Fig.6, this has been clarified in the text.

Line 413: The mentioned four observations, are outliers?

-A: Yes, a more detailed explanation had been included on lines 381-385:

"Looking at the low-salinity observations (~27 psu) collected at (136.4°W, 70.5°N) on 15th August 2011, marked by anti-triangles (Fig. 1b) near the Mackenzie River estuary, TP4 has a significant negative bias (< -4 psu) visible as the outliers above the dashed-black line in Fig. 11a. This hints to a lack of fresh water signatures from river discharge."

Line 536: In my opinion this is not a validation. Is a comparison. -A: Right, 'validation' is changed to 'evaluation'.

Line 61. Typo: MIRIAS should be MIRAS -A: It is corrected.

Technical corrections (Typos) Line 122: Typo: "in in Section" ("in" written twice) -A: It is corrected.

Line 133: Typo: "march-up" should be "match-up"? -A: Thanks for this point. It is corrected

Line 139: Correct address is http://bec.icm.csic.es -A: It is corrected.

Line 163: Typo: should be EASE instead of EASA -A: Here, it means an Equal-Area Scalable Earth Grid (EASE-Grid), and also changed in the revision.

Line 281: Typo: then should be than. The correct ending for the sentence should be "than the provided by BEC product"

-A: Thank you for this point. It is corrected

Line 317: Word SMOS is used twice. -A: It is corrected.

Line 552: The correct URL is bec.icm.csic.es -A: Thanks, it is corrected.

Anonymous Referee #2

The paper aims to quantify uncertainties of Arctic observation-based sea surface salinity to be included in the TOPAZ reanalysis. Two SMOS products are considered and compared against climatologies, observed data sets and reanalysis. This is an important problem in advancing in the data assimilation technics and improving the quality of CMEMS reanalyses. Anyway this study is not a significant step along that path. The paper has some unclear or incomplete reasoning. I do not feel that this research is ready to be published in OS. I do encourage resubmission after a much more detailed and careful investigation. -A: We thank the referee for the time spent and for the detailed revision of our manuscript. We appreciated very much for the comments which are all taken into account in the revised version of the manuscript.

My primary concerns are i) the research is poorly presented, with vital details missing -A: Thank you for this comment. We have improved the presentation of the work to help the reader understand. This evaluation has two parts: the first part is an intercomparison with reference to the TP4 reanalysis, the second part is an evaluation with respect to two types of in-situ datasets which are involved in the TOPAZ system and independent respectively. In the revision, the observed SSS by in situ near surface will be extended from no deeper than 5 m depth to 8 m depth, which will involve more observation samples for this evaluation.

ii) the BEC SMOS product selected from this study should actually be updated to version 2. -A: Yes. In the revision, we use the version 2.0 of BEC product to replace all the figure and the concerned analysis (also see the response to the same comment of Referee #1). *iii) the PHC data set is old, is included in WOA13 and assimilated in TOPAZ. It does not add much to the analysis*

-A: Thank you for this suggestion. The PHC dataset is one of the most important climatology in the Arctic Ocean, and still implemented widely in quantitative evaluation works (Carton et al., 2018, 2019). The PHC is based on the archive of observations primarily from the 1950s through the 1980s and so may have a somewhat cool climatology. In the current version of TOPAZ, the combined climatology of PHC and WOA13 are used as relaxation so that the quantitative comparison of two climatologies still could be helpful to reasonably reject this or not for the improving of the model relaxation process.

Carton, J.A., G.A. Chepurin, and L. Chen, 2018: SODA3: a new ocean climate reanalysis, *J. Clim.*, **31**, 6967-6983, doi:10.1175/JCLI-D-18-0149.1.

Carton, J.A., S.G. Penny, and E. Kalnay, 2019: Temperature and salinity variability in soda3, ECCO4r3, and ORAS5 ocean reanalyses, 1993-2015, *J. Clim.*, 32, 2277-2293, doi:10.1175/JCLI-D-18-0605.1.

iv) MOI is not a reanalysis. The CMEMS product MULTIOBS_GLO_PHY_REP_015_002 is a combination of four data set. I do define a reanalysis as a combination of ocean modeling, data assimilation scheme and observed data sets. I would rather include in this study a global CMEMS ocean/sea ice reanalysis to be compared with TOPAZ4.

-A: Thank you for this comment. We agree that evaluating more global reanalysis products in CMEMS would be very interesting, and give more knowledge of the uncertainties in the different model systems, but it would go beyond the initial aim of directing our next assimilation work.

As an objective analysis product MOB uses the multivariable optimal interpolation method and can be used as a special representative in reanalysis products just like Simple Ocean Data Assimilation (SODA, Carton et al., 2018) is often used for comparative analysis with respect to other traditional reanalysis products (Uotila et al., 2018).

So in this study, we choose to use these two representative types of reanalysis products in CMEMS to evaluate the new satellite SSS products.

v) The region of interested is the Arctic Ocean, but results are mostly related to the North Atlantic/Nordic Seas area.

-A: Thank you for this comment. In the current evaluation, the comparison in the Beaufort Sea has been presented referred to the independent SSS from BGEP and CLIVAR, which is directly linked to one of the main conclusions to support the BEC product. In addition, our forecast products more focus on the wide Arctic region (north of 60N), where our general interest is and discussed in this study.

In this study, the in situ SSS from CORA5.1 were used by the TOPAZ system either assimilated or filtered during pre-processing for QC. It results these dependent SSS from CORA5.1 primarily are distributed in the Nordic Sea as shown in that figure. There are in general few observations in Arctic, the additional reason is a strictly used limit for the SSS observations - near the surface no deeper than 5 m depth. In fact, if extending the limit to 8 m depth, more SSS observations extracted from Ice-Tethered Profiler (ITP) will be involved. The Fig. A2 shows the locations of the SSS from ITP in the three years. Clearly, the evaluation referred to this dataset would enrich our knowledge of the Arctic SSS uncertainty.



Fig. A2 The locations of the SSS observations extracted at the 8m depth from the ITP living more than 30 days during the years of 2011-2013.

In addition, compared with the in-situ SSS from CORA5.1, the scatterplots of the six SSS products have been added in the revision to investigate the uncertainties according to different areas in the Arctic Ocean, also see the lines of 326-353.

vi) Section 5 summarizes main results but a proper discussion to support the BEC SMOS and the "certain benefit (line 537) is missing. These points significantly detract from the conclusions of the study, make the conclusions much weaker than the present manuscript states.

-A: Thank you for this comment. The revision manuscript includes more discussion about this issue, with more consistence to the present results.

English need to be generally improved.

-A: Thanks for your comments. We will further improve the concerned parts.

Formatted: Header

Deleted: -Deleted: datasets

Deleted: Raj¹ Deleted: Bertino^{1, 2}

Deleted: Samuelsen^{1, 2} Deleted: Wakamatsu^{1, 2}

Deleted: and two CMEMS reanalyses

Formatted: Font: 16 pt, Font color: Text 1

Evaluation of Arctic Ocean surface salinities from SMOS

against a regional reanalysis and in situ data

Jiping Xie¹, Roshin P. <u>Raj²</u>, Laurent <u>Bertino²</u>, Annette <u>Samuelsen²</u>, and Tsuyoshi <u>Wakamatsu²</u>

1. Nansen Environmental and Remote Sensing Center, N5006 Bergen, Norway

2. <u>Nansen Environmental and Remote Sensing Center and Bjerknes Centre for</u> Climate Research, Bergen, <u>N5006 Bergen</u>, Norway

Correspondence to: Jiping. Xie (jiping.xie@nersc.no)

Deleted: ¶

1

1

1	Abstract	1
2	Recently two gridded Sea Surface Salinity (SSS) products that cover the Arctic Ocean	17
3	have been derived from the European Space Agency's (ESA) Soil Moisture and Ocean	
4	Salinity (SMOS) mission; one developed by the Barcelona Expert Centre (BEC) and	
5	the other developed by the Ocean Salinity Expertise Center of the Centre Aval de	
6	"Traitement des Données SMOS at IFREMER (CEC). The uncertainties of these two	
7	SSS products are guantified during the period of 2011-2013 against other SSS	
8	products: one data assimilative regional reanalysis, one data-driven reprocessing in	
9	the framework of the Copernicus Marine Environment Monitoring Services (CMEMS),	
10	two climatologies: the 2013 World Ocean Atlas (WOA) and the Polar science center	
11	Hydrographic Climatology (PHC), and in-situ datasets, both assimilated and	
12	independent. The CMEMS reanalysis comes from the TOPAZ4 system which	
13	assimilates a large set of ocean and sea-ice observations using an Ensemble Kalman	
14	Filter, (EnKF). Another CMEMS product is the Multi-OBservations reprocessing (MOB),	
15	a multivariate objective analysis combining in-situ data with satellite SSS. The monthly	$\left(\right)$
16	root mean squared deviations (RMSD) of both SMOS products, compared to the	
17	TOPAZ4 reanalysis, reach 1.5 psu in the Arctic summer, while in the winter months	/
18	the BEC SSS is closer to TOPAZ4 with a deviation of 0.5 psu. The comparison of CEC	
19	satellite SSS against in-situ data shows too fresh Atlantic waters in the Barents Sea.	
20	the Nordic seas, and in the northern North Atlantic Ocean, consistently with the	
21	abnormally fresh deviations against TOPAZ4. When compared against independent	/
22	in-situ data in the Beaufort Sea, the BEC product shows the smallest bias (<0.1 psu)	
23	in summer and the smallest RMSD (1.8 psu), although all six SSS products share a	
24	common challenge to represent fresher water masses (<24 psu). Along the Norwegian	- 2011
	common challenge to represent resher water masses (~24 psu). Along the Norwegian	
25	coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors	
25 26	coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors than TOPAZ4 and indicates the potential value of assimilating the satellite-derived	
25 26 27	coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors than TOPAZ4 and indicates the potential value of assimilating the satellite-derived salinity in this system.	
25 26 27 28 20	coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors than TOPAZ4 and indicates the potential value of assimilating the satellite-derived salinity in this system.	
25 26 27 28 29 30	coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors than TOPAZ4 and indicates the potential value of assimilating the satellite-derived salinity in this system.	
25 26 27 28 29 30 31	coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors than TOPAZ4 and indicates the potential value of assimilating the satellite-derived salinity in this system.	
25 26 27 28 29 30 31 32	coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors than TOPAZ4 and indicates the potential value of assimilating the satellite-derived salinity in this system.	
25 26 27 28 29 30 31 32 33	coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors than TOPAZ4 and indicates the potential value of assimilating the satellite-derived salinity in this system.	

Formattad	
	[1]
Deleted: ¶	[2]
Deleted: stratification of the upperrctic Ocean is	[4]
Formatted	[3]
Formatted	[5]
Deleted: , independently	
Deleted: in Spain	
Formatted	[6]
Formatted	[7]
Deleted: (CECOS)	
Formatted	[8]
Deleted: Traitemenent des Donnees SMOS (CATI	D <u>Ş)</u> [mg]
Formatted	[10]
Deleted: two	
Deleted: products providing the Arctic SSS	
Formatted	[11]
Formatted	[12]
Deleted: one global, and another regional product.	[13]
Formatted	[14]
Deleted: and ocean	
Formatted	[15]
Deleted: with	
Formatted	[16]
Deleted: , the global reanalysis combines in-situ a	nd [17]
Formatted	[18]
Deleted: ensemble optimal interpolation method. In	n thison
Formatted	[20]
Deleted: the climatology both World Ocean Atlas (WQA)1
Formatted	<u>[14 2]</u>
Deleted: datasets during 2011-2013. For the valida	<u>1221</u> ation
Formatted	[23]
Deleted: observations are divided in two: those the	n [23] at [24]
Formatted	<u>• [24]</u>
Deleted: -situ observations show largest disagreer	[23] nonte i
Formatted	[07]
Polated, summer SSS from the	[27]
Exercised	
	[28]
Deleted: nas	
Formatted	[29]
Deleted: - saline - bias (~0.6 psu) with the smalles	t.rqgt0]
Formatted	[31]
Deleted: of this product into the forthcoming Arctic	[32]
Formatted	[33]
Formatted	[34]
Formatted	[35]
Deleted: absolute deviation;	
Deleted: ¶	[36]

135 **1. Introduction**

The sea surface salinity (SSS) plays a key role in tracking processes in the global 136 137 water cycle through precipitation, evaporation, runoff, and sea-ice thermodynamics (Vialard and Delecluse, 1998; Sumner and Belaineh, 2005; Vancoppenolle et al., 138 139 2009; Yu, 2011). SSS is known to impact the oceanic upper mixing significantly (Latif 140 et al., 2000; de Boyer Montegut et al., 2004; Maes et al., 2006; Furue et al., 2018) 141 and via its effect on the surface layer density (Johnson et al, 2012). The SSS also 142 affects the decadal variability of hydrography in the upper waters of the North Atlantic (Reverdin et al., 1997). Using a coupled atmosphere-ocean model and an observed 143 144 SSS climatology dataset, Mignot and Frankignoul (2003) attributed the interannual 145 variability of the Atlantic SSS to two factors: anomalous Ekman advection and the 146 freshwater flux, Additionally, the increased melting of glaciers and sea-ice in the 147 Arctic (McPhee et al., 1998; Macdonald et al., 1999) leads to significant changes in 148 the salinity distribution and fresh water pathways (Steele and Ermold, 2004; Morison 149 et al., 2012). The freshwater flux is regarded as one of the least constrained 150 parameters in ocean models due to poorly known river discharge, precipitation, and 151 glacial/sea-ice melt (e.g., Tseng et al., 2016; Furue et al., 2018). In ocean models the 152 sea-surface freshwater flux is often adjusted directly or the SSS is restored to its 153 corresponding climatological value to avoid salinity drift. 154 155 Monitoring SSS from space is crucial for understanding the global water cycle and 156 the ocean dynamics, especially in the Arctic Ocean where our knowledge of the SSS 157 variability is limited due to non-homogenous and sparse in-situ data. The European 158 Space Agency's (ESA) Soil Moisture and Ocean Salinity (SMOS) satellite, launched 159 in November 2009, consists of the Microwave Imaging Radiometer using Aperture 160 Synthesis (MIRAS) instrument, a passive 2-D interferometric radiometer operating in 161 L-band (1.4 GHz, 21 cm), that measures the brightness temperature (BT) emitted 162 from the Earth, The L-band microwave is highly sensitive to water salinity, which 163 influences the dielectric constants in the sea, and is less susceptible to atmospheric 164 or vegetation-induced attenuation than higher frequency measurements (Font et al., 165 2010; Kerr et al., 2010; Mecklenburg et al., 2012). Committed to provide global 166 salinities averaged over 10-30 days with an accuracy of 0.1 psu in the open ocean, 167 ESA provides the MIRAS data into SMOS Level 1 (L1) and Level 2 (L2) products 168 through a set of sequential processors (Mecklenburg et al., 2012; ESA, 2017).

Deleted: to track hydrological

Formatted: Header

Deleted: de Boyer Montegut et al., 2004;

Deleted: dominance	
Deleted:) the	
Deleted: affects the thermohaline circulation	
Deleted: northern	
Deleted: Frankgnoul	

Deleted: ¶

	Increase in the freshwater content of the Arctic Ocean due to
-(Deleted:), a
1	Deleted: change in the global warming scenario, can leads to
-(Deleted: However, the
(Deleted: the small-scale features of
(Deleted: general, to avoid salinity drift in the
-(Deleted: ,
-(Deleted: by restoring
{	Deleted:

-(Deleted: MIRIAS
_(Deleted: to measure
_(Deleted: (Font et al.,
(Moved down [1]: 2010; Kerr et al.,
~(Deleted: 2010).
(Deleted: has
_(Moved (insertion) [1]
(Deleted: Mecklenburg et al. 2012) Since its operational

Deterd: Mecklenburg et al., 2012). Since its operational phase started in May 2010, SMOS provides the longest SSS record from space over the global ocean, even compared with the National Aeronautics and Space Administration's (NASA) Aquarius mission (between 2011 and 2015) and its follow-up SMAP (Soil Moisture Active and Passive, since 2015

Formatted: Header

o provide global salinities averaged an accuracy of 0.1 psu for open sible to interpreter the MIRAS

Ind Level 2 (L2) data through a set sors (Mecklenburg et al., 2012; 1 processing stage, the three L1A, L1B, and L1C are respectively calibrated engineering visibility, the postruction and multi-angular BT at e (TOA) Over oceans (TOA). Over oceans

of atmosphere and on…t the sea / ESA with swath-based format 016; ESA, 2017). Under...s a ...f the national agencies in France ly, two Level 3 (L3) data products illable, which are independently ean Salinity Expertise Center re Aval de nent des Donnees…onnées SMOS R and the Barcelona Expert tudies comprehensively investigate tities in the Arctic Ocean at same two SMOS ... These two SMOS .uccessfully used to local...gulhas salinity front ... [37]

Text 1, Not Highlight

these monitoring activities from [38] ice...ecessary step towards the 39] salinity, Uotila et al. (2018) . [40]

Formatted: Font color: Text 1
Deleted: of monthly salinity
Deleted: of
Formatted: Font color: Text 1
Formatted [41]
Formatted: Font color: Text 1
Deleted: (
Formatted: Font color: Text 1
Deleted:) shows a considerable spread among the e_{42}
Formatted: Font color: Text 1
Deleted: extraordinarily poor
Formatted: Font color: Text 1
Deleted: coverage
Formatted: Font color: Text 1
Deleted: domain. With the accumulated The SSS _[43]
Formatted: Font color: Text 1
Deleted: is derivedrom the TOPAZOPAZ4 [44]
Moved down [2]: Xie et al., 2017
Deleted:),a coupled ocean and sea-ice data [45]
Moved (insertion) [2]
Deleted: from The second is the CMEMS. This [46]

201	Υ		-1	Deleted: Committed to
202	Over the ocean, Level 2 products (L2OS) are comprised of three different ocean		0	over 10-30 days with ocean, ESA is respon
203	salinities, together with the BTs at the top of atmosphere and at the sea surface,		5	SMOS Level 1 (L1) ar
204	distributed by ESA with swath-based format (e.g., SMOS Team, 2016; ESA, 2017).		E	of sequential process ESA, 2017). In the L1
205	As a result of the efforts of the national agencies in France and Spain respectively,		C	corresponded to the c
206	two Level 3 (L3) data products of SSS are freely available, which are independently	/	t	he top of atmosphere
207	developed by the Ocean Salinity Expertise Center (CECOS) of the Centre Aval de	/	I و	Deleted: TOAhe top surface, distributed by
208	Traitement des Données SMOS at IFREMER and the Barcelona Expert Centre	$/\parallel$	((e.g., SMOS Team, 20 esult of the efforts at.
209	These two SMOS products have successfully resolved the Agulhas salinity front	//	a c	and Spain respectivel of SSS are freely avai
210	(D'Addezio et al., 2016) and proven useful for the estimating precipitation (Supply et	/	c (developed by the Oce
211	al., 2018). The work of Olmedo et al. (2018) quantitatively evaluate the accuracy of		Ì	Traitemenentraitem
212	the SMOS Arctic and sub-Arctic SSS to less than 0.35 psu, but this evaluation		Ċ	Centre (BEC). Few st
213	against Argo data was limited by the lack of data in the Arctic proper. The present		t	ime, although these t
214	study thus investigates the accuracy of these two SMOS SSS products in the Arctic		r	resolveesolved the
215	Ocean.		-(1	Formatted: Font color:
216	Y.		-(1	Deleted: In parallel to
217	A good estimate of surface salinity is a necessary step towards the knowledge of the		7	Deleted: practical choi
218	three-dimensional water mass properties, for which data assimilation and optimal	/	6	Delatad: Aratia Ocean
219	interpolation methods must be invoked. In a recent study, Uotila et al. (2018)			Formatted: Font color:
220	investigated the Arctic salinity in ten reanalysis products and found disagreements		G	Deleted: of monthly sa
221	within them regarding the seasonal cycle in the upper layer (0-100 m; Figure 12 of			Deleted: of
222	Uotila et al., 2018). Note that the full assessment of the Arctic SSS products has			Formatted: Font color:
223	been hindered by the extreme paucity of in-situ data in the Arctic. The SSS data from	M/	()	Formatted: Font color:
224	the SMOS mission should in principle allow the evaluation of salinity on a basin		Ì	Deleted: (
225	scale. In this study, we use two SSS products available from the Constraints Marine	וווו	U	Formatted: Font color:
225				Deleted:) shows a con
226	Environment Monitoring Service (CMEMS). The first is the regional Arctic CMEMS			Pormatted: Font color:
227	reanalysis (ARCTIC-REANALYSIS-PHYS-002-003) from the TOPAZ4 assimilation			Formatted: Font color:
228	system, which is a coupled ocean and sea-ice data assimilation system using the		Ī	Deleted: coverage
229	Ensemble Kalman filter (EnKF) to assimilate the various ocean and sea-ice		Q	Formatted: Font color:
230	observations (e.g., Xie et al., 2017). The second is the CMEMS multivariate optimal		J	Deleted: domain. With
231	interpolation reprocessing (MULTIOBS GLO PHY REP 015 002 Droghei et al.			Formatted: Font color:
032	2018) The latter product directly merges in situ data with satellite measurements			Moved down [2]: Xie e
2.52	including CMOC without the use of a model and is therefore a neuropeople and the	$/\parallel$		Deleted:),a coupled
233	Including Sivices without the use of a model and is therefore a reprocessing rather		G	Moved (insertion) [2]
234	than a reanalysis. There are four other global reanalysis products under CMEMS,		Y	Deleted: from The

L

394	but understanding well their differences requires an intimate knowledge of their		
395	setup, and is out of scope of the present study.		
396			
397	We assess the quantitative deviations of Arctic SSS among the two SMOS products		
398	and the two CMEMS products, together with two climatology datasets: WOA13		Deleted: optimally merg
399	(version 2.0 of World Ocean Atlas of 2013; Zweng et al., 2013) and the older PHC		CMEMS reanalysis pro
400	(Polar Science Center Hydrographic Climatology version 3.0; Steele et al., 2001). We		climatology datasets:
401	further extend the evaluation using available in-situ salinity observations during the		PHC (Polar Science Conversion 2.0) Steele et al
402	years 2011-2013 from different data sources. Can the evaluation against the in-situ][[evaluation using availa
403	data also shed light on the uncertainty of the SMOS products? Can it also give useful]	sources. Thean the
404	information needed for the assimilation of the SMOS SSS products into an Arctic	/	SMOS products toward
405	ocean forecast/reanalysis system?]	information needed for
406			systems in near future.
407	The paper is organized as follows: Section 2 describes all SSS products and the in-		Deleted: Inection 2,
408	situ datasets. The monthly mean SSS from these six products are intercompared and		are describedataset
409	monthly differences from the TOPAZ SSS are analyzed in Section 3. Section 4	///	theonthly deviations
410	evaluates the SSS products against in-situ data, which are divided between	//	4 illustrates the quantit
411	assimilated and independent data. A summary of this study is provided in Section 5.]	are divided into two se
412			assimilated into TOPA
413	2. Data description		
414	2.1 Sea surface salinity from SMOS		Deleted: comingrigin
415	The SSS retrieval from SMOS is subject to biases originating from various non-		contaminationson-ge so-called land-sea con
416	geophysical sources such as the so-called land-sea contamination and the latitudinal		biases likely mainly of instrument. A particula
417	biases, mainly caused by the thermal drift of the instrument. A particular challenge in	/ //	sea-ice edge because Based on different stat
418	the Arctic is the sea-ice edge because of ice-ocean contamination. Based on		marchpproaches, ma filtering flags, the CEC
419	different statistical approaches, match-up criteria, and SMOS data filtering flags, two]	have independently chain to produce the re
420	centers have developed separate processing chains producing a Level 3 SSS]	3 SSS product on a reg These two SSS product
421	product on a regular arid. These two SSS products are hereafter named respectively	/	respectively named
422	CEC and BEC in this study, evaluated during the three years of 2011-2013 (see]	Deleted: Thishe late
423	Table 1).		latitudes Oceans and in
424	• <u>The BEC product</u>		Deleted: June 2018).
425	The latest regional Arctic product (version 2.0) from BEC is available from	/	March 2019). The BEC from ESA L1B (v620) p
426	http://bec.icm.csie.es since December 2018 (last access: March 2019). The BEC	[/]	2016), and accumula

SSS product was generated from ESA L1B (v620) products, and accumulates salinity

Deleted: optimally merged observational data products.¶ In this paper, we assess the performance of ...he two CMEMS reanalysis products in comparison to the two SMOS SSS products...roducts, together with the ...wo climatology datasets: WOA13 (version 2.0 of World Ocean Atlas of 2013; Zweng et al., 2013) and the older PHC (Polar Science Center Hydrographic Climatology version 3.0; Steele et al., 2001). We further extend the evaluation using available in-situ salinity observations during the years of ...011-2013 from different data sources. The...an the evaluation against the in-situ data is ...Iso expected to...shed light on the uncertainty of SMOS products towards the reliable Arctic SSS monitoring program, which... Can it also give useful information needed for the assimilation of the SMOS SSS products int an Arctic ocean forecast/reanalysis

Formatted: Header

Deleted: In ...ection 2,...describes all the assessed ...SS products and reference...he in-situ data are described....atasets. The monthly means of...ean SSS from these six products are intercompared,...and the ...onthly deviations reference to...ifferences from the TOPAZ SSS are analyzed in in ...ection 3. Section 4 illustrates the quantitative evaluations of...valuates SSS products against the reference ...n-situ data, which are divided into two sets of observations based on whether the observations had been...etween assimilated into TOPAZ or not [48].

... [47]

developed in by...version 2.0) from BEC targeting high atitudes Oceans and in the Arctic Ocean,...s available rom http://cp34-bec.mima.csie.es[50]

Deleted: June 2018). ¶ March 2019). The BEC SSS product was generated from ESA L1B (v620) products (SMOS-BEC Team, 2016),... and accumulates the [51] Formatted: English (US)

Field Code Changed

526	data over 9 days with a spatial grid resolution of 25 km, With respect to its previous
527	version, a systematic bias in the retrieved salinity is corrected by computing the
528	SMOS climatology (the most probable value for a given lat-lon, incidence angle and
529	across-swath distance) which is substituted by a reference value from WOA13. In
530	addition, a temporal bias correction has been refined in this version using near-
531	surface Argo salinity to compute regional averages (see the details in Olmedo et al.
532	<u>2018).</u>
533	• <u>The</u> CEC product
534	The third version of LOCEAN SMOS SSS L3 maps (L3_DEBIAS_LOCEAN_v3) was
535	released by the CECOS in July 2018. Every 4 days, the SSS maps averaged over 9
536	days are released on ftp.ifremer.fr (last access: December 2018). This product uses
537	the Equal-Area Scalable Earth Grid (EASE-Grid) which has limited grid distortion and
538	a spatial resolution of 25km. Using a Bayesian retrieval approach (Kolodzejczyk et
539	al., 2016), the SMOS systematic errors in the vicinity of continents are discarded o
540	improve the product quality. Further, a 'de-biasing' method (Boutin et al., 2018) has
541	been applied in this version of the CEC product, in which the non-Gaussian
542	distribution of SSS is taken into account, refining the latitudinal correction at high
543	latitude, and preserving the naturally seasonal variability of SSS.
544	
545	2.2 Sea surface salinity from two CMEMS products
546	The <u>TOPAZ4</u> Arctic <u>MFC reanalysis</u>
547	" <u>TOPAZ4</u> uses the version 2.2 of Hybrid Coordinate Ocean Model (HYCOM,
548	Chassignet et al., 2003; Bertino and Lisæter, 2008) coupled with a simple
549	thermodynamic sea ice model (Drange and Simonsen, 1996) in which the elastic-
550	viscous-plastic rheology describes the sea ice dynamics (Hunke and Dukowicz,
551	1997). The model domain covers the Arctic Ocean and the north Atlantic Ocean with
552	a horizontal resolution of 12-16 km. In order to obtain an accurate and dynamically
553	consistent reanalysis in the Arctic Ocean, the deterministic EnKF (DEnKF; Sakov and
554	Oke, 2008) was implemented in TOPAZ with a dynamical ensemble of 100 members
555	all driven by perturbed 6-hourly atmosphere forcing from ERA interim (Simmons et
556	al., 2007). The perturbations of precipitations are following a log-normal probability
557	distribution and conserve the ensemble-average total precipitation.
558	Along the model lateral boundaries in the South Atlantic and in Bering Strait, the

temperature and salinity are relaxed to a combined climatology data from PHC and

Formatted: Header

Deleted: for the period of 2011-2013. Using a non-Bayesian approach

Deleted: of the L1B salinity data is debiased against

Deleted: SSS extrapolated from

Deleted: float at 7.5 m depth, which are provided by the Coriolis data center (<u>www.coriolis.eu.org</u>). For further processing detail, see

Formatted: English (US)

Deleted: (2016). The bias corrected data are spatiotemporally interpolated to the L3 binned maps. Then their anomaly is blended with WOA09 SSS climatology (Antonov et al., 2010) using optimal interpolation with 300 km influence radius to produce the final L3 regularly gridded, daily SSS product (OA L3 SSS). The OA L3 SSS maps are served daily on regular 25 km grids for an average period of 9 days.

Deleted: were

Deleted: of CATDS

Deleted: These

Deleted: with

Deleted: accumulation period at every 4 days are provided from 16th January 2010 to 25th December 2017. These products, using Equal-Area Scalable Earth (EASA) Grid in which pixels have a constant area and longitudes are equally spaced but not latitudes, have a spatial resolution of 25km freely available on FTP:

Field Code Changed

Deleted: Beginning from the ESA L1B products, the BTs are reconstructed under apodization window and interpolation procedure (Vergely and Boutin, 2017). Based on a semi-empirical ocean surface model developed internally, three different forward models in the L2 processors are implemented for the SSS retrieval and relevant geophysical parameters (SST, wind, etc.). Only one of these three SSSs from the L2 processors are used as L2OS on an EASE grid, similar to ESA L2OS (v622) products. Using the

Deleted: migrated to

 $\boldsymbol{Deleted:}$), an improved technique to correct systematic biases,...

Deleted: used

- Deleted: where
- Deleted: Gaussianity

Deleted: the

Deleted: reanalyses in

Deleted: reanalyses from TOPAZ

Deleted: TOPAZ

Deleted:). In the sea ice model,

Deleted:) was used to describe the ice dynamics.

Deleted: northern

Formatted: Header

608 WOA. The river discharges are treated as an additional mass and a negative salinity 609 flux. Near the surface, to avoid the salinity drift (Tseng et al., 2016; Furue et al., 2018), a weak relaxation to the same combined climatological SSS with 30 days 610 611 decay, is used as most ocean models, but restricted to the areas where the difference 612 to climatology is smaller than 0.5 psu. The EnKF assimilates various ocean and sea-613 ice observations (e.g., Xie et al., 2016, 2018) into a multivariate state update of the 614 HYCOM model. The understanding for the uncertainty of the TOPAZ4 SSS has been hindered by 615 poor coverage of in-situ data over the Arctic domain, although Xie et al. (2017) had 616 comprehensively assessed the <u>TOPAZ4</u> reanalysis during 1991-2013 against various 617 618 types of ocean and sea-ice observations. For the sake of brevity, the TOPAZ4 619 reanalysis SSS is named TP4 hereafter. 620 621 SSS from the Multi-OBservations dataset The CMEMS product of MULTIOBS_GLO_PHY_REP_015_002, combines the SSS 622 623 observations from in-situ and satellite data, using optimal interpolation (OI, 624 Buongiorno Nardelli et al., 2016; Verbrugge et al., 2018) at weekly interval on a 0.25° 625 x 0.25° regular grid. The main datasets used during the OI processing are; 1) the quality controlled in-situ data, COriolis dataset for Re-Analysis (CORA, Cabanes et 626 al., 2013) distributed through CMEMS: 2) the objectively analyzed SSS and SST data 627 generated from CORA, also distributed by CMEMS, which uses the WOA 2013 628 629 climatology as first guess and has been upscaled to the MOB grid as another first 630 guess of the multidimensional OL 3) The SMOS L3 binned (L3bin) data reprocessed by SMOS-BEC at 0.25° grid, although the previous version 1.0 of the product 631 632 mentioned above; 4) The daily Reynolds L4 AVHRR_OI Global blended SST product 633 on a 0.25° grid. This product is called MOB hereafter. 634 635 2.3 <u>Surface salinity</u> from in-situ data 636 The in-situ SSS data are acquired here from three quality-controlled datasets. The first data source is CORA from CMEMS (product id: 637 INSITU GLO TS REP OBSERVATIONS 013 001 b), also used in the MOB SSS. 638 CORA contains temperature and salinity profiles from various in-situ data sources 639

640 (Cabanes et al., 2013). Since 2013, the CORA dataset has been updated every year

Deleted: Near the northern model boundary, a barotropic inflow at the Bering Strait is imposed to involve the impact of Pacific water, which varies seasonally as indicated by observations. Due to the poor knowledge on the river discharge into the Arctic, a monthly climatology is calculated by the precipitation from the ERA interim (Simmons et al, 2007) averaged over 20 years, which was ingested to the Total Runoff Integrating Pathways (TRIP, Oki and Sud, 1998) hydrological model. In the model, the Deleted: exchange by Deleted: (Deleted:) Deleted: of other Deleted: adopted

Deleted: constrain Deleted: less Moved (insertion) [3] Deleted: In order to obtain a reliable and dynamically consistent reanalysis in the Arctic Ocean, the deterministic EnKF (DEnKF; Sakov and Oke, 2008) has Moved up [3]: various ocean and sea-ice observations Deleted: 2016, 2018) are assimilated into the HYCOM531 Deleted: full evaluation Deleted: TOPAZ Deleted: TOPAZ Deleted: The related Deleted: product from this reanalysis Deleted: here after Deleted: multivariable Optimal Interpolation Deleted: (Verbrugge et al., 2018) Deleted:) and covers the years of 1993-2017 Deleted: . This product available from [54] Deleted: as follow Deleted: (product id: [55] Deleted: the Deleted: analysis system Deleted: final Deleted: the Deleted: field for Deleted: Deleted: which are built separately for descending angs61 Deleted: is used Deleted: Over the same time period (2011-2013) [57] Deleted: Salinity near surface Deleted: Against the two SMOS products from and the sal Deleted:). Initially developed to supply in-situ data $in_{[59]}$

716	and includes all the Argo float profiles, moorings, gliders, Ice-Tethered Profilers (ITP;
717	Toole et al., 2011), XBT, CTD, and XCTD data. The latest version of the dataset,
718	CORA5.1, covers the period of 1950-2016. Figure 1a shows the distribution of SSS
719	(averaged over 0-8 m depth) observations from CORA5.1 (total 69,246 observations)
720	over the domain north of 52°N during the years 2011-2013
721	The second source of in-situ data is from the Beaufort Gyre Experiment Project
722	(BGEP, http://www.whoi.edu/website/beaufortgyre/background, last access: 14th
723	December 2018). In order to monitor the natural variabilities of the Beaufort Sea in
724	the Canada Basin, BGEP maintains moorings since 2003 and acquires in-situ
725	measurements over the Beaufort Sea region every summer. Symbols (anti-triangle,
726	square, and star) shown in Fig. 1b indicate the locations of valid SSS observations
727	obtained from BGEP. The in-situ dataset used in this study is obtained from the GO-
728	SHIP (the Global Ocean Ship-based Hydrographic Investigations Program, Talley et
729	al., 2017) database under the Climate Variability and Predictability Experiment
730	(CLIVAR). The SSS observations in the Beaufort Sea are extracted from
731	CLIVAR/GO-SHIP data with EXPOCODE (33HQ20111003 and 33HQ20121005, ref.
732	Mathis and Monacci, 2014), which are available from https://cdiac.ess-
733	dive.lbl.gov/ftp/oceans/CARINA/Healy/ (last access: 18th December 2018). All the
734	valid salinity profiles are averaged within the upper <u>8</u> m layer, in order to <u>match at</u>
735	best with the satellite SSS measurements. Contrarily to the CORA data, both BGEP
736	and CLIVAR data are independent from all the evaluated datasets.
737	
738	3. Intercomparison of monthly SSS <u>fields</u>
739	Prior to the intercomparison of different SSS products, all the gridded products from
740	satellite, reanalysis and climatology have been mapped on the same grid used in the
741	TP4 model by a "nearest neighbor" interpolation, To quantitatively evaluate the SSS

deviation in the Arctic, the bias and the root mean square deviation (RMSD) are

- 743 defined by
- 744

$\text{Bias} = \frac{1}{p} \sum_{i=1}^{p} (\mathbf{H}_{i} \mathbf{x}_{i}^{f} - \mathbf{s}_{i})$

$$\text{RMSD} = \sqrt{\frac{1}{p}\sum_{i=1}^{p}(H_i x_i^f - s_i)^2}$$

___(1) ___(2)

Formatted: Header

	Τ
eleted: Note that	
ormatted: Font color: Text 1	
eleted: profiles from CORA5.1 have been used in the	1
ormatted: Font color: Text 1	-
eleted: are 24249	<
ormatted: Font color: Text 1	<
ormatted: Font color: Text 1	
eleted: °	-
ormatted: Font color: Text 1	<
eleted: of	<
ormatted: Font color: Text 1	<
eleted: , and most of them are located in the norther \mathfrak{g}_2	≪ !L
ormatted: Font color: Text 1	-
eleted: sources	<
eleted: Aiming at monitoring	<
eleted: Gyre	<
eleted: is maintaining a set of observing system [63	≺ 1.
eleted: providing	4
eleted: observations	<
eleted: Gyre in	<
eleted: From the BGEP, the valid SSS observation \mathfrak{g}_{64}	
ormatted: Font color: Text 1	4
eleted: start)	<
ormatted: Font color: Text 1	<
eleted: the right panel of	
ormatted: Font color: Text 1	
eleted: 1. Last of all, we use	
ormatted: Font color: Text 1	
eleted: data from	
ormatted: Font: Arial, Font color: Text 1	
eleted: . (
eleted:))	
eleted: Specifically,	
eleted: 5	
eleted: near surface	
eleted: obtain	5
eleted: marched observations of	
eleted: for evaluation.	2
eleted: converted	2
eleted: grids as	2
eleted: method	
eleted: difference	
eleted:).	

789	Where p is the length of the time series, $\mathbf{x}_i^{\mathrm{f}}$ is the valid salinity from different sources
790	at the <i>i</i> th time, compared to the <u>reference</u> salinity field $\mathbf{s}_{i_{e}}\mathbf{H}_{i}$ is the observation
791	operator projecting \mathbf{x}_{i}^{t} onto \mathbf{s}_{i}

793	 Monthly mean comparison of SSS
794	Figure 2 shows the monthly mean Arctic SSS in March from the six products. Notable
795	differences in the two SMOS products, appear in the Nordic Seas, Barents Sea, and
796	around the Labrador Sea in the northern North Atlantic Ocean. At first sight, the
797	large-scale SSS features from SMOS products are similar to the other products.
798	However, the CEC <u>SSS</u> is fresher (as shown by the isolines of 35 psu) compared to
799	the BEC, TP4, MOB and both climatologies. The location of the sea-ice edge in the
800	two SMOS products match comparatively well with the TP4 reanalysis (Fig. 2a, d). In
801	sea-ice covered region_TP4 shows a gradual decrease in SSS from the European to
802	the American sector, with two minima near the Beaufort Sea and the East Siberian
803	Sea (ESS; Fig. 2b) consistently with the PHC (Fig. 2c). Those are unclear in the
804	MOB and WOA (Fig. 2e, f), especially the SSS minimum in the Beaufort Sea. The
805	Jatter two products also show artificial projection artefacts around the North Pole.
806	Figure 3 shows the corresponding SSS fields in September. In comparison to the
807	March situation, the BEC and CEC SSS in the Nordic Seas are both less saline,
808	indicated by the 35 psu isoline. The sea ice masking of the two SMOS products differ
809	considerably in the Canadian Basin and in the Arctic marginal seas. Although the
810	SSS of TP4, MOB, PHC and WOA agree relatively well in the northern Atlantic
811	Ocean, the discrepancies become dramatic in ice-covered areas. Below the ice or
812	near the sea-ice edge (denoted by the brown thick line in Fig. 2 and 3), TP4 and PHC
813	share common features, which can be explained by the model restoring to PHC. On /
814	the other hand, the MOB and WOA differ significantly in spite of WOA being used as $/$
815	input to the MOB. Short of a universal reference for Arctic SSS, the monthly mean
816	SSS deviations will be quantified using TP4 as a reference.
817	
818	Deviation analysis of monthly SSS referred to TP4
819	Figure 4 and Figure 5 show the deviations of the monthly mean SSS of the five
820	products with reference to the TP4 SSS in August and September respectively. In

821 August, the two SMOS products (Fig. 4a, c) show coherently negative deviations (~2

Formatted: Header

Deleted: 6	evaluated	times
------------	-----------	-------

1	Deleted: which isompared to the referredeference
l	salinity field s _i and [65]
•(Formatted: Font: Times New Roman, Bold
•(Deleted: if needs to project
1	Deleted: into

Formatted: Font color: Text 1

Deleted: means of ... ean Arctic SSS in March and reveals considerable...rom the six products. Notable differences in the two SMOS products. Notable differences are found in ... appear in the Nordic Seas, Barents Sea, and around the Labrador Sea in Northern ... he northern North Atlantic Ocean. In general, overall...t first sight, the large-scale SSS maps...eatures from SMOS products are consistent with SSS of ...imilar to the two reanalysis...ther products and the two climatology products, although the BEC SSS tends to be more saline than... However, the CEC. It...SSS is noticeable that the ...resher (as shown by the isolines of 35 psu) compared to the BEC, TP4, MOB and both climatologies. The location of the sea-ice edge in the two SMOS products marches...atch comparatively well with that of ...he TP4 reanalysis (Fig. 2a, d). Outside of the ...n sea-ice covered region in the Arctic (represented by the 15% sea ice concentration in Fig. 2) there is a good agreement between the subpolar SSS fields of the two reanalyses and the climatologies. Over the sea-ice covered region, the ... TP4 shows a gradual decrease in SSS from the sea-ice edge in ...uropean to the Nordic Seas...merican sector, with the...wo minima around...ear the Beaufort Sea and the East Siberian Sea (ESS; Fig. 2b), being consistent... consistently with the result in the ...HC (Fig. 2c). The features mentioned above, especially the minimal center in the Beaufort Sea,...hose are missing...nclear MOI...he MOB and WOA (Fig. 2e, f)...., especially the SSS minimum in the Beaufort Sea. The MOI and the WOA...atter two products also show commonly a potential ...rtificial projection issue [66] Deleted: As a contrast in summer, Fig. ...igure 3 shows the corresponding SSS fields in September respectively from the SMOS products, the reanalyses and the climatologies. Considerable differences in the two SMOS products are also found in Fig. 3 similar to that shown in Fig. 2. The SSS field from CEC is relatively fresher then the BEC.... In comparison to the climatologies, the BEC SSS reproduces a much better representation...arch situation, the BEC and CEC SSS

in the Nordic Seas are both less saline, indicated by the 35 psu isoline. The sea ice masking of the surface salinity in this region. As to the SSS from the reanalyses (TP4...wo SMOS products differ considerably in the Canadian Basin and MOI) and the climatologies (PHC and WOA), Fig. 3 shows a good agreement in the Northern...n the Arctic marginal seas. Although the SSS of TP4, MOB, PHC and WOA agree relatively well in the northern Atlantic Ocean. However... the discrepancies among them collectively emerge under the sea-ice cover in the Arctic. Over...ecome dramatic in ice- ... [67]

Deleted: (BEC, PHC, CEC, MOI, and WOA), referred to the TP4. The...

980	psu) in the marginal seas of the Beaufort Sea, the ESS, the Laptev Sea, and the		ſ	Deleted: along the sea-ice edge
981	Kara Sea. In the North Atlantic Ocean, away from the sea-ice edge, the deviation of		Í	Deleted: Highlighted
982	the BEC from TP4 is lower (bias less than 0.5 psu). Focusing on the Arctic domain		F	Formatted: Font color: Auto
983	(>60°N), the mean deviation of the BEC SSS is -0.87 psu and its root mean square is		ſ	Deleted: SSS deviation of BEC in August is about -0.5
984	1.75 psu. The CEC SSS shows considerable negative deviations over 1 psu in the		e	edge, the deviation of BEC has a slight positive bias
985	northern Atlantic, from north of Denmark Strait to the west coast of Ireland. This is		t	he CEC SSS, the averaged deviation is about -0.42
986	remarkably different from the BEC, and does not discern the subpolar from the		p d	su with RMSD about 1.73 psu. Notably clear negative leviations appear in both BEC and CEC products
987	subtropical waters there (Hátún et al., 2005). The deviations of MOB and the two		t	consistently along the sea-ice edge in the Beautort Sea, he ESS, the Laptev Sea and the Kara Sea. However,
988	climatology products are comparatively small in the open ocean of the northern		C	lear differences over the north Atlantic and Arctic
989	Atlantic (Fig. <u>4b, e). Near and below the sea-ice cover, the deviations are much</u>		0	leviations in the northern Atlantic with a minimum over
990	larger, particularly both the MOB and WOA show strong saline anomalies (> 1 psu) in		1 n	psu located at the north of Denmark Strait, it has elatively strong positive deviations near the coasts of
991	the Eurasian basin and low anomalies in the American basin.		T	The deviations in the northern Atlantic in MOI (Fig. 4d)
992	▲ ▲		a (Fig
993	In September, the SSS deviations of BEC, MOB, PHC and WOA show similar fresher			Moved down [4]: 4b, e).
994	patterns as in August, but the CEC deviations becomes surprisingly positive around	**********	I it	Deleted: However, over the sea-ice covered region and the surrounding sea waters, the differences are rather
995	the ice edge. The SSS deviation of CEC, averaged over the Arctic domain (>60°N),		s d	ignificant. The PHC has a relatively small negative leviation over the majority of the Arctic and north
996	swaps from -0.42 to 0.42 psu from one month to the next one. The seasonal		e A	Atlantic Oceans (Fig. 4b). However, around the sea-ice adge, the deviations are much larger. On the other
997	evolution of monthly SSS deviations from TP4 for all five remaining products.		h	and, MOI and WOA have strong positive deviations over the Eurasian basin (> 1 psu), with respective
998	averaged over the Arctic, are shown in Fig. 6. Among the five products, the MOB	annan an a		MSD of 4.21 and 3.29 psu in the whole Arctic region.
999	shows the strongest seasonality with the RMSD higher than 4 psu in July and August	10111111111111	2	2.96 and 2.28 psu respectively. The averaged SSS
000	(Fig. 6a), and close to 2 psu in winter. The spatially averaged deviation is much	10000000000000000000000000000000000000	A	leviation of PHC (Fig. 5b) becomes slightly less than in August mainly due to the positive deviations along the
001	fresher than TP4, over -2 psu in summer and -0.5 psu in winter (Fig. 6b). The	1122111221122	S	sea-ice edge in the marginal seas. Although the two SMOS SSS products from SMOS have the smallest
002	deviations of the two SMOS SSS show a relatively smaller seasonality (Fig. 6a)	12210200000	O	leviation among the five products (Fig. 5a, c) with [68]
002	During summer months, their RMSDs reach 1.5 nsu (Fig. 6a) in summer, and they	11111111111111111		Deleted: about -1.5 psu. This suggests the MOI SSS is
003	degraphic to 0.5 and 1.0 psu (for PEC and CEC respectively). Throughout the whole	1000000000	Q	uite close to the WOA in the Arctic domain. As for [69]
004	decrease to 0.5 and 1.0 psu (for BEC and CEC respectively). Throughout the whole	12112211221		Moved (insertion) [4]
005	year, the BEC RMSDs (Fig. 6a) are consistently smaller than that of CEC, and the	100000000	Ī	Deleted: deviation has a significant seasonality of the
006	seasonal cycles are different. This shows that the BEC SSS is closest to TP4,		I	nean deviations over -0.5 psu during summer and less
007	although it is overall fresher in the Summer.		L L	Cormatted: English (US)
008			Č	Deleted by
009	4. Evaluation against in-situ observations		T	Deleted: Referred to Eqs. 1-2, the quantitative
010	The misfits of the six SSS products from SMOS_CMEMS and climatologies are		Ì	Deleted: the
010		\leq	I	Deleted: the reanalyses
011	calculated as in Eqs. (1) and (2) against the pointwise in-situ observations described		T	Deleted: the

1012 in Section 2.3. For TP4, the SSS evaluation is conducted on the same model day as

Deleted: discrete Deleted: and BEC

Formatted: Header

11	111	the in-situ observations. Owing to the fact that the SSS from BEC, CEC and MOB are		Dele
11	112	averaged over either 9 days or one week (see Table 1), the product dates, at the		Dele
11	113	center of the averaging window lag 5 or 4 days compared to the observation date,		Dele
11	114	For PHC and WOA, the in-situ observations are sorted to monthly bins and evaluated		Dele
11	115	for each month. The quantitative evaluation is divided into two main sections starting		Dele
11	116	with dependent and then independent observations.		Dele
11	117			
11	118	4.1 Against SSS from CORA5.1		Dele thre
11	119	As shown in Fig. 1a, the distribution of SSS observations from CORA5.1 over the		area
11	120	Arctic is very inhomogeneous during the three years. Due to this, the evaluation of		Dele
11	121	the gridded SSS products against in-situ observations is restricted to the observation-		Her dom
11	122	rich regions. The SSS misfits bias and RMSD for the six products are reported in		peri whe
11	123	Table 2 according to the eight Arctic sub-regions defined previously (Figure 1a). The		mel
11	124	observations are displayed on scatterplots (Figure 7 and 8) to exhibit their		Dele
11	125	uncertainties for fresh and saline waters in different areas.	//	<i>In th</i> In th
11	126	<u>Central Arctic</u>		from avai
11	127	Figure 7 shows the SSS products compared with discrete observations in the central		97% from
11	128	Arctic (sub-regions S0, S1, S2, and S3). The observed SSS in S0 and S1 are mainly		SSS Ove
11	129	from the ITP at a minimal depth of 8 m. Around the North Pole (S0), where the		Nor con:
11	130	satellite SSS are absent, the TP4 reanalysis and MOB reprocessing show opposite		the Amo
11	131	biases: +0.48 psu and -0.52 psu respectively (Table 2). The two climatologies used		spa MO
11	132	by them, PHC and WOA respectively, also show opposite biases. Considering the		esp mos
11	133	latter climatologies, both SSS scatterplots shows a fresh bias for high salinity water		If or TP4
11	134	(>33 psu) and a saline bias for low salinity water (<31 psu).		app alor
11	135	In the Canadian basin (in S1), the two climatological SSSs show an obvious gap in		obvi Aga
11	136	comparison to the ITP observations. Comparing to the fresh in-situ SSS from 24 to		SMO
11	137	30 psu, the PHC has strong saline bias (from 2 to more than 5 psu). On the other		but sum
11	138	hand, the WOA shows both a fresh bias for relatively high salinity water (>28 psu)		RM (S6
11	139	and saline bias for fresher water (<26 psu). Owing to the different time periods (Table		the are
11	140	1) of the in-situ data they used, this result confirms the freshening of the Canadian		stro Figu
11	141	basin since in the 1990s (Morison et al., 2012).		devi Atla
11	142	In the S1 sub-region, the satellite SSS from BEC and CEC have only 20 and 42 data		loca proc
11	143	points for evaluation respectively. The resulting scatterplots show a significantly		8d) mini

Deleted: observing

Formatted: Header

Deleted: For CEC and MOI, the corresponding evaluation is made... Deleted: product

Deleted: nearest backwards in time to the observing

tes...

Deleted: bin

Deleted: in

Deleted: SSS observations from CORA5.1 during the three years are distributed unevenly over the pan-Arctic area. Due to the non-homogenous

Deleted: the observations

Deleted: limited to the observational-dense domains. Here, we specifically focus our evaluation over the two domains: the northern Atlantic Ocean during the entire period and the Beaufort Sea during summer seasons when the surface is exposed owing to the sea ice melting....

Formatted: Font: Italic, Font color: Auto

Deleted:

In the northern Atlantic Ocean and Nordic Seas¶ In the northern Atlantic Ocean including the sub-regions from S4 to S7 (Fig. 1a), 23626 salinity observations are available for this evaluation, corresponding to more than 97% of all valid observations over the Arctic domain from CORA5.1. Figure 7 shows the mean deviation of SSS for each product during the years of 2011-2013. Over the northern Atlantic oceans including the Norwegian Sea and the Greenland Sea, the considerable negative biases (<-0.16 psu) are shown in the products of CEC, PHC and WOA (Fig. 7c, d, f). Among of them, the CEC shows significantly high spatial variability. The SSS products of BEC, TP4 and MOI (Fig. a, b, e) have relatively small bias (<0.08 psu), especially the MOI shows the minimal deviations in most of this region. ¶

If only comparison of the SSS between the BEC and the TP4, the latter has two stronger positive biases appearing along the southern Norwegian coast and along the Greenland west coast, although it has obviously smaller bias than the BEC in the open seas. Against the Argo profiles from the Coriolis data center, SMOS-BEC Team (2016) found the RMSDs of the BEC SSS in the Arctic (>50°N) are mostly less than 0.4 psu, but also showing the interannual variability like in the summer of 2012 the RMSD close to 0.8 psu. The RMSDs of the BEC SSS in the Arctic (>50°N) are mostly less than 0.4 psu, but near the coast regions (S4 and S5 in Table 1) the RMSDs are over 1 psu. It further indicates the BEC quality has a strong dependency on the locations. ¶ Figure 8 shows the Root Mean Square (RMS) deviations of SSS for the all products over the northern Atlantic Ocean and the Nordic Seas. Averaged in the local domain, the maximal deviation among the six products can be found about 1.0 psu in the CEC (Fig. 8d) in which high spatial variability is also profound. The minimal deviation among them is found about 0.4 psu in

the MOI (Fig. 8e), in which similar magnitude of the RMSDs are distributed over the entire domain relatively

Formatted: Header

1284	positive salinity bias (>4 psu) for fresh waters (<27 psu). For relatively higher salinity	
1285	water (> 27 psu), the CEC has a stronger saline bias than the BEC.	
1286	In the Kara Sea (sub-region S2), the TP4 SSS has the smallest RMSD at 1.7 psu,	
1287	which is significantly smaller than other products. The scatterplot also shows a good	
1288	linear relationship between the TP4 and the in-situ SSS, while other products	
1289	generally show fresh biases, indicating that the SSS variability in the Kara Sea is well	
1290	captured by TP4. In the Barents Sea (sub-region S3), TP4 gives as well the smallest	
1291	misfit (RMSD: 0.34 psu; bias: -0.14 psu). The SSS scatterplots exhibits linear	
1292	relationships for all products except the CEC, which underestimates the Atlantic	
1293	water SSS.	
1294		
1295	Northern North Atlantic and Nordic Seas	
1296	Figure 8 shows the paired scatterplots of the six SSS products in the subpolar seas	
1297	from sub-regions S4 to S7 (see Fig. 1a). In S4 and S5, the bias of SSS products is	
1298	relatively small, less than 0.15 psu (Table 2), except for CEC in S4 and TP4 in S5,	
1299	both too saline by 0.2 psu. The scatterplots further indicate that low salinity waters	
1300	are too saline in all SSS products in S4 (<31 psu) and in S5 (<28 psu). Meanwhile,	
1301	the respective bias and RMSD of the SSS products are less than 0.1 psu and 0.43	
1302	psu respectively, except for the CEC in S6 and S7. The MOB SSS has the smallest	
1303	salinity bias. Among the eight regions compared here (S0 to S7), the SSS bias is	
1304	lowest in S6 (Irminger Sea).	
1305	Over the northern North Atlantic and the Nordic seas, Fig. 9 shows maps of the mean	
1306	SSS deviation for each product during the period 2011-2013. Considerable negative	
1307	biases (<-0.2 psu) are found in the CEC, whereas the MOB and WOA have the	
1308	smallest bias, less than 0.02 psu (Fig. 9 d, e, f). The SSS products from BEC, TP4	
1309	and PHC (Fig. 9 a, b, c) have slightly higher bias (~0.05 psu) in comparison to the	
1310	MOB and WOA. On average, the BEC bias is only -0.04 psu, much smaller than that	
1311	of the CEC (<-0.2 psu). Focusing on the BEC SSS, Fig 9a shows that while a fresh	
1312	bias dominates the Nordic Seas, the product is too saline in the northern North	
1313	Atlantic and the North Sea.	
1314	The inter-comparison of the biases against the in-situ data in Fig. 9a and 9b exhibits	Formatted: Font color: Text 1
1315	two strong positive biases of TP4 along the Norwegian coast and along the West	Deleted: the summer
1316	Greenland coast. Notably, the BEC has smaller bias along both coasts, although it	
1317	has a slightly saline bias offshore. This indicates potential benefits of the BEC SSS	

1319	for the TOPAZ system along the Norwegian and Greenland coasts, were it		
1320	successfully assimilated into the system. Figure 10 shows RMSDs of SSS for all the		
1321	products over the northern North Atlantic Ocean and the Nordic Seas. On average,		Deleted: are plotted in Fig. 9. The
1322	the largest uncertainty is found with the CEC (~1.0 psu; Fig. 10d), with RMSDs as		Formatted: Font: Italic
1323 1324	large as 1.5 psu in the Greenland Sea and the Barents Sea. The SSS RMSDs for the five other SSS products are much smaller (~0.5 psu).		Deleted: from in-situ dataange from 15 to 33 psu. The BEC SSS ranges from 24 to 31 psu with a bias of 0.65 psu and RMSD of 2.63 psu. On the same panel, the TP4 ranges from 26 to2 psu, with a bias of 2.73 psu and RMSD of 3.85 psu The range of BEC SSS is
1825			limited to 24 to 31 psu with a minor bias of 0.09 psu and a RMSD of 1.82 psu. On the other hand, the range of
1326	4.2 Independent SSS in the Beaufort Sea,		TP4 SSS is even shorter from 19 to 32 psu, with a large saline bias of 2 59 psu and a RMSD of 3 63 psu. The
1327	Independent in-situ data from BGEP and CLIVAR are used during the summer		linear regression coefficients for BEC and TP4 are
1328	months of 2011-2013 in the Beaufort Sea for the evaluation of the six SSS products		thatooking at the significant deviations of BEC and
1329	(Fig. 11). The in-situ SSS observations range from 15 to 32 psu, The range of BEC		attributed to the particular four observations around. [72]
1330	SSS is limited to 24 to 31 psu with a minor bias of 0.09 psu and a RMSD of 1.82 psu.		Formatted [73]
1331	On the other hand, the range of TP4 SSS is even shorter from 19 to 32 psu, with a		Deleted: collected, 15 th August 2011 of which locations are marked in Fig. 1bv anti-triangles.
1332	large saline bias of 2.59 psu and a RMSD of 3.63 psu. The linear regression		They become on the continental shelf(Fig. 1b) near estuary ofackenzie River, where the strong fresh
1333	coefficients for BEC and TP4 are 0, <u>57 and 0,07</u> respectively. <u>Looking at the Jow-</u>		water signature could be originated to
1334	salinity observations (~27 psu) collected at (136.4°W, 70.5°N) on 15th August 2011	\parallel	rangesSS climatologies, thethe range of FRC rangesSS climatology is only reaching from 254 to
1335	marked by anti-triangles, (Fig. 1b) near the Mackenzie River, estuary, TP4 has a		bias of 1.775 psu and RMSD of 3.1385 psu.
1336	significant negative bias (< -4 psu) visible as the outliers above the dashed-black line		basin, the deviations of theHC are quite
1337	in Fig. 11a. This hints to a lack of fresh water signatures from river discharge.		themaller. The strong positive bias in theP4 at these points mostly originated an then be partly
1338	The range of PHC SSS climatology is only reaching from 24 to 31 psu, similar to		attributed to the SSS relaxation inf the TOPAZ model
1339	TP4, with a saline bias of 1.65 psu and RMSD of 2.85 psu. Compared to the TP4		weak. The range of another climatology, he WOA
1340	deviation at the Makenzie River basin, the PHC saline bias is present, but smaller.		than the range of PHC. This contributes the minimal bias of the WOA about 0.02 psu among Among the
1341	The strong positive bias in TP4 at these points can then be partly attributed to the		six products, the WOA bias is the smallest (~0.02 psu)
1342	SSS relaxation of the TOPAZ model towards the PHC climatology, albeit rather		However, it should be noticedoted that the
1343	weak. The range of the WOA is much wider, from 12 to 31 psu, Among the six	/ //	wider unders very large for salinities lower than 24
1344	products, the WOA bias is the smallest (~0.02 psu) over the Beaufort Sea during all		RMSD over>3.0 psu for of both ofHC and WOA.
1345	three summers. However, it should be noted that the variability of in-situ observations		that the two climatologies have a big
1346	is very large for salinities lower than 24 psu, which contributes to the large RMSD		sea waterow salinity regions (<24 psu) dominated
1347	(≥3.0 psu) of both PHC and WOA. It confirms that the two climatologies have a		Deleted: The CEC SSS ranges from 18 psu to 34 psu
1348	sizable uncertainty over low salinity regions (<24 psu) in the Arctic Ocean.		which is significantly wider than the range of the BEC. The SSS bias ofhe CEC SSS ranges from 13 psu to
1349	The CEC SSS ranges from 13 psu to 34 psu, which is much wider than the range of	///	34 psu, which is aboutuch wider than the range of the BEC SSS. The saline bias of CEC is however larger at
1350	the BEC SSS. The saline bias of CEC is however larger at 2,38 psu and its RMSD is	///	2.78 psu and its RMSD is about quite large at 3.97 psu. Againuthermore, the CEC deviations from the in-
1351	about <u>quite large at 3,77</u> psu. Futhermore, the CEC deviations from the in-situ		situ observations become widerre larger in the range where the SSS is lessaters fresher than 247 psu.
10.50	the section of the section of the first section of the MOD section during the section of the sec	/	For the MOI, the satellite and in-situ datahe MOB

1852 observations are larger in waters fresher than 27 psu. The MOB combined product

13

Formatted: Header

[73]

d: The CEC SSS ranges from 18 psu to 34 psu is significantly wider than the range of the BEC. SS bias of ...he CEC SSS ranges from 13 psu to a, which is about...uch wider than the range of the SSS. The saline bias of CEC is however larger at psu and its RMSD is about quite large at 3.9...7 gain...uthermore, the CEC deviations from the inservations become wider...re larger in the range the SSS is less...aters fresher than 24...7 psu. e MOI, the satellite and in-situ data...he MOB combined product, a . [76]

1475	performs poorly with the largest negative bias (>5 psu) and an RMSD in excess of 8
1476	psu. <u>In contrast</u> to the other five SSS products, the anomalously fresh SSS observed
1477	around the point (140°W, 71°N) near the Mackenzie River estuary are represented
1478	by even fresher values of 12 psu in MOB, which may hint at an amplification of the
1479	anomalies.
1480	In order to characterize dependencies of the bias for the six SSS products against
1481	the in-situ data, their absolute differences are plotted as a function of observed SSS
1482	in Fig. <u>12</u> . In general, all products show considerable deviations with the maxima
1483	reaching 8 to 14 psu. While the absolute misfits of most of the SSS products
1484	monotonically increase towards lower salinity, the bias of MOB shows its peak
1485	around 20 psu shown in Fig. <u>12c</u> . The fourth-order <u>polynomial</u> curve function,
1486	$F(S) = p_1 S^4 + p_2 S^3 + p_3 S^2 + p_4 S + p_5 $ (3)
1487	is then <u>fitted</u> to the absolute bias for each of the SSS products, where S represents
1488	the in-situ salinity. The fitting coefficients p_1 to p_{5_1} for each product are listed in Table
1489	3. The norm residuals are displayed on each panel in Fig. 12 and clearly show that
1490	fitting for MOB has the largest uncertainty, while the minimal norm residuals are
1491	about 10 and 7 psu ² respectively for BEC and TP4. This suggests the derived fitting
1492	curves for BEC and TP4 have credible skill in charactering its error distribution as a
1493	function of the observed SSS. Both curves monotonically decrease towards the
1494	salinity higher than 28 (30) psu for BEC (TP4) and increase slightly afterwards. The
1495	absolute bias in TP4 is consistently larger than that in BEC. The fitted curves of PHC
1496	and WOA have the similar functional forms to TP4 and BEC, but with lower
1497	amplitudes.
1498	
1499	5. Conclusions
1500	To understand the uncertainties in the Arctic SSS, our study evaluates the two
1501	gridded SMOS SSS products (BEC and CEC), two CMEMS products (TP4 and
1502	M(OP) and two elimetric and weta (DUC and $M(OA)$ by their interpretation and

1502 MOB), and two climatology products (PHC and WOA) by their inter-comparison and

1503comparisons against both of dependent and independent in-situ datasets during the1504years of 2011-2013.

- 1505 The differences in the spatial coverage of the two SMOS SSS were clearly shown in
- 1506 the monthly mean (Fig. 2 and Fig.3), due to the different retrieval applied in these two
- 1507 <u>datasets</u>. The <u>spatial distributions of SSS</u> from TP4 and PHC are considerably close
- to each other, <u>mainly as for</u> the fact that the SSS in the TOPAZ model is relaxed

Formatted: Header

Deleted: is significant of more than 4...>5 psu) and the...n RMSD is more than 7...n excess of 8 psu. Contrast...n contrast to the other five SSS products, the anomalously fresh SSS observed around the point (140°W, 71°N) near the estuary of ...ackenzie River estuary are represented by further...ven fresher values of around...12 psu in the MOI _____ [77]]

Deleted: ...n order to characterize dependencies of the bias for the six SSS products against the in-situ data, their absolute biases...ifferences are paired ...lotted as a function of observed SSS in Fig. 10...2. In general, all products show considerable deviations by...ith the maxima reaching 8 to 14 psu. While the absolute misfits of the ...ost of the SSS products monotonically increase towards lower salinity range... the bias of MOI...OB shows its peak around 20 psu shown in Fig. 10....2c. The fourth-order polynominal[78]

Deleted: fit...itted to the absolute bias for each of the SSS products, where S represents the in-situ salinity. The fitting coefficients from... p_1 to p_5 , for each product are listed in Table 2.... The norm residuals printed...re displayed on each panel of ... n Fig. 10...2 and clearly show that fitting for MOI contains... OB has the largest uncertainty, while the minimal norm residuals no more than...re about 10 and 7 psu² are obtained...espectively for BEC and TP4. This suggests the derived fitting curves for BEC and TP4 have credible skill in charactering its error distribution as a function of the observed SSS. Both curves monotonically decrease towards the salinity greater...igher than 28 (30) psu for BEC (TP4) and increase slightly afterwards. The absolute bias in TP4 is consistently larger than that in BEC. Although with lower amplitudes, the ... he fitted curves of PHC and WOA have the similar functional forms of...o TP4 and BEC. Their relative relation of the fitted curves, PHC being consistently larger than WOA, is also similar to that between TP4 and BEC [79]

Deleted: means of SMOS SSS ...ean (Fig. 2 and Fig.3) clearly show the two SMOS products have equivalent data coverage in winter months but obviously different in summer months..., due to the different retrieval different BT filtering flags....n these two datasets. The salinity patterns...patial distributions of SSS from TP4 and PHC are considerably close to each other, which is consistent to [81]

1614	towards PHC at each time step. Relative to TP4, the SSS deviations of the four	
1615	products (BEC, MOB, WOA and PHC) in summer show similar magnitude over the	
1616	open waters, On the contrary, the CEC SSS shows a negative bias (<-1 psu) over	///
1617	the region extending from the Iceland towards the western side of Ireland (Fig. 4.5).	
1618	but clearly the BEC SSS has a slightly negative bias over the region. In general, the	///
1619	most significant differences in the SSS deviations relative to TP4 are found under the]
1620	sea-ice cover and in its surrounding marginal ice zones.	
1621	Furthermore, the intercomparison of the SSS products shows that the BEC SSS in	
1622	August and September (Fig. 4, 5) has consistent negative deviations along the sea-	
1623	ice edge in the Beaufort Sea and the Chukchi Sea, but the CEC <u>SSS has</u> opposite	\sum
1624	deviations in these two months. Thus, it may be arguable that the two SMOS	
1625	products would give rise to significantly different effects to the upper ocean state in	
1626	the <u>TOPAZ system if it to be assimilated into. Hence the SSS quantitative</u>	
1627	evaluations of two products for optimal selection or blending would be worth of	
1628	investigating further	
1629	Focusing on the wide Arctic domain (>60°N), the deviations of the five SSS products	
1630	relative to TP4 show diverse seasonal characteristics (Fig. 6). Although the SSS	
1631	products of BEC and CEC have the similar deviation of about 1.5 psu (Fig. 6a) in	
1632	summer, the BEC deviations in winter months are clearly lower (~0.5 psu). The	111
1633	deviations of MOB and WOA (Fig. 6a) varies from over 1.5 psu in winter to around 4	
1634	psu in summer, which suggests a considerable gap with the TP4. Consequently, the	
1635	intercomparison suggests that the BEC SSS has the most consistent pattern with the	
1636	TP4 <u>SSS</u> among all <u>other</u> SSS products.	
1637	Against the in-situ data from CORA5.1 which were used in both TP4 and MOB, the	
1638	quantitative evaluations of the six SSS products were investigated in the eight sub-	
1639	regions (Fig. 1a). It was divided into two parts: in the central Arctic Ocean; the	$\langle \rangle$
1640	northern North Atlantic Ocean and the Nordic Seas. Due to the limited coverage of	\mathbb{N}
1641	BEC and CEC in S1, the scatterplots (Fig. 7) show a positive saline bias (>4 psu) for	
1642	low salinity water (< 27 psu). However, the salinity bias of BEC is slightly reduced for	
1643	relative high salinity water (> 27 psu). In the Kara Sea and the Barents Sea, the TP4	k
1644	SSS has the minimal RMSD compared with others (Table 2). The BEC scatterplots in	
1645	S2 and S3 (Fig. 7) have similar distributions with respect to TP4.	C
1646	In the northern North Atlantic Ocean and the Nordic Seas (S6, S4, and S3; Fig. 8),	1

1647 the scatterplots of the CEC SSS show that it underestimates the Atlantic water

Formatted: Header

Deleted: to the...owards PHC SSS ...t each time step. The monthly SSS patterns of MOI are clearly close to that of WOA, and they both show some partial incompatibility near the North Pole owing to the map projection (shown as in Fig. 2). ¶ Relative to TP4, the TP4 ...SS, the...deviations of the four products (BEC, MOI...OB, WOA and PHC) in summer show similar magnitude over the open waters, but... On the contrary, the CEC SSS shows an obviously... negative bias (<-1 psu) over the region extending from the Iceland towards the western side of Ireland (Fig. 4, 5). This significant..., 5), but clearly the BEC SSS has a slightly negative bias of...ver the CEC should be paid further attention in future evaluation studies about this SSS product...egion. In general, the most significant differences among... the SSS deviations relative to the ...P4 are found under the Arctic ...ea-ice cover and in its surrounding marginal seas... [82]

Deleted: The...urthermore, the intercomparison of the SSS products shows that the BEC SSS in August and September (Fig. 4, 5) shows...as consistent negative deviations along the sea-ice edge in the Beaufort Sea and the Chukchi Sea, but the CEC along the ice edge shows the...SS has opposite deviations in these two months. This indicates special attention is necessary for selecting a suitable SMOS SSS product to be assimilated into an ocean and sea-ice forecasting system. The ... [83]

Formatted: Font color: Text 1	
Deleted: impacts to	
Formatted: Font color: Text 1	
Deleted: concerned ocean mixing	g so that the
Formatted: Font color: Text 1	
Deleted: worthy	
Formatted	[84]
Deleted: studying.	
Deleted: core	
Formatted: Font color: Text 1	
Deleted: °	
Formatted: Font color: Text 1	
Deleted: theP4 show the diver characteristics (Fig. 6). The MOI seasonality in which the RMSD . products of BEC and CEC have about 1.5 psu (Fig. 6a) in summe in winter months are clearly lowe deviations of MOB and WOA (Fig 1.5 psu in winter to overround second largest seasonality can b with the RMSD ranges from 1.5 p	selyiverse seasonal has the largest Ithough the SSS the similar deviation of er, the BEC deviations rr (-0.5 psu). The g. 6a) varies from over 4 psu in summer. The be found in the WOA psu to 3.5 psu. The [85]

Deleted: have been...ere used in the...oth TP4 and the MOI...OB, the quantitative evaluations of the six SSS products have been [86].

Deleted: , but in the sea-ice covered region they are hindered by the sparse observations in the Arctic. In the northern Atlantic Ocean domain, the MOI...(S6, S4, app

_	

1604	
1804	salinity, which also is consistent with the intercomparison results (low salinity
1805	deviation) shown in Fig. 4 and 5. The misfits of mean and RMSDs shown in Fig. 9
1806	and 10, suggest the CEC SSS has considerable uncertainty (RMSD of about 1 psu),
1807	especially in the Nordic Seas with obvious low salinity biases, On the other hand, the
1808	SSS uncertainties of the BEC are significantly lower in comparison to the CEC, but
1809	are equivalent compared with TP4 and PHC. Two notable regions where the BEC
1810	SSS has lower uncertainties referred to the in-situ observations than the TP4 are
1811	along the Norwegian coast and near the west coast of Greenland Island, It is
1812	reasonable to expect that they are the most beneficial region in the Nordic Seas if the
1813	BEC SSS is successfully assimilated into the TOPAZ system.
1814	Against independent in-situ observations from BGEP and CLIVAR, the SSS
1815	evaluation in the Beaufort Sea is performed in the summers of the three years.
1816	The linear regression against these independent SSS observations (Fig. 11)
1817	suggests the BEC SSS has the smallest RMSD of 1.8 psu with a positive bias of 0.1
1818	psu, and the CEC SSS has larger RMSD of about 3.8 psu with a larger positive bias
1819	of 2.4 psu (Fig. 11). On the other hand, the TP4 SSS also shows large RMSD of
1820	about 3.6 psu with large positive bias of 2.6 psu. They are smaller than MOB which
1821	has the RMSD of <u>8.2 psu and larger negative bias (-5.0 psu).</u> As for the two
1822	climatology products, the RMSDs of WOA and PHC both are more than 2.8 psu, but
1823	with significantly smaller bias in WOA. Overall, the large uncertainty found in the
1824	linear regression of all products is attributed to large product-observation mismatch
1825	against in-situ salinity data of less than 24 psu, which are observed over the
1826	continental shelf near the estuary of the Mackenzie River.
1827	In order to characterize the product-data misfits of all six products against in-situ
1828	data, a 4th order polynomial function is fitted to the absolute deviation as a function
1829	of observed salinity (Fig. <u>12</u>). The absolute deviations of most of the products except
1830	for <u>MOB</u> monotonically decrease as observed salinity increases. The norm residuals
1831	for <u>TP4</u> and <u>BEC</u> and are the smallest of 10.2 and 7.0, respectively, among all six
1832	products and the fitted curves give certain confidence in estimating the size of the
1833	error in the each SSS product. The fitted curve reaches its smallest value of less than
1834	1.0 psu at the in-situ salinity of 28 psu and 30 psu for BEC and TP4 respectively.
1835	Both the fitted curves for CEC and MOB have large norm residuals of 18.1 and 68.8
1836	psu ² respectively. Note that special attention must be paid in <u>usage of MOB</u> in the
	· · · · · · · · · · · · · · · · · · ·

Formatted	[88]
Deleted: along the south-west Norwe	egian coast and [89]
Deleted: Highlighting in the Beaufort	Sea, there are 193
Formatted	[91]
Deleted: 2.63	
Formatted	[92]
Deleted: 65	[22]
Formatted	[02]
Deleted: 9	[93]
Formatted	[04]
Deleted: 71	[94]
Formatted	[05]
	[93]
Formatted	
Deleted: 95	[96]
Deleted: 65	
Formatted	[97]
Deleted: a	
Formatted	[98]
Deleted: 73	
Formatted	[99]
Deleted: , but they	
Formatted	[100]
Deleted: obviously	
Formatted	[101]
Deleted: the corresponding misfits of	the MOI
Formatted	[102]
Deleted: 7.18	
Formatted	[103]
Deleted: with	
Formatted	[104]
Deleted: of -4.3	
Formatted	[105]
Deleted: .	
Formatted	[106]
Deleted: climatologies	L
Formatted	[107]
Deleted: the	[10/]
Formatted	[108]
Deleted: have BMSD	[108]
Formatted	[100]
Deleted: 3	[109]
Formatted	F1101
Polotod: small	[110]
Formetted	
Public di the	[111]
Deleted: the	
Pormatted	[112]
Deleted: a	
rormatted	[113]
Deleted: for	
Formatted	[114]
Deleted:	
Formatted	[115]
Deleted: , the absolute deviationso	f all six products 161

1948	Arctic Ocean due to its large negative bias and the RMSD in regions where the
1949	product is based on limited number of observations.
1950	Evaluations of the SSS products against TP4 product and in situ data conducted
1951	above suggest certain benefit can be expected in assimilating one of the SMOS
1952	salinity products, the BEC SSS, into the TOPAZ Arctic ocean analysis-forecast
1953	system. The knowledge of error structure in the SSS products provided in this study
1954	will assist to reasonably estimate the observation error for the SMOS product, which
1955	is required by a data assimilation system. We recommend that due to the poor spatial
1956	coverages of CORA in situ data in the Arctic Ocean, more data - especially from the
1957	Arctic Ocean marginal seas - should be compiled from independent data source for
1958	validating the SMOS SSS products. In addition, when comparing the two climatology
1959	products, PHC and WOA, the SSS scatterplots of the PHC in the central Arctic (Fig.
1960	7) show salinity bias for Jow saline water. Considering the different time periods of
1961	their compiled in-situ data sources (Table 1), it independently verifies that the
1962	freshening in the Canada Basin since 1990s is rather significant as discussed by
1963	Morison et al. (2012). Based on this evaluation, the next TOPAZ system will use the
1964	WOA to replace the PHC as the target relaxation field.
1965	
1966	Acknowledgement
1967	The authors acknowledge the support of CMEMS for the Arctic MFC, and partly
1968	funding from the European Space Agency in response to the Arctic+Salinity project
1969	(Dec 2018-June 2020). Grants of computing time (nn2993k and nn9481k) and
1970	storage (ns2993k) from the Norwegian Sigma2 infrastructures are gratefully
1971	acknowledged. The BEC SSS is produced by the Barcelona Expert Centre
1972	(bec.icm.csic.es) mainly funded by the Spanish National Program on Space. The
1973	CEC SSS is distributed by the Ocean Salinity Expertise Center (CECOS) of CATDS
1974	at IFREMER, France.

1975

1976 **Reference:**

1977	Bertino, L., and Lisæter, K. A.: The TOPAZ monitoring and prediction system for the Atlantic
1978	and Arctic Oceans, Journal of Operational Oceanography, 1(2), 15–19, doi:
1979	10.1080/1755876X.2008.11020098, 2008
1980	Boutin J., Vergely J.L., Marchand S., D'Amico F., Hasson A., Kolodziejczyk Nicolas, Reul

1981 Nicolas, Reverdin G., Vialard J.: New SMOS Sea Surface Salinity with reduced systematic

Formatted: Header

Deleted: RMSD, although its smallest misfits against CORA data in the northern Atlantic oceans among others
Deleted: Validation
Formatted: Font color: Text 1
Formatted: Font color: Text 1
Deleted: product like
Formatted: Font color: Text 1
Formatted: Font color: Text 1
Deleted: on
Formatted: Font color: Text 1
Deleted: earned
Formatted: Font color: Text 1
Deleted: help us
Formatted: Font color: Text 1
Formatted: Font color: Text 1
Deleted: Due
Formatted: Font color: Text 1
Deleted: the
Formatted: Font color: Text 1
Formatted: Font color: Text 1
Formatted: Font color: Text 1
Deleted: The newest SMOS product (Olmedo et al., 2018) that covers the years of 2010-2017 became available recently. Validation of the SMOS SSS product
Deleted: the longer period together with the extended in situ data is under preparation now
Deleted: step.
Deleted:

Deleted: www.smos-Deleted: .)

Deleted: Antonov, J., Seidov, D., Boyer, T., Locarnini, R., Mishonov, A., and Garcia, H.: World Ocean Atlas 2009, Volume 2: Salinity. NOAA Atlas NESDIS 69, NOAA, U.S. Government Printing Office, Washington D.C., 2010.¶

I	4	Formatted: Header
I		
2006	errors and improved variability. Remote Sensing of Environment, 214, 115-134,	
2007	<u>http://doi.org/10.1016/j.rse.2018.05.022,</u> 2018.	Field Code Changed
2008	Buongiorno Nardelli, B., Droghei, R., and Santoleri, R.: Multi-dimensional interpolation of	
2009	SMOS sea surface salinity with surface temperature and in situ salinity data. Remote	
2010	Sens. Environ. 180, 392–402. doi: 10.1016/j.rse.2015.12.052, 2016	
2011	Cabanes, C., A. Grouazel, K. von Schuckmann, M. Hamon, V. Turpin, C. Coatanoan, F. Paris,	
2012	S. Guinehut, C. Boone, N. Ferry, C. de Boyer Montégut, T. Carval, G. Reverdin, S.	
2013	Pouliquen, and P. Y. Le Traon: The CORA dataset: validation and diagnostics of in-situ	
2014	ocean temperature and salinity measurements. Ocean Science, 9, 1-	
2015	18, <u>http://www.ocean-sci.net/9/1/2013/os-9-1-2013.html</u> , doi:10.5194/os-9-1-2013, 2013	Field Code Changed
2016	Chassignet, E. P., Smith, L. T., and Halliwell, G. R.: North Atlantic Simulations with the Hybrid	
2017	Coordinate Ocean Model (HYCOM): Impact of the vertical coordinate choice, reference	
2018	pressure, and thermobaricity, J. Phys. Oceanogr., 33, 2504-2526. Doi:	
2019	http://dx.doi.org/10.1175/1520-0485(2003)033<2504:NASWTH>2.0.CO:2, 2003.	Field Code Changed
2020	de Boyer Montegut, C., G. Madec, A. Fischer, A. Lazar, and D. ludicone: Mixed Layer Depth	
2021	over the Global Ocean: An Examination of Profile Data and a Profile-Based Climatology.	
2022	J. Geophys. Res., 109 (C12003), 1–20, doi:10.1029/2004JC002378, 2004.	
2023	D'Addezio, J. M., and Subrahmanyam, B.: Sea surface salinity variability in the Agulhas	
2024	Current region inferred from SMOS and Aquarius. Remote Sensing of Environment, 180,	
2025	440–452. doi:10.1016/j.rse.2016.02.006, 2016	
2026	Drange, H. and Simonsen, K.: Formulation of air-sea fluxes in the ESOP2 version of MICOM,	
2027	Technical Report No. 125 of Nansen Environmental and Remote Sensing Center, 1996.	
2028	Droghei, R., Buongiorno Nardelli, B., and Santoleri, R.: A new global sea surface salinity and	
2029	density dataset from multivariate observations (1993-2016). Front. Mar. Sci. 5:84. doi:	
2030	10.3389/fmars.2018.00084, 2018	
2031	ESA, SMOS data products, available from	
2032	https://earth.esa.int/documents/10174/1854456/SMOS-Data-Products-Brochure (last	Field Code Changed
2033	access on 12 th December 2018), November 2017.	
2034	Font, J., Camps, A., Borges, A., Martín-Neira, M., Boutin, J., Reul, N., Kerr, Y. H., Hahne, A.,	
2035	and Mecklenburg, S.: SMOS: The challenging sea surface salinity measurement from	
2036	space, Proc. IEEE, 98(5), 649–665, DOI: <u>10.1109/JPROC.2009.2033096</u> , May 2010.	Field Code Changed
2037	Furue, R., Takatama, K., Sasaki, H., Schneider, N., Nonaka, M., and Taguchi, B.: Impacts of	
2038	sea-surface salinity in an eddy-resolving semi-global OGCM. Ocean Modelling, 122, 36-	
2039	56, doi:10.1016/j.ocemod.2017.11.004, 2018	Deleted: .
2040	Hátún, H., Sandø, A. B., Drange, H., Hansen, B., and Valdimarsson, H.: Influence of the	
2041	Atlantic Subpolar Gyre on the Thermohaline Circulation, Science, 309 (5742), 1841-1844,	
2042	doi:10.1126/science.1114777, 2005	
I		

Formatted: Header 2044 Hunke, E. C., and Dukowicz, J. K.: An elastic-viscous-plastic model for sea ice dynamics, J. 2045 27. Phys. Oceanogr.. 1849-1867 https://doi.org/10.1175/1520-Field Code Changed 2046 0485(1997)027<1849:AEVPMF>2.0.CO;2, 1997. 2047 Johnson, G. C., Schmidtko, S., and Lyman, J. M.: Relative contributions of temperature and 2048 salinity to seasonal mixed layer density changes and horizontal density gradients, J. 2049 Geophys. Res., 117, C04015, doi:10.1029/2011JC007651, 2012 Field Code Changed 2050 Kerr, Y. H., Waldteufel, P., Wigneron, J. P., Delwart, S., Cabot, F., Boutin, J., Escorihuela, M. 2051 J., Font, J., Reul, N., Gruhier, C., Juglea, S., Drinkwater, M. R., Hahne, A., Martín-Neira, 2052 M., and Mecklenburg, S.: The SMOS mission: New tool for monitoring key elements of the global water cycle, Proc. IEEE, 98(5), 666-687, doi:10.1109/JPROC.2010.2043032, 2010. 2053 Formatted: Font: 11 pt Field Code Changed 2054 Kolodziejczyk, N., Boutin, J., Vergely, J.-L., Marchand, S., Martin, N., and Reverdin, G.: Formatted: Font: 11 pt, English (US) 2055 Mitigation of systematic errors in SMOS sea surface salinity. Remote Sensing of Formatted: Font: 11 pt, English (US) 2056 Environment, 180, 164–177. doi:10.1016/j.rse.2016.02.061, 2016 2057 Latif, M., Roeckner, E., Mikolajewicz, U., and Voss R.: Tropical stabilization of the thermohaline 2058 circulation in a greenhouse warming simulation, J. Clim., 13: 1809-1813, 2000. 2059 Macdonald, R. W., Carmack, E. C., McLaughlin, F. A., Falkner, K. K., and Swift, J. H.: 2060 Connections among ice, runoff and atmospheric forcing in the Beaufort Gyre. Geophys. Res. Lett., 26, 2223-2226, 1999 2061 2062 Maes, C., Ando, K., Delcroix, T., Kessler, W. S., McPhaden, M. J., and Roemmich, 2063 D.: Observed correlation of surface salinity, temperature and barrier layer at the eastern 2064 edge of the western Pacific warm pool, Geophys. Res. Lett., 33, L06601, 2065 doi:10.1029/2005GL024772, 2006. Field Code Changed 2066 Mathis, J. T., and Monacci, N. M.: Carbon Dioxide and Hydrographic data obtained during the 2067 USCGC Healy Cruise HLY1203 in the Arctic Ocean (October 05 - 25, 2012). Available 2068 from http://cdiac.ess-dive.lbl.gov/ftp/oceans/CARINA/Healy/HLY-12-03/. Oak Ridge 2069 National Laboratory, US Department of Energy, Oak Ridge, Tennessee. doi: 2070 10.3334/CDIAC/OTG.CLIVAR 33HQ20121005, 2014. 2071 McPhee, M. G., Stanton, T. P., Morison, J. H. and Martinson, D. G.: Freshening of the upper 2072 ocean in the Arctic: is perennial sea ice disappearing? Geophys. Res. Lett. 25, 1729-1732, 2073 1998. 2074 Mecklenburg, S., Drusch, M., Kerr, Y. H., Font, J., Martín-Neira, M., Delwart, S., Buenadicha, 2075 G., Reul, N., Daganzo-Eusebio, E., Oliva, R., and Crapolicchio, R.: ESA's soil moisture 2076 and ocean salinity mission: Mission performance and operations. IEEE TGARS, 50(5), 2077 1354-1366, DOI: 10.1109/TGRS.2012.2187666, 2012 Field Code Changed 2078 Mignot, J., and Frankignoul, C.: On the interannual variability of surface salinity in the Atlantic. 2079 Climate Dynamics, 20(6), 555-565. doi:10.1007/s00382-002-0294-0, 2003

19

Formatted: Header 2080 Morison, J., Kwok, R., Peralta-Ferriz, C., Alkire, M., Rigor, I., Andersen, R., and Steele, M.: Formatted: Font: 11 pt 2081 Changing arctic ocean freshwater pathways. Nature, 481:66-70, 2012. 2082 Olmedo, E., Gabarró, C., González-Gambau, V., Martínez, J., Ballabrera-Poy, J., Turiel, A., Deleted: Martínez, J., Umbert, 2083 Portabella, M., Fournier, S., and Lee, T.: Seven Years of SMOS Sea Surface Salinity at Moved down [5]: M., Formatted: Font: 11 pt, Font color: Auto 2084 High Latitudes: Variability in Arctic and Sub-Arctic Regions. Remote Sensing. 2018; Formatted: Font: 11 pt, Font color: Auto 2085 10(11):1772, https://doi.org/10.3390/rs10111772, 2018. Deleted: Hoareau, N., Portabella, M., Ballabrera-Poy, 2086 Reverdin, G., Cayan, D., and Kushnir, Y.: Decadal variability of hydrography in the upper , and Turiel, A .: Improving time and space resolution of SMOS salinity maps using multifractal fusion. 2087 northern North Atlantic in 1948-1990. J. Geophys. Res., 102, 8505-8531, Remote Sens. Environ. 180, 246-263, https://doi.org/10.1016/j.rse.2016.02.038, 2016.¶ 2088 https://doi.org/10.1029/96JC03943, 1997 Olmedo, E., Formatted: English (US) 2089 Sakov, P., and Oke, P. R.: A deterministic formulation of the ensemble Kalman Filter: an Deleted: Oki, T. and Sud, Y. C.: Design of Total Runoff 2090 alternative to ensemble square root filters. Tellus A, 60(2), 361-371, doi:10.1111/j.1600-Integrating Pathways (TRIP)-A global river channel network, Earth Interact., 2, 1–37, doi:10.1175/1087-2091 0870.2007.00299.x, 2008. 3562(1998)002<0001:DOTRIP>2.3.CO;2, 1998. 2092 Simmons, A., Uppala, S., Dee, D., and Kobayashi, S.: ERA-Interim: New ECMWF reanalysis 2093 products from 1989 onwards. ECMWF Newsletter. 110. 25-35, doi:10.21957/pocnex23c6, 2094 2007. 2095 SMOS Team, SMOS L2 OS Algorithm Theoretical Baseline Document, ESA, Paris, France, 2096 SO-TN-ARG-GS-0007. version available 3.13. from 2097 https://earth.esa.int/documents/10174/1854519/SMOS L2OS-ATBD (last access: 12th Field Code Changed 2098 December 2018), 29th Arpil 2016. 2099 SMOS-BEC Team, Quality Report: Validation of SMOS-BEC experimental sea surface salinity 2100 products in the Arctic Ocean and high latitudes Oceans. Years 2011-2013, Barcelona 2101 Expert Centre, Spain. Technical note: BEC-SMOS-0007-QR version 1.0, available at 2102 http://bec.icm.csic.es/doc/BEC-SMOS-0007-QR.pdf (last access: 13th December 2018), Field Code Changed 2103 4th March 2016. 2104 Steele, M. and W. Ermold (2004) Salinity Trends on the East Siberian Shelves, Geophysical 2105 Research Letters, Vol. 31, L24308, doi:10.1029/2004GL021302, 2004. Steele, M., R. Morley, and W. Ermold, PHC: A global ocean hydrography with a high-quality 2106 2107 Arctic Ocean, Journal of Climate, 14, 2079-2087, 2001. 2108 Sumner, D., and Belaineh, G.: Evaporation, Precipitation, and Associated Salinity Changes at 2109 a Humid, Subtropical Estuary. Estuaries, 28(6), 844-855. Retrieved from 2110 http://www.jstor.org/stable/3526951, 2005. Field Code Changed 2111 Supply A, Boutin J, Vergely J-L, et al.: Precipitation Estimates from SMOS Sea-Surface 2112 Salinity. Q. J. R. Meteorol. Soc., 144 (Suppl. 1):103-119, https://doi.org/10.1002/qj.3110, Field Code Changed 2113 2018. 2114 Talley, L. D., Johnson, G. C., Purkey, S., Feely, R. A., and Wanninkhof, R.: Global Ocean 2115 Ship-based Hydrographic Investigations Program (GO-SHIP) provides key climate-2116 relevant deep ocean observations, US CLIVAR Variations, 15 (2), available from

Formatted: Header

I

2129 https://www.pmel.noaa.gov/pubs/PDF/tall4659/tall4659.pdf (last access: 19th December 2130 2018), 2017. 2131 Toole, J. M., Krishfield, R. A., Timmermans, M. J., and Proshutinsky, A.: The Ice-Tethered Moved (insertion) [5] Moved (insertion) [6] 2132 Profiler: Argo of the Arctic, Oceanography, 24, 126-135, doi:10.5670/oceanog.2011.64, Formatted: Font: 11 pt, Font color: Auto 2133 2011. 2134 Tseng, Y., Bryan, F. O., and Whitney, M. M.: Impacts of the representation of riverine 2135 freshwater input in the community earth system model. Ocean Modelling, 105, 71-86. 2136 doi:10.1016/j.ocemod.2016.08.002, 2016. 2137 Uotila, P., Goosse, H., Haines, K., Chevallier, M., Barthélemy, A., Bricaud, C., Carton, J., 2138 Fu'ckar, N., Garric, G., Iovino, D., Kauker, F., Korhonen, M., Lien, V. S., Marnela, M., 2139 Massonnet, F., Mignac, D., Peterson, A., Sadikn, R., Shi, L., Tietsche, S., Toyoda, T., Xie, 2140 J., and Zhang, Z.: An assessment of ten ocean reanalyses in the polar regions, Clim. Dynam.,https://doi.org/10.1007/s00382-018-4242-z, online first, 2018. 2141 2142 Vancoppenolle, M., Fichefet, T., and Goosse, H.: Simulating the mass balance and salinity of 2143 Arctic and Antarctic sea ice. 2. Importance of sea ice salinity variations. Ocean Modelling, 2144 27, 54-69. doi:10.1016/j.ocemod.2008.11.003, 2009. 2145 Verbrugge, N., Mulet, S., Guinehut, S., Buongiorno Nardelli, B., and Droghei, R.: Quality 2146 information document for global ocean multi observation products 2147 multiobs_glo_phy_rep_015_002, CMEMS-MOB-QUID-015-002, v1.0, available from http://cmems-resources.cls.fr/documents/QUID/CMEMS-MOB-QUID-015-002.pdf 2148 (last Field Code Changed

2149 access: 14th December 2018), February 2018.

2150 Vialard, J., and P. Delecluse (1998), An OGCM study for the TOGA decade: I. Role of salinity
2151 in the physics of the Western Pacific fresh pool, J. Phys. Oceanogr., 28, 1071–1088,
2152 Woodgate, R., Aagaard, K. and Weingartner, T.: Monthly temperature, salinity, and
2153 transport variability of the Bering Strait through flow. *Geophys. Res. Lett.*, 32, L04601,
2154 DOI: 10.1029/2004GL021880, 2005.

- Xie, J., Bertino, L., Counillon, F., Lisæter, K. A., and Sakov, P.: Quality assessment of the
 TOPAZ4 reanalysis in the Arctic over the period 1991–2013. *Ocean Science*, 13(1), 123–
 144. <u>http://doi.org/10.5194/os-13-123-2017</u>, 2017.
- Xie, J., Counillon, F., and Bertino, L.: Impact of assimilating a merged sea-ice thickness from
 CryoSat-2 and SMOS in the Arctic reanalysis, The Cryosphere, 12, 3671-3691,
 https://doi.org/10.5194/tc-12-3671-2018, 2018.
- Xie, J., Counillon, F., Bertino, L., Tian-Kunze, X., and Kaleschke, L.: Benefits of assimilating
 thin sea-ice thickness from SMOS into the TOPAZ system. *The Cryosphere*, 10, 2745–
 2761. <u>http://doi.org/10.5194/tc-10-2745-2016</u>, 2016.

Moved up [6]: L., Deleted: Vergely, J. -Deleted: and Boutin, J.: SMOS OS Level 3: The Algorithm Theoretical Basis Document (v300), available on. http://www.catds.fr/content/download/78841/1005020/f ile/ATBD L3OS v3.0.pdf?version=3. (last access: 14th December 2018), 5th May 2017.¶ Formatted: Normal, Justified, Indent: Left: 0 cm, Hanging: 0,75 cm Formatted: English (US)

I		 Formatted: Header
2172	Yu, L.: A global relationship between the ocean water cycle and near-surface salinity, J.	
2173	Geophys. Res., 116, C10025, doi:10.1029/2010JC006937, 2011.	
2174	Zweng, M. M., Reagan, J. R., Antonov J. I., Locarnini, R. A., Mishonov, A. V., Boyer, T. P.,	 Formatted: Font: 11 pt
2175	Garcia, H. E., Baranova, O. K., Johnson, D. R., Seidov, D., and Biddle, M. M.: World	
2176	Ocean Atlas 2013, Volume 2: Salinity, Levitus, S. (Ed.), Mishonov, A., Technical Ed.	
2177	NOAA Atlas NESDIS 74, 39pp, 2013.	 Formatted: Font: 11 pt

Page 1: [1] Formatted	Author	6/15/19 3:21:00 AM
Header		
Page 2: [2] Deleted	Author	6/15/19 3:21:00 AM
x		
Page 2: [3] Formatted	Author	6/15/19 3:21:00 AM
Page 2: [4] Deleted	Author	6/15/19 3:21:00 AM
▼		
A Page 2: [4] Deleted	Author	6/15/19 3:21:00 AM
		0,10,17,0121001101
v		
Page 2: [5] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [6] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [7] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [7] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [8] Formatted	Author	6/15/19 3:21:00 AM
Pattern: Clear		
Page 2: [9] Deleted	Author	6/15/19 3:21:00 AM
x		
Page 2: [10] Formatted	Author	6/15/19 3:21:00 AM
Font color: 1 ext 1		
Page 2: [11] Formatted	Author	6/15/19 3:21:00 AM
Font color: 1 ext 1		
Page 2: [12] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [13] Deleted	Author	6/15/19 3:21:00 AM
v		
A Page 2. [14] Formatted	Author	6/15/19 3·21·00 AM
Font color: Text 1	Autio	0/15/17 5.21.00 Alvi
Page 2: [14] Formatted	Authou	6/15/10 2.21.00 ANA
Font color: Text 1	Author	0/15/19 5:21:00 ANI
Page 2: [14] Formatted	Author	6/15/19 3:21:00 AM

Font color: Text 1		
Page 2: [15] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [16] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [17] Deleted	Author	6/15/19 3:21:00 AM
۲		
Fage 2: [18] Formatted	Author	6/15/19 3:21:00 AM
Page 2: [19] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 2: [20] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [21] Deleted	Author	6/15/19 3:21:00 AM
v		
<u>.</u>		1
Page 2: [22] Formatted	Author	6/15/19 3:21:00 AM
Font color: 1 ext 1		
Page 2: [23] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [23] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [24] Deleted	Author	6/15/19 3:21:00 AM
۲		
Page 2: [25] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [26] Deleted	Author	6/15/19 3:21:00 AM
۲		
Page 2: [27] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [28] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [29] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [30] Deleted	Author	6/15/19 3:21:00 AM

Page 2: [31] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 2: [32] Deleted	Author	6/15/19 3:21:00 AM
¥		
A		
Page 2: [33] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text I		
Page 2: [34] Formatted	Author	6/15/19 3:21:00 AM
Font: Bold		
Page 2: [35] Formatted	Author	6/15/19 3:21:00 AM
Justified, Don't adjust right inde between Latin and Asian text, D	ent when grid is defined, Line spaci Don't adjust space between Asian tex	ng: single, Don't adjust space tt and numbers
Page 2: [36] Deleted	Author	6/15/19 3:21:00 AM
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
¥.		
<u> </u>		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
V		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
v		
A		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
•		
۰		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
۷		
Page 1: [37] Delated	Author	6/15/10 3.21.00 AM
age 4: [5/] Deleted	Author	0/15/19 5:21:00 AM
۷		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
۷		
A		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
۷		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
Υ		

T

<u> </u>		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 4: [37] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 4: [38] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 4: [39] Deleted	Author	6/15/19 3:21:00 AM
v		
_		
Page 4: [39] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 4: [39] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 4: [40] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 4: [41] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 4: [41] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 4: [42] Delated	Author	6/15/19 3·21·00 AM
1 age 7. [72] Delettu	Autio	0/13/17 3.21.00 AM
Χ		
A Page 4: [43] Deleted	Author	6/15/19 3·21·00 AM
1 age 7. [75] Delettu	Autio	0/13/17 3.21.00 AM
v		
A	Author	6/15/19 3·21·00 AM
1 age 7. [75] Delettu	Autio	0/13/17 3.21.00 AM
v		
Page 4: [43] Deloted	Author	6/15/10 3·21·00 AM
	Autioi	0/15/17 5.21.00 AW
▼		
Page 4: [43] Delated	Author	6/15/10 2.21.00 AM
1 age 4: [45] Deleteu	Autilor	0/15/19 3:21:00 AM
v		
A		

Page 4: [43] Deleted	Author	6/15/19 3:21:00 AM
Υ		
A Page 4: [44] Deleted	Author	6/15/19 3:21:00 AM
۲		
Page 4: [44] Deleted	Author	6/15/19 3:21:00 AM
v		
A Page 4: [44] Deleted	Author	6/15/19 3:21:00 AM
V		
A Dago 4. [45] Deloted	Author	(15/10 2-21-00 AM
-	Author	0/15/19 5:21:00 AM
A		
Page 4: [45] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 4: [46] Deleted	Author	6/15/19 3:21:00 AM
Υ		
Page 4: [46] Deleted	Author	6/15/19 3:21:00 AM
۲		
A Page 4: [46] Deleted	Author	6/15/19 3:21:00 AM
۲		
A Page 4: [46] Deleted	Author	6/15/19 3·21·00 AM
Page 4: [46] Deleted	Author	6/15/19 3:21:00 AM
V		
Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
٧		
A Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
۷		
A Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
		0,10,17 012100 1111

Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
v		
A		
Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
v		
<u>.</u>		
Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
Υ		
Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
•		
▲		
Page 5: [47] Deleted	Author	6/15/19 3:21:00 AM
_		
×		
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
v		
A Page 5: [48] Deleted	Author	6/15/19 3·21·00 AM
Tage 5. [16] Detetu		0/15/19 0.21.00 / 11/1
Ψ		
A Page 5: [48] Deleted	Author	6/15/10 3·21·00 AM
Tage 5. [46] Deleteu	Aution	0/13/19 5.21.00 AM
▼		
A Daga 5: [48] Delated	Authou	6/15/10 2.21.00 AM
rage 5: [46] Deleted	Author	0/15/19 5:21:00 AM
v		
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
▼		
A		1
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
•		

Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
Υ		
A Page 5: [48] Deleted	Author	6/15/19 3·21·00 AM
1 age 3. [46] Deced		0/13/17 5.21.00 / 11/1
X		
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 5: [48] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
_		
×		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
_		
×		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
•		
· · · · · · · · · · · · · · · · · · ·		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
•		
^		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
v		
<u></u>		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
v		
	A	C/1E/10 2.31.00 AM
rage 5: [49] Deleted	Autnor	0/15/19 3:21:00 AM
v		
A Page 5: [40] Deleted	Author	6/15/10 2.21.00 AM
1 age 3. [47] Deleteu	Author	0/15/19 5:21:00 AM
v		

A		
Page 5: [49] Deleted	Author	6/15/19 3:21:00 AM
Page 5: [50] Deleted	Author	6/15/19 3:21:00 AM
,		
Page 5: [50] Deleted	Author	6/15/19 3:21:00 AM
r		
k		
Page 5: [50] Deleted	Author	6/15/19 3:21:00 AM
·		
Page 5: [50] Deleted	Author	6/15/19 3:21:00 AM
r		
Page 5: [51] Deleted	Author	6/15/19 3:21:00 AM
Page 5: [51] Deleted	Author	6/15/19 3:21:00 AM
Page 5: [51] Deleted	Author	6/15/19 3·21·00 AM
	Aumor	0/15/17 5.21.00 AM
Page 7: [52] Deleted	Author	6/15/19 3:21:00 AM
Page 7: [53] Deleted	Author	6/15/19 3:21:00 AM
r		
Page 7: [54] Deleted	Author	6/15/19 3:21:00 AM
,		
D 7. [55] D. 1.4]	A. (h	(115/10 2.21.00 AM
rage /: [55] Deleted	Autnor	6/15/19 3:21:00 AM
Dage 7. [54] Deleted	Author	6/15/10 2-21.00 AM
rage /: [50] Deleted	Author	0/15/19 5:21:00 AM
<i>.</i>		
Page 7. [57] Deleted	Author	6/15/19 3·21·00 AM
	Aumor	0/13/17 5.21.00 AM
ζ		
Page 7: [58] Deleted	Author	6/15/19 3·21·00 AM
r age / [50] Deleted		0/10/17 5.21.00 AM
ζ		
Page 7: [59] Deleted	Author	6/15/19 3·21·00 AM
age (1 [07] below		0/10/17 0.21.00 /11/1
Γ		

Page 8: [60] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 8: [61] Deleted	Author	6/15/19 3:21:00 AM
Υ		
Page 8: [62] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 8: [63] Deleted	Author	6/15/19 3:21:00 AM
Ψ		
A Page 8: [64] Deleted	Author	6/15/19 3·21·00 AM
Tage 0. [01] Deleteu	Author	0/15/19 0.21.00 AM
X		
Page 9: [65] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 9: [65] Deleted	Author	6/15/19 3:21:00 AM
Υ		
Page 9: [65] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
Ψ		
A Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
V		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
Ψ		
A Page 9: [66] Deleted	Author	6/15/19 3·21·00 AM
- age > 1001 Deleter	110101	0/ 10/ 17 0-21.00 / 11/1

<u> </u>		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
▼		
<u> </u>		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
Υ		
<u> </u>		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 9: [66] Deleted	Author	6/15/19 3·21·00 AM
		0/13/17 0.21.00 / 10/
Υ		
A Dago (); [66] Dolotod	Andhau	C/12/10 2.01.00 ANA
rage 9: [66] Deleted	Autnor	0/15/19 5:21:00 AM
v		
<u> </u>		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
v		
_ _		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
•		
·		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
Υ		
A	Author	6/15/10 2.21.00 AM
1 age 7. [00] Deteteu	Autio	0/15/19 5.21:00 AIVI
v		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
▼		
<u> </u>		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
•		
τ		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
v		

Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
▼		
A Dags 0: [66] Delated	Author	(/15/10 2-21-00 AM
Page 9: [66] Deleted	Autnor	6/15/19 3:21:00 AM
v		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
τ		
Page 9: [66] Deleted	Author	6/15/19 3:21:00 AM
τ		
A		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
v		
A		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
▼		
<u> </u>		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
۷		
<u> </u>		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
x		
<u> </u>		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
×		

A		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
Υ		
Page 0: [67] Deleted	Author	6/15/10 3.21.00 AM
rage 9: [07] Deleted	Author	0/15/19 5:21:00 AM
v		
A		
Page 9: [67] Deleted	Author	6/15/19 3:21:00 AM
Page 9: [67] Deleted	Author	6/15/19 3·21·00 AM
Tage 7. [07] Delettu	Autio	0/13/17 5.21.00 AW
Υ		
<u> </u>		
Page 10: [68] Deleted	Author	6/15/19 3:21:00 AM
_		
τ		
Page 10: [69] Deleted	Author	6/15/19 3:21:00 AM
Tuge Ioi [07] Deleteu		
τ		
A		
Page 10: [70] Deleted	Author	6/15/19 3:21:00 AM
•		
¥		
Page 11: [71] Deleted	Author	6/15/19 3:21:00 AM
1 ngo 111 [71] 2010104		
Υ		•
•		
Page 13: [72] Deleted	Author	6/15/19 3:21:00 AM
•		
Page 13: [72] Deleted	Author	6/15/19 3:21:00 AM
8 1 1		
▼		
A		
Page 13: [72] Deleted	Author	6/15/19 3:21:00 AM
•		
A		
Page 13: [72] Deleted	Author	6/15/19 3:21:00 AM
<u> </u>		
▼		
Page 13: [72] Deleted	Author	6/15/19 3:21:00 AM
V		
<u> </u>		
Page 13: [72] Deleted	Author	6/15/19 3:21:00 AM
Υ		
		_
Page 13: 72 Deleted	Author	6/15/19 3:21:00 AM

▼		
A		
Page 13: [72] Deleted	Author	6/15/19 3:21:00 AM
▼		
A Dage 12: [72] Formatted	Author	6/15/10 3.21.00 AM
rage 13: [75] Formatted	Autior	0/13/19 5.21.00 AM
Font color: Auto		
Page 13: [73] Formatted	Author	6/15/19 3:21:00 AM
Font color: Auto		
Page 13: [74] Deleted	Author	6/15/19 3:21:00 AM
A		
Page 13: [74] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 13. [74] Deleted	Author	6/15/19 3·21·00 AM
Tage 13. [74] Delettu		0/15/17 5.21.00 / 11/1
Χ		
A		
Page 13: [74] Deleted	Author	6/15/19 3:21:00 AM
•		
A		
Page 13: [74] Deleted	Author	6/15/19 3:21:00 AM
Page 13. [74] Deleted	Author	6/15/19 3·21·00 AM
Tage 15. [74] Detetu	Author	0/15/17 5.21.00 / 11/1
▼		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
V		
A		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
_		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
٧		
<u> </u>		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
-		
τ		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
rage tor [/o] beleted		0/10/17 0/#1000/11/1

Dama 12. [75] Dalatad	A 61	(15/10.2.21.00 AM
age 15: [/5] Deleted	Autnor	0/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Dama 12. [75] Dalatad	A 61	(/15/10 2-21.00 AM
rage 15: [75] Deleted	Autnor	0/15/19 5:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM

Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
Χ		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
•		
<u> </u>		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
· · · · · · · · · · · · · · · · · · ·		
•		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
▼		
A Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
-		
×		
Page 13: [75] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 13: [75] Deleted	Author	6/15/10 3·21·00 AM
Tage 13. [75] Detted	Author	0/13/17 5.21.00 /10/
×		
Page 13: [76] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 12: [76] Delated	Andhon	(/15/10.2.21.00 AM
rage 15: [70] Deleted	Author	0/15/19 5:21:00 AIM
v		
Page 13: [76] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 13: [76] Deleted	Author	6/15/19 3:21:00 AM
▼		
Page 13: [76] Deleted	Author	6/15/19 3:21:00 AM
Χ		
<u> </u>		
Page 13: [76] Deleted	Author	6/15/19 3:21:00 AM
•		

A Description 12: 17(1) Deleted	A41	(15/10 2-21-00 AM
rage 13: [70] Deleted	Author	0/15/19 5:21:00 AM
Ψ		
А. Раде 13: [76] Deleted	Author	6/15/19 3:21:00 AM
V		
Page 13: [76] Deleted	Author	6/15/19 3:21:00 AM
-		
▲		
Page 13: [76] Deleted	Author	6/15/19 3:21:00 AM
٧		
<u> </u>		
Page 14: [77] Deleted	Author	6/15/19 3:21:00 AM
τ		
A Deges 14 [77] Delets d	A +++++	C/1E/10 2.21.00 AN
rage 14: [//] Deleted	Autnor	0/15/19 3:21:00 AM
Υ		
A Page 14: [77] Deleted	Author	6/15/19 3·21·00 AM
Tuge In [7] Deteteu		
v		
Page 14: [77] Deleted	Author	6/15/19 3:21:00 AM
▲		
Page 14: [77] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 14: [77] Deleted	Author	6/15/19 3:21:00 AM
Ψ		
A Dage 14: [77] Deleted	Author	6/15/10 2:21:00 AM
rage 14: [//] Deleted	Author	0/15/19 5:21:00 AM
τ		
Page 14: [77] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
Y		
<u> </u>		
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
•		

	A	(15/10 2 21 00 AM
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
▼		
A Page 14: [78] Deleted	Author	6/15/19 3·21·00 AM
1 age 14. [76] Deleted	Author	0/15/17 5.21.00 / 10/
Ψ		
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
V		
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
•		
<u> </u>		
Page 14: [78] Deleted	Author	6/15/19 3:21:00 AM
٧		
<u> </u>		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
▼		
A		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
τ		
	A 41	(15/10 2 21 00 AM
Page 14: [79] Deleted	Author	0/15/19 3:21:00 AM
▼		
A Page 14. [79] Deleted	Author	6/15/19 3·21·00 AM
1 age 14. [75] Deteted	Author	0/15/17 5.21.00 / 10/
v		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
Υ		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
•		
_		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
•		

	A	(15/10 2 21 00 AM
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
▼		
A Page 14: [79] Deleted	Author	6/15/19 3·21·00 AM
1 age 14. [75] Deteted	Author	0/15/17 5.21.00 / 11/1
Ψ		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
V		
Page 14: [79] Deleted	Author	6/15/19 3:21:00 AM
-		
×		
Page 14: [80] Deleted	Author	6/15/19 3:21:00 AM
•		
<u> </u>		
Page 14: [80] Deleted	Author	6/15/19 3:21:00 AM
٧		
<u> </u>		
Page 14: [80] Deleted	Author	6/15/19 3:21:00 AM
▼		
A		
Page 14: [80] Deleted	Author	6/15/19 3:21:00 AM
τ		
	A 41	(15/10 2 21 00 AM
Page 14: [80] Deleted	Author	0/15/19 3:21:00 AM
▼		
A Page 14. [80] Deleted	Author	6/15/19 3·21·00 AM
Tage 14. [00] Deleted	Author	0/10/19 0/21:00 11/1
v		
Page 14: [80] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 14: [80] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 14: [80] Deleted	Author	6/15/19 3:21:00 AM
•		
_		
Page 14: [80] Deleted	Author	6/15/19 3:21:00 AM
•		

A Dage 14, [91] Delated	Author	(15/10 2-21.00 AM
rage 14: [61] Deleted	Author	0/15/19 5:21:00 AM
v		
Page 14: [81] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 14: [81] Deleted	Author	6/15/19 3:21:00 AM
v		
<u>.</u>		
Page 14: [81] Deleted	Author	6/15/19 3:21:00 AM
τ		
A	A 41	(15/10 2.21.00 AM
Page 14: [81] Deleted	Autnor	6/15/19 3:21:00 AM
Υ		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
v		
A Dage 15: [92] Delated	Author	(15/10 2.21.00 AM
Page 15: [82] Deleted	Autnor	6/15/19 3:21:00 AM
Υ		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
×		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
τ		
<u> </u>		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
v		
A Dago 15: [92] Dolatod	Author	6/15/10 2.21.00 AN
r age 15; [62] Deleteu	Author	0/15/19 5:21:00 AM
v		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
-		
×		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
•		

A		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
v		
<u> </u>		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
▼		
<u> </u>		
Page 15: [82] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 15: [83] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 15: [83] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 15: [83] Deleted	Author	6/15/19 3:21:00 AM
▼		
A	A 4L	(15/10 2-01-00 AM
Page 15: [85] Deleted	Autnor	6/15/19 3:21:00 AM
X		
A Dage 15: [94] Formattad	Author	(/15/10 2-21-00 AM
Fage 15: [64] Formatted	Author	0/15/19 5:21:00 AM
Page 15: [84] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 15: [85] Deleted	Author	6/15/19 3:21:00 AM
_		
v		
Page 15: [85] Deleted	Author	6/15/19 3:21:00 AM
-		
×		
Page 15: [85] Deleted	Author	6/15/19 3:21:00 AM
-		
×		
Page 15: [85] Deleted	Author	6/15/19 3:21:00 AM
•		

Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
Authon	(/15/10.2.21.00 AM
Autnor	0/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
A	(15/10.2.21.00 ANA
Autnor	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
A. 0	
Author	0/15/19 3:21:00 ANI
Author	6/15/19 3:21:00 AM
Author	6/15/19 3:21:00 AM
	Author Author

Page 16: [90] Deleted	Author	6/15/19 3:21:00 AM
v		
	A4b	(15/10.2.21.00 AM
Page 16: [90] Deleted	Autnor	6/15/19 3:21:00 AM
v		
Page 16: [91] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [91] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [91] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [92] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [93] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [93] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [94] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [95] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [96] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [96] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [97] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [98] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [99] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [100] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [101] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [102] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		

Page 16: [103] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [104] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [105] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [106] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [107] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [107] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [108] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [109] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [110] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [110] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [111] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [112] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [113] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [114] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [115] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [115] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [115] Formatted	Author	6/15/19 3:21:00 AM
Font color: Text 1		
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
v		
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM

Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
Daga 16. [116] Dalatad	Author	6/15/10 3·21·00 AM
age 10. [110] Deleteu	Aution	0/15/17 5.21.00 AW
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
	A. (1	(15/10 2.21.00 AM
rage 16: [116] Deleted	Autnor	6/15/19 5:21:00 AM
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
		0/10/17/0121000/12/1
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM
Page 16: [116] Deleted	Author	6/15/19 3:21:00 AM

Captions of Table and Figures:

Tabl	e 1. <u>Details o</u>	<u>f the six produ</u>	ucts evaluated during	<u>2011-2013.</u>	
<u>Product</u>	<u>Data</u> source	<u>Resolution</u>	<u>Provider</u>	Website or CMEMS id	<u>Releas</u> <u>year</u>
<u>BEC</u>	<u>SMOS</u>	<u>9 days; 25</u> <u>km</u>	Barcelona Expert Centre, Spain	http://bec.icm.csie.es	<u>2018</u>
<u>CEC</u>	<u>SMOS</u>	<u>9 days; 25</u> <u>km zonal</u>	<u>Ocean Salinity</u> Expertise Center, IFREMER	FTP: ftp.ifremer.fr	<u>2018</u>
<u>TP4</u>	<u>Reanalysis</u>	<u>Daily; 12~16</u> <u>km</u>	CMEMS	ARCTIC-REANALYSIS- PHYS-002-003	<u>2015</u>
<u>MOB</u>	<u>In situ +</u> <u>SMOS</u>	<u>7 days;</u> <u>1/4x1/4°;</u>	CMEMS	MULTIOBS GLO PHY REP _015_002	<u>2016</u>
<u>PHC</u>	<u>In situ</u> (1950-1994)	Monthly: <u>1x1°</u>	Polar Science Center, <u>University of</u> <u>Washington</u>	http://psc.apl.washington.edu	<u>2005</u>
<u>WOA</u>	<u>In situ</u> (1955~2012)	<u>Monthly;</u> <u>1/4x1/4°</u>	NODC, NOAA	https://www.nodc.noaa.gov/O <u>C5/woa13/</u>	<u>2013</u>

 Table 2. Misfits of SSS relative to in-situ CORA5.1 observations during 2011-2013 in each sub-region. Bold numbers denote the smallest error among the six products.

 Sub-region. Bold
 Bias (psu)
 RMSD (psu)

A												
Region	BEC	CEC	TP4	MOB	PHC	WOA	BEC	CEC	TP4	MOB	PHC	WOA <
S0	-	-	.48	52	<u>,48</u>	.11	-	-	1,25	1.78	1.28	,70
S1	4.03	3.18	<u>3.29</u>	<u>1.63</u>	<u>3.29</u>	42	4. <u>23</u>	3. <u>70</u>	<u>3.47</u>	2.22	3. <u>43</u>	<u>1.37</u>
S2	-1 <u>.76</u>	44	<u>97</u>	2.96	<u>-3.30</u>	- <u>2.93</u>	2.16	2. <u>57</u>	1.70	3. <u>68</u>	3.87	3.62
S3	<u>14</u>	70	.14	- <u>21</u>	<u>29</u>	- <u>25</u>	.45	1 <u>.17</u>	.34	. <u>42</u>	<u>.51</u>	.44
S4	<u>09</u>	- <u>,20</u>	<u>,12</u>	<u>,11</u>	<u>02</u>	.02	<u>.91</u>	1. <u>21</u>	<u>.89</u>	<u>_86</u>	<u>.94</u>	.84
S5	<u>07</u>	<u>.06</u>	.20	.01	.02	<u>.07</u>	1 <u>,47</u>	1, <u>52</u>	1, <u>42</u>	1, <u>44</u>	1. <u>39</u>	1.30
S6	01	<u>15</u>	.01	01	<u>09</u>	. <u>05</u>	<u>,25</u>	.66	. <u>14</u>	12	<u>,28</u>	.16 🖪
S7	.05	_34	_04	03	-23	03	.31	_88	.33	.22	_43	.27

Table 3. Optimal coefficients for the 4th order polynomial fit of the errors (see Eq. 3) as a function of in-situ SSS for each product.

		Residual	🚽 n situ 🖪				
Product	p1(x10-3)	p ₂	p ₃	p4	p ₅	<u>norm</u>	<u>sample</u>
BEC	0. <u>168</u>	-0. <u>016</u>	0, <u>614</u>	-11.345	<u>87.097</u>	7.03	<u>91</u>
CEC	0. <u>225</u>	<u>-0.033</u>	-1, <u>550</u>	<u>-29.886</u>	<u>205.179</u>	<u>18.13</u>	121
TP4	0.993	-0, <u>096</u>	<u>3.430</u>	<u>54.552</u>	<u>335.197</u>	<u>10.17</u>	<u>232</u>
MOB	-1. <u>080</u>	0.128	-5 <u>,469</u>	<u>99.824</u>	<u>-645.087</u>	<u>68.81</u>	<u>163</u>
PHC	<u>1.257</u>	-0. <u>120</u>	<u>4.235</u>	<u>-65.938</u>	<u>388.808</u>	<u>13.98</u>	<u>232</u>
WOA	-0, <u>121</u>	0, <u>010</u>	-0, <u>322</u>	<u>3.998</u>	<u>-10.847</u>	<u>38.91</u>	<u>232</u>

Formatted	[1]
Moved (insertion) [1]	
Formatted	[2]
Deleted: then-situ CORA5.1 observations from	[3]
Deleted: ¶	
Formatted	[4]
Formatted Table	[5]
Deleted: MOI	
Deleted: MOI	
Formatted	[6]
Moved down [2]:97	[0]
Formatted	[7]
Deleted: 44	[/]
Formatted	гот
Deleted: -1 12	[8]
Deletadi 28	
Deleted: 50	
Deleted: .52	
Formatted	[9]
Deleted: 64	
Formatted	[11]
Deleted: 43	
Deleted: 1.42	
Formatted	[10]
Formatted	[12]
Deleted: 95	
Deleted: 78	
Deleted: 4.11	
Formatted	[14]
Deleted: 4.39	
Deleted: 78	
(Moved (insertion) [3]	
Deleted: .56	
Deleted: 2.80	
Formatted	[12]
Deleted: 2.80	[13]
Deleted: -2	
Deleted: 75	
Deleted: ./5	
Deleted:17	
Formatted	[15]
Deleted: 2.04	
Formatted	[16]
Deleted: 42	
Deleted: .65	
Formatted	[17]
Moved (insertion) [2]	
Deleted: .70	
Deleted:74	[18]
Deleted: 1.77	
Moved up [3]: 1.37	
Formatted	[19]
Deleted: 1.81	
Formatted	. [20]
Deleted: 15	[20]
Deleted: 2.41	
Formatted	10.11
Deletede 97	[21]
Deleted: 3/	
Deleted: 2.90	
Formatted	[22]
Deleted: 2.71	
Deleted: .05	
Formatted	[23]

Deleted: 15



Fig. 1 (a): SSS locations of the in-situ observations north of 52°N in CORA5.1 during the years 2011-2013. <u>8 sub-regions divide the</u> Arctic Ocean, <u>with</u> the number of observations <u>indicated</u> in each region, (b): <u>Independent</u> SSS observations in the Beaufort Sea during the summer months of 2011-2013, from the BGEP (marked by anti-triangles, squares, and starts) and the CLIVAR (marked by triangles and crosses). <u>Different colors (red, black</u> and yellow) <u>indicate</u> the <u>years (2011, 2012 and 2013 resp.)</u>.



Fig. 2 Monthly SSS (unit: psu) in March from satellite products (BEC and CEC, *left column*), <u>reanalysis/reprocessing</u> (TP4 and <u>MOB</u>, *middle column*), and climatology (PHC and WOA, *right column*). White areas are masked by sea ice. The thick brown line represents the sea ice edge (15% concentration from <u>TP4</u>), and the black shaded isoline represents the <u>35 psu</u> salinity near the surface.



Deleted: of	_
Deleted: They are divided into	
Deleted: around	_
Deleted: and	
Deleted: are marked on the panel.	
Deleted: . They are collected	
Deleted:) respectively, and with different color in which the	_
Deleted: (_
Deleted: or	
Deleted: denotes	
Deleted: observations in	
Deleted: (
Deleted: or	
Deleted:).	
Deleted: ¶ ¶ [6	[2]

Deleted: reanalyzes	
Deleted: MOI	
Deleted: extent	
Deleted: TOPAZ4	
Deleted: of 35 psu	



concentration from TP4), the black lines represent ± 1 psu.

Formatted: Normal, Indent: Left: 0 cm





Fig. 7 Scatterplots of SSS compared to the CORA5.1 in-situ observations with respect to the SO-S3 regions in the Arctic. The diamonds (anti-triangles, stars, squares, circles, and triangles) represents the SSS from TP4 (BEC, PHC, WOA, MOB, and CEC respectively). The black (red) lines are the linear regressions of the blue (purple) dots in each panel, and the coefficient R² is indicated in the panel together with the number of observations in parentheses.





Fig. 8 Same as Fig. 7 but for the subpolar regions S4-S7.



Fig. 9 The mean deviation of SSS for the six datasets compared to in situ observations from CORA 5.1 during the three years of 2011-2013 in the northern North Atlantic and the Nordic seas. The SSS observations are distributed into the coarse grid cells of 9x9 grids in TP4, with a gray mask if the valid observations less than 10.





Fig. <u>10</u> The Root Mean Square deviation of SSS for six datasets compared to in situ observations from CORA 5.1 during the three years of 2011-2013 in the northern <u>North</u> Atlantic and <u>the</u> Nordic seas. The SSS observations are distributed into the coarse grid cells of 9x9 grids in TP4, with a gray mask if the valid observations less than 10.









Fig. <u>12</u> Scatterplots of SSS uncertainty compared to the in-situ observations in Beaufort Sea as a function of the observed salinity. The black dashed line represents the absolute deviation of <u>5</u> psu. <u>(a)</u> The diamond (anti-triangle) represents from TP4 (BEC) with blue (purple). <u>(b)</u> The star (square) from the climatology of PHC (WOA). <u>(c) The</u> circle (triangle) represents from <u>MOB</u> (CEC). The thick dashed curves are fitted by the fourth order polynomial function, and the norm residuals are marked on panel respectively.



Deleted: 3	
Deleted: Left:	
Deleted: Middle:	
Deleted: Right: the	
Deleted: MOI	

Page 24: [1] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Arial		
Page 24: [2] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold		
Page 24: [3] Deleted	Jiping Xie	6/15/19 3:27:00 AM
¥		
Page 24: [3] Deleted	Jiping Xie	6/15/19 3:27:00 AM
¥		
Page 24: [3] Deleted	Jiping Xie	6/15/19 3:27:00 AM
Y		
Page 24: [3] Deleted	Jiping Xie	6/15/19 3:27:00 AM
v		
Page 24: [3] Deleted	Jiping Xie	6/15/19 3:27:00 AM
•		
Page 24: [3] Deleted	Jiping Xie	6/15/19 3:27:00 AM
•		
Page 24: [4] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: 10,5 pt		
Page 24: [5] Formatted Table	Jiping Xie	6/15/19 3:27:00 AM
Formatted Table		
Page 24: [6] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Centered		
Page 24: [7] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold, Not Italic		
Page 24: [8] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold, Not Italic		
Page 24: [9] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Bold, Italic, Pattern: 15% (Auto Foreground, White Background)	
Page 24: [10] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Bold, Italic, Pattern: 15% (Auto Foreground, White Background)	
Page 24: [11] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold, Not Italic		
Page 24: [12] Formatted	Jining Xie	6/15/19 3:27:00 AM
Font: Bold. Italic. Pattern: 15% (Auto Foreground, White Background) High	light
Page 24. [13] Formatted		6/15/19 3·27·00 AM
left	orping Arc	0/15/17 5.27.00 AM

Jiping Xie	6/15/19 3:27:00 AM
White Background)	
Jiping Xie	6/15/19 3:27:00 AM
uto Foreground, White Background),	Highlight
Jiping Xie	6/15/19 3:27:00 AM
White Background), Highlight	
Jiping Xie	6/15/19 3:27:00 AM
Jiping Xie	6/15/19 3:27:00 AM
¥	
Jiping Xie	6/15/19 3:27:00 AM
Jiping Xie	6/15/19 3:27:00 AM
uto Foreground, White Background),	Highlight
Jiping Xie	6/15/19 3:27:00 AM
Jiping Xie	6/15/19 3:27:00 AM
uto Foreground, White Background)	
Jiping Xie	6/15/19 3:27:00 AM
Jiping Xie	6/15/19 3:27:00 AM
uto Foreground, White Background)	
Jiping Xie	6/15/19 3:27:00 AM
uto Foreground, White Background)	
Jiping Xie	6/15/19 3:27:00 AM
Jiping Xie	6/15/19 3:27:00 AM
Jiping Xie	6/15/19 3:27:00 AM
uto Foreground, White Background)	
Jiping Xie	6/15/19 3:27:00 AM
uto Foreground, White Background)	
Jiping Xie	6/15/19 3:27:00 AM
preground, White Background)	
Jiping Xie	6/15/19 3:27:00 AM
uto Foreground, White Background)	
Jiping Xie	6/15/19 3:27:00 AM
	Jiping Xie White Background) Jiping Xie Auto Foreground, White Background), Jiping Xie White Background), Highlight Jiping Xie Jiping Xie Jiping Xie Auto Foreground, White Background), Jiping Xie Jiping Xie Jiping Xie Auto Foreground, White Background) Jiping Xie

Font: Bold, Italic, Pattern: 15% (Auto Foreground, White Background)

A		
Page 24: [32] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Left		
Page 24: [33] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground	d, White Background)	
Page 24: [34] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Bold, Pattern: 15% (Auto	Foreground, White Background)	
Page 24: [35] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Bold, Pattern: 15% (Auto	Foreground, White Background)	
Page 24: [36] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold, Not Italic		
Page 24: [37] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold, Not Italic		
Page 24: [38] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground	d, White Background)	
Page 24: [39] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground	d, White Background)	
Page 24: [40] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Bold, Italic, Pattern: 15%	(Auto Foreground, White Background)
Page 24: [41] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Left, Tab stops: 0,31 cm, Cente	ered	
Page 24: [42] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground	d, White Background)	
Page 24: [43] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Bold, Italic, Pattern: 15%	(Auto Foreground, White Background)
Page 24: [44] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground	d, White Background)	
Page 24: [45] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground	d, White Background)	
Page 24: [46] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold, Not Italic		
Page 24: [47] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold, Not Italic		
Page 24: [48] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Bold, Italic, Pattern: 15%	(Auto Foreground, White Background)
Page 24: [49] Formatted	Jiping Xie	6/15/19 3:27:00 AM

Font: Bold, Italic, Pattern: 15% (Auto Foreground, White Background)

Т

Page 24: [50] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground,	White Background)	
Page 24: [51] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground,	White Background)	
Page 24: [52] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Pattern: 15% (Auto Foreground,	White Background)	
Page 24: [53] Formatted	Jiping Xie	6/15/19 3:27:00 AM
Font: Not Bold		
Page 24: [54] Deleted	Jiping Xie	6/15/19 3:27:00 AM
×		
Page 24: [54] Deleted	Jiping Xie	6/15/19 3:27:00 AM
X		
Page 24: [54] Deleted	Jiping Xie	6/15/19 3:27:00 AM
¥.		
Page 24: [54] Deleted	Jiping Xie	6/15/19 3:27:00 AM
· ·		
Page 24: [55] Formatted Table	Jiping Xie	6/15/19 3:27:00 AM
Formatted Table		
Page 24: [56] Deleted	Jiping Xie	6/15/19 3:27:00 AM
	۷	
Page 24: [57] Deleted	Jiping Xie	6/15/19 3:27:00 AM
	Ţ	
Page 24: [58] Deleted	Jiping Xie	6/15/19 3:27:00 AM
	X	
Page 24: [58] Deleted	Jiping Xie	6/15/19 3:27:00 AM
Page 24: [59] Deleted	Jiping Xie	6/15/19 3:27:00 AM
	X	
Page 24: [59] Deleted	Jiping Xie	6/15/19 3:27:00 AM
	X	
Page 24: [60] Deleted	Jiping Xie	6/15/19 3:27:00 AM
Dage 24. [(0] Dalate J	1:	(12/10 2.07.00 +34
rage 24: [60] Deleted	Jiping Xie	0/15/19 3:27:00 AM
Page 25. [61] Deleted	Jining Via	6/15/10 3.27.00 AM
	Jiping Alt	0/13/17 3.27.00 AIVI
Page 25: [62] Deleted	Jiping Xie	6/15/19 3:27:00 AM

Page 26: [63] Deleted	Jiping Xie	6/15/19 3:27:00 AM
Page 27: [64] Deleted	Jiping Xie	6/15/19 3:27:00 AM

¥...