

Dear Editor,

We thank the two reviewers for their critical and constructive comments on our research. Their comments have significantly improved our manuscript. The detailed responses to their comments are listed as following: the reviewer comments are in black and our response is in red.

Anonymous Referee #1

Although it is clarified in the text, it is not clear in figures 7, 8 and 11 the corresponding dataset for each provided value of R^2 ($R^2=X/Y$). Please, indicate the correspondence between X/Y and the datasets in the figure caption or by write X and Y numbers in a different color, according to each dataset, will help to the reader.

-A: Thanks for this comment. More explanation about R^2 is added in the caption like "... R^2 between the evaluated product and the in-situ SSS..." in Fig. 7, and same as in Fig.8 and Fig. 11.

Anonymous Referee #2

The authors did a satisfactory job in responding to my comments and rewriting their earlier draft. Minor revisions are needed and English might be improved, still.

-A: We thank the referee for the detailed evaluation of our manuscript and constructive suggestions. We appreciated this very much, all the comments are taken into account in the new revision. The English has been improved especially for the last parts of the paper.

Line 24: rephrase "all six SSS products share a common challenge to represent fresh water masses"

-A: Thank you for this comment. It is changed as

Line 21- 25: "When compared against independent in-situ data in the Beaufort Sea, the BEC product shows the smallest bias (<0.1 psu) in summer and the smallest RMSD (1.8 psu). The results also show that all six SSS products have a common challenge to represent fresh water masses (<24 psu) in the central Arctic."

Line 82: four SMOS products have been previously mentioned. Specify the two that are considered.

-A: Thank you for this comment. The more specific statement is added.

Line 81-83: “The present study thus investigates the accuracy of these two L3 SSS products from SMOS in the Arctic Ocean.”

Line 88: add ocean to reanalysis products

-A: Thanks for this point, it is added.

Line 89: Uotila et al. presented temperature and salinity fields in the Arctic. Do you refer here to the seasonal cycle of both variables? Why the ten reanalysis are so different in the Arctic salinity, and all probably wrong? Topaz is part of the inter-comparison, add a more specific comment on results by Uotila et al.

-A: Thank you for this comment. Here, we only refer to the salinity seasonal cycle in Uotila et al. (2018). Although most reanalysis products (seven in the ten reanalyses in Table 1 of Uotila et al., 2018) restored salinity to climatology, it should be noticed that different salinity datasets were used, which also reveals the lack of a universal SSS reference. So we add the related comment in the text.

Line 90-93: “Although most reanalysis products (seven out of ten reanalyses in Table 1 of Uotila et al., 2018) restored salinity to climatology, they did not use the same salinity climatology, which betrays the lack of a universal SSS reference.”

Line 112: “can it also give...” is here the evaluation against in situ data, the subject?

-A: Yes, it is. It is further corrected by “Can the evaluation against in-situ data also shed light on the uncertainties of the SMOS products?”

Line 210: BGEP is available from CMEMS too, as required by the title of section 2.2?

-A: Unfortunately not. This is why we list BGEP under Section 2.3, not 2.2. In this study, the in-situ observations from BGEP are directly downloaded from the website (<http://www.who.edu/>), and were not assimilated into TP4. The quantitative evaluation of SSS use that as one of the independent observations so we keep it in section 2.3.

Caption for Figure 1: only four sub-regions are in the Arctic Ocean, the others are located in the Nordic Seas and North Atlantic. Please add in the manuscript a clear

definition of the Arctic domain, North Atlantic domain. The two are often mistaken in the text.

-A: Thank you for this comment. In this study, the Arctic Ocean is limited to north of 60N. Here, considering the distributions of the valid in-situ observations from CORA5.1, the subregions are divided into 8 regions. Clearly, the subregions of S0-S4 are regarded as in the Arctic region, the other regions of S5-S7 are attributed into the northern North Atlantic.

So the caption for Fig. 1 has a change as “8 sub-regions divide the Arctic Ocean (S0-S4) and the northern North Atlantic Ocean (S5-S7), ...”

In additional, more statement about this issue is added in Section 4.1

Line 322-326: “In this study, the Arctic domain (>60N) is the core region for evaluation, divided into five sub-regions numbered from S0 to S4. It contains the central Arctic (sub-regions S0, S1, S2, and S3) and the Nordic Seas (S4). The regions from S5 to S7 are in the northern North Atlantic.”

Line 243: the 35 psu isoline marks the Atlantic water that does only marginally reach the Arctic ocean. I suggest to add a lower-salinity isoline to the plot to better highlight also the inflow within the Arctic, something between 33 and 34 psu for example

-A: Thank you this nice suggestion. We add the isoline of 33.6 psu and tuning the colorbar with a larger range as shown in the updated Fig. 2 and 3.

Figure 2: the minimum salinity is not clearly shown, the blue saturates at 30 psu? subplot e and f: are the salinity fields correct close to North Pole? It seems there is an issues in the interpolation at very high latitude for these two products

-A: Yes, the minimum salinity is not clearly shown in Fig.2 due to the colorbar is cut off when the salinity below 30 psu. So the colorbars in the new figures have been extended to represent fresh waters. In the central Arctic, the lower SSS in TP4 and PHC is around 30 psu, which is rather saline compared to that in MOB and WOA. Both suffer from interpolation artefacts due to their unfortunate regular lat-lon projection (singularity at the North Pole).

Line 252: The comparison is between BEC/CEC with all the other products, or BEC against CEC?

-A: The comparison is between BEC/CEC with all the other products, especially indicated by the dashed line of 35 psu in Fig. 3, they are both less saline.

Line 255: I do not see that the 4 products agree in the North Atlantic. Rephrase

-A: Thank you for this comment. The 4 products show the similar patterns by the dashed line (35 psu) in the North Atlantic and the Nordic seas. To avoid the misunderstanding, the text is changed at Line 257-259: "Although the SSS of TP4, MOB, PHC and WOA agree relatively well in the North Atlantic Ocean and the Nordic seas **as shown by the dashed lines of 35 psu, ...**"

Line 260: what is exactly a universal reference?

-A: Here a universal reference means a common reference to Arctic SSS analysis that can be consensually accepted or used in both spatial and temporal resolution and accuracy.

Line 267: the Beaufort Sea is almost all ice-covered in CEC. The area that you consider here is unclear.

-A: Thank you for this comment. In August, the CEC SSS appears a smaller area than BEC in the Beaufort Sea. For the two SMOS products, we only consider ice free pixels.

Line 268: CEC presents positive deviation in the Kara Sea close to coast line, probably due to the land-ocean interaction. Please add a line on that.

-A: In fact, we noticed the positive deviations of CEC near the coast line (not only in the Kara Sea) which are rather significant even compared with that in BEC.

A concerned comment is added as

Line 272-273: "A positive deviation of CEC is noticeable in the Kara Sea, which indicates the land-ocean interaction stronger than that in BEC."

Line 273: I suggest to add a line on the missing low salinity related to the polar water that travels southward from the Arctic

-A: Thank you for this suggestion. A comment is added as

Line 280-284: "For the BEC and CEC products that use different ice masks, the deviations are averaged outside their respective ice mask, not their intersection. Comparing the low salinity lines of 33.6 psu in Fig. 3a and 3d, it clearly shows the polar water southward from Arctic has a misinterpretation owing to the used ice mask."

Line 276: near and below the sea-ice cover reproduced by TP4?

-A: Yes, thank you this remind. This definition is added as Line 285-286: "Near and below the sea-ice cover reproduced by TP4 (the thick brown line in the figures), ..."

Line 285: how is sea ice cover treated in all products in computing the deviations in Fig 6?

-A: Figure 6 reveals the monthly deviations of the five SSS products referred to TP4, which is constrained at north of 60N without considering sea ice cover, although the two SMOS products only use ice free pixels. If averaging the deviations outside of ice cover (defined by 0.15 concentration in TP4), the monthly deviations of the five products referred to TP4 are shown in Fig. A as bellow. Clearly, the BEC and the CEC have similar deviation features like in Fig. 6, compared with other products except of the specific values.




Fig. A Monthly deviations in the Arctic Ocean (>60N; out of ice cover defined by TP4) of (a) the RMS and (b) the spatial average during the period 2011-2013 for the five SSS products referred to TP4. The anti-triangle (triangle, circle, star and square) line represents the SSS deviations from BEC (CEC, MOB, PHC and WOA respectively).

Line 293-294: is that evident in fig 6b?

-A: Referred to the TP4 SSS, the RMS deviation (Fig. 6a) of BEC has consistently smaller RMS compared with the other products. For the mean deviation (Fig. 6b), the same conclusion is evident for BEC except in the summer months. In summer, the SSS deviation of CEC clearly shows large deviations of opposite signs in Fig. 5c, which sums up to the smaller deviation compared to that in BEC in Fig. 6b.

Line 379: Rephrase. The range is larger, the salinity lower.

-A: Thank you for this point. It is revised as Line 392-393: "On the other hand, the range of TP4 SSS increases from 19 to 32 psu, with a larger saline bias of 2.59 psu and a RMSD of 3.63 psu."

Line 447: rewrite “if it to be assimilated into”

-A: Thanks for this point. The whole sentence is rephased as Line 456-458:” Thus, it seems that the two SMOS products would give rise to significantly different effects to the upper ocean state, were they assimilated.”

Evaluation of Arctic Ocean surface salinities from SMOS against a regional reanalysis and in situ data

Jiping Xie¹, Roshin P. Raj², Laurent Bertino², Annette Samuelsen², and
Tsuyoshi Wakamatsu²

1. Nansen Environmental and Remote Sensing Center, N5006 Bergen, Norway
2. Nansen Environmental and Remote Sensing Center and Bjerknes Centre for
Climate Research, Bergen, N5006 Bergen, Norway

Correspondence to: Jiping. Xie (jiping.xie@nersc.no)

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Abstract

Recently two gridded Sea Surface Salinity (SSS) products that cover the Arctic Ocean have been derived from the European Space Agency's (ESA) Soil Moisture and Ocean Salinity (SMOS) mission: one developed by the Barcelona Expert Centre (BEC) and the other developed by the Ocean Salinity Expertise Center of the Centre Aval de Traitement des Données SMOS at IFREMER (CEC). The uncertainties of these two SSS products are quantified during the period of 2011-2013 against other SSS products: one data assimilative regional reanalysis; one data-driven reprocessing in the framework of the Copernicus Marine Environment Monitoring Services (CMEMS); two climatologies; the 2013 World Ocean Atlas (WOA) and the Polar science center Hydrographic Climatology (PHC); and in-situ datasets, both assimilated and independent. The CMEMS reanalysis comes from the TOPAZ4 system which assimilates a large set of ocean and sea-ice observations using an Ensemble Kalman Filter (EnKF). Another CMEMS product is the Multi-OBServations reprocessing (MOB), a multivariate objective analysis combining in-situ data with satellite SSS. The monthly root mean squared deviations (RMSD) of both SMOS products, compared to the TOPAZ4 reanalysis, reach 1.5 psu in the Arctic summer, while in the winter months the BEC SSS is closer to TOPAZ4 with a deviation of 0.5 psu. The comparison of CEC satellite SSS against in-situ data shows too fresh Atlantic Water in the Barents Sea, the Nordic seas, and in the northern North Atlantic Ocean, consistently with the abnormally fresh deviations against TOPAZ4. When compared against independent in-situ data in the Beaufort Sea, the BEC product shows the smallest bias (<0.1 psu) in summer and the smallest RMSD (1.8 psu). The results also show that all six SSS products share a common challenge to represent fresh water masses (<24 psu) in the central Arctic. Along the Norwegian coast and at the southwestern coast of Greenland, the BEC SSS shows smaller errors than TOPAZ4 and indicates the potential value of assimilating the satellite-derived salinity in this system.

Keywords: Arctic Ocean; sea surface salinity; SMOS; reanalysis;

Formatted: Font color: Text 1

Deleted: ,

Deleted:),

Deleted: :

Deleted:),

Deleted: waters

Deleted:), although

Deleted: fresher

Deleted:).

Formatted: Font color: Text 1

1. Introduction

The sea surface salinity (SSS) plays a key role in tracking processes in the global water cycle through precipitation, evaporation, runoff, and sea-ice thermodynamics (Vialard and Delecluse, 1998; Sumner and Belaine, 2005; Vancoppenolle et al., 2009; Yu, 2011). SSS is known to impact the oceanic upper mixing significantly (Latif et al., 2000; de Boyer Montegut et al., 2004; Maes et al., 2006; Furue et al., 2018) and via its effect on the surface layer density (Johnson et al., 2012). The SSS also affects the decadal variability of hydrography in the upper waters of the North Atlantic (Reverdin et al., 1997). Using a coupled atmosphere-ocean model and an observed SSS climatology dataset, Mignot and Frankignoul (2003) attributed the interannual variability of the Atlantic SSS to two factors: anomalous Ekman advection and the freshwater flux. Additionally, the increased melting of glaciers and sea-ice in the Arctic (McPhee et al., 1998; Macdonald et al., 1999) leads to significant changes in the salinity distribution and fresh water pathways (Steele and Ermold, 2004; Morison et al., 2012). The freshwater flux is regarded as one of the least constrained parameters in ocean models due to poorly known river discharge, precipitation, and glacial/sea-ice melt (e.g., Tseng et al., 2016; Furue et al., 2018). In ocean models the sea-surface freshwater flux is often adjusted directly or the SSS is restored to its corresponding climatological value to avoid salinity drift.

Monitoring SSS from space is crucial for understanding the global water cycle and the ocean dynamics, especially in the Arctic Ocean where our knowledge of the SSS variability is limited due to non-homogenous and sparse in-situ data. The European Space Agency's (ESA) Soil Moisture and Ocean Salinity (SMOS) satellite, launched in November 2009, consists of the Microwave Imaging Radiometer using Aperture Synthesis (MIRAS) instrument, a passive 2-D interferometric radiometer operating in L-band (1.4 GHz, 21 cm), that measures the brightness temperature (BT) emitted from the Earth. The L-band microwave is highly sensitive to water salinity, which influences the dielectric constants in the sea, and is less susceptible to atmospheric or vegetation-induced attenuation than higher frequency measurements (Font et al., 2010; Kerr et al., 2010; Mecklenburg et al., 2012). Committed to provide global salinities averaged over 10-30 days with an accuracy of 0.1 psu in the open ocean, ESA provides the MIRAS data into SMOS Level 1 (L1) and Level 2 (L2) products through a set of sequential processors (Mecklenburg et al., 2012; ESA, 2017).

77

78 Over the ocean, Level 2 products (L2OS) are comprised of three different ocean
79 salinities, together with the BTs at the top of atmosphere and at the sea surface,
80 distributed by ESA with swath-based format (e.g., SMOS Team, 2016; ESA, 2017).

81 As a result of the efforts of the national agencies in France and Spain respectively,
82 two Level 3 (L3) data products of SSS are freely available, which are independently
83 developed by the Ocean Salinity Expertise Center (CECOS) of the Centre Aval de
84 Traitement des Données SMOS at IFREMER and the Barcelona Expert Centre.

85 These two SMOS products have successfully resolved the Agulhas salinity front
86 (D'Addezio et al., 2016) and proven useful for the estimating precipitation (Supply et
87 al., 2018). The work of Olmedo et al. (2018) quantitatively evaluate the accuracy of
88 the SMOS Arctic and sub-Arctic SSS to less than 0.35 psu, but this evaluation

89 against Argo data was limited by the lack of data in the Arctic proper. The present
90 study thus investigates the accuracy of these two L3 SSS products from SMOS in the
91 Arctic Ocean.

92

93 A good estimate of surface salinity is a necessary step towards the knowledge of the
94 three-dimensional water mass properties, for which data assimilation and optimal
95 interpolation methods must be invoked. In a recent study, Uotila et al. (2018)
96 investigated the Arctic salinity in ten ocean reanalysis products and found

97 disagreements within them regarding the seasonal cycle in the upper layer (0-100 m;

98 Figure 12 of Uotila et al., 2018). Although most reanalysis products (seven out of ten
99 reanalyses in Table 1 of Uotila et al., 2018) restored salinity to climatology, they did
100 not use the same salinity climatology, which betrays the lack of a universal SSS

101 reference. Note that the full assessment of the Arctic SSS products has been

102 hindered by the extreme paucity of in-situ data in the Arctic. The SSS data from the
103 SMOS mission should in principle allow the evaluation of salinity on a basin scale. In
104 this study, we use two SSS products available from the Copernicus Marine

105 Environment Monitoring Service (CMEMS). The first is the regional Arctic CMEMS
106 reanalysis (ARCTIC-REANALYSIS-PHYS-002-003) from the TOPAZ4 assimilation

107 system, which is a coupled ocean and sea-ice data assimilation system using the
108 Ensemble Kalman filter (EnKF) to assimilate the various ocean and sea-ice
109 observations (e.g., Xie et al., 2017). The second is the CMEMS multivariate optimal
110 interpolation reprocessing (MULTIOBS_GLO_PHY_REP_015_002, Droghei et al.,

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: SMOS

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: 2018).

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

2018). The latter product directly merges in-situ data with satellite measurements including SMOS without the use of a model and is therefore a reprocessing rather than a reanalysis. There are four other global reanalysis products under CMEMS, but understanding well their differences requires an intimate knowledge of their setup, and is out of scope of the present study.

We assess the quantitative deviations of Arctic SSS among the two SMOS products and the two CMEMS products, together with two climatology datasets: WOA13 (version 2.0 of World Ocean Atlas of 2013; Zweng et al., 2013) and the older PHC (Polar Science Center Hydrographic Climatology version 3.0; Steele et al., 2001). We further extend the evaluation using available in-situ salinity observations during the years 2011-2013 from different data sources. Can the evaluation against in-situ data also shed light on the uncertainties of the SMOS products? Can it also give useful information needed for the assimilation of the SMOS SSS products into an Arctic ocean forecast/reanalysis system?

The paper is organized as follows: Section 2 describes all SSS products and the in-situ datasets. The monthly mean SSS from these six products are intercompared and monthly differences from the TOPAZ SSS are analyzed in Section 3. Section 4 evaluates the SSS products against in-situ data, which are divided between assimilated and independent data. A summary of this study is provided in Section 5.

2. Data description

2.1 Sea surface salinity from SMOS

The SSS retrieval from SMOS is subject to biases originating from various non-geophysical sources such as the so-called land-sea contamination and the latitudinal biases, mainly caused by the thermal drift of the instrument. A particular challenge in the Arctic is the sea-ice edge because of ice-ocean contamination. Based on different statistical approaches, match-up criteria, and SMOS data filtering flags, two centers have developed separate processing chains producing a Level 3 SSS product on a regular grid. These two SSS products are hereafter named respectively CEC and BEC in this study, evaluated during the three years of 2011-2013 (see Table 1).

- *The BEC product*

Formatted: Font color: Text 1

Deleted: the

Formatted: Font color: Text 1

Deleted: uncertainty

Formatted: Font color: Text 1

149 The latest regional Arctic product (version 2.0) from BEC is available from
150 <http://bec.icm.csie.es> since December 2018 (last access: March 2019). The BEC
151 SSS product was generated from ESA L1B (v620) products, and accumulates salinity
152 data over 9 days with a spatial grid resolution of 25 km. With respect to its previous
153 version, a systematic bias in the retrieved salinity is corrected by computing the
154 SMOS climatology (the most probable value for a given lat-lon, incidence angle and
155 across-swath distance) which is substituted by a reference value from WOA13. In
156 addition, a temporal bias correction has been refined in this version using near-
157 surface Argo salinity to compute regional averages (see the details in Olmedo et al.,
158 2018).

- 159 • *The CEC product*

160 The third version of LOCEAN SMOS SSS L3 maps (L3_DEBIAS_LOCEAN_v3) was
161 released by the CECOS in July 2018. Every 4 days, the SSS maps averaged over 9
162 days are released on <ftp.ifremer.fr> (last access: December 2018). This product uses
163 the Equal-Area Scalable Earth Grid (EASE-Grid) which has limited grid distortion and
164 a spatial resolution of 25km. Using a Bayesian retrieval approach (Kolodziejczyk et
165 al., 2016), the SMOS systematic errors in the vicinity of continents are discarded o
166 improve the product quality. Further, a 'de-biasing' method (Boutin et al., 2018) has
167 been applied in this version of the CEC product, in which the non-Gaussian
168 distribution of SSS is taken into account, refining the latitudinal correction at high
169 latitude, and preserving the naturally seasonal variability of SSS.

171 2.2 Sea surface salinity from two CMEMS products

- 172 • *The TOPAZ4 Arctic MFC reanalysis*

173 TOPAZ4 uses the version 2.2 of Hybrid Coordinate Ocean Model (HYCOM,
174 Chassignet et al., 2003; Bertino and Lisæter, 2008) coupled with a simple
175 thermodynamic sea ice model (Drange and Simonsen, 1996) in which the elastic-
176 viscous-plastic rheology describes the sea ice dynamics (Hunke and Dukowicz,
177 1997). The model domain covers the Arctic Ocean and the [North](#) Atlantic Ocean with
178 a horizontal resolution of 12-16 km. In order to obtain an accurate and dynamically
179 consistent reanalysis in the Arctic Ocean, the deterministic EnKF (DEnKF; Sakov and
180 Oke, 2008) was implemented in TOPAZ with a dynamical ensemble of 100 members
181 all driven by perturbed 6-hourly atmosphere forcing from ERA interim (Simmons et

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: north

Formatted: Font color: Text 1

al., 2007). The perturbations of precipitations are following a log-normal probability distribution and conserve the ensemble-average total precipitation. Along the model lateral boundaries in the South Atlantic and in Bering Strait, the temperature and salinity are relaxed to a combined climatology data from PHC and WOA. The river discharges are treated as an additional mass and a negative salinity flux. Near the surface, to avoid the salinity drift (Tseng et al., 2016; Furue et al., 2018), a weak relaxation to the same combined climatological SSS with 30 days decay is used as most ocean models, but restricted to the areas where the difference to climatology is smaller than 0.5 psu. The EnKF assimilates various ocean and sea-ice observations (e.g., Xie et al., 2016, 2018) into a multivariate state update of the HYCOM model. The understanding for the uncertainty of the TOPAZ4 SSS has been hindered by poor coverage of in-situ data over the Arctic domain, although Xie et al. (2017) had comprehensively assessed the TOPAZ4 reanalysis during 1991-2013 against various types of ocean and sea-ice observations. For the sake of brevity, the TOPAZ4 reanalysis SSS is named TP4 hereafter.

- *SSS from the Multi-Observations dataset*

The CMEMS product of MULTIOBS_GLO_PHY_REP_015_002 combines the SSS observations from in-situ and satellite data, using optimal interpolation (OI, Buongiorno Nardelli et al., 2016; Verbrugge et al., 2018) at weekly interval on a $0.25^{\circ} \times 0.25^{\circ}$ regular grid. The main datasets used during the OI processing are: 1) the quality controlled in-situ data, CORIOLIS dataset for Re-Analysis (CORA, Cabanes et al., 2013) distributed through CMEMS; 2) the objectively analyzed SSS and SST data generated from CORA, also distributed by CMEMS, which uses the WOA 2013 climatology as first guess and has been upscaled to the MOB grid as another first guess of the multidimensional OI; 3) The SMOS L3 binned (L3bin) data reprocessed by SMOS-BEC at 0.25° grid, although the previous version 1.0 of the product mentioned above; 4) The daily Reynolds L4 AVHRR_OI Global blended SST product on a 0.25° grid. This product is called MOB hereafter.

2.3 Surface salinity from in-situ data

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

215 The in-situ SSS data are acquired here from three quality-controlled datasets. The
216 first data source is CORA from CMEMS (product id:
217 INSITU_GLO_TS_REP_OBSERVATIONS_013_001_b), also used in the MOB SSS.
218 CORA contains temperature and salinity profiles from various in-situ data sources
219 (Cabanes et al., 2013). Since 2013, the CORA dataset has been updated every year
220 and includes all the Argo float profiles, moorings, gliders, Ice-Tethered Profilers (ITP;
221 Toole et al., 2011), XBT, CTD, and XCTD data. The latest version of the dataset,
222 CORA5.1, covers the period of 1950-2016. Figure 1a shows the distribution of SSS
223 (averaged over 0-8 m depth) observations from CORA5.1 (total 69,246 observations)
224 over the domain north of 52°N during the years 2011-2013.

225 The second source of in-situ data is from the Beaufort Gyre Experiment Project
226 (BGEP, <http://www.whoi.edu/website/beaufortgyre/background>, last access: 14th
227 December 2018). In order to monitor the natural variabilities of the Beaufort Sea in
228 the Canada Basin, BGEP maintains moorings since 2003 and acquires in-situ
229 measurements over the Beaufort Sea region every summer. Symbols (anti-triangle,
230 square, and star) shown in Fig. 1b indicate the locations of valid SSS observations
231 obtained from BGEP. The in-situ dataset used in this study is obtained from the GO-
232 SHIP (the Global Ocean Ship-based Hydrographic Investigations Program, Talley et
233 al., 2017) database under the Climate Variability and Predictability Experiment
234 (CLIVAR). The SSS observations in the Beaufort Sea are extracted from
235 CLIVAR/GO-SHIP data with EXPOCODE (33HQ20111003 and 33HQ20121005, ref.
236 Mathis and Monacchi, 2014), which are available from [https://cdiac.ess-](https://cdiac.ess-dive.lbl.gov/ftp/oceans/CARINA/Healy/)
237 [dive.lbl.gov/ftp/oceans/CARINA/Healy/](https://cdiac.ess-dive.lbl.gov/ftp/oceans/CARINA/Healy/) (last access: 18th December 2018). All the
238 valid salinity profiles are averaged within the upper 8 m layer, in order to match at
239 best with the satellite SSS measurements. Contrarily to the CORA data, both BGEP
240 and CLIVAR data are independent from all the evaluated datasets.

242 3. Intercomparison of monthly SSS fields

243 Prior to the intercomparison of different SSS products, all the gridded products from
244 satellite, reanalysis and climatology have been mapped on the same grid used in the
245 TP4 model by a “nearest neighbor” interpolation. To quantitatively evaluate the SSS
246 deviation in the Arctic, the bias and the root mean square deviation (RMSD) are
247 defined by

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

$$\text{Bias} = \frac{1}{p} \sum_{i=1}^p (\mathbf{H}_i \mathbf{x}_i^f - \mathbf{s}_i) \quad (1)$$

$$\text{RMSD} = \sqrt{\frac{1}{p} \sum_{i=1}^p (\mathbf{H}_i \mathbf{x}_i^f - \mathbf{s}_i)^2} \quad (2)$$

Where p is the length of the time series, \mathbf{x}_i^f is the valid salinity from different sources at the i th time, compared to the reference salinity field \mathbf{s}_i . \mathbf{H}_i is the observation operator projecting \mathbf{x}_i^f onto \mathbf{s}_i .

Formatted: Font color: Text 1

• Monthly mean comparison of SSS

Formatted: Font color: Text 1

Figure 2 shows the monthly mean Arctic SSS in March from the six products. Notable differences in the two SMOS products appear in the Nordic Seas, Barents Sea, and around the Labrador Sea. At first sight, the large-scale SSS features from SMOS products are similar to the other products. However, the CEC SSS is fresher (as shown by the isolines of 35 psu) compared to the BEC, TP4, MOB and both climatologies. The location of the sea-ice edge in the two SMOS products match comparatively well with the TP4 reanalysis (Fig. 2a, d). In sea-ice covered region, TP4 shows a gradual decrease in SSS from the European to the American sector, with two minima near the Beaufort Sea and the East Siberian Sea (ESS; Fig. 2b) consistently with the PHC (Fig. 2c). Those are unclear in the MOB and WOA (Fig. 2e, f), especially the SSS minimum in the Beaufort Sea. The latter two products also show artificial projection artefacts around the North Pole.

Deleted: in the northern North Atlantic Ocean.

Formatted: Font color: Text 1

Figure 3 shows the corresponding SSS fields in September. In comparison to the March situation, the BEC and CEC SSS in the Nordic Seas are both less saline, indicated by the 35 psu isoline. The sea ice masking of the two SMOS products differ considerably in the Canadian Basin and in the Arctic marginal seas. Although the SSS of TP4, MOB, PHC and WOA agree relatively well in the North Atlantic Ocean as shown by the dashed lines of 35 psu, the discrepancies become dramatic in ice-covered areas. Below the ice or near the sea-ice edge (denoted by the brown thick line in Fig. 2 and 3), TP4 and PHC share common features, which can be explained by the model restoring to PHC. On the other hand, the MOB and WOA differ significantly in spite of WOA being used as input to the MOB. Short of a universal reference for Arctic SSS, the monthly mean SSS deviations will be quantified using TP4 as a reference.

Deleted: northern

Formatted: Font color: Text 1

Formatted: Font color: Text 1

• *Deviation analysis of monthly SSS referred to TP4*

Figure 4 and Figure 5 show the deviations of the monthly mean SSS of the five products with reference to the TP4 SSS in August and September respectively. In August, the two SMOS products (Fig. 4a, c) show coherently negative deviations (~2 psu) in the marginal seas of the Beaufort Sea, the ESS, the Laptev Sea, and the Kara Sea. A positive deviation of CEC is noticeable in the Kara Sea, which indicates the land-ocean interaction stronger than that in BEC. In the North Atlantic Ocean, away from the sea-ice edge, the deviation of the BEC from TP4 is lower (bias less than 0.5 psu). Focusing on the Arctic domain (>60°N), the mean deviation of the BEC SSS is -0.87 psu and its root mean square is 1.75 psu. The CEC SSS shows considerable negative deviations over 1 psu in the North Atlantic, from north of Denmark Strait to the west coast of Ireland. This is remarkably different from the BEC, and does not discern the subpolar from the subtropical waters there (Hátún et al., 2005). For the BEC and CEC products that use different ice masks, the deviations are averaged outside their respective ice mask, not their intersection. Comparing the low salinity lines of 33.6 psu in Fig. 3a and 3d, it clearly shows the polar water southward from Arctic has a misinterpretation in CEC owing to the used ice mask. The deviations of MOB and the two climatology products are comparatively small in the open ocean of the North Atlantic (Fig. 4b, e). Near and below the sea-ice cover, reproduced by TP4 (the thick brown line in the figures), the deviations are much larger, particularly both the MOB and WOA show strong saline anomalies (> 1 psu) in the Eurasian basin and low anomalies in the American basin.

In September, the SSS deviations of BEC, MOB, PHC and WOA show similar fresher patterns as in August, but the CEC deviations becomes surprisingly positive around the ice edge. The SSS deviation of CEC, averaged over the Arctic domain (>60°N), swaps from -0.42 to 0.42 psu from one month to the next one. The seasonal evolution of monthly SSS deviations from TP4 for all five remaining products, averaged over the Arctic, are shown in Fig. 6. Among the five products, the MOB shows the strongest seasonality with the RMSD higher than 4 psu in July and August (Fig. 6a), and close to 2 psu in winter. The spatially averaged deviation is much fresher than TP4, over -2 psu in summer and -0.5 psu in winter (Fig. 6b). The deviations of the two SMOS SSS show a relatively smaller seasonality (Fig. 6a).

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: northern

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: northern

Formatted: Font color: Text 1

Deleted: ,

Formatted: Font color: Text 1

Formatted: Font color: Text 1

During summer months, their RMSDs reach 1.5 psu (Fig. 6a) in summer, and they decrease to 0.5 and 1.0 psu (for BEC and CEC respectively). Throughout the whole year, the BEC RMSDs (Fig. 6a) are consistently smaller than that of CEC, and the seasonal cycles are different. This shows that the BEC SSS is closest to TP4, although it is overall fresher in the Summer.

4. Evaluation against in-situ observations

The misfits of the six SSS products from SMOS, CMEMS and climatologies are calculated as in Eqs. (1) and (2) against the pointwise in-situ observations described in Section 2.3. For TP4, the SSS evaluation is conducted on the same model day as the in-situ observations. Owing to the fact that the SSS from BEC, CEC and MOB are averaged over either 9 days or one week (see Table 1), the product dates at the center of the averaging window lag 5 or 4 days compared to the observation date. For PHC and WOA, the in-situ observations are sorted to monthly bins and evaluated for each month. The quantitative evaluation is divided into two main sections starting with dependent and then independent observations.

4.1 Against SSS from CORA5.1

As shown in Fig. 1a, the distribution of SSS observations from CORA5.1 over the Arctic is very inhomogeneous during the three years. Due to this, the evaluation of the gridded SSS products against in-situ observations is restricted to the observation-rich regions. The SSS misfits bias and RMSD for the six products are reported in Table 2 according to the eight Arctic sub-regions defined previously (Figure 1a). In this study, the Arctic domain (>60°N) is the core region for evaluation, divided into five sub-regions numbered from S0 to S4. It contains the central Arctic (sub-regions S0, S1, S2, and S3) and the Nordic Seas (S4). The regions from S5 to S7 are in the northern North Atlantic. The observations are displayed on scatterplots (Figure 7 and 8) to exhibit their uncertainties for fresh and saline waters in different areas.

• Central Arctic

Figure 7 shows the SSS products compared with discrete observations in the central Arctic. The observed SSS in S0 and S1 are mainly from the ITP at a minimal depth of 8 m. Around the North Pole (S0), where the satellite SSS are absent, the TP4 reanalysis and MOB reprocessing show opposite biases: +0.48 psu and -0.52 psu respectively (Table 2). The two climatologies used by them, PHC and WOA

Formatted: Font color: Text 1

Deleted: (sub-regions S0, S1, S2, and S3).

Formatted: Font color: Text 1

respectively, also show opposite biases. Considering the latter climatologies, both SSS scatterplots shows a fresh bias for high salinity water (>33 psu) and a saline bias for low salinity water (<31 psu). In the Canadian basin (in S1), the two climatological SSSs show an obvious gap in comparison to the ITP observations. Comparing to the fresh in-situ SSS from 24 to 30 psu, the PHC has strong saline bias (from 2 to more than 5 psu). On the other hand, the WOA shows both a fresh bias for relatively high salinity water (>28 psu) and saline bias for fresher water (<26 psu). Owing to the different time periods (Table 1) of the in-situ data they used, this result confirms the freshening of the Canadian basin since in the 1990s (Morison et al., 2012). In the S1 sub-region, the satellite SSS from BEC and CEC have only 20 and 42 data points for evaluation respectively. The resulting scatterplots show a significantly positive salinity bias (>4 psu) for fresh waters (<27 psu). For relatively higher salinity water (> 27 psu), the CEC has a stronger saline bias than the BEC. In the Kara Sea (sub-region S2), the TP4 SSS has the smallest RMSD at 1.7 psu, which is significantly smaller than other products. The scatterplot also shows a good linear relationship between the TP4 and the in-situ SSS, while other products generally show fresh biases, indicating that the SSS variability in the Kara Sea is well captured by TP4. In the Barents Sea (sub-region S3), TP4 gives as well the smallest misfit (RMSD: 0.34 psu; bias: -0.14 psu). The SSS scatterplots exhibits linear relationships for all products except the CEC, which underestimates the Atlantic water SSS.

- *Northern North Atlantic and Nordic Seas*

Figure 8 shows the paired scatterplots of the six SSS products in the subpolar seas from sub-regions S4 to S7 (see Fig. 1a). In S4 and S5, the bias of SSS products is relatively small, less than 0.15 psu (Table 2), except for CEC in S4 and TP4 in S5, both too saline by 0.2 psu. The scatterplots further indicate that low salinity waters are too saline in all SSS products in S4 (<31 psu) and in S5 (<28 psu). Meanwhile, the respective bias and RMSD of the SSS products are less than 0.1 psu and 0.43 psu respectively, except for the CEC in S6 and S7. The MOB SSS has the smallest salinity bias. Among the eight regions compared here (S0 to S7), the SSS bias is lowest in S6 (Irminger Sea).

386 Over the northern North Atlantic and the Nordic seas, Fig. 9 shows maps of the mean
 387 SSS deviation for each product during the period 2011-2013. Considerable negative
 388 biases (<-0.2 psu) are found in the CEC, whereas the MOB and WOA have the
 389 smallest bias, less than 0.02 psu (Fig. 9 d, e, f). The SSS products from BEC, TP4
 390 and PHC (Fig. 9 a, b, c) have slightly higher bias (~0.05 psu) in comparison to the
 391 MOB and WOA. On average, the BEC bias is only -0.04 psu, much smaller than that
 392 of the CEC (<-0.2 psu). Focusing on the BEC SSS, Fig 9a shows that while a fresh
 393 bias dominates the Nordic Seas, the product is too saline in the northern North
 394 Atlantic.

395 The inter-comparison of the biases against the in-situ data in Fig. 9a and 9b exhibits
 396 two strong positive biases of TP4 along the Norwegian coast and along the West
 397 Greenland coast. Notably, the BEC has smaller bias along both coasts, although it
 398 has a slightly saline bias offshore. This indicates potential benefits of the BEC SSS
 399 for the TOPAZ system along the Norwegian and Greenland coasts, were it
 400 successfully assimilated into the system. Figure 10 shows RMSDs of SSS for all the
 401 products over the northern North Atlantic Ocean and the Nordic Seas. On average,
 402 the largest uncertainty is found with the CEC (~1.0 psu; Fig. 10d), with RMSDs as
 403 large as 1.5 psu in the Greenland Sea and the Barents Sea. The SSS RMSDs for the
 404 five other SSS products are much smaller (~0.5 psu).

406 4.2 Independent SSS in the Beaufort Sea

407 Independent in-situ data from BGEP and CLIVAR are used during the summer
 408 months of 2011-2013 in the Beaufort Sea for the evaluation of the six SSS products
 409 (Fig. 11). The in-situ SSS observations range from 15 to 32 psu. The range of BEC
 410 SSS is limited to 24 to 31 psu with a minor bias of 0.09 psu and a RMSD of 1.82 psu.
 411 On the other hand, the range of TP4 SSS increases from 19 to 32 psu, with a larger
 412 saline bias of 2.59 psu and a RMSD of 3.63 psu. The linear regression coefficients
 413 for BEC and TP4 are 0.57 and 0.07 respectively. Looking at the low-salinity
 414 observations (~27 psu) collected at (136.4°W, 70.5°N) on 15th August 2011, marked
 415 by anti-triangles (Fig. 1b) near the Mackenzie River estuary, TP4 has a significant
 416 negative bias (< -4 psu) visible as the outliers above the dashed-black line in Fig.
 417 11a. This hints to a lack of fresh water signatures from river discharge.
 418 The range of PHC SSS climatology is only reaching from 24 to 31 psu, similar to
 419 TP4, with a saline bias of 1.65 psu and RMSD of 2.85 psu. Compared to the TP4

Deleted: and the North Sea

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: is even shorter

Formatted: Font color: Text 1

Deleted: large

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

deviation at the Makenzie River basin, the PHC saline bias is present, but smaller. The strong positive bias in TP4 at these points can then be partly attributed to the SSS relaxation of the TOPAZ model towards the PHC climatology, albeit rather weak. The range of the WOA is much wider, from 12 to 31 psu. Among the six products, the WOA bias is the smallest (~ 0.02 psu) over the Beaufort Sea during all three summers. However, it should be noted that the variability of in-situ observations is very large for salinities lower than 24 psu, which contributes to the large RMSD (> 3.0 psu) of both PHC and WOA. It confirms that the two climatologies have a sizable uncertainty over low salinity regions (< 24 psu) in the Arctic Ocean. The CEC SSS ranges from 13 psu to 34 psu, which is much wider than the range of the BEC SSS. The saline bias of CEC is however larger at 2.38 psu and its RMSD is about quite large at 3.77 psu. Furthermore, the CEC deviations from the in-situ observations are larger in waters fresher than 27 psu. The MOB combined product performs poorly with the largest negative bias (> 5 psu) and an RMSD in excess of 8 psu. In contrast to the other five SSS products, the anomalously fresh SSS observed around the point (140°W , 71°N) near the Mackenzie River estuary are represented by even fresher values of 12 psu in MOB, which may hint at an amplification of the anomalies.

In order to characterize the dependency of the bias on the SSS values for the six SSS products, we used the in-situ data, plotting their absolute differences as a function of observed SSS in Fig. 12. In general, all products show considerable deviations as high as 8 to 14 psu. While the absolute misfits of most SSS products increase monotonically with lower salinity, the bias of MOB shows a peak around 20 psu (Fig. 12c). A fourth-order polynomial function,

$$F(S) = p_1 S^4 + p_2 S^3 + p_3 S^2 + p_4 S + p_5, \quad (3)$$

is then fitted to the absolute bias for each SSS product, where S represents the in-situ salinity. The fitting coefficients, p_1 to p_5 , are listed in Table 3 for each product. The norm residuals are displayed on each panel in Fig. 12 and clearly show that the fitting for MOB has the largest uncertainty, while the minimal norm residuals are about 10 and 7 psu^2 respectively for BEC and TP4. This suggests the derived fitting curves for BEC and TP4 have relatively credible skill characterizing the error distribution as a function of the observed SSS. Both curves decrease with increasing salinity above 28 (30) psu for BEC (TP4) and increase slightly afterwards. The

- Formatted ... [1]
- Deleted: dependencies
- Formatted ... [2]
- Deleted: against
- Formatted ... [3]
- Deleted: are plotted
- Formatted: Font color: Text 1
- Deleted: with the maxima reaching
- Formatted: Font color: Text 1
- Deleted: of the
- Formatted ... [4]
- Deleted: increase towards
- Formatted: Font color: Text 1
- Deleted: its
- Formatted: Font color: Text 1
- Deleted: shown in
- Formatted: Font color: Text 1
- Deleted: . The
- Formatted: Font color: Text 1
- Deleted: curve
- Formatted: Font color: Text 1
- Formatted ... [5]
- Deleted: of the
- Formatted: Font color: Text 1
- Deleted: products
- Formatted: Font color: Text 1
- Deleted: for each product
- Formatted: Font color: Text 1
- Deleted: .
- Formatted ... [6]
- Deleted: in
- Formatted: Font color: Text 1
- Deleted: its
- Formatted: Font color: Text 1
- Deleted: monotonically
- Formatted: Font color: Text 1
- Deleted: towards the
- Formatted: Font color: Text 1
- Deleted: higher than
- Formatted: Font color: Text 1

494 absolute bias in TP4 is consistently larger than that in BEC. The fitted curves of PHC
495 and WOA have similar functional forms to TP4 and BEC, but with lower amplitudes.

497 5. Conclusions

498 To understand the uncertainties in the Arctic SSS, our study evaluates two gridded
499 SMOS SSS products (BEC and CEC), two CMEMS products (TP4 and MOB), and
500 two climatology products (PHC and WOA) by mutual inter-comparison and
501 comparisons against both dependent and independent in-situ datasets during the
502 years 2011-2013.

503 The differences in spatial coverage of the two SMOS SSS were shown in the monthly
504 mean (Fig. 2 and Fig.3), due to the different retrievals applied in these two datasets.

505 The spatial distributions of SSS from TP4 and PHC are close to each other, due to
506 the relaxation of TOPAZ model towards PHC. Relative to TP4, the SSS deviations of
507 the four products (BEC, MOB, WOA and PHC) in summer show similar magnitude
508 over open waters. On the contrary, the CEC SSS shows a negative bias (<-1 psu)
509 over the region extending from Iceland towards the western side of Ireland (Fig. 4, 5),
510 but the BEC SSS has a slightly but clear negative bias over the region. In general,
511 the most significant differences in the SSS deviations relative to TP4 are found under
512 the sea-ice cover and in its surrounding marginal ice zones.

513 Furthermore, the intercomparison of the SSS products shows that the BEC SSS in
514 August and September (Fig. 4, 5) has consistent negative deviations along the sea-
515 ice edge in the Beaufort Sea and the Chukchi Sea, but the CEC SSS has opposite
516 deviations in these two months. Thus, it seems that the two SMOS products would
517 give rise to significantly different effects to the upper ocean state, were they
518 assimilated.

519 Focusing on the wider Arctic domain (>60°N), the deviations of the five SSS products
520 relative to TP4 show diverse seasonal characteristics (Fig. 6). Although the BEC and
521 CEC SSS products show similar deviations of 1.5 psu (Fig. 6a) in summer, the BEC
522 deviations in winter are clearly lower (~0.5 psu). The deviations of MOB and WOA
523 (Fig. 6a) vary from over 1.5 psu in winter to around 4 psu in summer, so all are in
524 considerable disagreement with TP4. Consequently, our intercomparison suggests
525 that the BEC SSS has more consistent pattern with the TP4 SSS among the SSS
526 products compared here.

- Deleted: the
- Formatted ... [7]
- Deleted: the
- Formatted ... [8]
- Deleted: their
- Formatted ... [9]
- Deleted: of
- Formatted ... [10]
- Deleted: of
- Formatted ... [11]
- Deleted: the
- Formatted ... [12]
- Deleted: clearly
- Formatted ... [13]
- Deleted: retrieval
- Formatted ... [14]
- Deleted: considerably
- Formatted ... [15]
- Deleted: mainly as for the fact that the SSS in the
- Formatted ... [16]
- Deleted: is relaxed
- Formatted ... [17]
- Deleted: at each time step
- Formatted ... [18]
- Deleted: the
- Formatted ... [19]
- Deleted: the
- Formatted ... [20]
- Deleted: clearly
- Formatted ... [21]
- Deleted: may be arguable...eems that the two SMOS
- Formatted ... [23]
- Deleted: wide
- Formatted ... [24]
- Deleted: of BEC and CEC have the
- Formatted ... [25]
- Deleted: deviation
- Formatted ... [26]
- Deleted: about
- Formatted ... [27]
- Deleted: months
- Formatted ... [28]
- Deleted: varies
- Formatted ... [29]
- Deleted: which suggests a
- Formatted ... [30]
- Deleted: gap
- Formatted ... [31]
- Deleted: the
- Formatted ... [32]
- Deleted: the
- Formatted ... [33]
- Deleted: the most
- Formatted ... [34]
- Deleted: all other
- Formatted ... [35]

593 The in-situ data from CORA5.1, which were used in both TP4 and MOB, has been
 594 used for evaluation of the six SSS products in eight sub-regions (Fig. 1a). These
 595 were divided into two parts: the central – seasonally ice covered - Arctic Ocean and
 596 the open ocean areas (the northern North Atlantic Ocean and the Nordic Seas). Due
 597 to limited coverage of BEC and CEC in S1, the scatterplots (Fig. 7) show a positive
 598 saline bias (>4 psu) for low salinity water (< 27 psu). However, the salinity bias of
 599 BEC is slightly reduced for relatively higher salinity water (> 27 psu). In the Kara Sea
 600 and the Barents Sea, the TP4 SSS has minimal RMSD compared with others (Table
 601 2). The BEC scatterplots in S2 and S3 (Fig. 7) are similar to TP4.
 602 In the northern North Atlantic Ocean and the Nordic Seas (Fig. 8), the scatterplots of
 603 the CEC SSS show that it underestimates the Atlantic water salinity, which is also
 604 consistent with the intercomparison results (low salinity deviation) shown in Fig. 4
 605 and 5. The misfits of mean and RMSDs shown in Fig. 9 and 10, suggest the CEC
 606 SSS has considerable uncertainty (RMSD of about 1 psu), especially in the Nordic
 607 Seas with an obvious low salinity bias. By comparison, the SSS uncertainties of BEC
 608 are significantly lower than CEC, and are equivalent to both TP4 and PHC. Two
 609 notable regions, where the BEC SSS has lower uncertainties than TP4 against the
 610 in-situ observations, are along the Norwegian coast and near the west coast of
 611 Greenland. It is reasonable to expect that they should benefit the most if the BEC
 612 SSS were successfully assimilated into the TOPAZ system.
 613 Against independent in-situ observations from BGEP and CLIVAR, the SSS
 614 evaluation in the Beaufort Sea is performed in three successive summers. The linear
 615 regression against these independent SSS observations (Fig. 11) shows that the
 616 BEC SSS has the smallest RMSD of 1.8 psu with a positive bias of 0.1 psu, and the
 617 CEC SSS has larger RMSD of about 3.8 psu with a larger positive bias of 2.4 psu
 618 (Fig. 11). On the other hand, the TP4 SSS also shows large RMSD of about 3.6 psu
 619 with large positive bias of 2.6 psu. These are smaller than MOB which has the RMSD
 620 of 8.2 psu and a larger negative bias (-5.0 psu). As for the two climatology products,
 621 the RMSDs of WOA and PHC are both above 2.8 psu, but with significantly smaller
 622 bias in WOA. More specifically, the poor fit of all products is attributed to large
 623 product-observation mismatches against in-situ salinity observations below 24 psu,
 624 which are located over the continental shelf near the estuary of the Mackenzie River.
 625 In order to characterize the product-data misfits of all six products against in-situ
 626 data, a 4th order polynomial is fitted to the absolute deviation as a function of the

- Deleted: Against the
- Formatted ... [36]
- Deleted: the quantitative evaluations
- Formatted ... [37]
- Deleted: were investigated
- Formatted ... [38]
- Deleted: the
- Formatted ... [39]
- Deleted: It was
- Formatted ... [40]
- Deleted: in
- Formatted ... [41]
- Deleted: ;
- Formatted ... [42]
- Deleted: .
- Formatted ... [43]
- Deleted: the
- Formatted ... [44]
- Deleted: relative high
- Formatted ... [45]
- Deleted: the
- Formatted ... [46]
- Deleted: have
- Formatted ... [47]
- Deleted: distributions with respect
- Formatted ... [48]
- Deleted: S6, S4, and S3;
- Formatted ... [49]
- Deleted: is
- Formatted ... [50]
- Deleted: biases. On the other hand
- Formatted ... [51]
- Deleted: the
- Formatted ... [52]
- Deleted: in comparison to the
- Formatted ... [53]
- Deleted: but
- Formatted ... [54]
- Deleted: compared with
- Formatted ... [55]
- Deleted: referred to
- Formatted ... [56]
- Deleted: than the TP4
- Formatted ... [57]
- Deleted: Island
- Formatted ... [58]
- Deleted: are
- Formatted ... [59]
- Deleted: beneficial region in the Nordic Seas
- Formatted ... [60]
- Deleted: is
- Formatted ... [61]
- Deleted: the
- Formatted ... [62]
- Deleted: of the three years. ¶
- Formatted ... [63]
- Deleted: function
- Formatted ... [64]
- Formatted ... [65]

711 observed salinity (Fig.12). The absolute deviations of most of the products except the
 712 MOB decrease monotonically with increasing salinity. The norm residuals for TP4
 713 and BEC and are the smallest among all six products with 10.2 and 7.0, respectively.
 714 The fitted curve reaches its smallest value of below 1.0 psu for an in-situ salinity of
 715 28 psu and 30 psu for BEC and TP4 respectively. Both the fitted curves for CEC and
 716 MOB have large norm residuals of 18.1 and 68.8 psu² respectively. Note that special
 717 attention must be paid in usage of MOB in the Arctic Ocean due to a large negative
 718 bias and high RMSD in regions where the product is based on a limited number of
 719 observations.
 720 The above evaluations suggest that certain benefit can be expected in assimilating
 721 the BEC SSS into the TOPAZ Arctic ocean analysis-forecast system. The knowledge
 722 of the error structure in the SSS products provided in this study will serve as an input
 723 to the observation error for the SMOS product, as required by data assimilation. The
 724 poor spatial coverages of CORA in situ data in the Arctic Ocean beg for more data -
 725 especially from the Arctic Ocean marginal seas - to be compiled from independent
 726 data source to validate the SMOS SSS products. In addition, when comparing the
 727 two climatology products, PHC and WOA, the SSS scatterplots of the PHC in the
 728 central Arctic (Fig. 7) reveal a saline bias for low salinity waters. Considering that
 729 PHC does not include the two more recent decades of data (Table 1), this confirms
 730 that the freshening in the Canadian Basin since the 1990s is rather significant as
 731 discussed by Morison et al. (2012). Based on this, the next TOPAZ system will use
 732 WOA in replacement of PHC as target relaxation data.

734 Acknowledgement

735 The authors acknowledge the support of CMEMS for the Arctic MFC and funding
 736 from the European Space Agency project Arctic+Salinity. Grants of computing time
 737 (nn2993k and nn9481k) and storage (ns2993k) from the Norwegian Sigma2
 738 infrastructures are gratefully acknowledged. The BEC SSS is provided by the
 739 Barcelona Expert Centre (bec.icm.csic.es), Spain. The CEC SSS is distributed by the
 740 Ocean Salinity Expertise Center (CECOS) of CATDS at IFREMER, France.

742 Reference:

Formatted	... [66]
Deleted: for	
Formatted	... [67]
Deleted: decrease as observed	
Formatted	... [68]
Deleted: increases	
Formatted	... [69]
Deleted: of	
Formatted	... [70]
Deleted: , among all six products and the fitted curves	711
Formatted	... [72]
Deleted: less than	
Formatted	... [73]
Deleted: at the	
Formatted	... [74]
Deleted: its	
Formatted	... [75]
Deleted: the	
Formatted	... [76]
Deleted: Evaluations of the SSS products against TP4	717
Formatted	... [78]
Deleted: show salinity	
Formatted	... [79]
Deleted: saline water.	
Formatted	... [80]
Deleted: different time periods	
Formatted	... [81]
Deleted: their compiled in-situ	
Formatted	... [82]
Deleted: sources	
Deleted: it independently verifies	
Formatted	... [83]
Formatted	... [84]
Deleted: Canada	
Formatted	... [85]
Deleted: evaluation	
Formatted	... [86]
Deleted: the	
Formatted	... [87]
Deleted: to replace the	
Formatted	... [88]
Deleted: the	
Formatted	... [89]
Deleted: field	
Formatted	... [90]
Deleted: partly	
Formatted	... [91]
Deleted: in response to the	
Formatted	... [92]
Deleted: project (Dec 2018-June 2020).	
Formatted	... [93]
Deleted: produced	
Formatted	... [94]
Deleted:) mainly funded by the Spanish National	741
Formatted	... [96]

830 Bertino, L., and Lisæter, K. A.: The TOPAZ monitoring and prediction system for the Atlantic
831 and Arctic Oceans, *Journal of Operational Oceanography*, 1(2), 15–19, doi:
832 10.1080/1755876X.2008.11020098, 2008

833 Boutin J., Vergely J.L., Marchand S., D'Amico F., Hasson A., Kolodziejczyk N., Reul N.,
834 Reverdin G., Vialard J.: New SMOS Sea Surface Salinity with reduced systematic errors
835 and improved variability. *Remote Sensing of Environment*, 214, 115-134,
836 <http://doi.org/10.1016/j.rse.2018.05.022>, 2018.

837 Buongiorno Nardelli, B., Droghei, R., and Santoleri, R.: Multi-dimensional interpolation of
838 SMOS sea surface salinity with surface temperature and *in situ* salinity data. *Remote Sens.*
839 *Environ.* 180, 392–402. doi: 10.1016/j.rse.2015.12.052, 2016

840 Cabanes, C., A. Grouazel, K. von Schuckmann, M. Hamon, V. Turpin, C. Coatanoean, F. Paris,
841 S. Guinehut, C. Boone, N. Ferry, C. de Boyer Montégut, T. Carval, G. Reverdin, S.
842 Pouliquen, and P. Y. Le Traon: The CORA dataset: validation and diagnostics of in-situ
843 ocean temperature and salinity measurements. *Ocean Science*, 9, 1-
844 18, <http://www.ocean-sci.net/9/1/2013/os-9-1-2013.html>, doi:10.5194/os-9-1-2013, 2013

845 Chassignet, E. P., Smith, L. T., and Halliwell, G. R.: North Atlantic Simulations with the Hybrid
846 Coordinate Ocean Model (HYCOM): Impact of the vertical coordinate choice, reference
847 pressure, and thermobaricity, *J. Phys. Oceanogr.*, 33, 2504-2526. Doi:
848 [http://dx.doi.org/10.1175/1520-0485\(2003\)033<2504:NASWTH>2.0.CO;2](http://dx.doi.org/10.1175/1520-0485(2003)033<2504:NASWTH>2.0.CO;2), 2003.

849 de Boyer Montegut, C., G. Madec, A. Fischer, A. Lazar, and D. Iudicone: Mixed Layer Depth
850 over the Global Ocean: An Examination of Profile Data and a Profile-Based Climatology.
851 *J. Geophys. Res.*, 109 (C12003), 1–20, doi:10.1029/2004JC002378, 2004.

852 D'Addezio, J. M., and Subrahmanyam, B.: Sea surface salinity variability in the Agulhas
853 Current region inferred from SMOS and Aquarius. *Remote Sensing of Environment*, 180,
854 440–452. doi:10.1016/j.rse.2016.02.006, 2016

855 Drange, H. and Simonsen, K.: Formulation of air-sea fluxes in the ESOP2 version of MICOM,
856 Technical Report No. 125 of Nansen Environmental and Remote Sensing Center, 1996.

857 Droghei, R., Buongiorno Nardelli, B., and Santoleri, R.: A new global sea surface salinity and
858 density dataset from multivariate observations (1993–2016). *Front. Mar. Sci.* 5:84. doi:
859 10.3389/fmars.2018.00084, 2018

860 ESA, SMOS data products, available from
861 <https://earth.esa.int/documents/10174/1854456/SMOS-Data-Products-Brochure> (last
862 access on 12th December 2018), November 2017.

863 Font, J., Camps, A., Borges, A., Martín-Neira, M., Boutin, J., Reul, N., Kerr, Y. H., Hahne, A.,
864 and Mecklenburg, S.: SMOS: The challenging sea surface salinity measurement from
865 space, *Proc. IEEE*, 98(5), 649–665, DOI: 10.1109/JPROC.2009.2033096, May 2010.

Formatted: Font: Italic

Formatted: Font color: Text 1

Deleted: Nicolas,

Formatted: Font color: Text 1

Deleted: Nicolas,

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

868 Furue, R., Takatama, K., Sasaki, H., Schneider, N., Nonaka, M., and Taguchi, B.: Impacts of
869 sea-surface salinity in an eddy-resolving semi-global OGCM. *Ocean Modelling*, 122, 36–
870 56, doi:10.1016/j.ocemod.2017.11.004, 2018

871 Hátún, H., Sandø, A. B., Drange, H., Hansen, B., and Valdimarsson, H.: Influence of the
872 Atlantic Subpolar Gyre on the Thermohaline Circulation, *Science*, 309 (5742), 1841-1844,
873 doi:10.1126/science.1114777, 2005

874 Hunke, E. C., and Dukowicz, J. K.: An elastic-viscous-plastic model for sea ice dynamics, *J.*
875 *Phys. Oceanogr.*, 27, 1849-1867, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0485(1997)027<1849:AEVPMF>2.0.CO;2)
876 [0485\(1997\)027<1849:AEVPMF>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<1849:AEVPMF>2.0.CO;2), 1997.

877 Johnson, G. C., Schmidtko, S., and Lyman, J. M.: Relative contributions of temperature and
878 salinity to seasonal mixed layer density changes and horizontal density gradients, *J.*
879 *Geophys. Res.*, 117, C04015, doi:10.1029/2011JC007651, 2012

880 Kerr, Y. H., Waldeufel, P., Wigneron, J. P., Delwart, S., Cabot, F., Boutin, J., Escorihuela, M.
881 J., Font, J., Reul, N., Gruhier, C., Juglea, S., Drinkwater, M. R., Hahne, A., Martín-Neira,
882 M., and Mecklenburg, S.: The SMOS mission: New tool for monitoring key elements of the
883 global water cycle, *Proc. IEEE*, 98(5), 666–687, doi:10.1109/JPROC.2010.2043032, 2010.

884 Kolodziejczyk, N., Boutin, J., Vergely, J.-L., Marchand, S., Martin, N., and Reverdin, G.:
885 Mitigation of systematic errors in SMOS sea surface salinity. *Remote Sensing of*
886 *Environment*, 180, 164–177. doi:10.1016/j.rse.2016.02.061, 2016

887 Latif, M., Roeckner, E., Mikolajewicz, U., and Voss R.: Tropical stabilization of the thermohaline
888 circulation in a greenhouse warming simulation, *J. Clim.*, 13: 1809–1813, 2000.

889 Macdonald, R. W., Carmack, E. C., McLaughlin, F. A., Falkner, K. K., and Swift, J. H.:
890 Connections among ice, runoff and atmospheric forcing in the Beaufort Gyre. *Geophys.*
891 *Res. Lett.*, 26, 2223–2226, 1999

892 Maes, C., Ando, K., Delcroix, T., Kessler, W. S., McPhaden, M. J., and Roemmich,
893 D.: Observed correlation of surface salinity, temperature and barrier layer at the eastern
894 edge of the western Pacific warm pool, *Geophys. Res. Lett.*, 33, L06601,
895 doi:10.1029/2005GL024772, 2006.

896 Mathis, J. T., and Monacci, N. M.: Carbon Dioxide and Hydrographic data obtained during the
897 USCGC Healy Cruise HLY1203 in the Arctic Ocean (October 05 - 25, 2012). Available
898 from <http://cdiac.ess-dive.lbl.gov/ftp/oceans/CARINA/Healy/HLY-12-03/>. Oak Ridge
899 National Laboratory, US Department of Energy, Oak Ridge, Tennessee. doi:
900 10.3334/CDIAC/OTG.CLIVAR_33HQ20121005, 2014.

901 McPhee, M. G., Stanton, T. P., Morison, J. H. and Martinson, D. G.: Freshening of the upper
902 ocean in the Arctic: is perennial sea ice disappearing? *Geophys. Res. Lett.* 25, 1729–1732,
903 1998.

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

904 Mecklenburg, S., Drusch, M., Kerr, Y. H., Font, J., Martín-Neira, M., Delwart, S., Buenadicha,
 905 G., Reul, N., Daganzo-Eusebio, E., Oliva, R., and Crapolicchio, R.: ESA's soil moisture
 906 and ocean salinity mission: Mission performance and operations. *IEEE TGARS*, 50(5),
 907 1354–1366, DOI: [10.1109/TGRS.2012.2187666](https://doi.org/10.1109/TGRS.2012.2187666), 2012

908 Mignot, J., and Frankignoul, C.: On the interannual variability of surface salinity in the Atlantic.
 909 *Climate Dynamics*, 20(6), 555–565. doi:10.1007/s00382-002-0294-0, 2003

910 Morison, J., Kwok, R., Peralta-Ferriz, C., Alkire, M., Rigor, I., Andersen, R., and Steele, M.:
 911 Changing arctic ocean freshwater pathways. *Nature*, 481:66–70, 2012.

912 Olmedo, E., Gabarró, C., González-Gambau, V., Martínez, J., Ballabrera-Poy, J., Turiel, A.,
 913 Portabella, M., Fournier, S., and Lee, T.: Seven Years of SMOS Sea Surface Salinity at
 914 High Latitudes: Variability in Arctic and Sub-Arctic Regions. *Remote Sensing*. 2018;
 915 10(11):1772, <https://doi.org/10.3390/rs10111772>, 2018.

916 Reverdin, G., Cayan, D., and Kushnir, Y.: Decadal variability of hydrography in the upper
 917 northern North Atlantic in 1948–1990. *J. Geophys. Res.*, 102, 8505–8531,
 918 <https://doi.org/10.1029/96JC03943>, 1997

919 Sakov, P., and Oke, P. R.: A deterministic formulation of the ensemble Kalman Filter: an
 920 alternative to ensemble square root filters. *Tellus A*, 60(2), 361-371, doi:10.1111/j.1600-
 921 0870.2007.00299.x, 2008.

922 Simmons, A., Uppala, S., Dee, D., and Kobayashi, S.: ERA-Interim: New ECMWF reanalysis
 923 products from 1989 onwards. *ECMWF Newsletter*, 110. 25-35, doi:10.21957/pocnex23c6,
 924 2007.

925 SMOS Team, SMOS L2 OS Algorithm Theoretical Baseline Document, ESA, Paris, France,
 926 SO-TN-ARG-GS-0007, version 3.13, available from
 927 https://earth.esa.int/documents/10174/1854519/SMOS_L2OS-ATBD, (last access: 12th
 928 December 2018), 29th April 2016.

929 SMOS-BEC Team, Quality Report: Validation of SMOS-BEC experimental sea surface salinity
 930 products in the Arctic Ocean and high latitudes Oceans. Years 2011-2013, Barcelona
 931 Expert Centre, Spain. Technical note: BEC-SMOS-0007-QR version 1.0, available at
 932 <http://bec.icm.csic.es/doc/BEC-SMOS-0007-QR.pdf> (last access: 13th December 2018),
 933 4th March 2016.

934 Steele, M. and W. Ermold, Salinity Trends on the East Siberian Shelves, *Geophysical*
 935 *Research Letters*, Vol. 31, L24308, doi:10.1029/2004GL021302, 2004.

936 Steele, M., R. Morley, and W. Ermold, PHC: A global ocean hydrography with a high-quality
 937 Arctic Ocean, *Journal of Climate*, 14, 2079-2087, 2001.

938 Sumner, D., and Belaine, G.: Evaporation, Precipitation, and Associated Salinity Changes at
 939 a Humid, Subtropical Estuary. *Estuaries*, 28(6), 844-855. Retrieved from
 940 <http://www.jstor.org/stable/3526951>, 2005.

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: (2004)

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

942 Supply A, Boutin J, Vergely J-L, et al.: Precipitation Estimates from SMOS Sea-Surface
 943 Salinity. *Q. J. R. Meteorol. Soc.*, 144 (Suppl. 1):103–119, <https://doi.org/10.1002/qj.3110>,
 944 2018.

945 Talley, L. D., Johnson, G. C., Purkey, S., Feely, R. A., and Wanninkhof, R.: Global Ocean
 946 Ship-based Hydrographic Investigations Program (GO-SHIP) provides key climate-
 947 relevant deep ocean observations, *US CLIVAR Variations*, 15 (2), available from
 948 <https://www.pmel.noaa.gov/pubs/PDF/tall4659/tall4659.pdf> (last access: 19th December
 949 2018), 2017.

950 Toole, J. M., Krishfield, R. A., Timmermans, M.-L., and Proshutinsky, A.: The Ice-Tethered
 951 Profiler: Argo of the Arctic, *Oceanography*, 24, 126–135, doi:10.5670/oceanog.2011.64,
 952 2011.

953 Tseng, Y., Bryan, F. O., and Whitney, M. M.: Impacts of the representation of riverine
 954 freshwater input in the community earth system model. *Ocean Modelling*, 105, 71–86.
 955 doi:10.1016/j.ocemod.2016.08.002, 2016.

956 Uotila, P., Goosse, H., Haines, K., Chevallier, M., Barthélemy, A., Bricaud, C., Carton, J.,
 957 Fučkar, N., Garric, G., Iovino, D., Kauker, F., Korhonen, M., Lien, V. S., Marnela, M.,
 958 Massonnet, F., Mignac, D., Peterson, A., Sadikn, R., Shi, L., Tietsche, S., Toyoda, T., Xie,
 959 J., and Zhang, Z.: An assessment of ten ocean reanalyses in the polar regions, *Clim.*
 960 *Dynam.*, <https://doi.org/10.1007/s00382-018-4242-z>, online first, 2018.

961 Vancoppenolle, M., Fichefet, T., and Goosse, H.: Simulating the mass balance and salinity of
 962 Arctic and Antarctic sea ice. 2. Importance of sea ice salinity variations. *Ocean Modelling*,
 963 27, 54–69. doi:10.1016/j.ocemod.2008.11.003, 2009.

964 Verbrugge, N., Mulet, S., Guinehut, S., Buongiorno Nardelli, B., and Droghei, R.: Quality
 965 information document for global ocean multi observation products
 966 multiobs_glo_phy_rep_015_002, CMEMS-MOB-QUID-015-002, v1.0, available from
 967 <http://cmems-resources.cls.fr/documents/QUID/CMEMS-MOB-QUID-015-002.pdf> (last
 968 access: 14th December 2018), February 2018.

969 Vialard, J., and P. Delecluse: An OGCM study for the TOGA decade: I. Role of salinity in the
 970 physics of the Western Pacific fresh pool, *J. Phys. Oceanogr.*, 28, 1071–1088, 1998.

971 Woodgate, R., Aagaard, K. and Weingartner, T.: Monthly temperature, salinity, and transport
 972 variability of the Bering Strait through flow. *Geophys. Res. Lett.*, 32, L04601, DOI:
 973 10.1029/2004GL021880, 2005.

974 Xie, J., Bertino, L., Counillon, F., Lisæter, K. A., and Sakov, P.: Quality assessment of the
 975 TOPAZ4 reanalysis in the Arctic over the period 1991–2013. *Ocean Science*, 13(1), 123–
 976 144. <http://doi.org/10.5194/os-13-123-2017>, 2017.

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: (1998),

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: .

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

979 Xie, J., Counillon, F., and Bertino, L.: Impact of assimilating a merged sea-ice thickness from
 980 CryoSat-2 and SMOS in the Arctic reanalysis, *The Cryosphere*, 12, 3671-3691,
 981 <https://doi.org/10.5194/tc-12-3671-2018>, 2018.

982 Xie, J., Counillon, F., Bertino, L., Tian-Kunze, X., and Kaleschke, L.: Benefits of assimilating
 983 thin sea-ice thickness from SMOS into the TOPAZ system. *The Cryosphere*, 10, 2745–
 984 2761. <http://doi.org/10.5194/tc-10-2745-2016>, 2016.

985 Yu, L.: A global relationship between the ocean water cycle and near-surface salinity, *J.*
 986 *Geophys. Res.*, 116, C10025, doi:10.1029/2010JC006937, 2011.

987 Zweng, M. M., Reagan, J. R., Antonov J. I., Locarnini, R. A., Mishonov, A. V., Boyer, T. P.,
 988 Garcia, H. E., Baranova, O. K., Johnson, D. R., Seidov, D., and Biddle, M. M.: World
 989 Ocean Atlas 2013, Volume 2: Salinity, Levitus, S. (Ed.), Mishonov, A., Technical Ed.
 990 NOAA Atlas NESDIS 74, 39pp, 2013.

991

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Italic, Font color: Text 1

Formatted: Font color: Text 1

[illegible]

Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [5] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [6] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [6] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 14: [6] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 15: [7] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 15: [8] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 15: [9] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 15: [10] Formatted	Authors	8/13/19 9:35:00 AM

Font color: Text 1

Page 15: [11] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [12] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [13] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [14] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [15] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [16] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [17] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [18] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [19] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [20] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [21] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [21] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [22] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼.....

Page 15: [22] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼.....

Page 15: [23] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [24] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [24] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [24] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [25] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [26] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [27] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [28] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [29] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [30] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [31] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [32] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [33] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [34] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [35] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 15: [35] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [36] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [36] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [37] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [38] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [39] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [40] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [41] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [41] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [42] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [43] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [44] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [45] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [46] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [47] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [48] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [49] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [49] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [50] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [50] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [51] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [52] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [53] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [54] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [55] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		
Page 16: [55] Formatted	Authors	8/13/19 9:35:00 AM
Font color: Text 1		

Page 16: [56] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [57] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [58] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [59] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [60] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [61] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [62] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [63] Deleted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

▼ Page 16: [64] Formatted	Authors	8/13/19 9:35:00 AM
---------------------------	---------	--------------------

Font color: Text 1

Page 16: [65] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [66] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [67] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [67] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [68] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [69] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [70] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [71] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼

Page 17: [72] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [73] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [74] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [75] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [76] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [76] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼

Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼

Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼

Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼

Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼

Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼

Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼

Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼		
Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM

▼		
Page 17: [77] Deleted	Authors	8/13/19 9:35:00 AM

▼		
Page 17: [78] Formatted	Authors	8/13/19 9:35:00 AM

Font color: Text 1

Page 17: [79] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [80] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [80] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [81] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [82] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [83] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [84] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [85] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [85] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [86] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [87] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [88] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [89] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [90] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [91] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [92] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [93] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [94] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Page 17: [95] Deleted	Authors	8/13/19 9:35:00 AM
-----------------------	---------	--------------------

▼.....

Page 17: [96] Formatted	Authors	8/13/19 9:35:00 AM
-------------------------	---------	--------------------

Font color: Text 1

Captions of Table and Figures:

Table 1. Details of the six products evaluated during 2011-2013.

Product	Data source	Resolution	Provider	Website or CMEMS id	Release year
BEC	SMOS	9 days; 25 km	Barcelona Expert Centre, Spain	http://bec.icm.csie.es	2018
CEC	SMOS	9 days; 25 km zonal	Ocean Salinity Expertise Center, IFREMER	FTP: ftp.ifremer.fr	2018
TP4	Reanalysis	Daily; 12~16 km	CMEMS	ARCTIC-REANALYSIS-PHYS-002-003	2015
MOB	In situ + SMOS	7 days; 1/4x1/4°	CMEMS	MULTIOBS_GLO_PHY_REP_015_002	2016
PHC	In situ (1950-1994)	Monthly; 1x1°	Polar Science Center, University of Washington	http://psc.apl.washington.edu/	2005
WOA	In situ (1955-2012)	Monthly; 1/4x1/4°	NODC, NOAA	https://www.nodc.noaa.gov/O_C5/woa13/	2013

Table 2. Misfits of SSS relative to in-situ CORA5.1 observations during 2011-2013 in each sub-region. Bold numbers denote the smallest error among the six products.

Region	Bias (psu)						RMSD (psu)					
	BEC	CEC	TP4	MOB	PHC	WOA	BEC	CEC	TP4	MOB	PHC	WOA
S0	-	-	.48	-.52	.48	-.11	-	-	1.25	1.78	1.28	.70
S1	4.03	3.18	3.29	1.63	3.29	.42	4.23	3.70	3.47	2.22	3.43	1.37
S2	-1.76	-.44	-.97	2.96	-3.30	-2.93	2.16	2.57	1.70	3.68	3.87	3.62
S3	-.14	-.70	-.14	-.21	-.29	-.25	.45	1.17	.34	.42	.51	.44
S4	-.09	-.20	.12	.11	-.02	.02	.91	1.21	.89	.86	.94	.84
S5	-.07	.06	.20	.01	.02	.07	1.47	1.52	1.42	1.44	1.39	1.30
S6	-.01	.15	.01	-.01	-.09	.05	.25	.66	.14	.12	.28	.16
S7	.05	.34	.04	-.03	-.23	-.03	.31	.88	.33	.22	.43	.27

Table 3. Optimal coefficients for the 4th order polynomial fit of the errors (see Eq. 3) as a function of in-situ SSS for each product.

Product	F(p ₁ , p ₂ , p ₃ , p ₄ , p ₅ , s)					Residual norm	In situ samples
	p ₁ (x10 ⁻³)	p ₂	p ₃	p ₄	p ₅		
BEC	0.168	-0.016	0.614	-11.345	87.097	7.03	91
CEC	0.225	-0.033	-1.550	-29.886	205.179	18.13	121
TP4	0.993	-0.096	3.430	-54.552	335.197	10.17	232
MOB	-1.080	0.128	-5.469	99.824	-645.087	68.81	163
PHC	1.257	-0.120	4.235	-65.938	388.808	13.98	232
WOA	-0.121	0.010	-0.322	3.998	-10.847	38.91	232




Fig. 1 (a): SSS locations of the in-situ observations north of 52°N in CORA5.1 during the years 2011-2013. 8 sub-regions divide the Arctic Ocean (S0-S4) and the northern North Atlantic Ocean (S5-S7), with the number of observations indicated in each region. **(b):** Independent SSS observations in the Beaufort Sea during the summer months of 2011-2013 from the BGEF (marked by anti-triangles, squares, and starts) and the CLIVAR (marked by triangles and crosses). Different colors (red, black and yellow) indicate the years (2011, 2012 and 2013 resp.).





Fig. 2 Monthly SSS (unit: psu) in March from satellite products (BEC and CEC, left column), reanalysis/reprocessing (TP4 and MOB, middle column), and climatology (PHC and WOA, right column). White areas are masked by sea ice. The thick brown line represents the sea ice edge (15% concentration from TP4), and the black shaded isolines represent the salinities of 33.6 and 35 psu near the surface.



Deleted:

Deleted: s

Deleted: the

Deleted: salinity




Fig. 3 Similar to previous figure in September.





Fig. 4 Deviations of monthly SSS (unit: psu) in August for (a) BEC; (b) PHC; (c) CEC; (d) MOB; and (e) WOA relative to TP4. The thick brown line represents sea ice edge (15% concentration from TP4), the black lines represent ± 1 psu.



Deleted:




Fig. 5 Same as previous for September.




Fig. 6 Monthly deviations in the Arctic Ocean (>60N) of (a) the RMS and (b) the spatial average during the period 2011-2013 for the five SSS products referred to TP4. The anti-triangle (triangle, circle, star and square) line represents the SSS deviations from BEC (CEC, MOB, PHC and WOA respectively).




Fig. 7 Scatterplots of SSS compared to the CORA5.1 in-situ observations with respect to the S0-S3 regions in the Arctic. The diamonds (anti-triangles, stars, squares, circles, and triangles) represents the SSS from TP4 (BEC, PHC, WOA, MOB, and CEC respectively). The black (red) lines are the linear regressions of the blue (purple) dots in each panel, and the coefficient R^2 between the evaluated product and the in-situ SSS is indicated in the panel together with the number of observations in parentheses.




Fig. 8 Same as Fig. 7 but for the subpolar regions S4-S7.




Fig. 9 The mean deviation of SSS for the six datasets compared to in situ observations from CORA 5.1 during the three years of 2011-2013 in the northern North Atlantic and the Nordic seas. The SSS observations are distributed into the coarse grid cells of 9x9 grids in TP4, with a gray mask if the valid observations less than 10.




Fig. 10 The Root Mean Square deviation of SSS for six datasets compared to in situ observations from CORA 5.1 during the three years of 2011-2013 in the northern North Atlantic and the Nordic seas. The SSS observations are distributed into the coarse grid cells of 9x9 grids in TP4, with a gray mask if the valid observations less than 10.




Fig. 11 Scatterplots of SSS compared to the in-situ observations in Beaufort Sea during the summer months of 2011-2013: (a) The diamond (anti-triangle) represents the SSS from TP4 (BEC) with blue (purple), and the linear regression is denoted by the dashed black (red) line. (b) The star (square) from the climatology of PHC (WOA). (c) The circle (triangle) represents from MOB (CEC). The coefficient R^2 is the squared linear relationship between the evaluated product and the in-situ SSS, and the misfits also shown on the panels.




Fig. 12 Scatterplots of SSS uncertainty compared to the in-situ observations in Beaufort Sea as a function of the observed salinity. The black dashed line marks 5 psu. (a) The diamonds (anti-triangles) represent TP4 (BEC) in blue (purple). (b) The stars (squares) are the PHC (WOA) climatology. (c) The circles (triangles) represent the MOB (CEC). The thick dashed curves are fitted by a fourth order polynomial, and the norm residuals are marked on each panel respectively.

Deleted: represents

Deleted: the absolute deviation of

Deleted: from

Deleted: climatology of

Deleted:) represents from