



**National
Oceanography Centre**
NATURAL ENVIRONMENT RESEARCH COUNCIL

Joseph Proudman Building
6 Brownlow Street
Liverpool
Merseyside, L3 5DA
United Kingdom

Tel: +44 (0) 151 795 4800
Fax: +44 (0) 151 795 4801

www.noc.ac.uk

Editor, Ocean Science

26 June 2018

Dear Sir,

Paper Submitted to Ocean Science

Many thanks for the reviewer comments on our paper submitted to Ocean Science, which appeared in OS Discussions, entitled "The Nodal Dependence of Long-Period Ocean Tides in the Drake Passage" (os-2018-50). They were very useful. I have uploaded our replies to the comments and these are included again below

We have made a number of small changes to the text following the comments, and a few others of our own. The only major change has been to Section 3.2 which discusses Vernadsky data. That has been rewritten and expanded, and there are extra figures to do with that (Figures 3c,d and 8c,h). We hope there are not now too many figures, but most of them can be printed fairly small. In addition, we have added a Table 1 to summarise the many numbers given in the text.

Also below is a Word comparison file which shows where changes have been made. The Supplementary Information file is not changed.

If there are problems, I can be contacted at plw@noc.ac.uk.

Many thanks for your help with this.

Yours sincerely

Philip L. Woodworth

Philip Woodworth



INVESTOR IN PEOPLE

Reviewer 1

We are grateful for the time that both reviewers spent on this paper. The comments of Reviewer 1 are given below followed by our replies. The page and line numbers refer to the version in OS Discussions.

P2 131: maybe add the reference Lyard et al 2006

Done

P4 125: OK, but in this high latitude region, the ocean response to atmospheric pressure can be significantly different from IB + effects of wind not negligible => might need to use a model forced by the atmosphere (at least a barotropic model for high frequencies) to remove correctly this non tidal variability. Have you done this test ?

We are simply saying here that sea level variability due to air pressure changes are largely compensated for in the BP records by air pressure itself, at the timescales we are interested in, thanks to the inverse barometer. That automatically removes a lot of the non-tidal variability from the records. We believe that is all correct. We agree that there are still other processes in the ocean (caused by the wind or whatever) that result in non-tidal variability – those are investigated using the NEMO model data discussed on page 12 which the reviewer commented on (so, yes, the ‘test was done’).

P4 129: “low-frequency process” : what are the frequencies concerned ? annual/semi annual only or some other components ?

This was badly worded. Instead of ‘primarily a low-frequency process’ we now say ‘a slow monotonic process’. We have also added an extra reference (Polster et al., 2009) and the three references now given should be adequate to give the reader an idea of the problems of drift in deep pressure sensor data.

P6 17: have you considered the same length of record for each BPR ? if not, can you estimate the impact of the different lengths of record on the harmonic estimation of M_f , M_m , M_t ? this impact is likely not negligible and should be considered in the discussion.

This is related to the question for page 11 line 15, see below. Almost all deployments were annual ones apart from those that used the MYRTLE instruments. If the lengths had been very different, then we agree that the lengths and noise contents of each record would determine differently the size of the uncertainties for amplitude and phase (Eq 5). However, we cannot see that being a problem in itself. We have worked with the computed formal errors from each regression and a bigger issue with this approach, as the text discusses, is the assumption of white noise for the computed errors.

P6 129: sentence not clear. Please rephrase.

The sentence should now be clearer.

P7 115 : add ref to eq 4

The same references apply as for M_2 in Eq [3]. We have referenced Doodson and Warburg (1941) here again.

P7 133: “28.4 +/-1.4_” : what about the sign? Do you obtain the same sign as in eq 4 ?

It obviously has the same sign as in Eq 4, as can be seen from Figure 5(a). We have added a few words to say that.

P8 14: add a sentence like “this N-S difference is likely explained by the dynamic response of the ocean at this frequency” : see the spatial patterns of FES2014 showed in supplementary materials.

We have not done this. This would be getting ahead of things. The spatial variations (dynamic response) are discussed in detail in the next section using the FES2014 model.

P8 115: add ref to eq 8

The same references apply as for M2 in Eq [3]. We have referenced Doodson and Warburg (1941) here again.

P8 121: have you tried to fit cos or sin ?

Let us explain what we did again. A first step using the data from each deployment is to find the individual amplitudes and phase lags determined using Eq [5], as explained in Section 2. A second step is that the set of all the values of amplitude are parameterised as a cosine peaking at year 2006.5 when $N=0.0$, as explained in Section 3.1. For example, that is easy to see for Mf from Eq 4. In the case of Mm the cosine should in theory be upside-down because of the negative sign in Eq 8 – such an upside-down shape was indeed obtained for Mm in Figure 6(a) although its small amplitude meant that it looks almost like a straight line – see the text on page 8 line 16.

For phase lags we fit a sine constrained to be zero at 2006.5 instead of a cosine (see end of page 7) as you can see from Eq 4 that the nodal variation in the equilibrium tide for Mf or Mt is expected to vary like $\sin(N)$.

We don't understand why the reviewer points to page 8, line 21 in this context as that sentence makes it clear that there is expected to be no nodal variation in phase lag for Mm (Eq 8), and indeed there is no evident nodal variation in the data, so we just show in Figure 6(b) a straight line at the average value of 177.3° .

P8 130-31: mean value 0.43 is smaller than in eq 4. Please explain

We don't understand the comment. The 0.43 mbar is the mean value of Mt amplitude. If the reviewer is asking why this is smaller than the 1.043 in Eq. 4, then the latter is the nodal factor 'f' which is a dimensionless multiplier of the amplitude in Eq 2 (the lines following explain why that is not exactly 1.0 in this case). Eq 4 applies to both Mf and Mt to a good approximation by the way, the average amplitude of Mf is of course the larger.

P8 133: “which follows from the larger average amplitudes in the second half of the data” : not clear, please explain

To be more complete, the text says “which follows from the larger average amplitudes in the second half of the data (Figure 7a)”. So please take a look at Figure 7a again – the nodal

variation means that the amplitudes tend to be larger in the second half. However, we agree there is maybe a problem with saying ‘average amplitudes’ which we have changed to ‘larger amplitudes on average’.

P9 117: “individual uncertainties approximately five times larger than for the BPRs”: how do you explain this point ?

Because the non-tidal background is much larger in a coastal tide gauge record due to air pressures and winds than in a BP record from deeper water – this is the reviewer’s own point above referring to page 4 line 25. See the next two points regarding spectra and DAC corrections to the tide gauge data. Section 3.2 has been rewritten and extended.

P9 124: “the superiority of BP measurements” : this point is not clearly demonstrated here. Need a spectrum of TG as in figure 3 + see next point.

The point is demonstrated clearly by the ‘five times larger’ in the previous comment and from inspection of the error bars in Figures 5 and 8. However, we agree that it is important to include additional spectra for the TG data, we should have done that before, please see our answer to the next point and to Reviewer 2 point 4.

P9 125-27: clearly modelling the non tidal variability should improve the results, you should make the test. You can use the Dynamic Atmospheric Correction (which is a barotropic modelling) to check this impact (the data are available on line on the AVISO website) or use NEMO as in page 12.

Many thanks for this comment which is similar to that of Reviewer 2 point 4. We have modified and expanded the discussion of Vernadsky data using DAC corrections in Section 3.2.

P10 111: add references for FES2014

The web site (dated 2018) from which FES2014 can be downloaded is referenced on page 10 as FES2014 (2018) and given in the reference list. This is admittedly a strange way of referring to something, with a date in the model name which is not the date it was obtained. If there is a better way of referencing the model we would be happy to use it.

P11 115: “typically 1-year long records” : for BP different lengths have been used isn’t it ?

Not really. On page 4 (section 2) we explained that almost all the BPR records are from annual deployments (apart from the MYRTLE records). These redeployments happened at almost the same time of year, constrained by the schedules of the BAS ships, resulting in roughly 1-year records. The exact dates for each one can be found from the PSMSL web site.

P11 125-26: comparison is ambiguous: did you choose the 185_ contour because this is the closest to the observed average phase lag ? or do you really take the geometrically mid-passage contour ? need to clarify

We don’t see how the comparison is ambiguous. These lines of text are just mentioning a general comparison of the average phase lag for Mm reported in Section 3.1, and given that there were a similar number of deployments north and south, compared to the average phase lag at mid-Passage from the FES2014 model anyone would conclude by inspecting

Supplementary Figure 2(d). We have added '(Section 3.1)' following the mention of 177° , which makes it a little clearer where the numbers come from. We have also added '(Section 3.1)' where Mf average phase lag is mentioned on page 10, and for Mt following the reviewer's comment on page 12 line 26 below.

P11 129-30: indeed for 92-99, Mf amplitudes are smaller for south deployments : : : is this N-S difference small enough to be not significant ?

There is a misunderstanding here. Page 11, lines 29-30 are discussing Mm and not Mf. They make the point that the corresponding Mf amplitudes are not so different to later ones. We suspect that the earlier and lower Mm values referred to are to do with being further east at the south end of the F-S line. This would be another artefact of our having to deal with a data set that has both spatial and temporal dependence.

P12 17: "use of 5day values of BP": is it a running 5 days average ? why not using 1-day as what is done on BP measurements ?

5-day values are a standard NEMO product, see Hughes et al. (2018). Anyway, here we are discussing Mm for which 5-day sampling should be just about adequate.

P12 110-11: " : : : correlations were weaker in the north : : : " : can you explain more ?

Not really. The weaker correlation with NEMO in the north is almost certainly to do with the higher eddy variability as demonstrated by Sheen et al. (2014). Although NEMO has eddies in it, it is unreasonable to expect a model such as that to explain all the details of that variability. It does a better job in the south. The different character of variability north and south is demonstrated in Figure 3 of Hughes et al. (2018), see also Supplementary Figure 3 in the present paper. This is already explained in the text.

P12 124: why do you use different names for Mt/Mtm ?

We don't. Mtm is mentioned only once, to point out that the constituent has that name in the FES2014 model. Otherwise we call it Mt throughout. You will often find tidal constituents having different names in the literature when they have been studied by different people through the years.

P12 126: same comment as for Mm, see above.

We have added '(Section 3.1)' after the mention of 197° to make it clearer.

P12 130: "similar to that obtained above for figure 7a" : the estimation for figure 7a are not shown in the text above ? : : : to add

We see the problem here with the word 'above'. We meant a couple of pages above in section 3.1. The wording has been made clearer.

P12 132: idem for estimations on figure 7b

Ditto

P13 13: likely true for old versions of tidal packages : : :

It is not a question of old versions of packages as such. All of them use different sets of constituents depending on the lengths of records analysed etc. The fact of the matter is that the mid-latitude heritage of much tidal research (e.g. Darwin/Doodson/Cartwright) meant that Mt was not normally included in the standard sets, although there is no real reason why it could not have been.

P13 128: “should be separable from Mf : : : given a year of data”: have you performed some tests ? using a long time series and then a one year time series to be able to say that ?

There are no tests necessary. We have mentioned the periods of the additional constituents MSf, MSm and MSt and those of our main three Mf, Mm and Mt, and you can check that the pairs are all separable within a year by the Rayleigh criterion. Now, of course, measurement errors can make this procedure more uncertain. So we have added ‘In principle’ to this sentence.

P13 129 : you mean removing these small constituents using an ocean model and then analyzing the studies frequencies ? but ocean models might not be enough accurate for such small constituents : : : please clarify.

We have changed the text to read ‘ocean tide models’ instead of ‘ocean models’ which may have been misleading. Inference of smaller constituents is a standard procedure in tidal analyses. The details can be left to whoever in the future does the analysis.

P14 13: + this point might also explain the different behaviours of BP and TG ?

We are not sure about this. We never combine BP and TG data in any fit, and within the TG fit alone the same ‘k’ factor will apply.

P14 18: “our determination of Mm”: why not other components Mf, Mt ? please explain

Because Mm is the lowest of the three in frequency and the spectrum is red (or pink). Then please see the previous sentences.

P15 1 17: “stacks of records” : please explain

‘Stacks’ are when many records are combined into an overall fit. It is a technique often used in seismology, for example, and was used by Trupin and Wahr (1990) to look at long-period aspects of tide gauge records. Anyone interested can read that paper. We are not digressing into a discussion of that here.

P16 A2: you get these formulae from eq 2 and A1 ?

Yes. That is correct.

P17 111: how do you choose $R=0.414$?

We did not choose this value ourselves. It is the value of the amplitude of the main sideband in the tidal potential. However, we agree this was unclear and have added the Cartwright and Tayler (1971) and Cartwright and Edden (1973) references again.

P18 15: It is not clear why you choose to use simplified formulae in this paper ? explain please.

Because they are quite adequate for the simple nodal variations we are looking at here, especially given the uncertainties in the data. These simple forms are also the ones that normally appear in text books on tides. But there are applications such as Ray and Egbert (2012) where you have to take more complete ones. They are never completely correct, however, as they are trying to provide simple algebraic expressions of combinations of multiple sidebands in the tidal potential.

P18 113: $R=0.065$?

Yes. Again, this is a value that comes from the amplitudes for the sidebands of Mm in the tidal potential. The Cartwright tables (Cartwright and Tayler, 1971) were already referred to but we have added Cartwright and Edden (1973) to be more complete. We hope this is clearer now.

Legend of Figure 5: "one standard error" : please give a bit more details.

The caption has been expanded.

Technical corrections:

P1 1 16: replace by "while the phase difference for Mm"

Done. Many thanks.

P2 127: replace by "seems to be a good theory"

Done

P4 118: replace has -> have

Not done. Bottom pressure (BP) is singular.

P13 122: replace will -> may

In our opinion, 'will' is better as these other constituents are bound to be present to some extent and there is no 'may' or 'perhaps' about it.

Reviewer 2

We are grateful for the time that both reviewers spent on this paper. The comments of Reviewer 2 are given below followed by our replies. The page and line numbers refer to the version in OS Discussions.

1. P1 31: ' $t=0$,' => ' $t=0$. Eq[1] is further modified to;'

Thanks but we think it reads ok as it is.

2. P6 6-8: Are the daily mean values suitable for resolving Mt (9.13 days)? Using daily means might be one of reasons for large error bars in Fig7. Though there are some discussions on the complications of this analysis approach, it will be better if authors can further discuss(/investigate) the dependence of long-period tides, especially Mt, on data temporal resolution.

We understand this comment but daily means (Nyquist period of 2 days) should be adequate for study of a cycle with a period of 9 days, irrespective of the small signals and large error bars in Figure 7. It was anyway convenient for us to use daily means which were a product of the Weighing the Ocean project. We don't feel a discussion suggested by the Reviewer's last sentence is warranted.

3. P7 4-7, Fig4 and other places in text: why is non-tidal variability larger in the south side as however there are more eddies in the north as authors mentioned in P12 5? MJO (intraseasonal) is taken as one potential contributor but it seems to me that there are still some significant features at longer timescales (e.g. in Fig4b between 350day and 450day, between 510day and 610 day). A bit more explanations/speculations are suggested to add here.

The Reviewer has misunderstood our purpose in showing Figure 4(a,b). Mf is the largest of the long-period tides discussed here and it varies a lot over the nodal cycle, hence the ratio of tidal to non-tidal variability varies over the cycle. We wanted to include Figure 4(a,b) as examples of that variability, when the tidal component was large and small respectively. The two plots (a,b) were not intended to show north/south differences in variability as such. We have added some words to make that clear.

As we mentioned, there is a lot of non-tidal variability due to eddies etc. in the north (Sheen et al., 2014) but also some in south. This results in spectra parameterised as shown in the Supplementary Material Figure 3 and discussed in Hughes et al. (2018). The MJO was mentioned in particular because the timescale of its variability is not too different from the Mm period. The features pointed out in Figure 4b are indeed interesting – they are presumably associated with rapid (non-tidal) ocean variability of some kind, there are odd features like that in many BP records that we have not investigated in detail. Fortunately, the stationary signals of the tide are fairly immune to such things.

4. Subsection 3.2: It compares the long-period tides derived from BPR and also the tide gauge record. How the power spectral distributions differ between those two kinds of records? If BPR has advantages of resolving long-period tides over tide gauge data, due to less non-tidal variability, one may expect there are more noises close to the 3 constituents' frequencies (Fig3). Is this true? It's good to show this merit.

The Reviewer is right, and this point relates to two by Reviewer 1 (P9 124 and P9 125-27). We have added spectra for Vernadsky for comparison. We have also considerably extended the discussion of Vernadsky data in Section 3.2 by using DAC corrections, and have added extra words to the Conclusions. Many thanks for suggesting we do this, which we should have done before.

5. P8 1-3 and Fig5b: It's worth proposing some explanations why such north-south differences are observed here, when this is not expected from the theory.

At this point in the paper, the different phase lags north and south demonstrate spatial variation unexpected from the Equilibrium tide, that (if correct) would have to be explained

by dynamical tidal differences. That is what we investigate further in the discussion of Section 4 by making use of the FES2014 model which, if you read on, are explained well by the model.

6. Fig 6&7: small amplitudes and large error bars make it difficult to detect the nodal cycle. It seems to me that error bars are slightly larger in the 1st decade. Is this related to BPR data density used? It'll be good if the data availability (after QC) of such 45 BPR records is provided.

It is true that error bars are slightly larger in the first decade, but it depends which plot you mean. For example, for the amplitude of Mt (Figure 7a) they are much the same, but the amplitudes themselves are larger in the second part which results in the errors on the phases (Figure 7b) being smaller. See also our reply to Reviewer 1 (P8 133). But in general this comment is right, we have no simple explanation, presumably it depends on the mix of locations (e.g. the F-S line in the early years), ocean variability changes in time etc.

Most of the BPR records were re-QC'd as part of the Weighing the Ocean project and are available on the PSMSL web site. All of them were inspected for possible glitches in the time series. We have put some wording in the Acknowledgements for anyone who would like copies of the data.

7. Author used FES2014 model to discuss the spatial variation of long-period tide parameters. FES models are assimilated by satellite altimeter data, which to me however have some limitations at high latitudes. Is this a noticeable concern here?

We don't think so. The orbit of TOPEX and its follow-on missions was designed to include the Drake Passage, and there is now over 25 years of precise altimetry, so there has been plenty of data for assimilation. Anyway much of the non-equilibrium dynamics of the long-period tides can be modelled from first principles without assimilation (some references are given in the paper). We suspect that FES2014 is a very good model for the long-period tides. Anyway, it is certainly the best available for use here.

The nodal dependence of long-period ocean tides in the Drake Passage

Philip L. Woodworth¹, Angela Hibbert¹

¹National Oceanography Centre, Joseph Proudman Building, 6 Brownlow Street, Liverpool L3 5DA, United Kingdom

Correspondence to: Philip L. Woodworth (plw@noc.ac.uk)

Abstract. Almost three decades of bottom pressure recorder (BPR) measurements at the Drake Passage, and 31 years of hourly tide gauge data from Vernadsky station on the Antarctic Peninsula, have been used to investigate the temporal and spatial variations in this region of the three main long-period tides Mf, Mm and Mt (in order of decreasing amplitude, with periods of a fortnight, a month and third of a month respectively). The amplitudes of Mf and Mt, and the phase lags for all three constituents, vary over the nodal cycle (18.61 years) in essentially the same way as in the equilibrium tide, so confirming the validity of Doodson's 'nodal factors' for these constituents. The amplitude of Mm is found to be essentially constant, and so inconsistent at the three-sigma level from the $\pm 13\%$ (or $\sim \pm 0.15$ mbar) anticipated variation over the nodal cycle, which can probably be explained by energetic non-tidal variability in the records at monthly timescales and longer. The north-south differences in amplitude for all three constituents are consistent with those in a modern ocean tide model (FES2014), as are those in phase lag for Mf and Mt, while the phase difference for Mm is smaller than in the model. BPR measurements are shown to be considerably superior to conventional coastal tide gauge data in such tidal studies, thanks owing to the lower/larger proportion of non-tidal variability in the latter. However, correction of the tide gauge records for non-tidal variability results in the uncertainties in nodal parameters being reduced by a factor of two (for Mf at least) to a magnitude comparable (approximately twice) to those obtained from the BPR data.

1 Introduction

The ocean tide at each location is usually represented as a combination of harmonic constituents with frequencies corresponding to those of lines in the tidal potential (Cartwright and Tayler, 1971; Cartwright and Edden, 1973). The major lunar constituents are always accompanied by sidebands separated in frequency by $\pm 1/18.61$ cycles per year, 18.61 years being the nodal (or draconic) period of regression in the mean longitude of the lunar ascending node (Doodson and Warburg, 1941). The most efficient way of accounting for the sidebands in a harmonic expansion is via the use of 'nodal factors' f and u , whereby the simple representation of a single constituent:

$$H \cos(\omega t + A - G) \quad [1]$$

in which ω is the angular frequency of the constituent, H and G are its amplitude and phase lag, and A is its astronomical argument at time $t = 0$, is modified to:

$$f H \cos(\omega t + A + u - G) \quad [2]$$

where f and u are time-dependent functions of the longitude of the ascending node (N). For example, in the tidal potential (or equilibrium tide), the main lunar semidiurnal tide (M2) has nodal factors:

$$f = 1.0 - 0.037 \cos(N), u = -2.1^\circ \sin(N) \quad [3]$$

retaining only terms in $\cos(N)$ and $\sin(N)$, and neglecting smaller terms depending on $\cos(2N)$ etc. (Doodson, 1928; Doodson and Warburg, 1941; Pugh and Woodworth, 2014).

Because the frequencies of the sidebands are similar to the constituent's central frequency, it is usually assumed that the response of the ocean at the sidebands and at the central frequency will be in proportion to that given in the tidal potential i.e. that the same admittance will apply. However, nodal factors different to expectations from the tidal potential (or equilibrium tide) have been found at many locations, at least for semidiurnal tides.

For example, smaller values of f for M2 were found around the UK by Amin (1983, 1985) and were explained as being a consequence of non-linear frictional damping. Similar findings were obtained from measurements of mean tidal range around the UK by Woodworth et al. (1991). Differences from the expected nodal factors were found in data from the west coast of Australia (Amin, 1993) and the Bay of Fundy and Gulf of Maine (Ku et al., 1985; Ray, 2006; Müller, 2011). Feng et al. (2015) found differences for both semidiurnal and diurnal tides at locations along the coast of China. In a survey of long-term changes in the amplitudes and phase lags of the four main tidal constituents around the world (M2, S2, O1 and K1), Woodworth (2010) pointed to many locations where differences in f from those expected from the equilibrium tide were evident.

Turning to the long-period tides, all of the long-period constituents of the equilibrium tide have amplitudes proportional to $\left(\frac{1}{3} - \sin^2(\text{latitude})\right)$ with no zonal dependence. The amplitudes are twice as large at the poles as at the equator; they are 180° out-of-phase between high and low latitudes; and they have zero amplitude at 35° N/S. Proudman (1960) suggested that, at least for the longest of the long-period tides (the 18.61 year nodal tide), the tide in the real ocean should be a close approximation of its equilibrium form, and that still seems to [be](#) a good theory (Woodworth, 2012). However, tidal modelling and observations by tide gauges and satellite altimetry have demonstrated that the long-period tides with shorter periods in the real ocean, such as Mf and Mm with periods of approximately a fortnight and a month respectively, have significant spatial variations from their equilibrium form (Wunsch et al., 1997; Mathers and Woodworth, 2001; Egbert and Ray, 2003; [Lyard et al., 2006](#); Ray and Egbert, 2012; Ray and Erofeeva, 2014).

Although the spatial variations of the long-period tides are now much better understood, it is also of interest to consider whether their temporal (nodal) variability conforms to expectations. The Mf constituent (period 13.66 days) is particularly interesting in this regard. In the equilibrium tide, Mf is the largest of the long-period tides and has very large nodal variations:

$$f = 1.043 + 0.414 \cos(N), u = -23.7^\circ \sin(N) \quad [4]$$

[\(Doodson and Warburg, 1941\)](#). Why the first term in f is not identically 1.0 (for Mf and for many other constituents) arises from the way that Doodson (1928) combined sideband constituents in order to provide simple functions in terms of N only. Doodson's nodal parameterisations, especially those for Mf, are discussed in the Appendix.

As far as we know, the magnitude of this temporal variability for the long-period tides in the real ocean has never been verified properly. In principle, one would have expected that

the relatively large amplitude and short period of Mf would have enabled the temporal variation of its amplitude and phase lag to be estimated reliably from two decades of tide gauge data. However, there is always non-tidal background variability at fortnightly timescales to contend with. Most research on long-period tides in tide gauge records has been focused on regions such as the low-latitude Pacific, where the non-tidal background is much less than at higher latitudes (e.g. Miller et al., 1993). However, the long-period tides are also small in these regions (i.e. centimetric, see Figure 5 of Ray and Egbert, 2012). These studies of Pacific data were primarily concerned with establishing how the non-equilibrium aspects of Mf and Mm varied spatially, rather than temporally (Wunsch, 1967). Even though some long tide gauge records exist at high latitudes (e.g. northern Norway or Canada), where long-period amplitudes are larger, the relatively high background of non-tidal sea level variability means that it is difficult to make an accurate determination of the long-period tides without also modelling the non-tidal background (e.g. Crawford, 1982).

In this paper, we report on the temporal variations of the amplitudes and phase lags of Mf, Mm and Mt (period of one third of a month) at the Drake Passage to see if they are consistent with equilibrium expectations. These are the three long-period tides, in order of decreasing amplitude in the equilibrium tide, that one is likely to be able to extract from records of about one year. The Drake Passage is at a sufficiently high latitude that any long-period tides should be larger than in most parts of the ocean. In addition, our investigation is based on the use of measurements of bottom pressure (BP) obtained over almost three decades, instead of on conventional coastal tide gauge data. It will be seen that bottom pressure recorders (BPRs) are inherently more suitable for providing long-period tidal information than coastal tide gauges. However, as a comparison of different measurement techniques, we also make use of 31 years of hourly sea level data from the nearby Vernadsky station, which has the longest tide gauge record in Antarctica.

2 Bottom pressure recorder data and methods

Cartwright (1999, chapter 13) provides a history of the development of BPRs, primarily by groups in Germany, France, USA and UK. Cartwright himself and colleagues from the National Oceanography Centre (NOC, as it is now called) made extensive use of BPRs in sets of ‘pelagic’ (pertaining to the open sea) tidal measurements, first in waters around the UK, and then throughout the Atlantic Ocean (Cartwright et al., 1988; Spencer and Vassie, 1997). The same equipment was also used in international studies of non-tidal ocean processes (e.g. Cartwright et al., 1987), culminating in the late 1980s in the deployment of BPRs at Drake Passage in order to monitor fluctuations in the transport of the Antarctic Circumpolar Current (ACC) as part of the World Ocean Circulation Experiment (Woodworth et al., 1993).

Most of the BPR deployments were made on the north and south sides of Drake Passage in order to measure changes in the pressure gradient between them. Bottom landers based on the ‘Mk.IV’ or similar designs were used in most cases (Figure 1a, Spencer and Vassie, 1997). Over half of the deployments took the form of recoveries and redeployments on an annual basis at depths around 1000 metres, providing records of 15-minute average bottom pressure typically one year long. The other half of the deployments were made at greater depths, between 2000-4000 metres. Three deployments were made using the longer duration Multi Year Return Time Level Equipment (MYRTLE) instrument that provided BP records approximately 4 years long (Figure 1b). The measurement programme was terminated in 2016, resulting in a BP data set spanning almost three decades.

The measurements have been used in studies of ACC variability, as reviewed by Meredith et al. (2011). BP measured at the south side of the Drake Passage has been shown to

be particularly useful as a monitor of fluctuations in ACC transport (Hughes et al., 2003; Hibbert et al., 2010). The data have even proved to be useful in studies of tsunami travel times (Rabinovich et al., 2011). As regards tides, the data have been employed in validation studies of models of the semidiurnal and diurnal tides observed by satellite altimetry (Ray, 2013).

BP has advantages in tidal studies over sea level recorded by conventional tide gauges at the coast. An obvious factor is that BPRs can be deployed in deep water off-shore (pelagic), at some distance from where storm surges and other shallow-water processes are largest. Another factor is that much of the sea level variability due to air pressure changes is compensated automatically by air pressure itself in the bottom pressure measurement (the inverse barometer effect). As a result of these two factors, BP records tend to have a smaller percentage of non-tidal variability (or ‘noise’) than do tide gauge records.

The main disadvantages of a BP record are instrumental drift (also known as ‘creep’) and the absence of a geodetic datum. Fortunately for tidal studies, creep is ~~primarily a low-frequency~~ slow, monotonic process that tends not to impact upon the determination of high-frequency components of the record, such as the semidiurnal and diurnal tides (Watts and Kontoyiannis, 1990; Spencer and Vassie, 1997); Polster et al., 2009). Instrumental drift does tend to preclude the reliable observation of annual and semiannual tides in BPR data. However, these long-period tides are not of lunar origin and so are not the concern of the present investigation. The absence of a datum is an important factor when it is required to combine individual yearly records into longer, continuous records. Unless overlapping records are available, from which the datum of one deployment can be related to that of another, then techniques such as ‘end point matching’ have to be employed (e.g. Meredith et al., 2004). Although we make use of one such combined record below, in order to demonstrate clearly the existence of long-period tides in the data, combinations of records are not required for most of the present study in which we analyse the records from each deployment separately.

Almost all the Drake Passage BPR data obtained by NOC since 1992 have been re-analysed recently as part of a Natural Environment Research Council (NERC) project called ‘Weighing the Ocean’. (Several NOC deployments in the centre of the Passage and in the Scotia Sea were not included.) Data were subjected to a new set of quality control that identified any suspect measurements and corrected as far as possible for timing uncertainties and instrumental drift. The processed data, consisting of records from 35 individual deployments, are available on the web site of the Permanent Service for Mean Sea Level (PSMSL) (<http://www.psmsl.org>). Ten other records were added from deployments before 1992 at Drake Passage and from the Falkland-Signy (F-S) line (Woodworth et al., 1996). These earlier records can be obtained from <http://www.ntsif.org/files/acclaimdata/bprs/>.

Figure 2 shows the locations of the 45 deployments, many of which were at essentially the same positions and so overlap on the map. The 35 locations with reanalysed data from the PSMSL web site include those on the north and south sides of the Passage south of the Falkland Islands, and that of the first of the three MYRTLE deployments close to Signy Island. The two other MYRTLE deployments were also made on the south side, but in more central positions. The ten earlier deployments include the western-most north-south pair and those on the F-S line to the east.

We have treated the data from each deployment as a separate record, with record lengths from 296 to 1470 days. Each record of BP was first subjected to a tidal analysis consisting of typically 57 semidiurnal, diurnal and higher-frequency constituents, the exact number of constituents depending on the record length. However, importantly, long period tides were not included in the tidal analysis. Residuals of the analysis were interpolated to hourly values, and simple arithmetic averaging of the 24 hourly residuals each day provided

the time series of daily mean values of BP that are discussed below. For full details of the data processing, see http://www.pmsl.org/data/bottom_pressure/processing_procedures.php.

The evidence for long-period tides in the BP data is demonstrated clearly in Figure 3(a). In this case, the individual time series of daily mean BP were de-measured and de-trended and, when daily values were available from more than one location on the same day (i.e. from both north and south sides of the Passage), they were averaged, so providing a continuous, composite time series spanning over 26 years. Figure 3(a) shows the resulting power spectrum which indicates clearly the presence of Mm (period 27.55 days), MSf (14.77 days), Mf (13.66 days) and Mt (9.13 days). Such tidal signals are obviously less well resolved when analysing records individually (Figure 3(b)). In this case, we are dealing with records of different lengths, at times when the relative proportions of each long-period component (primarily Mf) will be different, and when there will be different proportions of tidal and non-tidal variability. Consequently, spectra were produced for each individual record, normalised to have unit energy in the long-period tidal band (0.02-0.15 cpd), and then averaged into bins of 0.005 cpd, so providing a spectrum ‘typical’ of an individual record. It can be seen that Mf, and to a lesser extent Mm, are still present, while Mt is less well resolved, and MSf cannot be seen above the background.

MSf is an interesting constituent that occurs for two reasons. It is partly a long-period tide in its own right, with an amplitude in the equilibrium tide 8.7% that of Mf, and with variations in amplitude through the nodal cycle of $\pm 14\%$. It is also partly an interaction constituent (see below), with a nodal variation of $\pm 3.7\%$ as for M2 in Equation 3. However, its generally low amplitude suggests that verification of its nodal variation in real data will be much harder than for Mm and Mt, and we have not considered MSf in detail further.

In order to study the time-dependence of the long-period tides, their amplitudes and phase lags were determined for each deployment record independently, by means of a regression of the daily means of BP in terms of three harmonics with periods of Mf, Mm and Mt plus a linear trend. The three periods are so different that the amplitudes and phase lags determined for each harmonic are almost the same whether the regression includes all three constituents or each one individually. This procedure assumes that the amplitudes and phase lags of each harmonic do not change during the record i.e.

$$h(t) = H_i \cos(\omega t - \varphi_i) \quad [5]$$

where $h(t)$ is BP for a particular harmonic that is a function of time t measured from the start of 1988, $\omega = \left(\frac{2\pi}{\text{period}}\right)$ radians per day, and H_i and φ_i are the amplitude and phase lag from the regression for deployment i . Therefore, one can investigate how H_i varies as a function of the central date of each record (T_i), and similarly from Equations 2 and 5 one can relate:

$$\varphi_i + A = G - u_i \quad [6]$$

where the variation of φ_i as a function of T_i can be described by an oscillation (u_i) around the average phase lag (G). A is the astronomical argument for the harmonic constituent concerned at the start of 1988. If the start of that year is defined by GMT (UT), then G will be the constituent’s Greenwich phase lag.

The regression is made using the G02CGF function of the Numerical Algorithms Group (NAG) library (<https://www.nag.co.uk>). This results in the determined amplitudes for the three harmonics having the same standard errors, while standard errors on each phase lag are defined by the standard error on the amplitude divided by the amplitude itself (times $360^\circ/2\pi$). The same standard error for each amplitude arises from an assumption of white noise in the residuals of the regression. Consequently, they may be potentially estimated too

low (see below). However, inspection of the magnitude of the scatter in the plots below points relative to the nodal fits in Figures 5-8, compared to their individual formal errors, demonstrates suggests that the standard errors will have been estimated fairly reliably.

3 Results for Mf, Mm and Mt

3.1 BPR data

In this section, we discuss findings for Mf, Mm and Mt obtained from the BPR data. Figure 4(a) shows an example of one of the records of daily mean BP and the result of a regression fit in terms of the three harmonics. In fact, this is a particularly good example of a record from the north side of the Drake Passage with a relatively small proportion of non-tidal variability, at a time (2008-9) when the amplitude of Mf was larger than average. It serves to make the point that information on the amplitude and phase lag of Mf can be extracted reliably from such records.

Figure 2 shows that the deployments in the Drake Passage took place over a large area. However, we can take advantage of the fact that the spatial scale of variation in Mf, and of other long-period tides, is also large (e.g. see Figure 5 of Ray and Egbert, 2012 and the discussion of the FES2014 model in Section 4). Consequently, as a first approximation, all of the values of H_i and G_i from the many deployments can be considered as having been obtained at the same location.

Figure 5(a,b) presents the amplitude and phase lag of Mf respectively, obtained from the harmonic analysis of each record. The amplitude units are mbar which can be taken as being approximately equivalent to cm of seawater. A clear nodal (18.61 year) variation can be seen in the amplitudes (Figure 5a), with the red line showing a fit in terms of $\cos(N)$, constrained to peak when $N = 0$ at 2006.5. The red line passes equally well through the black and blue points, representing deployments on the north and south sides of Drake Passage respectively

The mean amplitude in the fit is 2.18 ± 0.04 mbar, and the amplitude of the nodal variation is 0.93 ± 0.06 mbar, or 43 ± 3 % of the mean value, with the sign expected from Equation 4. This may be compared to the 40% expected from the equilibrium tide (i.e. $0.414/1.043$ in Equation 4). These and other findings reported below are summarised in Table 1.

If the real Mf had a spatial variation similar to its counterpart in the equilibrium tide, one could adjust the measured amplitudes for the difference in latitude of the various deployments (an equilibrium long-period tide has no variation zonally). Consequently, if the amplitudes in Figure 5(a) are multiplied by:

$$\left(\frac{1}{3} - \sin^2(\text{reference latitude})\right) / \left(\frac{1}{3} - \sin^2(\text{latitude})\right) \quad [7]$$

where a reference latitude of 58°S is chosen in the middle of the Drake Passage, then one obtains Supplementary Figure 1. There is a larger scatter about the fit than in Figure 5(a), with a chi-square three times as large. Most of the north-side values (black) are now systematically larger than the south-side values (blue), a result which is inconsistent with Mf amplitudes having the same latitude dependence as in the equilibrium tide.

Figure 5(b) shows the variation in phase lag obtained from each record i.e. the variation in values of $(\varphi_i + A)$ or $(G - u_i)$. The red line shows a fit in terms of $\sin(N)$, with the nodal variation constrained to be 0° when $N = 0$. The mean value in the fit is $191.9 \pm 1.0^\circ$

while the amplitude of the sinusoidal variation is $28.4 \pm 1.4^\circ$, which is a little larger than equilibrium tide expectations (Equation 4). The black and blue points are clearly separated, indicating a phase lag on the south side of the Drake Passage $22 \pm 2^\circ$ larger than on the north side (obtained by weighting the individual observed phase lags minus the fitted phase lag by the reciprocal of the square of the standard error on the phase lag). Once again, this is inconsistent with the equilibrium tide, in which both sets would have a phase lag of 180° at these latitudes.

The next largest long-period tide one can investigate is Mm. This represents more of a challenge, with a longer period (27.55 days) and an amplitude in the equilibrium tide that is approximately half that of Mf. In addition, it has a nodal variation in its equilibrium amplitude that is about a third that of Mf:

$$f = 1.0 - 0.130 \cos(N), u = 0.0 \quad [8]$$

(Doodson and Warburg, 1941). Figure 4(b) shows an example of a BP record from the south side of the Drake Passage, at a time (1999-2000) when the amplitude of Mf was much less than in Figure 4(a), indicating that Mm can be readily identified by eye at such times. Therefore, we can have some confidence in the harmonic fitting. (To be clear, Figures 4(a,b) are not to be taken as examples of north-south differences, rather than differences in the relative proportions of tidal and non-tidal variability in all BP time series at different epochs.)

Figure 6(a) shows the observed variation in Mm amplitude with no obvious differences between north and south side values. Once again, the red line shows a cosine fit to the amplitude values. The mean amplitude is 1.34 ± 0.04 mbar. However, the amplitude of the cosine is close to zero at 0.00 ± 0.06 mbar, or $0.1 \pm 4.2\%$ of the mean value (with the correct negative sign of Equation 8). This is much less than the 13% expected from the equilibrium tide, so there is approximately a 3-sigma difference between measurements and expectations.

The individual phase lags obtained for Mm (Figure 6b) are similar on each side of the Drake Passage. However, they have large uncertainties. Weighting each phase lag as for Mf above gives a south-north difference of $2 \pm 3^\circ$. They have no evident nodal variation, as suggested by Equation 8. Therefore, in this case, instead of a nodal fit the red line in Figure 6(b) indicates the median phase lag of $177.3 \pm 4.4^\circ$. This value is consistent with equilibrium expectations for a long-period tide at this latitude.

The third long-period tide to be investigated is Mt (period 9.13 days). This is the next largest long-period tide in the equilibrium tide, with an amplitude about one third that of Mm and one sixth that of Mf, and with a nodal variation in f and u similar to that for Mf in Equation 4. In this case, the amplitudes are so small that the contribution of Mt to the BP time series is not readily apparent by eye, such as in Figure 4(a,b), although Mt is undoubtedly present as shown by Figure 3(a,b). Therefore, in this case, one has to rely on the formal uncertainties provided by the regression fits.

Figure 7(a) shows the amplitudes obtained for Mt, which are similar on the north and south sides of the Drake Passage, with a mean value of 0.43 ± 0.04 mbar. The red line indicates a nodal variation with an amplitude of 0.12 ± 0.06 mbar, or $28 \pm 13\%$ of the mean value, which is consistent with f in Equation 4 within the uncertainties. Figure 7(b) shows the estimated phase lags from the analysis of each record. Phase lags have smaller uncertainties after 2001, which follows from the larger ~~average~~ on average amplitudes in the second half of the data (Figure 7a). They have an average value of $197.3 \pm 5.0^\circ$. A weighted fit indicates phase lags $22 \pm 9^\circ$ larger on the south side. A sinusoidal fit to all of the phase lag values considered together results in an amplitude of $30 \pm 7^\circ$, consistent with Equation 4.

3.2 Vernadsky data

Vernadsky on the west coast of the Antarctic Peninsula (Figure 2) has the longest tide gauge record in Antarctica. The base is now operated by the National Antarctic Scientific Center of Ukraine. A float gauge was installed at the base (then called Faraday) at around the time of the International Geophysical Year (1957-58). Monthly mean sea levels are available from the PSMSL starting in 1958, while hourly values from March 1984 to December 2014 can be obtained from the Global Extreme Sea Level Analysis (GESLA) data set (<http://www.gesla.org>, Woodworth et al., 2017).

Vernadsky tide gauge data have been used in several studies of ACC variability alongside the information from the Drake Passage BPRs (Hughes et al., 2003; Woodworth et al., 2006). For present purposes, Vernadsky data enable an interesting comparison to be made on how much better Mf can be observed in BP measurements than in coastal tide gauge data. It might be supposed that Vernadsky data would have an advantage in being all from the same location, rather than at different positions for the BPR deployments. On the other hand, a coastal tide gauge record will clearly contain a considerable amount of non-tidal variability due to storm surges etc.

Each year of hourly data from Vernadsky has been Figure 3(c) shows the spectrum of sea level variability at Vernadsky. Comparison with Figure 3(a) demonstrates an order of magnitude larger amount of non-tidal background in Figure 3(c), with only Mf observed clearly, only a hint of Mm, and Mt hidden within the background. Each year of hourly data from Vernadsky was analysed in a similar way as described for the BP measurements, providing daily values of sea level from which estimates of Mf amplitude and phase lag have been were obtained. ~~(We(Given the high noise levels at Mm and Mt frequencies in Figure 3(c), we considered Mm and Mt similar analyses for them to be below noise level in these one-year records unfeasible.)~~ Figure 8(a) shows the amplitude values, which have individual uncertainties approximately five times larger than for the BPRs in Figure 5(a). The mean amplitude in the plot is 2.90 ± 0.25 cm (and so the Mf harmonic constant would have an amplitude of $2.90/1.043 = 2.78$ cm). This is larger than for the nearby BPRs. The nodal cycle shown in red has an amplitude of 1.20 ± 0.36 cm, or 41 ± 12 % of the mean value, almost exactly the same as for the BPRs and again consistent with expectations from Equation 4. Phase lag (Figure 8(b)) is also consistent with the BP data, in having an average of $184.9 \pm 4.7^\circ$. Within the large scatter from year to year, a nodal variation with an amplitude of $22.1 \pm 7.5^\circ$ can be just about discerned. (Five years of data with phase lags outside the plot limits were not used in this nodal fit.)

~~Comparisons~~ Therefore, comparisons of Figures 5 and 8 ~~underline the point we wish to make regarding demonstrate~~ the superiority of BP measurements ~~to compared to coastal tide gauge records in~~ long-period tidal studies. ~~It is possible that modelling of, unless the non-tidal variability in the Vernadsky records in terms of a response to winds and air pressures could result in a more clearly identified Mf, but one doubts if it could ever background in the latter can be equally as good modelled efficiently.~~ Crawford (1982) provides an earlier example of an attempt at such modelling in Canadian tide gauge data.

Fortunately, Dynamic Atmospheric Correction (DAC) data sets are now available which provide estimates of the sea level response to air pressures and winds every 6 hours on a 0.25° global grid. Estimates are based on the use of a high-resolution barotropic model for high-frequency variability (timescales less than 20 days) and the assumption of the inverse barometer response for longer timescales. Details are available from the Archive, Validation et Interprétation des données des Satellites Océanographiques (AVISO) web site (<https://www.aviso.altimetry.fr>). Carrère and Lyard (2003) demonstrated how effective such modelling could be in estimating non-tidal variability in tide gauge records.

Figure 3(d) shows the spectrum of sea level variability at Vernadsky once the DAC correction has been applied. Complete years of DAC corrections are available for 1993-onwards. Therefore, they have been employed for the 22 years 1993-2014 only. Comparison

to Figure 3(c) shows that most of the background has been modelled effectively, down to a level little greater than that for the BPRs in Figure 3(a), and that Mf, Mm and Mt can now all be clearly identified above the background. Figure 8(c-h) contains a set of analyses of nodal variations for Mf amplitude and phase lag (c,d), Mm (e,f) and Mt (g,h), all based on the DAC-corrected data for 1993-2014. In the case of Mf (Figures 8c,d), the mean amplitude is 2.59 ± 0.13 cm (and so the Mf harmonic amplitude is $2.59/1.043 = 2.49$ cm). The nodal cycle red has an amplitude of 1.05 ± 0.19 cm, or $41 \pm 7\%$, about the same as for the uncorrected data in Figure 8(a). Average phase lag (Figure 8d) has a value of $207.7 \pm 2.9^\circ$ (approximately 23° larger than for the uncorrected data), and now a clear nodal cycle can be seen with an amplitude of $23.4 \pm 4.0^\circ$ (without the need to reject any values for being outside plot limits).

In the case of Mm (Figure 8e), the average amplitude is 1.47 ± 0.13 cm, while the nodal fit has an amplitude of $10 \pm 13\%$ of the average but with the opposite sign expected from Equation 8. This finding is similar to the difficulty of explaining Mm amplitude from the BPR data in Figure 6(a) reported above, and discussed further in the following section. Phase lag for Mm (Figure 8f) has an average value of $174.0 \pm 6.8^\circ$, with no evident nodal variation as expected from Equation 8. The average amplitude of Mt (Figure 8g) is 0.57 ± 0.13 cm with a nodal variation of $13 \pm 34\%$ of the mean, while Figure 8h shows an average phase lag of $232.6 \pm 12.4^\circ$ and a nodal amplitude of $47.3 \pm 19.2^\circ$.

Overall, one can see the benefit of using the DAC corrections. The non-tidal variability in the sea level spectrum is much reduced, and nodal variations in all three long-period tides can now be investigated more reliably. Mf and Mt amplitudes and phase lags, and Mm phase lag, are generally consistent with equilibrium expectations, Mm amplitude being an exception to be discussed further below. All the above Vernadsky findings are summarised in Table 1.

4 Discussion

Some of the findings of the previous section are consistent with expectations from the equilibrium tide, while those that are not require explanation.

As mentioned above, the long-period tides in the equilibrium tide have simple spatial distributions in amplitude and phase, with north-south variations only. However, their spatial distributions in the real ocean are now known to depart considerably from equilibrium expectations, with larger departures at shorter period (e.g. see Figure 2 of Ray and Erofeeva, 2014). These differences are most evident when contrasting the Pacific, Atlantic and Indian Ocean low- and mid-latitude basins.

If one considers Mf in particular, atlases of this constituent have been available for many years, notably since the data assimilation numerical modelling of Schwiderski (1982). More recent co-tidal distributions for Mf have been obtained from altimeter measurements and models by Kantha et al. (1998, Figure 7), Mathers and Woodworth (2001, Plate 4) and Egbert and Ray (2003, Figure 1). These are consistent with Mf phase lag increasing when travelling south down the Pacific coast of South America, with the 180° contour around the Drake Passage, and with a complicated amphidromic pattern in the South Atlantic to the NE of the Falklands. More recent studies have included the development of the FES2004 ocean tide model, which also showed these features (Lyard et al., 2006, Figure 2), with roughly the same Mf amplitude on both sides of the Drake Passage and larger phase lag on the south side than north side.

FES2014 (Finite Element Solution 2014) is the latest in the series of state-of-the-art global ocean tide models provided by French groups. It provides elevations and currents (amplitude and phase) and tidal loading information for 34 tidal constituents on a global $1/16^\circ \times 1/16^\circ$ grid. FES2014 (2018) provides more detailed information.

Supplementary Figure 2(a,b) shows the Mf amplitude and phase lag for Mf at the Drake Passage from the FES2014 model. Some points of consistency with our findings are as

follows. First, the model has much the same amplitude over the whole area (~ 2 cm), and phase lags are essentially zonal, largely justifying our decision to combine amplitudes and phase lags from all deployments in Figure 5, and the subsequent discussion in terms of north- and south-side values.

Second, we found the amplitudes for Mf to be similar on the north and south sides of Drake Passage (Figure 5a), but phase lags were shown to be $22 \pm 2^\circ$ larger for the southern deployments (Figure 5b). The latter is qualitatively consistent with Supplementary Figure 2(b). Third, the 192° average phase lag for Mf from all the BPRs taken together (Section 3.1, Figure 5b) is consistent with the $\sim 190^\circ$ contour in mid-Passage in Supplementary Figure 2(b). ~~On In addition, the other hand, the 185° phase lag at Mf harmonic constants estimated above for Vernadsky is a little lower than the $\sim 200^\circ$ one would infer from Supplementary Figure using DAC-corrected data (24.49 cm amplitude and 208° phase lag) are similar to those in FES2014 (2.41 cm and 202° respectively). (FES2014 amplitudes and phase lags for Mm and Mt (1.31 cm and 190° and 0.42 cm and 211° respectively) are all consistent with DAC-corrected Vernadsky findings to within ~ 1 or ~ 2 standard deviations for amplitudes and phase lags respectively).~~

If a tide model such as FES2014 was perfect, then any differences in observed amplitudes and phase lags due to the different deployment locations could be removed by relating each set of findings to those which would have been obtained at a reference location, using an admittance relationship:

$$H_i^{ref} = H_i \frac{HM^{ref}}{HM_i} \quad [9]$$

where H_i is the measured amplitude for deployment i , HM_i and HM^{ref} are the model amplitudes at the deployment and reference point locations respectively, and H_i^{ref} is the inferred amplitude at the reference point. Similarly,

$$\varphi_i^{ref} = \varphi_i + \varphi M^{ref} - \varphi M_i \quad [10]$$

where φ_i is the measured phase lag for deployment i , φM_i and φM^{ref} are the model phase lags at the deployment and reference point locations respectively, and φ_i^{ref} is the inferred phase lag at the reference point. If the model represented the spatial dependence of the tide correctly, then H_i^{ref} and φ_i^{ref} should have only a temporal dependence.

Figure 5(c) shows the resulting model-adjusted values of Mf amplitude, using a reference point location of 57° W, 58° S, demonstrating satisfactory consistency between values north and south. That was already the case in Figure 5(a), and the similarity of Figures 5(a) and (c) reflects the uniformity of amplitude in the model in this area. The nodal fit in red shows a cosine with an amplitude of $43 \pm 3\%$ of the mean, which is identical to that in Figure 5(a). For phase lag, Figure 5(d) demonstrates a considerable improvement on Figure 5(b), with values north and south in agreement (weighted south-north difference of $0 \pm 2^\circ$). In addition, the nodal fit in red has an amplitude of $23.4 \pm 1.4^\circ$, which is closer to Equation 4 than for Figure 5(b).

Consequently, the temporal variation of Mf can be seen from Figure 5 to conform closely to expectations from its equilibrium form shown by Equation 4. Mf has the largest amplitude of the long-period tides we have investigated, which together with its relatively short period compared to the typically 1-year long records, means that it is the best resolved. Our finding of consistency with equilibrium expectations parallels an observation regarding

fortnightly variations in the solid earth in a study of polar motion data by Ray and Egbert (2012), who concluded that a similar admittance applied to Mf and its nodal sideband (see also earlier work by Gross, 2009). As explained above, the same admittance for a central frequency and its sidebands indicates that the nodal factors of the equilibrium tide apply equally as well to the tide in the real ocean (or solid earth).

Turning to Mm, its spatial variation in FES2014 is shown in Supplementary Figure 2(c,d). Once again, amplitudes are much the same over the whole area, and phase lag contours are roughly zonal. However, in this case, Supplementary Figure 2(d) indicates a north-south gradient of phase lag about half that for Mf in Supplementary Figure 2(b). Our observation of a small south-north difference of $2 \pm 3^\circ$ is qualitatively consistent with the smaller gradient in the model (a south-north difference of $\sim 10^\circ$). The observed average phase lag of 177° for Mm from all deployments combined (Section 3.1) is a little lower than the $\sim 185^\circ$ contour in mid-Passage in Supplementary Figure 2(d).

Figure 6(a) shows that Mm amplitudes for the first decade are lower in the south, but they become more equal to the northern ones thereafter. One may note that five of the six deployments with particularly low amplitudes before 1994 are from the F-S line. However, some kind of general amplitude bias in these early deployments is unlikely, given that their corresponding amplitudes for Mf are consistent with later ones (Figure 5a). Overall, Figure 6(a) does not provide evidence for a temporal dependence of Mm amplitude similar to that of Equation 8. However, identifying a nodal signal of only ~ 0.15 mbar is clearly a challenge given the uncertainties. At least, the absence of any evidence for nodal variation in Mm phase lag (Figure 6b) is consistent with Equation 8.

A nodal variation in amplitude of ~ 0.15 mbar might be technically within the resolution of the BPR measurements if Mm was accompanied by only a limited amount of non-tidal variability on similar (monthly) timescales. Monthly timescales are more comparable to processes associated with ACC variability. Sheen et al. (2014) showed that eddy kinetic energy is more intense in the north of Drake Passage, where the main fronts and their meanders occur. However, eddy activity also occurs in the south. In addition, variability in BP in this region has a contribution on 30-70 day timescales from the Madden-Julian Oscillation (Matthews and Meredith, 2004), which could potentially impact on our determination of Mm.

An attempt was made to reduce the amount of non-tidal variability in the records with the use of 5-day values of BP from the Nucleus for European Modelling of the Ocean (NEMO) $1/12^\circ$ ocean circulation model for 1988-2012 (Hughes et al., 2018), with the aim of better resolving any nodal tidal signals, particularly that for Mm. The model BP was found to have a high correlation with measured non-tidal BP for most of the southern deployments, while correlations were weaker in the north, as Sheen et al. (2014) would suggest. However, subtraction of the model values from the measurements resulted in little change in the determined Mm amplitudes and phase lags.

FES2014 model adjustments for Mm from Equations 9 and 10 result in Figure 6(c,d). Figure 6(c) confirms similar amplitudes north and south, and the nodal fit gives an amplitude of $0.8 \pm 4.3\%$ of the mean value, a little larger than that from Figure 6(a), but still 3-sigma away from expected in Equation 8. As for phase lag (Figure 6d), the weighted south-north difference is now $-11 \pm 3^\circ$, as can be readily observed by eye. This indicates that the model over-corrects for spatial variation in phase lag. This suggests that the difference in Mm phase lag across the real Drake Passage is less than in the model.

One might have expected the detection of Mt to be easier than that for Mm, thanks to its shorter period, even though it has a much smaller amplitude. Figure 7(a) shows an average amplitude of 0.43 mbar, with little evidence for differences between values north and south, while Figure 7(b) indicates an average phase lag of $\sim 197^\circ$, and some evidence for phase lags about 22° larger in the south than in the north. The temporal variations in Mt amplitude and

phase lag in Figure 7(a,b) are consistent with equilibrium expectations within the large uncertainties for this small constituent.

Supplementary Figure 2(e,f) gives the corresponding information for Mt from the FES2014 model. (This constituent is called Mtm in the model.) Supplementary Figure 2(e) shows an amplitude of ~0.4 cm over most of the area, while Supplementary Figure 2(f) shows a meridional gradient for phase lag similar to that obtained from the BPRs. The observed mean phase lag of 197° (Section 3.1) is consistent with the mid-Passage contour in Supplementary Figure 2(f).

If one applies the FES2014 model adjustments from Equations 9 and 10 to the observed amplitudes and phase lags for Mt, then one obtains Figure 7(c,d). This procedure results in apparent improvements as for Mf. Figure 7(c) is much the same as Figure 7(a), with similar amplitudes north and south. The nodal fit in Figure 7(c) has an amplitude of $30.531 \pm 14.0\%$ which is similar to that obtained above in Section 3.1 for Figure 7(a). For phase lag, Figure 7(d) demonstrates an improvement on Figure 7(b) with a weighted south-north difference of $-5 \pm 9^\circ$, consistent with zero difference. The nodal fit shows an amplitude of $23 \pm 7.0 \pm 6.9^\circ$, closer to Equation 4 than the value for Figure 7(b) obtained in Section 3.1.

As an aside, one can mention that Mt is to some extent a ‘forgotten constituent’. It is represented in harmonic expansions of the tidal potential (Doodson, 1921; Cartwright and Tayler, 1971) as a line with Doodson number 0,3,0,-1,0,0 (or 085.455 in Doodson’s notation) with one major nodal sideband (0,3,0,-1,1,0). However, Doodson did not usually refer to it explicitly in his own papers (e.g. Doodson, 1928), and it is not included in the standard sets of harmonic constituents used in tidal analysis packages (e.g. Bell et al., 1996), even though Figure 3 shows that it is resolvable at higher latitudes, at least in BPR data. One supposes that the reason for lack of interest in this constituent by previous tidal analysts has been due to its smaller amplitude at low- and mid-latitudes and to the generally higher level of noise in tide gauge records.

There are several complications we are aware of in the above analyses. One is that, when measurements are combined from different locations, then the observed tidal amplitudes should be adjusted for spatial-variations in water density, latitude-dependent variations in acceleration due to gravity, and depth-dependent compressibility of sea water. However, these will be at the ~1% level (Ray, 2013) and so are much less than other uncertainties.

A second complication concerns whether imperfections in our tidal analyses and subsequent averaging of the BP residuals into daily means of BP could have aliased residual components of the main diurnal and semi-diurnal tides into frequencies similar to those of the three long-period tides. We do not believe this is an important issue. All of the tidal analyses were subjected to quality control to check that tidal and non-tidal components of the records were separated efficiently. However, any residual tidal signals would then have been considerably reduced by the daily averaging. For example, the amplitude of any residual M2 would have been reduced to approximately 3.5% of its original value and aliased into the period of MSf (14.77 days). Consequently, while it is possible that aliasing could have contributed to some of the MSf in Figure 3(a), we believe most of that to be real. Furthermore, it is hard to see how the observed Mf, Mm and Mt could have been affected to any significant extent by aliasing. In principle, residuals of the tiny constituents OP2, Lambda2 and SNK2 could be aliased into Mf, Mm and Mt respectively, although reduced to negligibility by the daily averaging. Lambda2 is included explicitly in the tidal analysis. The other two are interaction constituents (see below) and do not appear as significant lines in the tidal potential (Cartwright and Tayler, 1971).

A further complication is that there will be other constituents present in the data (i.e. genuine and not-aliased ones) with a similar period to Mf, Mm or Mt. We have ignored this complication for present purposes as the other constituents are likely to be small. In the case

of Mf, the other main constituent will be MSf. MSf has an amplitude 9% of that of Mf in the equilibrium tide, which is similar to that found in the composite BPR record (Figure 3a). Similarly, MSm (period 31.81 days) has an amplitude of 19% of Mm in the equilibrium tide, and MSt (period 9.56 days) is 19% of Mt. However, there is little evidence for significant amounts of either in Figure 3(a). Again in principle, these other constituents should be separable from Mf, Mm and Mt given a year of data. One might imagine a more sophisticated harmonic expansion in future work in which information on these and other constituents is inferred from ocean tide models.

Another complication is that observations of the three long-period tides considered here can contain contributions from nonlinear interactions between shorter-period tides. For example, the difference between K1 and O1 frequencies is identical to that of Mf, and so their interaction can contribute to the observed Mf. K2 and M2 interactions can also contribute. Similarly, M2 and S2 can provide an interaction with the same frequency as MSf, which is similar to that of Mf. N2 and M2 interaction can contribute to Mm. An interaction will have an f and u determined by the product of the individual f and u values of the two short-period tides involved (see Table 4.4 of Pugh and Woodworth, 2014). Therefore, interaction nodal factors will be different to those of the long-period tide. This complication is primarily an issue for shallow waters, rather than the deeper ocean areas of the Drake Passage where our BPR measurements are located. Nevertheless, it should be possible to estimate the contributions from such interactions using tide modelling.

A final complication relates to all BP spectra having a continuous non-tidal background in addition to a tidal line spectrum (e.g. Figure 3). The background will tend to increase the amplitudes calculated for each tidal constituent (see Appendix B of Munk and Cartwright, 1966 and discussion in Wunsch, 1967). We simply note that this aspect would impact primarily on our determination of Mm. Another issue to do with the background is that it is not white noise. As mentioned above, this could lead to the errors in the harmonic analysis regressions being underestimated (e.g. Williams, 2003). In fact, the background spectra for all of the 45 BPR deployments are similar and can be parameterised reasonably well by a (frequency^k) dependence where $k \sim -1.5$ (Supplementary Figure 3). This suggests similar biases in estimated errors for each constituent for each deployment. Such biases, as long as they are similar in each case, should not significantly affect the fits to determined parameters from all deployments in Figures 5-7.

5 Conclusions

If one has several decades (or at least 19 years) of good tide gauge (or, in theory, BPR) data available for a tidal analysis then, if background noise levels allow, it should be possible to avoid having to consider the nodal sidebands as perturbations of the main harmonic via the use of ‘nodal factors’ f and u . Instead, one can treat them as independent constituents and make an explicit determination of their amplitudes and phase lags. Examples of such analyses of long records include those of Amin (1983) and Foreman and Neufeld (1991).

However, in practice most tidal analyses are made using one or several years of data, for which assumptions are required for f and u . The drawbacks of this approach have been recognised for many years, but primarily for the semidiurnal and diurnal constituents. As far as we know, the question of whether the variation of the long-period tides through the nodal cycle differs from equilibrium expectations has never been investigated properly.

In this paper, we have used data from 45 separate BPR deployments in the Drake Passage, and 31 years of hourly tide gauge data from the Vernadsky station in Antarctica, to estimate how well the nodal variation of the amplitudes and phase lags of Mf, Mm and Mt compare to expectations from the equilibrium tide. Our analysis uses simple harmonic expansions of daily values of BP or sea level at each location.

The combined data set provides information on how the amplitudes and phase lags of each constituent vary between the north and south sides of the Passage. The measurements indicate that amplitudes are similar throughout the region, which is consistent with a state-of-the-art ocean tide model (FES2014, 2018). Phase lags for Mf and Mt are $\sim 20^\circ$ larger in the south than in the north, which is also consistent with the model. However, the observed south-north difference in Mm phase lag is consistent with zero, compared to $\sim 10^\circ$ in the model. In fact, the Mm difference is probably consistent with the model given the uncertainties, and at least the BPR data and FES2014 are in agreement on indicating a smaller meridional gradient for Mm phase lag than for the other two constituents. Any detailed differences for all the long-period tides may be understood better by future modelling.

However, our main interest is in the temporal variability of the long-period tides. The variation of the amplitudes and phase lags of Mf and Mt in the BPR data have been found to be consistent with those suggested by the equilibrium tide within their uncertainties. To a great extent this is an expected finding given that, as explained in the Introduction, the long-period tides are closer to equilibrium than the diurnal and semi-diurnal tides and the frequencies of the nodal sidebands are close to that of the central line. Nevertheless, this is a reassuring finding for tidal analysts who might now (in this region at least) be able to employ f and u for the long-period tides as anticipated. The variation in phase lag of Mm (or rather its non-variation) is also consistent with equilibrium expectations. The absence of an expected 13% variation in the amplitude of Mm (Equation 8) at 3-sigma level (or possibly less if, as explained above, our uncertainties were slightly underestimated) is probably due to the background of non-tidal variability in the ocean circulation in this energetic area and/or in our inability to account adequately for spatial variations in Mm amplitude with the use of FES2014.

Our study has shown clearly that BPR data have ~~many advantages over conventional tide gauge measurements in long-period tidal studies such as this~~ advantages over conventional tide gauge measurements in long-period tidal studies such as this. Section 3.2 showed that, when Vernadsky coastal tide gauge data were corrected for non-tidal variability, then a major improvement in identification of the long-period tides results (e.g. reduction in the uncertainties for Mf in Table 1 by a factor of two). However, the Drake Passage BPRs, which were located in deeper water where the inverse barometer-related sea level variations are compensated automatically by BP itself, have still provided more accurate estimates of nodal variation, in spite of different locations for deployments. Table 1 demonstrates that the uncertainties for Vernadsky Mf, even when DAC-corrected, are still double those of the FES2014-corrected BPRs. (A similar conclusion can be obtained from inspection of the uncertainties displayed in Figures 5(c,d) and 8(c,d)). Nevertheless, it is the case that there is a lot more tide gauge data available for study worldwide than BPR data (Woodworth et al., 2017). Therefore, an obvious recommendation following from the present work is that tide gauge data be investigated more completely in order to investigate whether the temporal variation of long-period tides conforms to equilibrium expectations, perhaps by employing ‘stacks’ of records, as has been used previously to investigate other long-period components of tide gauge records (e.g. Trupin and Wahr, 1990-), with DAC-type corrections applied to each record.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements. The programme of Drake Passage bottom pressure measurements was led by scientists from the National Oceanography Centre including Ian Vassie, Mike Meredith, Chris Hughes and Miguel Ángel Morales Maqueda. The bottom pressure recorders

were designed, constructed and deployed by Bob Spencer, Peter Foden, Jeff Pugh, Steve Mack, Geoff Hargreaves and others. The help of the British Antarctic Survey with the deployments is much appreciated. Data sets were processed by David Blackman ~~and~~, Philip Axe and Angela Hibbert, and may be obtained from the web sites mentioned in Section 2 or from the authors of this paper. The NERC ‘Weighing the Ocean’ project (Grant No. NEW344022) was led by Mark Tamisiea and Chris Hughes. Loren Carrère is thanked for advice on the FES2014 ocean tide model which is distributed by AVISO (~~<http://www.aviso.altimetry.fr/>~~ <https://www.aviso.altimetry.fr/>). The DAC data sets may also be obtained from AVISO. NEMO ocean model data and advice on this study were provided by Chris Hughes. We are grateful to Richard Ray and Simon Williams for comments on aspects of this work, and to Xiangbo Feng and a second reviewer for their detailed remarks. Part of this work was funded by UK Natural Environment Research Council National Capability funding. Some figures were generated using the Generic Mapping Tools (Wessel and Smith, 1998).

Appendix: The accuracy of Doodson's nodal factors

The formulae for f and u presented in Doodson (1928) and Doodson and Warburg (1941) are more complicated than those in Equations 3, 4 or 8, in that they include additional terms depending on the cosines and sines of $2N$ and $3N$. However, the ones we have used are adequate for the present paper. It is useful to explain where they come from.

Imagine a constituent of unit amplitude described schematically by $\cos(\omega t)$, where for simplicity we have ignored the A and G in Equation 2. Consider the constituent as having a single important nodal sideband with an amplitude R which is less than 1, and an angular frequency $\omega + n$, where $n = \left(\frac{2\pi}{18.61 \text{ years}}\right)$ is the angular frequency of the nodal angle $N' = -N$ (Doodson, 1921). This $\omega + n$ situation represents Mf and its sidebands. Mt and K2 can be represented similarly. M2 has its single important sideband at $\omega - n$. Although most lunar constituents have one sideband that is much larger than the other, there are some constituents for which the amplitudes of the sidebands are approximately the same, such as Mm, see below.

Therefore, in the example of Mf, we can express the total tide as:

$$\cos(\omega t) + R\cos((\omega + n)t) = [1 + R\cos(nt)]\cos(\omega t) - [R\sin(nt)]\sin(\omega t) \quad [\text{A1}]$$

The nodal factor for amplitude (f) can then be expressed by:

$$f^2 = 1 + R^2\cos^2(nt) + 2R\cos(nt) + R^2\sin^2(nt) = 1 + R^2 + 2R\cos(nt)$$

$$f = \sqrt{1 + R^2} \sqrt{1 + \frac{2R\cos(nt)}{1 + R^2}} \quad [\text{A2}]$$

We can expand the second square-root by a Maclaurin series:

$$f = \sqrt{1 + R^2} \left[1 + \left(\frac{2R}{1 + R^2}\right) \frac{\cos(nt)}{2} - \left(\frac{2R}{1 + R^2}\right)^2 \frac{\cos^2(nt)}{8} \text{ etc.} \right] \quad [\text{A3}]$$

from which the second term provides the nodal time dependence of f i.e. $\frac{R\cos(nt)}{\sqrt{1+R^2}}$. When R is very small this is simply $R\cos(nt)$. (In the case of M2, for which the sideband is at $\omega - n$, then it becomes $-R\cos(nt) = -0.037 \cos(nt)$, as in Equation 3.) However, the main sideband of Mf has a much larger R value of 0.414; (Cartwright and Tayler, 1971; Cartwright and Edden, 1973), from which Equation A3 gives a time dependence of $0.382 \cos(nt)$. As can be seen from Equation 4, Doodson ignored the complication of the denominator and taken $R\cos(nt)$ to apply for Mf also.

The first and third terms provide the time-independent part of f for which Doodson took the time-average value of the third term. When R is very small, then the sum of the first and third terms can be approximated by:

$$\left(1 + \frac{R^2}{2}\right) - \left(\frac{R^2}{4}\right) = 1 + \frac{R^2}{4} \quad [\text{A4}]$$

from which one obtains 1.0004 for M2 (Doodson, 1928). When R is larger, we would have:

$$\sqrt{1 + R^2} - \frac{R^2}{4(1 + R^2)^{\frac{3}{2}}} \quad [A5]$$

which gives a value for Mf of 1.0485 given that $R = 0.414$. However, once again, Doodson appears to have assumed the small R approximation of Equation A4, giving the 1.043 in Equation 4.

From Equations 2 and A1, we can express the nodal factor for phase lag as:

$$u = \tan^{-1} \left(\frac{R \sin(nt)}{1 + R \cos(nt)} \right) \quad [A6]$$

and from the Maclaurin series $\tan^{-1}x = x - \frac{x^3}{3} + \frac{x^5}{5}$ etc. for $-1 < x < 1$, this gives $u = R \sin(nt)$ if the denominator is taken to be 1.0 for small values of R . Once again, this is clearly an acceptable approximation for M2. However, Doodson also used this approximation for Mf, resulting in the $u = 0.414 \sin(nt)$ radians or $23.7^\circ \sin(nt) = -23.7^\circ \sin(N)$ as in Equation 4.

As a test of whether these approximations by Doodson matter, Figure 9(a,b) shows the values of f and u that one obtains for Mf by calculating them rigorously using Equations A2 and A6, or by using Doodson's formulae. It can be seen that Doodson's values of f and u are good approximations, with standard deviations of the differences between the red and blue curves of 0.03 and 3.6° respectively. Therefore, they can be adopted reliably for analysis of generally noisy tide gauge or BPR data. However, in other tidal applications they may not be adequate. For example, Ray and Egbert (2012) made a study of fortnightly variations in earth rotation. When the nodal sidebands of Mf were treated rigorously, and additional double-nodal and double-perigean sidebands were included (i.e. sidebands with angular speeds which differ from that of the main line by the angular speeds of $2N'$ and $2p$ respectively, where p is the angle of lunar perigee), then improvements were obtained over the Doodson descriptions of f and u we have used here, which in turn improved upon their interpretation of high-precision length of day information.

As mentioned above, the formulae for f and u presented in Doodson (1928) and Doodson and Warburg (1941) are more complicated than the simplified ones discussed here. For example, his values for Mf include the double-nodal terms considered by Ray and Egbert (2012) (but not the double-perigean ones), and these more complete expressions will have been included in most tidal analysis and prediction software packages.

Finally, we can refer to Mm which has two nodal sidebands with amplitudes that are the same to within 1%, and that have opposite sign to that of the Mm central line in the harmonic expansion of the tidal potential (Cartwright and Tayler, 1971); [Cartwright and Edden, 1973](#)). The total tide can then be expressed as:

$$\cos(wt) - R \cos((w + n)t) - R \cos((w - n)t) = [1 - 2R \cos(nt)] \cos(wt) \quad [A8]$$

It is straightforward to see that in this case when $R = 0.065$ that $f = 1 - 0.130 \cos(N)$ and $u = 0.0$ as shown in Equation 8.

A more complicated discussion of Mm would include its other sidebands. Mm has Doodson number 0,1,0,-1,0,0. Its main double-perigean sideband 0,1,0,1,0,0 (i.e. differing by $2p$ from the main line) has an amplitude ~5% of Mm itself (as does the double-perigean sideband of Mf), while there is a component 0,1,0,1,1,0 (i.e. differing by $2p + N'$ from the

main line). There is even a 3rd-degree single-perigean component 0,1,0,0,0 (i.e. differing by p from the main line). The overall nodal factors f and u can then be obtained via:

$$f \cos(u) = 1.0 - 0.130 \cos(N) - 0.0535 \cos(2p) - 0.0216 \cos(2p - N) - 0.0551 \sin(p)$$

$$f \sin(u) = -0.0535 \sin(2p) - 0.0216 \sin(2p - N) + 0.0551 \cos(p)$$

[A9]

where the amplitudes of each term are taken from Cartwright and Tayler (1971) and that of the 3rd-degree term is evaluated at 58°S. Figure 10 indicates the simple nodal components of f and u as described by Equation 8 (or A8) by thin black and blue lines respectively. The overall values after combining all components in Equation A9 are shown by the thick lines. (This would pre-suppose that both 2nd and 3rd-degree long-period tides have a near-equilibrium behaviour. The overall values if one were to include only 2nd-degree components are shown in Supplementary Figure 4.) Equation 8 can be seen to be a good approximation of the overall f and u in spite of the other sidebands.

References

- Amin, M.: On perturbations of harmonic constants in the Thames Estuary, *Geophys. J. Roy. Astr. Soc.*, 73, 587-603, doi:10.1111/j.1365-246X.1983.tb03334.x, 1983.
- Amin, M.: Temporal variations of tides on the west coast of Great Britain, *Geophys. J. Roy. Astr. S.*, 82, 279-299, doi:10.1111/j.1365-246X.1985.tb05138.x, 1985.
- Amin, M.: Changing mean sea level and tidal constants on the west coast of Australia, *Mar. Freshwater Res.*, 44(6), 911-925, doi:10.1071/MF9930911, 1993.
- Bell, C., Vassie, J.M., and Woodworth, P.L.: The Tidal Analysis Software Kit (TASK Package). TASK-2000 version dated December 1998. National Oceanography Centre, Liverpool, 1996.
- Carrère, L., and Lyard, F.: Modeling the barotropic response of the global ocean to atmospheric wind and pressure forcing - comparisons with observations, *Geophys. Res. Lett.*, 30, 1275, doi:10.1029/2002GL016473, 2003.
- Cartwright, D.E., and Tayler, R.J.: New computations of the tide-generating potential, *Geophys. J. Roy. Astr. S.*, 23, 45-74, doi:10.1111/j.1365-246X.1971.tb01803.x, 1971.
- Cartwright, D.E., and Edden, A.C.: Corrected tables of tidal harmonics, *Geophys. J. Roy. Astr. S.*, 33, 253-264, doi:10.1111/j.1365-246X.1973.tb03420.x, 1973.
- Cartwright, D.E., Spencer, R., and Vassie, J.M.: Pressure variations on the Atlantic equator, *J. Geophys. Res.*, 92(C1), 725-741, doi:10.1029/JC092iC01p00725, 1987.
- Cartwright, D.E., Spencer, R., Vassie, J.M., and Woodworth, P.L.: The tides of the Atlantic Ocean, 60N to 30S, *Philos. T. R. Soc. Lond., A*, 324, 513-563, doi:10.1098/rsta.1988.0037, 1988.
- Cartwright, D.E.: Tides: a scientific history. Cambridge University Press: Cambridge, 292pp, 1999.
- Crawford, W.R.: Analysis of fortnightly and monthly tides, *Int. Hydrogr. Rev.*, LIX, 131-141, 1982.
- Doodson, A.T.: The harmonic development of the tide-generating potential, *Philos. T. R. Soc. Lond., A*, 100, 305-329, doi:10.1098/rspa.1921.0088, 1921.
- Doodson, A.T.: VI. The analysis of tidal observations, *Philos. T. R. Soc. Lond., A*, 227, 223-279, doi:10.1098/rsta.1928.0006, 1928.
- Doodson, A.T., and Warburg, H.D.: Admiralty Manual of Tides, His Majesty's Stationery Office, 270pp, 1941.
- Egbert, G.D., and Ray, R.D.: Deviation of long-period tides from equilibrium: kinematics and geostrophy, *J. Phys. Oceanogr.*, 33, 822-839, doi:10.1175/1520-0485(2003)33<822:DOLTFE>2.0.CO;2, 2003.
- Feng, X., Tsimplis, M.N., and Woodworth, P.L.: Nodal variations and long-term changes in the main tides on the coasts of China, *J. Geophys. Res. Oceans*, 120, doi:10.1002/2014JC010312, 2015.
- FES2014: Description of the FES2014 ocean tide model, obtained from <https://www.avisio.altimetry.fr/en/data/products/auxiliary-products/global-tide-fes/description-fes2014.html>. Downloaded January 2018.
- Foreman, M.G.G., and Neufeld, E.T.: Harmonic tidal analyses of long series, *Int. Hydrogr. Rev.*, LXVIII, 85-108, 1991.
- Gross, R.S.: An improved empirical model for the effect of long-period ocean tides on polar motion, *J. Geod.*, 83, 635-644, doi:10.1007/s00190-008-0277-y, 2009.
- Hibbert, A., Leach, H., Woodworth, P.L., Hughes, C.W., and Roussenov, V.M.: Quasi-biennial modulation of the Southern Ocean coherent mode, *Q. J. R. Meteor. Soc.*, 136, 755-768, doi:10.1002/qj.581, 2010.
- Hughes, C.W., Woodworth, P.L., Meredith, M.P., Stepanov, V., Whitworth, T., and Pyne A.R.: Coherence of Antarctic sea levels, Southern Hemisphere Annular Mode, and

- flow through Drake Passage, *Geophys. Res. Lett.*, 30(9), 1464, doi:10.1029/2003GL017240, 2003.
- Hughes, C.W., Williams, J., Blaker, A., Coward, A., and Stepanov, V.: A window on the deep ocean: The special value of ocean bottom pressure for monitoring the large-scale, deep-ocean circulation, *Prog. Oceanogr.*, 161, 19–46, doi:10.1016/j.pocean.2018.01.011, 2018.
- Kantha, L.H., Stewart, J.S., and Desai, S.D.: Long-period lunar fortnightly and monthly ocean tides, *J. Geophys. Res.*, 103 (C6), 12639–12647, doi: 10.1029/98JC00888, 1998.
- Ku, L.F., Greenberg, D.A., Garrett, C.J.R., and Dobson, F.W.: Nodal modulation of the lunar semidiurnal tide in the Bay of Fundy and Gulf of Maine, *Science*, 230(4721), 69–71, doi:10.1126/science.230.4721.69, 1985.
- Lyard, L., Lefevre, F., Letellier, T., and Francis, O.: Modelling the global ocean tides: modern insights from FES2004, *Ocean Dynam.*, 56: 394–415, doi:10.1007/s10236-006-0086-x, 2006.
- Mathers, E.L., and Woodworth, P.L.: Departures from the local inverse barometer model observed in altimeter and tide gauge data and in a global barotropic numerical model, *J. Geophys. Res.*, 106, C4, 6957–6972, doi:10.1029/2000JC000241, 2001.
- Matthews, A.J., and Meredith, M.P.: Variability of the Antarctic circumpolar transport and the Southern Annular Mode associated with the Madden-Julian Oscillation, *Geophys. Res. Lett.*, 31, L24312, doi:10.1029/2004GL021666, 2004.
- Meredith, M.P., Woodworth, P.L., Hughes, C.W., and Stepanov, V.: Changes in the ocean transport through Drake Passage during the 1980s and 1990s, forced by changes in the Southern Annular Mode, *Geophys. Res. Lett.*, 31, L21305, doi:10.1029/2004GL021169, 2004.
- Meredith, M.P., Woodworth, P.L., Chereskin, T.K., Marshall, D.P., Allison, L.C., Bigg, G.R., Donohue, K., Heywood, K.J., Hughes, C.W., Hibbert, A., Hogg, A. McC. Johnson, H.L., King, B.A., Leach, H., Lenn, Y-D., Morales Maqueda, M.A., Munday, D.R., Naveira Garabato, A.C., Provost, C., and Sprintall, J.: Sustained monitoring of the Southern Ocean at Drake Passage: past achievements and future priorities, *Rev. Geophys.*, 49, RG4005, doi:10.1029/2010RG000348, 2011.
- Miller, A.J., Luther, D.S., and Hendershott, M.C.: The fortnightly and monthly tides: resonant Rossby waves or nearly equilibrium gravity waves? *J. Phys. Oceanogr.*, 23, 879–897, doi:10.1175/1520-0485(1993)023<0879:TFAMTR>2.0.CO;2, 1993.
- Müller, M.: Rapid change in semi-diurnal tides in the North Atlantic since 1980, *Geophys. Res. Lett.*, 38, L11602, doi:10.1029/2011GL047312, 2011.
- Munk, W.H., and Cartwright, D.E.: Tidal spectroscopy and prediction, *Philos. T. R. Soc. Lond., A*, 259, 533–581, doi:10.1098/rsta.1966.0024, 1966.
- [Polster, A., Fabian, M., and Villinger, H.: Effective resolution and drift of Paroscientific pressure sensors derived from long-term seafloor measurements, *Geochem. Geophys. Geosyst.*, 10, Q08008, doi:10.1029/2009GC002532, 2009.](#)
- Proudman, J.: The condition that a long-period tide shall follow the equilibrium law, *Geophys. J. R. Astr. S.*, 3, 244–249, 10.1111/j.1365-246X.1960.tb00392.x, 1960.
- Pugh, D.T., and Woodworth, P.L.: *Sea-level science: Understanding tides, surges, tsunamis and mean sea-level changes*. Cambridge: Cambridge University Press. ISBN 9781107028197. 408pp, 2014.
- Rabinovich, A.B., Woodworth, P.L., and Titov, V.V.: Deep-sea observations and modeling of the 2004 Sumatra tsunami in Drake Passage, *Geophys. Res. Lett.*, 38, L16604, doi:10.1029/2011GL048305, 2011.
- Ray, R.D.: Secular changes of the M2 tide in the Gulf of Maine, *Cont. Shelf Res.*, 26(3), 422–427, doi:10.1016/j.csr.2005.12.005, 2006.

- Ray, R.D., and Egbert, G.D.: Fortnightly Earth rotation, ocean tides and mantle anelasticity, *Geophys. J. Int.* 189, 400–413, doi: 10.1111/j.1365-246X.2012.05351.x, 2012.
- Ray, R.D.: Precise comparisons of bottom-pressure and altimetric ocean tides, *J. Geophys. Res. Oceans*, 118, 4570-4584, doi:10.1002/jgrc.20336, 2013.
- Ray, R. D., and Erofeeva, S.Y.: Long-period tidal variations in the length of day, *J. Geophys. Res. Solid Earth*, 119, 1498–1509, doi:10.1002/2013JB010830, 2014.
- Schwiderski, E.W.: Global ocean tides. Part X. The fortnightly lunar tide (M_f). Atlas of Tidal Charts and Maps. Report TR 82-151. Naval Surface Weapons Center, Dahlgren, VA, 1982.
- Sheen, K.L., Naveira Garabato, A.C., Brearley, J.A., Meredith, M.P., Polzin, K.L., Smeed, D.A., Forryan, A., King, B.A. Sallée, J-B., St. Laurent, L., Thurnherr, A.M., Toole, J.M., Waterman, S.N., and Watson, A.J.: Eddy-induced variability in Southern Ocean abyssal mixing on climatic timescales, *Nat. Geosci.*, 7, 577-582, doi:10.1038/NGEO2200, 2014.
- Spencer, R., and Vassie, J.M.: The evolution of deep ocean pressure measurements in the U.K, *Prog. Oceanogr.*, 40, 423-435, doi:10.1016/S0079-6611(98)00011-1, 1997.
- Trupin, A., and Wahr, J.: Spectroscopic analysis of global tide gauge sea level data, *Geophys. J. Int.*, 100, 441-453, doi:10.1111/j.1365-246X.1990.tb00697.x, 1990.
- Watts, D.R., and Kontoyiannis, H.: Deep-ocean bottom pressure measurement: Drift removal and performance, *J. Atmos. Ocean. Tech.*, 7, 296-306, doi:10.1175/1520-0426(1990)007<0296:DOBPMD>2.0.CO;2, 1990.
- Wessel, P., and Smith, W.H.F.: New, improved version of generic mapping tools released, *EOS Transactions of the American Geophysical Union*, 79, 579-579, 1998.
- Williams, S.D.P.: The effect of coloured noise on the uncertainties of rates estimated from geodetic time series, *J. Geod.*, 76, 483–494, doi:10.1007/s00190-002-0283-4, 2003.
- Woodworth, P.L., Shaw, S.M., and Blackman, D.B.: Secular trends in mean tidal range around the British Isles and along the adjacent European coastline, *Geophys. J. Int.*, 104, 593-610, doi:10.1111/j.1365-246X.1991.tb05704.x, 1991.
- Woodworth, P.L.: A review of recent sea-level research, *Oceanogr. Mar. Biol.*, 31, 87-109, 1993.
- Woodworth, P.L., Vassie, J.M., Hughes, C.W., and Meredith, M.P.: A test of the ability of TOPEX/POSEIDON to monitor flows through the Drake Passage, *J. Geophys. Res.*, 101, C5, doi:10.1029/96JC00350, 1996.
- Woodworth, P.L., Hughes, C.W., Blackman, D.L., Stepanov, V.N., Holgate, S.J., Foden, P.R., Pugh, J.P., Mack, S., Hargreaves, G.W., Meredith, M.P., Milinevsky, G., and Fierro Contreras, J.J.: Antarctic peninsula sea levels: a real time system for monitoring Drake Passage transport, *Antarct. Sci.*, 18(3), 429-436, doi:10.1017/S0954102006000472, 2006.
- Woodworth, P.L.: A survey of recent changes in the main components of the ocean tide, *Cont. Shelf Res.*, 30, 1680-1691, doi:10.1016/j.csr.2010.07.002, 2010.
- Woodworth, P.L.: A note on the nodal tide in sea level records, *J. Coastal Res.*, 28, 316-323, doi:10.2112/JCOASTRES-D-11A-00023.1, 2012.
- Woodworth, P.L., Hunter, J.R. Marcos, M., Caldwell, P., Menéndez, M., and Haigh, I.: Towards a global higher-frequency sea level data set, *Geosci. Data J.*, 3, 50-59, doi:10.1002/gdj3.42, 2017.
- Wunsch, C.: The long-period tides, *Rev. Geophys. Space Phys.*, 5, 447-475, doi:10.1029/RG005i004p00447, 1967.
- Wunsch, C., Haidvogel, D.B., Iskandarani, M., and Hughes, R.: Dynamics of the long-period tides, *Prog. Oceanogr.*, 40, 81-108, doi:10.1016/S0079-6611(97)00024-4, 1997.

Table 1. A summary of estimates of nodal variation parameters for each long-period tide from Drake Passage BPR and Vernadsky tide gauge data. The last column shows the corresponding value in the Equilibrium Tide and appropriate equation number in the text.

<u>Long-Period Tide and Nodal Parameter</u>	<u>Amplitude of Variation</u>	<u>Section in the Paper where Discussed</u>	<u>Corresponding Amplitude in Equilibrium Tide [Equation Number in text] (and see references given in the text)</u>
<u>Mf 'f'</u>	<u>43 ± 3%</u>	<u>BPR data discussed in Section 3.1</u>	<u>40% [4]</u>
<u>Mf 'u'</u>	<u>28.4 ± 1.4°</u>		<u>23.7° [4]</u>
<u>Mm 'f'</u>	<u>0.1 ± 4.2%</u>		<u>13% [8]</u>
<u>Mm 'u'</u>	<u>No evident variation, see text</u>		<u>Zero [8]</u>
<u>Mt 'f'</u>	<u>28 ± 13%</u>		<u>40% [4]</u>
<u>Mt 'u'</u>	<u>30 ± 7°</u>		<u>23.7° [4]</u>
<u>Mf 'f'</u>	<u>43 ± 3%</u>	<u>BPR data, adjusted for deployments being at different locations using the FES2014 model, discussed in Section 4</u>	<u>As above for each long-period tide</u>
<u>Mf 'u'</u>	<u>23.4 ± 1.4°</u>		
<u>Mm 'f'</u>	<u>0.8 ± 4.3%</u>		
<u>Mm 'u'</u>	<u>No evident variation, see text</u>		
<u>Mt 'f'</u>	<u>31 ± 14%</u>		
<u>Mt 'u'</u>	<u>23.0 ± 6.9°</u>		
<u>Mf 'f'</u>	<u>41 ± 12%</u>	<u>Vernadsky tide gauge data spanning 1984-2014, discussed in Section 3.2</u>	<u>As above for each long-period tide</u>
<u>Mf 'u'</u>	<u>22.1 ± 7.5°</u>		
<u>Mf 'f'</u>	<u>41 ± 7%</u>	<u>Vernadsky tide gauge data spanning 1993-2014 with non-tidal variability removed using the</u>	<u>As above for each long-period tide</u>

		<u>DAC model</u>	
<u>Mf 'u'</u>	<u>23.4 ± 4.0°</u>		
<u>Mm 'f'</u>	<u>10 ± 13% Note opposite sign to that predicted by the Equilibrium Tide, see text</u>		
<u>Mm 'u'</u>	<u>No evident variation, see text</u>		
<u>Mt 'f'</u>	<u>13 ± 34%</u>		
<u>Mt 'u'</u>	<u>47.3 ± 19.2°</u>		