

Topic Editor Decision: Publish subject to minor revisions (review by editor) (10 May 2019)
by John M. Huthnance

Comments to the Author:

Dear Authors

Thank-you again for your revised manuscript. Both referees have seen it and as a result I am asking for minor modifications prior to publication in Ocean Science.

I am not sure whether you have seen the comments so I am copying them below with a few “editorial” comments from myself. Most are from Referee 1 who is an expert in tides (as well as a “native” English speaker) so please do address these. Referee 2 endorses a couple of points. There is also a comment about figure 9 forwarded via Referee 1 (you might guess where it comes from).

Referee 1

Comments on resubmission of 'Tidal variability in the Hong Kong region'
by Devlin et al. (Ocean Science)

This paper is a resubmission of an earlier and longer paper submitted to OSD which discussed tidal changes in the Hong Kong region, and less convincingly, in the South China Sea. I am pleased that the authors considered my suggestion to shorten it and focus on Hong Kong.

-Thank you very much for all your constructive comments in the previous revision!

I have read it again carefully and I have no doubt that the analysis has been done well at a technical level. However, my main concern is that the text does not read well at all. I have made some suggestions on rewording below.

-Thank you for these new comments, we will pay close attention to these suggestions as well as giving all parts of the manuscript a careful edit for readability.

The second concern, which I mentioned last time, is that the parameters used (TACs and delta-HATs) are simple ones but they are non-standard in tidal literature (unless you are familiar with Devlin's previous papers). They have to be explained therefore. But the paper assumes that the reader has read, or now wants to read, the previous papers. I insist that the paper have an Appendix wherein these two parameters are explained adequately.

-OK, we have added an appendix to better explain these terms. Thank you for your patience and understanding. As mentioned previously, we realize now that the term “delta-HAT” may not have been the best choice as it may confuse some people, but we still wanted to try and keep consistent with previous papers. We hope that our logic in the Appendix will be able to satisfy all readers.

As I mentioned last time, I don't have a problem with the TAC parameter. However, I really don't like the name 'approximate delta-HAT' which goes against all common use of the term 'HAT' in tidal literature. As I understand it, it reflects the maximum level that would be obtained in a year from a chosen set of time-dependent amplitudes and phases extracted by the admittance method. So please spell this out in the Appendix.

-Yes, absolutely! We have carefully thought about the definition and meaning of our δ -HAT terminology and have also endeavored to explain previous definitions of HAT as compared to our " δ -HAT". We hope our efforts are now easier to understand!

Detailed comments, many trivial to do with the text:

38 .. tidal variability in the 31-year period 1986-2016.

-Done.

40/41 ... locations, time series of approximations of the parameter delta-HAT, computed from combinations of the major tidal constituents, are found to be highly sensitive ..

-Fixed

44 - individual tides --> individual tidal constituents

-Fixed

49 - as important --> important in combination with

-Fixed

I really don't think tidal changes will ever be as important as MSL rise but as you say the tidal changes will add to the problem

-Agreed, thanks for the input.

81 - additional shorter-term

-Fixed

82 - and would imply that flood risk

-Done

85 - considered as a substantial complement to

-Fixed

89 - critical interest if all such factors are undergoing change.

-Done

At this point you should refer to the Appendix.

-Ok, we refer to the Appendix here.

95 - as a proxy for what can be described as changes.

-OK, fixed.

99 - was seen

-OK, fixed.

mmm should be mm m

-OK.

101 - ok so you have tidal changes and MSL in the extremes, what about non-tidal and non-MSL changes like storm surges?

-Thanks, this is a good comment. We have added a bit of text here to explain better that storm surge was not considered in the cited study here (or in the present study), though we do say that since tides and storm surge are both long-wave processes, they may be due to similar reasons, and knowledge of one part of the water level spectrum may instruct about other parts of the spectrum.

106 - I have never seen the word metropolises used in English (although it is correct I think). I would replace 'urban metropolises' with 'areas'

-Ok, I have changed this. Actually, I have also seen the word "Megalopolis" used to describe the ultra-populated areas of the planet (such as the Pearl River Delta which includes approx. 100 million people. But we do not use this term here.

two extensives in this sentence

-We changed the 2nd instance.

109 SCS --> South China Sea (SCS)

-OK, fixed!

You define this acronym only at the moment in figure 2 caption

-Should be fixed by the previous comment.

114-116 - this sentence needs rewording. It reads like you do something by doing the something.

-I have now cut down this sentence to be more direct and clearer.

122 - the longest record in Table 1 is 1954-2016 which is 63 and not 65 years

-Fixed.

123 - you have not yet explained to the reader that station names are accompanied by station codes, so I would drop (QB) here and see below.

-OK, I agree with your comments below about explaining station codes, so we will go forward with this explanation, but will try to use full station names the majority of instances.

129 environment --> geographical setting

-OK, fixed

131 .. including station name and station code, latitude and the ranges of the data records used in this study.

-Fixed

Then add here:

For brevity, we often refer to stations below by their station codes rather than their full names e.g. QB for Quarry Bay.

-Done.

You use station codes a lot in the text which seems unnecessary to me when it would be much clearer to the reader to use the place names. There is no space shortage here. So, if you were to remove them all and replace with the names then the above sentence would not be needed.

-We decided to keep the station codes on the first figure and tables, but will try to spell out the place name in all other locations.

133-137 - I know what you mean here but this sentence is rather a mouthful. Can you split into two? Also the sentence at line 143 'The tidal potential' should come earlier.

-Done, and done!

142 - 'effects are eliminated'. Why? I don't understand this. They would be eliminated only in an analysis of 18.6 years and would well and truly still be present in a 1-year analysis.

-Sorry for the confusing statement. We have changed it to better explain the process as:

“Nodal variabilities are typically present with similar strengths in both the observed tidal record and in the tidal potential. Therefore, when the observed data (harmonically analyzed in one-year windows) is divided by the potential (also analyzed in one year windows, nodal effects are mostly cancelled in the resulting admittance time series.”

-We have also moved this statement down to come after the definition of admittance so it makes more sense.

142 - 'may not always hold true'. You could refer to Amin's papers.

-I added the citation for Amin here.

144 start a new para at 'The result'

-Done

146 -.. analysis window (e.g. at mid-year) ..

-Done

157 - more apparent in the data sets used here

-Fixed

159 - ... MSL variability (Appendix 1). With the use of the TACs we determine ...

-Fixed

164 .. (delta-HAT) (see Appendix 1).

Start a new para.

-OK, done.

167 - mmm should be mm m

-Fixed, sorry about the careless mistake!

174 - The use of a window of a year in a harmonic analysis

-OK

181-183 This sentence could come earlier where you mention the Appendix

-I moved it up and changed the text there slightly

183 start new para at 'For the'

-Done

Here you start using codes in the text. I have no idea what TPK means and I can't be bothered referring all the time to Figure 1

-After re-reading closely, I can see your point. We spell out the full station names now.

30 years should be 31 (1986-2016)

-Fixed

.. determinations (Table 1).

-OK

136 .. in the TAC values over time .. [although I am not sure I understand this. I would reword this and simply say that as the TAC could change over time you have adopted a common epoch for the work]

-Fixed

188 twiddle 12-30 --> 12-31 (Table 1).

-Fixed

199 - I was not provided with the supplement

-Sorry, I thought it was uploaded, but I will instead add an appendix to the main text.

203 - I believe Victoria Harbour is usually spelt with a 'u'. It would be good to show it on Figure 1.

-You are right, HK uses British English, and I knew that because I live here and can see the harbor from my rooftop. But I think I over-applied the American English standard in the text.

.. all other gauges except .. moderately negative ...

-OK, fixed.

216 - you add the acronyms in the header at line 200 so do the same in the header here

-OK, thanks!

218 discrete --> particular

-OK

drop 'In Hong Kong' Five stations ...

229 - HK --> Hong Kong.

-Sorry, force of habit from living here in "HK"

233 - drop 'and we report ..'. Irrelevant.

-OK, thanks, fixed!

234 - OT --> overrules (OT)

-This is now irrelevant due to the removal of Figure 9(c) at the Editor's suggestion.

238 drop 'an additional'

-Dropped.

241-242 reword: .. Therefore, all MSL values reported here are given relative to the HKPD for the epoch 1965-1985.

-Fixed

246 drop 'drastically'

-OK

249 correlated to --> correlated with

-OK

254 though --> although

-OK

263 - I don't understand the sentence 'The TACs'. What does it mean that they are present? You mean they are large or what? And you don't actually show delta-HATs but you do show the TACs of the approximate delta-HATs. This all needs rewording.

-I think I have fixed it to be clearer, I hope you agree!

270 The spatial similarity in the ..

-OK, fixed

284 - with processes at other frequencies, such as at

-Fixed.

292 varies with

-OK, fixed.

293 with the spring-neap

-Fixed

294 drop 'away'

-We refined this sentence.

307 forcing of the tides

-OK, fixed

315 - some records are of shorter length and/or have many gaps, making ..

-Fixed

320 as was

-Fixed

328 perform --> employ

-OK, fixed

three-dimensional numerical ocean models to simulate the changing impacts on

-OK, fixed

330 comma before 'to better'

-OK, fixed

As I mentioned last time, one limitation of this study is the possibility of instrumental changes in the tide gauges. You don't even mention what sort of gauges they are or what changes there might have been.

-Thanks for this comment! This is important to explain, and we apologize for not doing it before! We have, fortunately, a good working relationship with the relevant authorities at the Hong Kong Observatory, and after a request for information and a short delay, they led us to a webpage on their site that gives the official yearly government reports about all Hong Kong tide gauges, their history, locations, and instruments (<https://www.hko.gov.hk/publica/pubsmo.htm>). Four of six of the HKO gauges (Quarry Bay, Tai Po Kau, Tsim Bei Tsui, and Waglan Island) are sea level pressure transducer types of gauges, and the other two (Shek Pik and Tai Miu Wan) are pneumatic type tide gauges. The Quarry Bay gauge was updated from a float type gauge recently (2017), and the Tai Po Kau gauge was also updated from a float gauge in 2006. Neither of these times correspond to any obvious anomalies in the tidal admittance records (the large changes at Tai Po Kau predate this by a few years), so we conclude that the instrumental changes were not a factor in the observed variability. Finally, all gauges operated by the HK Marine Department were all set up in 2004 as sea level pressure transducers. We now give this information in the Methods (Sec 2.1), and reprise the gauge changes in the Discussion (Sect 4.3).

351 drop hyphen in sea level. There should be a hyphen only when used as an adjective e.g. sea-level rise.

-Fixed here and elsewhere.

353 can be positively reinforced by what? by MSL changes?

- We have changed this sentence to read:

“The δ -HATs and D1/D2 TACs results illustrate that the tidal variability of multiple constituents can may be positively additive, and may reinforce MSL changesd at some locations”

354 agitate --> aggravate

-Fixed

360-364 these web addresses should have http or https

-Fixed

368 interest --> interests

-Fixed

383/422 - you could just call it SCS if has been defined in the text

-We decided to spell it out since I don't refer to it too often in the revised manuscript

figure 1/2 - add extra names as mentioned above

-Done

figure 2 - red on dark blue is not good. I would add China.

-We tried a different color scheme and increased the fonts

Taiwan Strait and Luzon are very small and unreadable when printed in A4

-Fonts are increased and some labels moved to be clearer

Figure 3 etc. - mention again that red/blue is +/-.

-OK

Black marks indicate TACs which are not significantly different from zero.

-OK

figure 7 - as I understand it this is not delta-HAT but the TAC computed from the approximate delta-HAt time series made from 4 constituents. Right?

-Yes, and we say this better in the caption now.

Please reword the caption to make that clear and not read as jargon. Also you have not defined what delta-HAT4 means in the text.

-Thanks, we admit that this caption didn't read smoothly. We have attempted to spell this out better as:

“Figure 7 The tidal anomaly correlation computed from the combination of the four largest tidal constituent amplitudes (given by the detrended sum of the $M_2 + S_2 + K_1 + O_1$) as a proxy for the change in the approximate highest astronomical tide (δ -HAT) relative to detrended MSL in Hong Kong, with the marker size showing the relative magnitude according to the legend, in units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers indicate TACs which are not significantly different from zero. “

-The delta-HAT₄ was a mistake on the figures and has been cleared up

455 sea-level --> sea level

-Fixed here and elsewhere

649 giving the station names and station codes, ... year of the available records, as well as the range of data analysed ..

-Fixed

652 - and O1 over the period 1986-2016.

-Fixed

656 - TACs over the period 1986-2016.

-Fixed

337 TAC --> TACs

-Fixed

Referee 2

L99: mmm-1 is quite confusing; please add a space where appropriate.

-We have fixed this here and elsewhere, it was a careless mistake.

LL162: what is delta-hat? On L162 it seems to be “change in the highest astronomical tide (δ -HAT)”, whereas on L163-164, we're told that “...the full tidal range (δ -HAT).” I assume it is the astronomical tide, but this could be clearer.

-Reviewer #1 had the same comments, and was more insistent on this being explained better, so we have added an Appendix that I hope clear up the confusion!

Forwarded comment

John - you will have had my review through the system. I didn't pick up on them not resolving the overtides as xxx commented on. See below. I don't know if he will reply to you himself.

“Re:Devlin... I just glanced at it. I noticed that they didn't really learn anything from my comment. Figure 9 still has annual estimates of tides like MO3 which are affected by close frequencies, but now added together with a bunch of other compound tides, perhaps in the hope that if you add together a bunch of dubious time series you'll end up with something legit.”

-Thanks for the comments. I realize now that the OT plot in Fig 9(c) doesn't add much information, and only confuses things. So, I have removed it and only kept the other three parts.

Editor comments.

Regarding Figure 9 and overtides, I am not expert (Referee 1 and the commenter are) but I do know that most of your subscript-4 (and subscript-3?) constituents are from non-linearity, whereas M6 is mainly from friction on the M2 flow. There might even be some forcing from higher-order astronomical terms. So I do doubt the value of figure 9c. You don't conclude much from it and I think any gain in understanding would need grouping of the constituents according to their origin (non-linearity, friction, . .).

-I have re-done Figure 9 to be only 3 parts, removing the OT plot in 9(c). I agree that the OT plots should not be there, and I should have removed it last revision since I removed other OT related material.

Line 310. “multiple tides” -> “multiple tidal constituents”?

-OK fixed!

Final Note to Editor about authorship

-Dr. Huthnance, thank you once again for your help, patience and understanding with our manuscript! I wanted to add a final note about the authorship of this paper. Last revision I had mentioned some affiliation changes, and in this (hopefully final) version I want to clarify and confirm these changes for the final publication.

All of this paper's authors have now officially moved to our new positions in China. So, the affiliation order should be:

-First:

Department of Geography and the Environment,
Jiangxi Normal University, Nanchang, Jiangxi, China

Second:

Institute of Space and Earth Information Science, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

And third (only for the first two authors, Devlin and Pan):

Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen, Guangdong, China

Please instruct us if there is anything else we need to do to confirm and validate these correct affiliations!

And again, Thank you for all your help!

Tidal variability in the Hong Kong region

Adam T. Devlin

Department of Geography and the Environment, Jiangxi Normal University.

Nanchang, Jiangxi, China

Institute of Space and Earth Information Science, The Chinese University of Hong Kong, Shatin,

Hong Kong SAR, China

Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen, Guangdong, China

Jiayi Pan*

Department of Geography and the Environment, Jiangxi Normal University.

Nanchang, Jiangxi, China

Institute of Space and Earth Information Science, The Chinese University of Hong Kong, Shatin,

Hong Kong SAR, China

~~*Department of Geography and the Environment, Jiangxi Normal University.*~~

~~*Nanchang, Jiangxi, China*~~

Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen, Guangdong, China

Hui Lin

Department of Geography and the Environment, Jiangxi Normal University.

Nanchang, Jiangxi, China

Institute of Space and Earth Information Science, The Chinese University of Hong Kong, Shatin,

Hong Kong SAR, China

~~*Department of Geography and the Environment, Jiangxi Normal University.*~~

~~*Nanchang, Jiangxi, China*~~

* - Corresponding author

32
33 Second re-submission to *Ocean Science*

34 March-May 2019

35
36 **Abstract**

37
38 Mean sea level (MSL) is rising worldwide, and correlated changes in ocean tides are also
39 occurring. This combination may influence future extreme sea levels, possibly increasing
40 coastal inundation and nuisance flooding events in sensitive regions. Analyses of a set of tide
41 gauges in Hong Kong reveal complex tidal behavior. Most prominent in the results are strong
42 correlations of MSL variability to tidal variability over the 31-year period of 1986-2016~~tidal~~
43 ~~variability~~; these tidal anomaly correlations (TACs) express the sensitivity of tidal amplitudes
44 and phases (M_2 , S_2 , K_1 , O_1) to MSL fluctuations and are widely observed across the Hong
45 Kong region. At a few important harbor locations, time series of approximations of the
46 parameter δ -HAT, computed from combinations of the major tidal constituents, are found to
47 be highly sensitive ~~locations, combined tidal variability that can approximate changes in the~~
48 ~~highest astronomical tide (δ -HAT) is highly sensitive~~ to MSL variability which may further
49 increase local flood levels under future MSL rise. Other open-water locations in Hong Kong
50 only show TACs for some individual tidal constituents but not for combined tidal
51 amplitudes, suggesting that the dynamics in enclosed harbor areas may be partially
52 frequency-dependent and related to resonance or frictional changes. We also observe positive
53 correlations of the fluctuations of diurnal (D_1) tides to semidiurnal (D_2) tides at most
54 locations in the region which may lead to further amplified tidal ranges under MSL. Overall,
55 it is shown that tidal changes in the Hong Kong coastal waters may be important in
56 combination with as important as MSL rise in impacting future total water levels.

62

63

64

65

66

67


68 **1. Introduction**

69 Ocean tides have long been thought of as a stationary process, as they are driven by
70 the gravitational forcing of the Sun and Moon whose motions are complex but highly
71 predictable (Cartwright and Tayler, 1971). Yet, long-term changes in the tides have been
72 observed recently on regional (Ray, 2006; Jay et al., 2009; Zaron and Jay, 2014; Rasheed and
73 Chua, 2014; Feng et al., 2015; Ross et al., 2017) and worldwide spatial scales (Woodworth,
74 2010; Müller, et al. 2011; Haigh et al., 2014; Mawdsley et al., 2015), concurrent with long-
75 term global mean sea level (MSL) rise (Church and White, 2006; 2011). Since gravitational
76 changes are not the reason, the tidal changes are likely related to terrestrial factors such as
77 changes in water depth which can alter friction (Arbic et al, 2009), coastal morphology and
78 resonance changes of harbor regions (Cartwright, 1972; Bowen and Gray, 1972; Amin, 1983;
79 Vellinga et al., 2014; Jay et al., 2011; Chernetsky et al., 2010, Familkhalili & Talke, 2016), or
80 stratification changes induced by increased upper-ocean warming (Domingues et al., 2008;
81 Colosi and Munk, 2006; Müller, 2012; Müller et al., 2012), all of which are also related to
82 ~~sea level~~sea level rise.

83 Tides can also exhibit short-term variability correlated to short-term fluctuations in
84 MSL (Devlin et al., 2014; 2017a; 2017b). These variabilities may influence extreme water
85 level events, such as storm surge or nuisance flooding (Sweet and Park, 2014; Cherqui et al.,
86 2015; Mofstakhari et al., 2015; 2017; Ray and Foster, 2016; Buchanan et al., 2017). Such
87 short-term extreme events are obscured when only considering long-term linear trends. Any
88 significant additional shorter-term positive correlation between tides and ~~sea level~~sea level
89 fluctuations may amplify this variability ~~and would imply that flood risk and implies that~~
90 ~~flood risk~~ based only on the superposition of present-day tides and surge onto a higher
91 baseline ~~sea level~~sea level will be inaccurate in many situations. The analysis of the

92 correlations between tides and sea level at a local or regional scale can indicate locations
93 where tidal evolution should be ~~considered as a substantial complement to~~ ~~considered a~~
94 ~~substantial modification to sea level~~ sea level rise. Moreover, since storm surge is a long wave,
95 factors affecting tides can also alter storm surge (Familkhalil and Talke, 2016; Arns et al.,
96 2017). Hong Kong is often subject to typhoons, with some recent storms yielding
97 anomalously high storm surges, so this issue is of critical interest if all such factors are
98 undergoing change ~~critical interest~~.

99 Recent works surveyed tidal anomaly correlations (TACs) at multiple locations in the
100 Pacific, a metric that quantifies the sensitivity of tides to short-term ~~sea level~~ sea level
101 fluctuations (Devlin et al., 2014; Devlin et al., 2017a), finding that over 90% of tide gauges
102 analyzed exhibited some measure of correlation in at least one tidal component. In a related
103 work (Devlin et al., 2017b), the combined TACs of the four largest tidal components was
104 calculated as a proxy for what can be described as changes ~~as a proxy for the changes~~ in the
105 highest astronomical tide (δ -HAT), with 35% of gauges surveyed exhibiting a sensitivity of
106 δ -HATs to ~~sea level~~ sea level fluctuations of at least ± 50 mm under a 1-m ~~sea level~~ sea level
107 change ($\sim 5\%$). A step-by-step description of the TAC and δ -HAT methods, including the
108 details of the calculations of the regressions and statistics can be found in the supplementary
109 materials of Devlin et al. (2017a and 2017b), and in this paper we summarize the meaning
110 and interpretations of the TACs and the δ -HATs in the Appendix.

111 A recent paper performed ~~similar a similar~~ analyses approach in the Atlantic Ocean, 
112 finding comparable results to the Pacific (Devlin et al., 2019). ~~The Comparing all worldwide~~
113 locations found that the greatest (positive) δ -HAT response ~~in the global ocean was~~ seen in
114 Hong Kong ($+650 \text{ mm m}^{-1}$). A probability distribution function analysis revealed that an
115 extreme sea level exceedance which includes tidal changes can be nearly double ($+150$ mm)
116 that which only considers MSL exceedance alone ($+78$ mm) over the past 50 years (Devlin et
117 al., 2017b). However, this approach did not consider water level extremes due to non-tidal or
118 non-MSL factors, such as storm surge, which may further complicate extreme water levels.
119 This ~~Yet, even without storm surge included, it was demonstrated~~ demonstrates that the non-
120 stationarity of tides can be a significant contributor to total (non-storm) water levels in this
121 region and warrants closer examination. Furthermore, tides and storm surge are both long-
122 wave processes and may be sensitive to the same forcing factors, so the behavior of tides may
123 be a possible instructor of the future behavior of storm surge events.

Formatted: Indent: First line: 1.27 cm

124 Hong Kong and the Pearl River Delta (PRD) region contains many densely-populated
125 ~~urban metropolises~~ areas with extensive coastal infrastructure and ~~extensive significant and~~
126 ~~continuous~~ recent land reclamation projects. ~~Sea level~~ Sea level rise in the region has
127 exhibited a variable rate in the region over the past 50 years (Li and Mok, 2012; Ip and Wai,
128 1990), but a common feature of all sea level records in the ~~SCS~~ South China Sea (SCS) is a
129 steep increase in the late 1990s with a subsequent decrease in the early 2000s, followed by a
130 sustained increase to the present day. In addition to this variable MSL behavior, there are
131 also anomalous tidal events observed at gauges in semi-enclosed harbor regions during the
132 late 1990s and early 2000s (shown and discussed below), corresponding to times of both
133 rapidly changing sea level and aggressive land reclamation. In this study, we ~~document the~~
134 ~~fine scale variability of tidal behavior and variability in the Hong Kong waters in response to~~
135 ~~MSL variability by performing~~ perform a spatial and temporal analysis of tidal sensitivity to
136 MSL variations in Hong Kong using the tidal anomaly correlation (TAC) method at 12
137 closely-located tide gauges.

138 **2. Methods**

139 *2.1 Data sources*

140 A set of 12 tide gauge records in the Hong Kong region were provided by the Hong
141 Kong Observatory (HKO) and the Hong Kong Marine Department (HKMD), spanning from
142 12 to ~~63~~ 5 years in length, including two gauges that are “historical” (i.e., no longer
143 operational). The longest record is the North Point/Quarry Bay (~~QB~~) tide gauge, located in
144 Victoria Harbor, established originally in 19524 and relocated from North Point to Quarry
145 Bay in 1986. The datums were adjusted and quality controlled by HKO to provide a
146 continuous record (Ip and Wai, 1990). Another long and continuous record is located at Tai
147 Po Kau (~~TPK~~) inside Tolo Harbor. Gauge locations in Hong Kong are shown in Figure 1,
148 with the gauges from HKO indicated by green markers, gauges from HKMD by light blue,
149 and historical (non-operational) gauges by red. Four of six of the HKO gauges (Quarry Bay,
150 Tai Po Kau, Tsim Bei Tsui, and Waglan Island) are sea level pressure transducer types of
151 gauges, and the other two (Shek Pik and Tai Miu Wan) are pneumatic type tide gauges. The
152 Quarry Bay gauge was updated from a float type gauge recently (2017), and the Tai Po Kau
153 gauge was also updated from a float gauge in 2006, and all gauges operated by the HK
154 Marine Department were all set up in 2004 as sea level pressure transducers
155 (<https://www.hko.gov.hk/publica/pubsmo.htm>).

156 Figure 2 shows the environment-geographical setting of the South China Sea, with the
157 location of Hong Kong indicated by the red box. Table 1 lists the metadata for all locations,
158 including station name and station code, latitude and the ranges of the data records used in
159 this study, including latitude, longitude, overall record length, and data used in this study.

Formatted: Indent: First line: 1.27 cm

160 2.2 Tidal admittance calculations

161 Our investigations of tidal behavior use a tidal admittance method. ~~An~~ The tidal
162 admittance is the unitless ratio of an observed tidal constituent to the corresponding tidal
163 constituent in the astronomical tide generating force expressed as a potential, V . This
164 potential can then be, divided by the acceleration due to gravity, g , to yield $Z_{pot}(t) = V/g$, with
165 units of length that can be compared to tidal elevations, $Z_{obs}(t)$, ~~via harmonic analysis~~.
166 Yearly harmonic analyses are performed on both $Z_{obs}(t)$ and $Z_{pot}(t)$ at each location, using the
167 R_T_TIDE package for MATLAB (Leffler and Jay, 2009), a robust analysis suite based on
168 T_TIDE (Pawlowicz, 2002). The tidal potential is determined based on the methods of
169 Cartwright and Tayler (1971). ~~Nodal and other low frequency astronomical variabilities are~~
170 ~~typically present with similar strengths in both the observed tidal record and in $Z_{pot}(t)$, but~~
171 ~~their effects are mostly eliminated in yearly analyzed admittance time series. This may not~~
172 ~~always hold true in shallow water areas but does seem to valid for the locations and tides~~
173 ~~analyzed in Hong Kong. The tidal potential is determined based on the methods of Cartwright~~
174 ~~and Tayler (1971).~~

175 The result from a single harmonic analysis of $Z_{obs}(t)$ or $Z_{pot}(t)$ determines an amplitude,
176 A , and phase, θ , at the central time of the analysis window for each tidal constituent, with
177 error estimates. A moving analysis window (e.g., at mid-year) produces an annual time-
178 series of amplitude, $A(t)$, and phase, $\theta(t)$, with the complex amplitude, $\mathbf{Z}(t)$, given by:

Formatted: Indent: First line: 1.27 cm

$$179 \quad \mathbf{Z}(t) = A(t)e^{i\theta(t)} \quad (1)$$

180 The tidal admittance (\mathbf{A}) and phase lag (\mathbf{P}) are formed using Eqs. (2) and (3)

$$181 \quad \mathbf{A}(t) = abs \left| \frac{\mathbf{Z}_{obs}(t)}{\mathbf{Z}_{pot}(t)} \right| , \quad (2)$$

$$182 \quad \mathbf{P}(t) = \theta_{obs}(t) - \theta_{pot}(t) \quad (3)$$

183 Nodal variabilities are typically present with similar strengths in both the observed
184 tidal record and in the tidal potential. Therefore, when the observed data (harmonically

Formatted: Indent: First line: 1.27 cm

185 analyzed in one-year windows is divided by the potential (also analyzed in one-year
186 windows), nodal effects are mostly constrained in the resulting admittance time series. This
187 may not always hold true in shallow-water areas (Amin, 1983) but does seem to valid for the
188 locations and tides analyzed in Hong Kong. The harmonic analysis procedure also provides
189 an annual MSL time-series. For each resultant dataset (MSL, **A** and **P**), the mean and trend
190 are removed from the time series to allow direct comparison of their co-variability. The
191 magnitude of the long-term trends is typically much less than the magnitude of the short-term
192 variability, which more apparent in the data sets used here ~~has previously been shown to be~~
193 ~~more apparent~~ (Devlin et al., 2017a; 2017b).

194 Tidal sensitivity to ~~sea-level~~sea level fluctuations is quantified using tidal anomaly
195 correlations (TACs), the relationships of detrended tidal variability to detrended MSL
196 variability (see Appendix). With the use of the TACs we determine the sensitivity of the
197 amplitude and phase of individual constituents (M_2 , S_2 , K_1 , O_1) to ~~sea-level~~sea level
198 perturbations at the yearly-analyzed scale. We also consider a proxy for the change in the
199 approximate highest astronomical tide (δ -HAT; see Appendix for details). The approximate
200 δ -HAT reflects the maximum tide-related water level that would be obtained in a year from a
201 combination of time-dependent amplitudes and phases ~~by combining the yearly analyzed~~
202 ~~time-series~~ of the four largest tidal amplitudes (M_2 , S_2 , K_1 , and O_1) extracted by the
203 admittance method, typically ~75% of the full tidal ~~range~~range (δ -HAT).

204 The detrended time series of the year-to-year change of the δ -HATs are compared to
205 detrended yearly MSL variability in an identical manner as the TACs, and both are expressed
206 in units of millimeter change in tidal amplitude per 1-meter fluctuation in ~~sea-level~~sea level
207 (~~mm/mm~~ m^{-1}). These units are adopted for convenience, though in practice, the observed
208 fluctuations in MSL are on the order of ~0.25 m. The phase TACs are reported in units of
209 degree change per 1-meter fluctuation in ~~sea-level~~sea level. The TAC methodology can also
210 be used to examine correlations between different parts of the tidal spectrum. We additionally
211 examine the sensitivity of combined diurnal (D_1 ; $K_1 + O_1$) tidal amplitudes to semidiurnal (D_2 ;
212 $M_2 + S_2$) tidal amplitudes (D_1/D_2 TACs). The units of the D_1/D_2 TACs are dimensionless
213 (i.e., mm/mm), and statistics are calculated as above.

214 The use of a window of a year in a harmonic analysis ~~The definition of the year~~
215 ~~window used for harmonic analysis~~ may have an influence on the value of the TAC or δ -
216 HAT, e.g. calendar year (Jan-Dec) vs. water year (Oct-Sep). To provide a better estimate of

Formatted: Indent: First line: 1.27 cm

217 the overall correlations for all data we take a set of determinations of the correlations using
218 twelve distinct year definitions (i.e., one-year windows running from Jan-Dec, Feb-Jan, ...,
219 Dec-Jan.). We take the average of the set of significant determinations (i.e., p -values of <
220 0.05) as the magnitude of the TAC or δ -HAT. For an estimate of the confidence interval of
221 the TAC or δ -HAT, the interquartile range (middle 50% of the set) is used. ~~A step-by-step
222 description of the TAC and δ -HAT methods, including the details of the calculations of the
223 regressions and statistics can be found in the supplementary materials of Devlin et al. (2017b).~~

224 For the very long record stations (e.g., ~~QB-Quarry Bay~~ and Tai Po Kau), we only
225 consider the past ~~310~~ years for TAC and δ -HAT determinations (Table 1). ~~Comparative
226 analyses in other studies have shown that any longer of a regression analysis may obscure
227 changes in the TAC over time~~The TAC values may change over time, so ~~using the past 30
228 years is a good window that standardizes results with~~we adopt a common epoch to better
229 match the rest of the Hong Kong tide gauge networks, which are typically ~12-310 years long.
230 Finally, we highlight some anomalous tidal events observed at certain Hong Kong gauges,
231 and discuss the temporal evolution of the tidal characteristics in Hong Kong.

Formatted: Indent: First line: 1.27 cm

232 3. Results

233 The individual TACs for amplitude and phase in Hong Kong are discussed first,
234 followed by the δ -HATs and the D_1/D_2 TACs. In all figures, significant positive results will
235 be reported by red markers, significant negative results by blue markers, and insignificant
236 values are shown as black markers. The relative size of the markers will indicate the relative
237 magnitude of the TAC or δ -HAT according the legend scale on each plot. All numerical
238 results for the major amplitude TACs (M_2 , S_2 , K_1 , and O_1) are listed in Table 2, and the δ -
239 HATs and D_1/D_2 TACs are listed in Table 3. Phase TACs of the individual constituents are
240 reported in Table S1 of the supplement.

241 3.1 Tidal anomaly correlations (TACs)

242 The strongest positive M_2 TACs are seen at Quarry Bay ($+218 \pm 37$ mm m^{-1}), and at
243 Tai Po Kau ($+267 \pm 42$ mm m^{-1}), with a smaller positive TAC seen at Shek Pik (Figure 3).
244 In the waters west of Victoria Harbour, all other gauges except Kwai Chung exhibit moderate
245 negative TACs. The semidiurnal phase TACs in Hong Kong (shown in the Supplementary
246 materials, Figure S1) show an earlier M_2 tide under higher MSL at ~~QB-Quarry Bay~~ and ~~TPK
247 Tai Po Kau~~ and a later tide west of Victoria Harbour. The S_2 results in Hong Kong (Figure 4)
248 show that only ~~QB-Quarry Bay~~ and ~~TPK-Tai Po Kau~~ have significant amplitude TAC values

249 (though smaller than M_2), and the S_2 phase TACs in Hong Kong (Figure S2) also show an
250 earlier tide at [QB-Quarry Bay](#) and [TPK-Tai Po Kau](#) under higher MSL.

251 The diurnal TACs in Hong Kong generally exhibit a larger-magnitude and more
252 spatially-coherent response than semidiurnal TACs. Like M_2 , the strongest K_1 values in Hong
253 Kong (Fig 5) are seen at [QB-Quarry Bay](#) ($+220 \pm 15 \text{ mm m}^{-1}$) and [TPK-Tai Po Kau](#) ($+190 \pm$
254 68 mm m^{-1}). The O_1 results in Hong Kong (Fig 6) are like the M_2 results, showing positive
255 TACs at [QB-Quarry Bay](#) ($+146 \pm 11 \text{ mm m}^{-1}$) and [TPK-Tai Po Kau](#) ($+100 \pm 25 \text{ mm m}^{-1}$), and
256 strongly negative TACs west of [QB-Quarry Bay](#). However, unlike the semidiurnal
257 constituents, the phase TACs for K_1 are mostly insignificant in Hong Kong (Figure S3), and
258 O_1 phase TACs (Figure S4) are only significant at [QB-Quarry Bay](#).

259 3.2 Combined tidal variability (~~δ -HATs~~) and tidal co-variability

260 The TACs are widely observed in Hong Kong, but the δ -HATs are only of
261 significance at ~~discrete-particular~~ locations (Figure 7). ~~In Hong Kong, five~~Five stations
262 exhibit significant δ -HAT values, with [QB-Quarry Bay](#) and [TPK-Tai Po Kau](#) having very
263 large positive magnitudes ($+665 \pm 85 \text{ mm m}^{-1}$ and $+612 \pm 210 \text{ mm m}^{-1}$, respectively), and
264 Shek Pik having a lesser magnitude of $+138 \pm 47 \text{ mm m}^{-1}$. Conversely, Ma Wan and Chi Ma
265 Wan exhibit moderate negative δ -HAT values, ($\sim -100 \text{ mm m}^{-1}$). The remainder of gauges
266 (which are mainly open-water locations) have statistically insignificant results for the
267 combined tidal amplitudes, even where some large individual TACs were observed. This
268 shows that the combined tidal amplitude effect as expressed by the δ -HATs is most important
269 in semi-enclosed harbors. The D_1/D_2 TACs are also important in Hong Kong and are seen at
270 almost every location. All significant D_1/D_2 TACs results are positive (Figure 8), and at most
271 locations the correspondence is nearly 1-to-1, indicating that a change in D_1 can yield a
272 nearly-identical magnitude change in D_2 , and vice-versa. Smaller magnitude relations are
273 seen in the western areas of the [HK-Hong Kong](#) region.

274 3.3 Anomalous tidal events at Hong Kong harbor locations

275 The overall temporal behavior of the tidal spectrum at enclosed harbor locations in
276 Hong Kong (Quarry Bay and Tai Po Kau) is especially interesting, ~~and we report here what is~~
277 ~~observed~~. In Figure 9, the time series of water level spectrum components are shown for [QB](#)
278 [Quarry Bay](#) and [TPK-Tai Po Kau](#), presenting the D_1 ($K_1 + O_1$) band (a), the D_2 ($M_2 + S_2$) band
279 (b), ~~the OT ($M_4 + M_6 + MK_3 + MO_3 + MS_4 + MN_4 + SN_4$) band~~ (c) and mean ~~sea level~~
280 ~~level~~ (MSL) (c~~d~~), given as normalized amplitudes with mean values shown in the legends.

281 The magnitude of MSL is given in relation to the Hong Kong Chart Datum as defined by the
282 Hong Kong Observatory. The Chart Datum is defined as ~~an additional~~ 0.146 m below the
283 Hong Kong Principal Datum (HKPD). The HKPD determined for the years 1965-1983 was
284 approximately 1.23 m below MSL. The HKPD has been recently re-determined using data
285 from 1997-2015 to be 1.30 m below MSL. Therefore, all MSL values reported here are given
286 relative to the HKPD for the epoch 1965-1985. Therefore, all MSL values are given in
287 relation to the sum of both values, so 1.376 m for the early years, and 1.446 m for the later
288 years (www.hko.gov.hk).

289 Some very notable features of the tides are clear. At QB-Quarry Bay, the early part of
290 the record shows nearly constant tidal amplitudes in D₁, while D₂ amplitudes show a slight
291 decrease, and MSL exhibits a slight positive trend. In the late 1980s, however, both D₁ and
292 D₂ increase ~~drastically~~ until around the year 2003, at which time both tidal bands undergo a
293 rapid decrease of amplitude of ~15%, sustaining this diminished magnitude for about five
294 years before increasing nearly as rapidly. The OT band shows a sustained increase over the
295 historical record, but many of the fluctuations around the trend are negatively anti-correlated
296 ~~to~~ with the perturbations in D₁ and D₂, and during the times of diminished major tides, the
297 OTs increase by about +20%. The MSL record is also highly variable at QB-Quarry Bay,
298 with a nearly zero trend during the increase in tides seen in the 1980s, followed by a strong
299 increase from ~1993-2000, and then a steep decrease concurrent with the time of diminished
300 tides before increasing again. The gauge at TPK-Tai Po Kau shows a similar tidal behavior,
301 although the timing and magnitudes are different. The increase in D₁ and D₂ at TPK-Tai Po
302 Kau in the 1980s is much larger and peaks earlier than QB-Quarry Bay, reaching a maximum
303 around 1996, and then decreasing around 1998, about five years before the drop at QB-Quarry
304 Bay. Both locations experience an absolute minimum around 2007 in D₂, but the D₁
305 minimum at TPK leads the QB-Quarry Bay minimum by a few years. These observed
306 anomalies are only observed at these two gauges; other locations in Hong Kong did not
307 reveal similar behavior.

308 **4. Discussion**

309 *4.1 Summary of observed tidal variability*

310 This survey has identified several types of tidal variability in Hong Kong. The
311 individual TACs ~~of individual tides are present~~ are significant at many Hong Kong locations,
312 while the TACs of the approximate δ -HATs appear to be more locally important, as the

313 strongest responses are mainly concentrated at specific locations (e.g., ~~QB~~Quarry Bay and
314 ~~TPK~~Tai Po Kau). The M_2 response (Fig 3) is negative at gauges just west of Quarry Bay and
315 positive at Shek Pik, with a similar pattern seen for the O_1 TACs (Fig 6). Conversely, the K_1
316 TAC results are generally positive (Fig 5). At both ~~QB~~Quarry Bay and ~~TPK~~Tai Po Kau, the
317 positive reinforcements of individual tidal fluctuations lead to very large δ -HATs, though
318 moderately negative δ -HATs are seen near ~~QB~~Quarry Bay at ~~CMW~~Chi Ma Wan and Ma
319 Wan (Fig 7). The spatial ~~connections~~similarity in the semi-enclosed center harbor regions
320 suggest a connected mechanism; this area is where most recent Hong Kong coastal
321 reclamation projects have occurred, including the construction of a new island for an airport,
322 shipping channel deepening and other coastal morphology changes. Such changes in water
323 depth and coastal geometry strongly suggest a relation to frictional or resonance mechanisms.

324 The D_1/D_2 TAC relations (Fig 8) are a more regionally-relevant phenomenon, being
325 significant nearly everywhere in Hong Kong. The majority of significant D_1/D_2 TACs are
326 positive, with most being nearly 1-to-1 (i.e., a ~ 1 -mm change in D_1 will yield a ~ 1 -mm
327 change in D_2), confirmed by the close similarity of temporal tidal trends of the D_1 and D_2
328 tidal bands in Hong Kong (Fig 9). This aspect of tidal variability in Hong Kong may be
329 related to the dynamics near the Luzon Strait, where large amounts of baroclinic conversion
330 in both D_1 and D_2 tides may tend to couple the variabilities (Jan et al., 2007; 2008; Lien et al.,
331 2015; Xie et al., 2008; 2011; 2013). The D_1 and D_2 internal tides may interact with each
332 other as well as with ~~processes at~~ other frequencies, such as ~~at~~ the local inertial frequency, f ,
333 via parametric subharmonic instability (PSI) interactions (McComas and Bretherton, 1977;
334 MacKinnon and Winters, 2005; Alford, 2008; Chinn et al., 2012), a form of resonant triad
335 interactions (Craik, 1985). The low-mode baroclinic energy can travel great distances, being
336 enhanced upon arrival at the shelf and leading to the further generation of baroclinic energy.
337 In the western part of Hong Kong, the D_1/D_2 relationships are less than 1 to 1 (~ 0.33 to ~ 0.25
338 at TBT and LOP, respectively). This may be partially influenced by effects of the Pearl River,
339 which discharges part of its flow along the Lantau Channel. The flow of the river is highly
340 seasonal and ejects a freshwater plume at every ebb tide that varies ~~by-with~~ prevailing wind
341 conditions and ~~by-with~~ the spring-neap cycle (Pan et al, 2014). The plumes may affect
342 turbulence and mixing in the region ~~and can dissipate and can dissipate tidal~~ energy ~~away the~~
343 ~~tidal bands~~, which may “decouple” the correlated response of D_1 and D_2 seen in the rest of
344 the Hong Kong coastal waters.

345 *4.2 Effects of local dynamics on tidal variability*

346 Hong Kong has had a long history of land reclamation to accommodate an ever-
347 growing infrastructure and population, including the building of a new airport island (Chep
348 Lap Kok), new land connections, channel deepening to accommodate container terminals,
349 and many bridges, tunnels, and “new cities”, built on reclaimed land. All of these may have
350 changed the resonance and/or frictional properties of the region. Tai Po Kau has also had
351 some land reclamation projects that have changed the coastal morphology and may have
352 modulated the tidal response. Both locations also show coherent D_1/D_2 TACs, as well as
353 having the largest positive δ -HATs, and large tidal anomalies (Figure 9). Other locations in
354 Hong Kong did not show such extreme variations, so these variations appear to only be
355 amplified in harbor areas. Decreases in friction associated with ~~sea-level~~ sea level rise may
356 lead to higher forcing ~~of the tides~~ tides, and those changes may also be amplified by the close
357 correlations of D_1 and D_2 variability or local harbor development which may further decrease
358 local friction. Hence, a small change in friction due to a small ~~sea-level~~ sea level change may
359 induce a significant change in tidal amplitudes. The positive reinforcement of multiple ~~tides~~
360 tidal constituent correlated with regional ~~sea-level~~ sea level adjustments may amplify the risks
361 of coastal inundation and coastal flooding, as evidenced by the gauges that had the largest δ -
362 HAT values.

363 *4.3 Limitations of this study and future steps*

364 The ~~inventory of tide gauges in Hong Kong~~ analysis of tides in the Hong Kong tide
365 gauge network revealed new dynamics and spatial connections in the area. However, some
366 records are of shorter length and/or have many gaps, making some gauges are of short length
367 and/or riddled with data gaps, making a full analysis of the area problematic. For example,
368 the Tsim Bei Tsui (~~TBT~~) gauge covers a long period, but there are significant gaps in the
369 record, which complicated our analysis. This gauge is located within a harbor region (Deep
370 Bay), bordered to the north by Shenzhen, PRC, which has also grown and developed its
371 coastal infrastructure in past decades, therefore, one might expect similar dynamics ~~are as~~
372 was seen at ~~QB~~ Quarry Bay and ~~TPK~~ Tai Po Kau. While there were moderately significant
373 D_1/D_2 correlations at ~~TBT~~ Tsim Bei Tsui, no significant TACs or δ -HATs were observed.
374 The large anomalies seen at ~~QB~~ Quarry Bay and ~~TPK~~ Tai Po Kau around 2000 are suggested
375 by the data at ~~TBT~~ Tsim Bei Tsui, but some data is missing around this time, making any
376 conclusions speculative. The Deep Bay region is ecologically sensitive, being populated by
377 extensive mangrove forests which may be disturbed by rapidly changing sea levels (Zhang et
378 al., 2018), so accurate determination of future ~~sea-level~~ sea levels is of utmost importance to

379 the vitality of these important ecosystems. Future studies ~~employing~~considering highly-
380 accurate digital elevation models will ~~perform~~employ simple analytical models as well as
381 high resolution ~~three-dimensional numerical ocean models to simulate the changing impacts~~
382 ~~on three dimensional models to simulate changing~~ coastlines under a variety of ~~sea level~~sea
383 level, tidal forcing, and anthropogenic change scenarios (historical and future), to better
384 understand the tidal dynamics in Hong Kong, and to try to separate the relative importance of
385 local and regional effects. ~~Lastly, we quickly mention the instrumental changes at two of the~~
386 ~~HKO gauges. The Quarry Bay gauge was updated from a float type gauge recently (2017),~~
387 ~~and the Tai Po Kau gauge was also updated from a float gauge in 2006. Neither of these~~
388 ~~times correspond to any obvious anomalies in the tidal admittance records (the large changes~~
389 ~~at Tai Po Kau predate this by a few years at least, and are consistent before and after the~~
390 ~~gauge change), so we conclude that the instrumental changes were not a factor in the~~
391 ~~observed variability.~~

392 5. Conclusions

393 This study has presented new information about the tidal variability in Hong Kong,
394 based on observations of a set of closely-located tide gauges in Hong Kong. The TACs,
395 D_1/D_2 relations, δ -HATs, and the anomalous events in tidal amplitudes seen at the Quarry
396 Bay and Tai Po Kau gauges show an amplified tidal response to MSL fluctuations in these
397 harbor regions as opposed to more open-water locations, where individual TAC were
398 sometimes significant, but not as much for the δ -HATs. The reason for the observed behavior
399 may be due to changing friction or resonance induced by coastal engineering projects that are
400 only significant at highly local (i.e., individual harbor) scales. Alternatively, the observed
401 behavior could be related to regional ~~SCS-South China Sea~~ changes due to climate change
402 (such as increased upper-ocean warming and/or regional stratification and internal tide
403 generation) may also be a factor. It is difficult to separate the local engineering changes from
404 regional climatic changes without closer investigations. However, even without exact
405 knowledge of the relevant mechanisms, these anomalies do suggest that a pronounced change
406 in tidal properties occurred around the year 2000 in Hong Kong, with the effect being most
407 pronounced at gauges in semi-enclosed harbors. Overall, the tidal variability in Hong Kong
408 documented here may have significant impacts on the future of extreme sea level in the
409 region, especially if the strong positive reinforcements hold or increase in coming decades.
410 Short-term inundation events, such as nuisance flooding, may be amplified under scenarios of
411 higher ~~sea level~~sea levels that lead to corresponding changes in the tides, which may amplify

412 small changes in water levels and/or reductions in friction due to harbor improvements. The
413 δ -HATs and D₁/D₂ TACs results illustrate that the tidal variability of multiple constituents
414 ~~can may~~ be ~~positively-additive, and may~~ reinforce ~~MSL changes~~ at some locations, which
415 may further ~~agitate-aggravate~~ coastal flooding under MSL future rise. Since tides and storm
416 surge are both long-wave processes, the locations of strong tidal response may also
417 experience an exaggerated storm surge in the near future.

418

419 **Code availability** All code employed in this study was developed using MATLAB, version
420 R2011B. All code and methods can be provided upon request.

421 **Data Availability** The data used in this study from the Hong Kong Observatory (HKO;

422 <http://www.hko.gov.hk>) and the Hong Kong Marine Department (HKMD;

423 <http://www.mardep.gov.hk/en/home.html>) was provided upon request, discussion of

424 intentions of use, and permission from the appropriate agency supervisors. Data used from

425 the University of Hawaii Sea Level Center (UHSLC; <http://www.uhslc.soest.hawaii.edu>) is

426 publicly available.

427

428 Appendix

429 A1. Tidal Anomaly Correlations (TACs)

430 Tidal admittances are constructed as described above, employing the use of the tidal
431 potential and Eqs. (2) and (3) to constrain the nodal variation present in the observed tidal
432 amplitudes and phases. Our primary interest in this paper is the interannual to decadal
433 variations and not the long-term trends in mean values. Therefore, we first remove the long-
434 term trends and mean values using the MATLAB “detrend” function. The detrended time-
435 series of residual variations in A and P, and the residual variations in MSL, can now be
436 examined for coherence, using scatter plots, cross-correlations, and regression statistics. We
437 define the tidal anomaly correlation (TAC) as the slope between detrended tidal properties
438 (amplitude and phase) and detrended MSL, expressed as the millimeter change in tidal
439 amplitude per meter of sea level rise (mm m⁻¹). The same approach is used with the phase
440 difference time-series to provide phase anomaly trends, with the trends expressed as degree
441 change in tidal phase per meter of sea level rise (deg m⁻¹). The errors of the TAC
442 determinations are defined as the 95% confidence interval (CI) of the linear trend

Field Code Changed

Field Code Changed

Field Code Changed

443 determination. Trends are deemed significant if the signal-to-noise ratio (SNR) of the linear
444 trend to the associated error is greater than 2.0.

445 **A2. Approximate change in the highest astronomical tide (δ -HAT)**

446 We also construct a “proxy” quantity as an approximate change in the highest
447 astronomical tide (δ -HAT) using an extension of the TAC method. To do this, we combine
448 the tidal admittance amplitudes of the (typically) four largest astronomical tides (M_2 , S_2 , K_1 ,
449 and O_1), then detrend the resultant combined time series as above. Next, we perform a
450 similar scatterplot and regression approach against the detrended MSL time series as was
451 done with the TACs. The benefit of this approach is to give a clear picture of the overall
452 changes in tides related to sea level changes. Some locations may show that the variability in
453 multiple tidal constituents partially “cancel” each other (e.g., semidiurnal tides may have a
454 large positive tendency compared to MSL variability while diurnal tides may have a large
455 negative tendency, resulting in an offsetting of variabilities under MSL changes, and a
456 smaller overall magnitude δ -HAT), while other locations may show a “reinforced” variability
457 (e.g., both diurnal and semidiurnal tides have positive tendencies compared to MSL changes,
458 resulting in an amplified δ -HAT). Thus, the accurate interpretation of the δ -HAT is that it
459 reflects the maximum tide-related water level that would be obtained in a given analysis
460 period (here, one year) from a chosen set of time-dependent amplitudes extracted from the
461 admittance method.

462 Two details about the δ -HAT parameter should be noted here. First, only the
463 amplitude of the tidal admittance can be combined in this manner, as combining the phase
464 variability of multiple frequencies may be inaccurate at worst, and at best is not very helpful.
465 Second, we acknowledge that the use of the term “ δ -HAT” may be somewhat confusing, as
466 previous literature about tidal analysis uses the term “Highest Astronomical Tide” (HAT) to
467 denote the highest water level that can be expected to occur under average meteorological
468 conditions due purely to astronomical forcing in a given epoch. This typical period is 19
469 years, which considers the full nodal cycle. This definition of HAT does not reflect the
470 highest *possible* water level at a given location, since storm surge or other “non-average”
471 meteorological conditions may amplify water levels far above this level on a shorter time
472 scale than a 19-year determination can reveal. The intention behind our chosen nomenclature
473 of the “approximate change in the highest astronomical tide” (δ -HAT) attempts to expand on
474 this concept by considering the “full” tidal variability (not strictly true since the 4 largest

475 tides are only about 75% of the full tidal range, but these tidal components are nearly always
476 stable in one year analyses, so it is a dependable and easily comparable metric) at timescales
477 shorter than a nodal period (~19 years), but longer than a storm surge (~2-5 days) or other
478 meteorological anomalies. Furthermore, our interest is the changes in tidal components that
479 is not due to astronomy or to meteorology. Rather, we show possible changes to tide-related
480 water level modifications due to MSL modifications, which may be important on seasonal to
481 decadal time scales, induced by mechanisms associated with global climate change (e.g.,
482 steric sea level rise due to ice melt, thermal sea level rise due to upper-ocean warming), or to
483 more local effects (such as rapid harbor modifications or land reclamation that adjusts tidal
484 resonance at a particular location).

485 The changes shown by the δ -HATs are important to consider, since a full
486 understanding of the changes in all components and timescales of the tides may better instruct
487 future coastal planning and engineering. The δ -HAT method used here can give important
488 information about possible future water level inundation in coastal locations that are not
489 storm-related, such as nuisance flooding (or, sometimes called “sunny day flooding”). These
490 may be obscured by longer-term analyses of the classical HAT (i.e., 19 years) if changes are
491 more rapid (i.e., year-to-year or season-to-season). However, it should also be reiterated that
492 a good understanding of changes in tides due to changing background water levels may also
493 be instructive about future storm surge related inundation at a location; both tides and storms
494 are long wave processes, so changes in one aspect of water level variability (i.e., a large
495 positive δ -HAT) may also indicate future increase in storm surge levels at the same location.

496

497 **Author Contributions** ATD did all analyses, figures, tables, the majority of writing, and
498 complied the manuscript. JP provided editing, insight, guidance, and direction to this study.
499 HL provided critical insight and helpful input.

500 **Competing Interests** The authors declare they have no competing interests.

501 **Acknowledgements** This work is supported by The National Basic Research Program of
502 China (2015CB954103), the National Natural Science Foundation of China (project
503 41376035), the General Research Fund of Hong Kong Research Grants Council (RGC)
504 (CUHK 14303818, 402912, and 403113), the Hong Kong Innovation and Technology Fund
505 under the grants (ITS/259/12 and ITS/321/13), and the direct grants of the Chinese University

506 of Hong Kong. The authors also thank the Hong Kong Observatory. In addition to sharing
507 their data archive, they were also a part of the discussions that led to this paper.

508

509

510 **FIGURE CAPTIONS:**

511 **Figure 1** Tide gauge locations in Hong Kong used in this study. Green markers indicate
512 active gauges provided by the Hong Kong Observatory (HKO), light blue markers indicate
513 gauges provided by the Hong Kong Marine Department (HKMD), and red markers indicate
514 historical gauges (once maintained by HKO) that are no longer operational.

515 **Figure 2** Location of Hong Kong in the South China Sea (SCS), given by the red box, with
516 some major oceanographic features labelled. Depth is given by the color bar, in units of
517 meters.

518 **Figure 3** Tidal anomaly correlations (TACs) of detrended M_2 amplitude to detrended MSL in
519 Hong Kong, with the marker size showing the relative magnitude according to the legend, in
520 units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers
521 indicate TACs which are not significantly different from zero. ~~insignificant TACs.~~

522 **Figure 4** Tidal anomaly correlations (TACs) of detrended S_2 amplitude to detrended MSL in
523 Hong Kong, with the marker size showing the relative magnitude according to the legend, in
524 units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers
525 indicate TACs which are not significantly different from zero. ~~Black marks indicate~~
526 ~~insignificant TACs.~~

527 **Figure 5** Tidal anomaly correlations (TACs) of detrended K_1 amplitude to detrended MSL in
528 Hong Kong, with the marker size showing the relative magnitude according to the legend, in
529 units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers
530 indicate TACs which are not significantly different from zero. ~~Black marks indicate~~
531 ~~insignificant TACs.~~

532 **Figure 6** Tidal anomaly correlations (TACs) of detrended O_1 amplitude to detrended MSL in
533 Hong Kong, with the marker size showing the relative magnitude according to the legend, in
534 units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers
535 indicate TACs which are not significantly different from zero. ~~Black marks indicate~~
536 ~~insignificant TACs.~~

537 **Figure 7** The tidal anomaly correlation computed from the combination of the four largest
538 tidal constituent amplitudes (given by the detrended sum of the $M_2 + S_2 + K_1 + O_1$) as a proxy
539 for the change in the approximate highest astronomical tide (δ -HAT) relative to detrended
540 MSL in Hong Kong, with the marker size showing the relative magnitude according to the
541 legend, in units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black
542 markers indicate TACs which are not significantly different from zero. ~~Figure 7 The change~~
543 ~~in the highest astronomical tide (δ HAT), given by the detrended sum of the $M_2 + S_2 + K_1 +$~~
544 ~~O_1 amplitudes to detrended MSL in Hong Kong, with the marker size showing the relative~~

545 magnitude according to the legend, in units of mm m^{-1} . Black marks indicate insignificant
 546 TACs. ▲

547 **Figure 8** The OT TACs; the relations of detrended diurnal tidal amplitude sum (D_1 ; $K_1 + O_1$)
 548 to detrended semidiurnal tidal amplitude sum (D_2 ; $M_2 + S_2$) in Hong Kong, with the marker
 549 size showing the relative magnitude according to the legend, in dimensionless units. Red/blue
 550 markers indicate positive/negative TACs, and black markers indicate TACs which are not
 551 significantly different from zero. Black marks indicate insignificant TACs.

552 **Figure 9** Time series of water level spectrum components at the Quarry Bay (QB; blue) and
 553 Tai Po Kau (TPK; red) tide gauges in Hong Kong, showing the D_1 band (a), the D_2 band (b),
 554 the OT band (c) and mean ~~sea level~~ sea level (MSL) (d). Components are plotted as a function
 555 of normalized amplitudes to show relative variability, with mean values given in the legend.

556

557


558

559

560


561

562 **FIGURES:**




563

564 **Figure 1** Tide gauge locations in Hong Kong used in this study. Green markers indicate
565 active gauges provided by the Hong Kong Observatory (HKO), light blue markers indicate
566 gauges provided by the Hong Kong Marine Department (HKMD), and red markers indicate
567 historical gauges (once maintained by HKO) that are no longer operational.



568
569 **Figure 2** Location of Hong Kong in the South China Sea (SCS), given by the red box, with
570 some major oceanographic features labelled. Depth is given by the color bar, in units of
571 meters.


572



573

574 **Figure 3** Tidal anomaly correlations (TACs) of detrended M_2 amplitude to detrended MSL in
 575 Hong Kong, with the marker size showing the relative magnitude according to the legend, in
 576 units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers
 577 indicate TACs which are not significantly different from zero. Black marks indicate
 578 insignificant TACs.

Formatted: English (United States)




579

580 **Figure 4** Tidal anomaly correlations (TACs) of detrended S_2 amplitude to detrended MSL in
 581 Hong Kong, with the marker size showing the relative magnitude according to the legend, in

Formatted: Left

582 units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers
583 indicate TACs which are not significantly different from zero. Black marks indicate
584 insignificant TACs.


Formatted: English (United States)



585


586 **Figure 5** Tidal anomaly correlations (TACs) of detrended K_1 amplitude to detrended MSL in
587 Hong Kong, with the marker size showing the relative magnitude according to the legend, in
588 units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers
589 indicate TACs which are not significantly different from zero. Black marks indicate
590 insignificant TACs.

Formatted: English (United States)



591

592 **Figure 6** Tidal anomaly correlations (TACs) of detrended O_1 amplitude to detrended MSL in
 593 Hong Kong, with the marker size showing the relative magnitude according to the legend, in
 594 units of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers
 595 indicate TACs which are not significantly different from zero. Black marks indicate
 596 insignificant TACs.



597
 598 **Figure 7** ~~The change in the tidal anomaly correlation computed from the combination of the~~
 599 ~~four largest tidal constituent amplitudes (given by the detrended sum of the $M_2 + S_2 + K_1 +$~~
 600 ~~O_1) as a proxy for the change in the approximate highest astronomical tide (δ -HAT); ~~given by~~~~
 601 ~~the detrended sum of the $M_2 + S_2 + K_1 + O_1$ amplitudes relative~~ to detrended MSL in Hong
 602 Kong, with the marker size showing the relative magnitude according to the legend, in units
 603 of mm m^{-1} . Red/blue markers indicate positive/negative TACs, and black markers indicate
 604 TACs which are not significantly different from zero. Black marks indicate insignificant
 605 TACs.


Formatted: Left

Formatted: English (United States)

Formatted: Centered

Formatted: Left


Formatted: English (United States)



606

607 **Figure 8** The Θ_{D_1/D_2} TACs; the relations of detrended diurnal tidal amplitude sum (D_1 ; K_1
 608 + O_1) to detrended semidiurnal tidal amplitude sum (D_2 ; $M_2 + S_2$) in Hong Kong, with the
 609 marker size showing the relative magnitude according to the legend, in dimensionless units.
 610 Red/blue markers indicate positive/negative TACs, and black markers indicate TACs which
 611 are not significantly different from zero. Black marks indicate insignificant TACs.

Formatted: Left



612

613 **Figure 9** Time series of water level spectrum components at the Quarry Bay (QB; blue) and
 614 Tai Po Kau (TPK; red) tide gauges in Hong Kong, showing the D₁ band (a), the D₂ band (b),
 615 ~~the OT band (c)~~ and mean ~~sea level~~ sea level (MSL) (c). Components are plotted as a
 616 function of normalized amplitudes to show relative variability, with mean values given in the
 617 legend.

618

619

620

621

622

623

624

625

626

627

628

629

630

631 **REFERENCES:**

632 Alford, M. H.: Observations of parametric subharmonic instability of the diurnal internal tide
633 in the South China Sea. *Geophysical Research Letters*, 35, L15602, 2008.

634 doi:10.1029/2008GL034720

635 Amin, M.: On perturbations of harmonic constants in the Thames Estuary. *Geophysical*

636 *Journal of the Royal Astronomical Society*. 73(3): 587-603. doi:10.1111/j.1365-

637 246X.1983.tb03334.x, 1983.

638 Arbic, B.K., Karsten, R.H., Garrett, C.: On tidal resonance in the global ocean and the back-

639 effect of coastal tides upon open-ocean tides. *Atmosphere-Ocean* 47(4), 239–266.

640 doi:10.3137/OC311.2009, 2009.

641 Arns, A., Dangendorf, S., Jensen, J., Bender, J., Talke, S.A., & Pattiaratchi, C.: [Sea-level](#)[Sea](#)

642 [level](#) rise induced amplification of coastal protection design heights. *Nature: Scientific*

643 *Reports*, 7, 40171. doi:10.1038/srep40171, 2017.

644 Bowen, A. J., & Gray, D. A.: The tidal regime of the River Thames; long-term trends and

645 their possible causes. *Phil. Trans. R. Soc. Lond. A*, 272(1221), 187-199.

646 doi:10.1098/rsta.1972.0045, 1972.

647 Buchanan, M. K., Oppenheimer, M., & Kopp, R. E.: Amplification of flood frequencies with

648 local sea level rise and emerging flood regimes. *Environmental Research Letters*, 12(6),

649 064009. doi:10.1088/1748-9326/aa6cb3, 2017.

650 Cartwright, D.E., & Tayler, R.J.: New computations of the tide-generating potential. *Geophys.*

651 *Journal of the Royal Astronomical Society*, 23, 45-74. doi: 10.1111/j.1365-

652 246X.1971.tb01803.x, 1971.

653 Cartwright, D.E.: Secular changes in the oceanic tides at Brest, 1711–1936. *Geophysical*

654 *Journal International*, 30(4), 433-449. doi:10.1.1.867.2468, 1972.

655 Chernetsky, A. S., Schuttelaars, H. M., & Talke, S. A.: The effect of tidal asymmetry and

656 temporal settling lag on sediment trapping in tidal estuaries. *Ocean Dynamics*, 60(5), 1219-

657 1241.. doi: 10.1007/s10236-010-0329-8, 2010.

658 Cherqui, F., Belmeziti, A., Granger, D., Sourdril, A., & Le Gauffre, P.: Assessing urban
659 potential flooding risk and identifying effective risk-reduction measures. *Science of the Total*
660 *Environment*, 514, 418-425, 2015.

661 Chinn, B. S., Girton, J. B., & Alford, M. H.: Observations of internal waves and parametric
662 subharmonic instability in the Philippines archipelago. *Journal of Geophysical Research:*
663 *Oceans*, 117(C5). doi:10.1029/2011JC007392, 2012.

664 Church, J. A., & White, N. J.: A 20th century acceleration in global sea-level
665 rise. *Geophysical research letters*, 33(1). doi:10.1029/2005GL024826, 2006.

666 Church, J. A., & White, N. J.: ~~Sea level~~Sea level rise from the late 19th to the early 21st
667 century. *Surveys in geophysics*, 32(4-5), 585-602. doi 10.1007/s10712-011-9119-1, 2011.

668 Colosi, J. A., & Munk, W.: Tales of the venerable Honolulu tide gauge. *Journal of physical*
669 *oceanography*, 36(6), 967-996. doi:10.1175/JPO2876.1, 2006.

670 Craik, A.D.D.: Wave Interactions and Fluid Flows. Cambridge Univ. Press, Cambridge, U. K.,
671 ISBN: 978-0521368292, 1985.

672 Devlin, A. T., Jay, D. A., Talke, S. A., & Zaron, E.: Can tidal perturbations associated with
673 sea level variations in the western Pacific Ocean be used to understand future effects of tidal
674 evolution? *Ocean Dynamics*, 64(8), 1093-1120. doi:10.1007/s10236-014-0741-6, 2014.

675 Devlin, A. T., Jay, D. A., Zaron, E. D., Talke, S. A., Pan, J., & Lin, H.: Tidal variability
676 related to sea level variability in the Pacific Ocean. *Journal of Geophysical Research:*
677 *Oceans*, 122(11), 8445-8463. doi:10.1002/2017JC013165, 2017.

678 Devlin, A. T., Jay, D. A., Talke, S. A., Zaron, E. D., Pan, J., & Lin, H.: Coupling of sea level
679 and tidal range changes, with implications for future water levels. *Scientific Reports*, 7(1),
680 17021. doi:10.1038/s41598-017-17056-z, 2017.

681 Devlin, A. T., Zaron, E. D., Jay, D. A., Talke, S. A., & Pan, J.: Seasonality of Tides in
682 Southeast Asian Waters. *Journal of Physical Oceanography*. doi: 10.1175/JPO-D-17-0119.1,
683 2018.

684 Devlin, A. T., Pan, J., & Lin, H.: Extended spectral analysis of tidal variability in the North
685 Atlantic Ocean. *Journal of Geophysical Research: Oceans*, 124(1), 506-526, 2019.

686 Domingues, C. M., Church, J. A., White, N. J., Gleckler, P. J., Wijffels, S. E., Barker, P. M.,
687 & Dunn, J. R.: Improved estimates of upper-ocean warming and multi-decadal ~~sea level~~
688 ~~sea level~~ rise. *Nature*, 453(7198), 1090. doi:10.1038/nature07080, 2008.

689 Haigh, I. D., Wijeratne, E. M. S., MacPherson, L. R., Pattiaratchi, C. B., Mason, M. S.,
690 Crompton, R. P., & George, S.: Estimating present day extreme water level exceedance
691 probabilities around the coastline of Australia: tides, extra-tropical storm surges and mean sea
692 level. *Climate Dynamics*, 42(1-2), 121-138. doi: 10.1007/s00382-012-1652-1, 2014.

693 Familkhalili, R., & Talke, S. A.: The effect of channel deepening on tides and storm surge: A
694 case study of Wilmington, NC. *Geophysical Research Letters*, 43(17), 9138-9147.
695 doi:10.1002/2016GL069494, 2016.

696 Fang, G., Kwok, Y. K., Yu, K., & Zhu, Y.: Numerical simulation of principal tidal
697 constituents in the South China Sea, Gulf of Tonkin and Gulf of Thailand. *Continental Shelf*
698 *Research*, 19(7), 845-869. doi: 10.1016/S0278-4343(99)00002-3, 1999.

699 Feng, X., Tsimplis, M. N., & Woodworth, P. L.: Nodal variations and long-term changes in
700 the main tides on the coasts of China. *Journal of Geophysical Research: Oceans*, 120(2),
701 1215-1232. doi:10.1002/2014JC010312, 2015.

702 Ip, S.F. and Wai, H.G.: *An application of harmonic method to tidal analysis and prediction in*
703 *Hong Kong*. Royal Observatory, 1990.

704 Jan, S., Chern, C. S., Wang, J., & Chao, S. Y.: Generation of diurnal K₁ internal tide in the
705 Luzon Strait and its influence on surface tide in the South China Sea. *Journal of Geophysical*
706 *Research: Oceans*, 112(C6). doi:10.1029/2006JC004003, 2007.

707 Jan, S., Lien, R. C., & Ting, C. H.: Numerical study of baroclinic tides in Luzon
708 Strait. *Journal of Oceanography*, 64(5), 789. doi:10.1007/s10872-008-0066-5, 2008.

709 Jay, D. A. (2009). Evolution of tidal amplitudes in the eastern Pacific Ocean. *Geophysical*
710 *Research Letters*, 36(4). doi: 10.1029/2008GL036185

711 Leffler, K. E., & Jay, D. A.: Enhancing tidal harmonic analysis: Robust (hybrid L1/L2)
712 solutions. *Continental Shelf Research*, 29(1), 78-88. doi: 10.1016/j.csr.2008.04.011, 2009.

713 Li, K. W., & Mok, H. Y.: Long term trends of the regional sea level changes in Hong Kong
714 and the adjacent waters. In *Asian And Pacific Coasts 2011* (pp. 349-359).
715 doi:10.1142/9789814366489_0040, 2012.

716 Lien, R. C., Tang, T. Y., Chang, M. H., & d'Asaro, E. A.: Energy of nonlinear internal waves
717 in the South China Sea. *Geophysical Research Letters*, 32(5). doi:10.1029/2004GL022012,
718 2005.

719 MacKinnon, J. A., & Winters, K. B.: Subtropical catastrophe: Significant loss of low-mode
720 tidal energy at 28.9°. *Geophysical Research Letters*, 32(15). doi:10.1029/2005GL023376,
721 2005.

722 Mawdsley, R. J., Haigh, I. D., & Wells, N. C.: Global changes in mean tidal high water, low
723 water and range. *Journal of Coastal Research*, 70(sp1), 343-348. doi:10.2112/SI70-058.1,
724 2014.

725 McComas, C. H., & Bretherton, F. P.: Resonant interaction of oceanic internal
726 waves. *Journal of Geophysical Research*, 82(9), 1397-1412. doi:10.1029/JC082i009p01397,
727 1977.

728 Moftakhari, H. R., AghaKouchak, A., Sanders, B. F., Feldman, D. L., Sweet, W., Matthew,
729 R. A., & Luke, A.: Increased nuisance flooding along the coasts of the United States due to
730 sea level rise: Past and future. *Geophysical Research Letters*, 42(22), 9846-9852.
731 doi:10.1002/2015GL066072, 2015.

732 Moftakhari, H. R., AghaKouchak, A., Sanders, B. F., & Matthew, R. A.: Cumulative hazard:
733 The case of nuisance flooding. *Earth's Future*, 5(2), 214-223. doi:10.1002/2016EF000494,
734 2017.

735 Müller, M., Arbic, B. K., & Mitrovica, J. X.: Secular trends in ocean tides: Observations and
736 model results. *Journal of Geophysical Research: Oceans*, 116(C5) doi:
737 10.1029/2010JC006387, 2011.

738 Müller, M., Cherniawsky, J. Y., Foreman, M. G. G., & Storch, J. S.: Global M₂ internal tide
739 and its seasonal variability from high resolution ocean circulation and tide
740 modeling. *Geophysical Research Letters*, 39(19). doi:10.1029/2012GL053320, 2012.

741 Müller, M.: The influence of changing stratification conditions on barotropic tidal transport
742 and its implications for seasonal and secular changes of tides. *Continental Shelf Research*, 47,
743 107-118. doi: 10.1016/j.csr.2012.07.003, 2012.

744 Pan, J., Gu, Y. and Wang, D.: Observations and numerical modeling of the Pearl River plume
745 in summer season, *Journal of Geophysical Research*, 119, doi:10.1002/2013JC009042, 2014.

746 Pawlowicz, R., Beardsley, B., & Lentz, S.: Classical tidal harmonic analysis including error
747 estimates in MATLAB using T_TIDE. *Computers & Geosciences*, 28(8), 929-937.
748 doi:10.1016/S0098-3004(02)00013-4, 2002.

749 Rasheed, A. S., & Chua, V. P.: Secular trends in tidal parameters along the coast of
750 Japan. *Atmosphere-Ocean*, 52(2), 155-168. doi:10.1080/07055900.2014.886031, 2014.

751 Ray, R. D.: Secular changes of the M₂ tide in the Gulf of Maine. *Continental shelf*
752 *research*, 26(3), 422-427. doi: 10.1016/j.csr.2005.12.005, 2006.

753 Ray, R. D., & Foster, G.: Future nuisance flooding at Boston caused by astronomical tides
754 alone. *Earth's Future*, 4(12), 578-587. doi:10.1002/2016EF000423, 2016.

755 Ross, A. C., Najjar, R. G., Li, M., Lee, S. B., Zhang, F., & Liu, W.: Fingerprints of Sea Level
756 Rise on Changing Tides in the Chesapeake and Delaware Bays. *Journal of Geophysical*
757 *Research: Oceans*, 122(10), 8102-8125. doi:10.1002/2017JC012887, 2017.

758 Sweet, W. V., & Park, J.: From the extreme to the mean: Acceleration and tipping points of
759 coastal inundation from sea level rise. *Earth's Future*, 2(12), 579-600, 2014.

760 Vellinga, N. E., Hoitink, A. J. F., van der Vegt, M., Zhang, W., & Hoekstra, P.: Human
761 impacts on tides overwhelm the effect of sea level rise on extreme water levels in the Rhine-
762 Meuse delta. *Coastal Engineering*, 90, 40-50. doi: 10.1016/j.coastaleng.2014.04.005, 2014.

763 Woodworth, P. L.: A survey of recent changes in the main components of the ocean
764 tide. *Continental Shelf Research*, 30(15), 1680-1691. doi: 10.1016/j.csr.2010.07.002, 2010.

765 Xie, X. H., Chen, G. Y., Shang, X. D., & Fang, W. D.: Evolution of the semidiurnal (M₂)
766 internal tide on the continental slope of the northern South China Sea. *Geophysical Research*
767 *Letters*, 35(13). doi:10.1029/2008GL034179, 2008.

768 Xie, X. H., Shang, X. D., van Haren, H., Chen, G. Y., & Zhang, Y. Z.: Observations of
769 parametric subharmonic instability-induced near-inertial waves equatorward of the critical
770 diurnal latitude. *Geophysical Research Letters*, 38(5). doi:10.1029/2010GL046521, 2011.

771 Xie, X., Shang, X., Haren, H., & Chen, G.: Observations of enhanced nonlinear instability in
772 the surface reflection of internal tides. *Geophysical Research Letters*, 40(8), 1580-1586.
773 doi:10.1002/grl.50322, 2013.

774 Zaron, E. D., & Jay, D. A.: An analysis of secular change in tides at open-ocean sites in the
775 Pacific. *Journal of Physical Oceanography*, 44(7), 1704-1726. doi:10.1175/JPO-D-13-0266.1,
776 2014.

777 Zhang, H., Wang, T., Liu, M., Jia, M., Lin, H., Chu, L. M., & Devlin, A. T.: Potential of
778 Combining Optical and Dual Polarimetric SAR Data for Improving Mangrove Species
779 Discrimination Using Rotation Forest. *Remote Sensing*, 10(3), 467. doi: 10.3390/rs10030467,
780 2018.

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805
806
807
808

809 **TABLES:**

810 **Table 1** Metadata for all tide gauge locations, giving the station names and station codes
811 giving latitude/longitude, year of the available records, as well as the range of data
812 analyzed and start year/end year of the full record, as well as of data analyzed in this study.

Station	Latitude	Longitude	Start Year	End Year	Number of years used
Quarry Bay (QB)	22.27° N	114.21° E	1954	2016	310 (1986-2016)
Tai Po Kau (TPK)	22.42° N	114.19° E	1963	2016	310 (1986-2016)
Tsim Bei Tusi (TBT)	22.48° N	114.02° E	1974	2016	310 (1986-2016)
Chi Ma Wan (CMW)	22.22° N	114.00° E	1963	1997	36 (1963-1997)
Cheung Chau (CHC)	22.19° N	114.03° E	2004	2016	12 (2004-2016)
Lok On Pai (LOP)	22.35° N	114.00° E	1981	1999	18 (1981-1999)
Ma Wan (MW)	22.35° N	114.06° E	2004	2016	12 (2004-2016)
Tai Miu Wan (TMW)	22.26° N	114.29° E	1996	2016	20 (1996-2016)
Shek Pik (SP)	22.21° N	113.89° E	1999	2016	17 (1999-2016)
Waglan Island (WAG)	22.17° N	114.30° E	1995	2016	21 (1995-2016)
Ko Lau Wan (KLW)	22.45° N	114.34° E	2004	2016	12 (2004-2016)
Kwai Chung (KC)	22.31° N	114.12° E	2004	2016	12 (2004-2016)

813

814 **Table 2** Amplitude TACs for M₂, S₂, K₁, and O₁ for the period of 1986-2016. All values
815 given are in units of millimeter change in tidal amplitude for a 1-meter fluctuation in sea-
816 level/sea level (mm m⁻¹). Statistically significant positive values are given in bold italic text.

Station	M ₂ TAC	S ₂ TAC	K ₁ TAC	O ₁ TAC
Quarry Bay (QB)	+218 ± 37	+85 ± 16	+220 ± 15	+146 ± 11
Tai Po Kau (TPK)	+267 ± 42	+98 ± 17	+190 ± 68	+100 ± 25
Tsim Bei Tusi (TBT)	+7 ± 80	-10 ± 15	+32 ± 22	+24 ± 22
Chi Ma Wan (CMW)	-58 ± 11	-7 ± 5	-18 ± 8	-37 ± 10
Cheung Chau (CHC)	-63 ± 20	-22 ± 35	+69 ± 48	+50 ± 92
Lok On Pai (LOP)	-81 ± 24	-18 ± 8	+8 ± 32	-24 ± 12
Ma Wan (MW)	-68 ± 4	+1 ± 25	+52 ± 4	-62 ± 21
Tai Miu Wan (TMW)	+22 ± 59	-1 ± 9	+10 ± 22	+3 ± 8
Shek Pik (SP)	+62 ± 29	+11 ± 18	+70 ± 4	+28 ± 17
Waglan Island (WAG)	+1 ± 21	+3 ± 6	+9 ± 7	-9 ± 8
Ko Lau Wan (KLW)	-46 ± 39	-11 ± 17	+29 ± 65	+60 ± 57
Kwai Chung (KC)	-90 ± 46	-10 ± 29	-91 ± 226	-202 ± 161

817
818
819
820
821

822 **Table 3** The δ -HAT and D_1/D_2 TACs for the period of 1986-2016. The δ -HAT values given
823 are in units of millimeter change in tidal amplitude for a 1-meter fluctuation in ~~sea level~~
824 level (mm m⁻¹). D_1/D_2 TACs are in unitless ratios (i.e., mm mm⁻¹) Statistically significant
825 values are given in bold italic text.

<i>Station</i>	δ -HAT	D_1/D_2
<i>Quarry Bay (QB)</i>	<i>+665 ± 82</i>	<i>+1.08 ± 0.05</i>
<i>Tai Po Kau (TPK)</i>	<i>+612 ± 210</i>	<i>+1.01 ± 0.04</i>
<i>Tsim Bei Tusi (TBT)</i>	+56 ± 117	<i>+0.37 ± 0.02</i>
<i>Chi Ma Wan (CMW)</i>	<i>-119 ± 19</i>	<i>+0.74 ± 0.19</i>
<i>Cheung Chau (CHC)</i>	-12 ± 42	+0.81 ± 1.03
<i>Lok On Pai (LOP)</i>	-114 ± 45	<i>+0.26 ± 0.05</i>
<i>Ma Wan (MW)</i>	<i>-91 ± 73</i>	+0.57 ± 1.02
<i>Tai Miu Wan (TMW)</i>	+42 ± 100	<i>+1.04 ± 0.20</i>
<i>Shek Pik (SP)</i>	<i>+138 ± 37</i>	<i>+0.89 ± 0.06</i>
<i>Waglan Island (WAG)</i>	+3 ± 31	<i>+1.11 ± 0.17</i>
<i>Ko Lau Wan (KLW)</i>	-66 ± 47	<i>+1.31 ± 0.62</i>

826