

CO₂ effects on diatoms: A Synthesis of more than a decade of ocean acidification experiments with natural communities

Lennart. T. Bach^{1,2*}, Jan. Taucher¹

¹Biological Oceanography, GEOMAR, Helmholtz Centre for Ocean Research, Kiel, Germany

²Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

*Corresponding Author lennart.bach@utas.edu.au

Keywords: Ocean acidification, climate change, mesocosm, literature review, food web, phytoplankton, Bacillariophyceae

Short summary

Diatoms are a group of phytoplankton species responsible for ~20% of primary production on Earth. Ocean acidification (OA) could influence diatoms but the key question is if they become more or less important within marine food webs. Here we synthesize OA experiments with natural communities and found: Diatoms are more likely to be positively than negatively affected by OA and larger species may profit more than smaller ones. This has important implications for ecosystem services diatoms provide.

Abstract

Diatoms account for ~40% of marine primary production and are considered to be key players in the biological carbon pump. Ocean acidification (OA) is expected to affect diatoms primarily by changing the availability of CO₂ as a substrate for photosynthesis

or through altered ecological interactions within the marine food web. Yet, there is little consensus how entire diatom communities will respond to increasing CO₂. To address this question, we synthesized the literature from over a decade of OA-experiments with natural diatom communities to uncover: 1) if and how bulk diatom communities respond to elevated CO₂ with respect to abundance or biomass; 2) if shifts within the diatom communities could be expected and how they are expressed with respect to taxonomic affiliation and size structure. We found that bulk diatom communities responded to high CO₂ in ~60 % of the experiments and in this case more often positively (56 %) than negatively (32 %; 12 % did not report the direction of change). Shifts among different diatom species were observed in 65 % of the experiments. Our synthesis supports the hypothesis that high CO₂ particularly favors larger species as 12 out of 13 experiments which investigated cell size found a shift towards larger species. Unraveling winners and losers with respect to taxonomic affiliation was difficult due to a limited database. We conclude that OA-induced changes in diatom competitiveness and assemblage structure may alter key ecosystem services due to the pivotal role diatoms play in trophic transfer and biogeochemical cycles.

1. Introduction

The global net primary production (NPP) of all terrestrial and marine autotrophs amounts to approximately 105 petagrams (Pg) of carbon per year (Field et al., 1998). Marine diatoms, a taxonomically diverse group of cosmopolitan phytoplankton, were estimated to contribute up to 25 % (26 Pg C year⁻¹) to this number, which is more than the annual primary production in any biome on land (Field et al., 1998; Nelson et al., 1995; Tréguer and De La Rocha, 2013). Thus, diatoms are likely the most important single taxonomic group of primary producers on Earth and any change in their prevalence relative to other phytoplankton taxa could profoundly alter marine food web structures and thereby affect

ecosystem services such as fisheries or the sequestration of CO₂ in the deep ocean (Armbrust, 2009; Tréguer et al., 2018).

The most conspicuous feature of diatoms is the formation of a silica shell, which is believed to primarily serve as protection against grazers (Hamm and Smetacek, 2007; Pančić and Kiørboe, 2018). Since the formation of this shell requires dissolved silicate, diatoms are often limited by silicon as a nutrient rather than by nitrogen or phosphate (Brzezinski and Nelson, 1996). However, when dissolved silicate is available, diatoms benefit from their high nutrient uptake and growth rates, allowing them to outcompete other phytoplankton and form intense blooms in many ocean regions (Sarhou et al., 2005).

Diatoms display an enormous species richness, with recent estimates accounting for so far undiscovered diatoms (including freshwater) being in the range of 20,000 – 100,000 species (Guiry, 2012; Mann and Vanormelingen, 2013). Sournia et al. (1991) derive a number between 1400 – 1800 of described marine diatoms based on microscopy while *Tara* Oceans reported ~4700 operational taxonomic units from genetic samples distributed over all major oceans except the North Atlantic and North Pacific (Malviya et al., 2016). Known diatom taxa span a size range of several orders of magnitude (<5 µm up to a few mm) with a wide range of morphologies and life strategies, e.g. single cells and cell chains, pelagic and benthic habitats (Armbrust, 2009; Mann and Vanormelingen, 2013; Sournia et al., 1991). Accordingly, they should not be treated as one functional group, but rather as a variety of subgroups occupying different niches.

It is well recognized that the global importance of diatoms as well as their diversity in morphology and life style is tightly linked to the functioning of pelagic food webs and elemental cycling in the oceans. For example, iron enrichment experiments in the

Southern Ocean found that a shift in diatom community composition from thick- to thin-shelled species (“persistence strategy” vs. “boom-and-bust strategy”) can enhance carbon and alter nutrient export via sinking particles (Assmy et al., 2013; Smetacek et al., 2012). This may not only affect element fluxes locally but enhance nutrient retention within the Southern Ocean and reduce productivity in the north which underlines how important diatom community shifts can be on a global scale (Boyd, 2013; Primeau et al., 2013; Sarmiento et al., 2004). Likewise, the cell size of diatoms can play an important role in transferring energy to higher trophic levels, as the dominance of larger species is generally considered to reduce the length of the food chain and lead to higher trophic transfer efficiency (Sommer et al., 2002). Consequently, understanding impacts of global change on diatom community composition is crucial for assessing the sensitivity of biogeochemical cycles and ecosystem services in the world oceans.

It has become evident that the sensitivity of diatoms to increasing $p\text{CO}_2$ is highly variable, likely being related to specific traits such as cell size or the carbon fixation pathway, as well as interactions with other environmental factors such as nutrient stress, temperature or light (Gao et al., 2012; Hoppe et al., 2013; Wu et al., 2014). However, it is still rather unclear how these species-specific differences in CO_2 sensitivities manifest themselves on the level of diatom communities. This knowledge gap motivated us to compile the presently available experimental data in order to reveal common responses of diatom communities to high CO_2 and thereby assess potential scenarios of shifts in diatom community composition under ocean acidification.

2. Literature investigation

2.1. Approach

Our original intention was to conduct a classical meta-analysis, which would have yielded

the benefit of a quantitative measure of diatom responses to OA, expressed as an overall effect size (i.e. combined magnitude) such as the response ratio. However, our literature analysis revealed a large variability in experimental pCO₂ ranges as well as measured response variables, which cannot be directly compared among each other (e.g. microscopic cell counts, pigment concentrations, genetic tools). These limitations impede data aggregation as required for a classical meta-analysis. Furthermore, experimental setups differed widely in terms of other environmental factors such as temperature, light, and nutrient concentrations, all of which are known to modulate potential responses to pCO₂ (Boyd et al., 2018), thereby further complicating data aggregation for meta-analysis. Therefore, we chose an alternative, semi-quantitative approach where diatom responses to increasing CO₂ are grouped in categories (see section 2.2) and also allows to account for differences in experimental setups, e.g. with respect to container volume. While this approach excludes the determination of effect size, it provides an unbiased insight on the direction of change of potential CO₂ effects.

2.2. Data compilation

We explored the response of diatom assemblages to high CO₂ (low pH) by searching the literature for relevant results with Google Scholar (December 15, 2017) using the following search query: diatom OR Bacillariophyceae AND "ocean acidification" OR "high CO₂" or "carbon dioxide" OR "elevated CO₂" OR "elevated carbon dioxide" OR "low pH" OR "decreased pH". The first 200 results were inspected and considered to be relevant when they were published in peer-reviewed journals, contained a description of the relevant methodological details, a statistical analysis or at least a transparent description of variance and uncertainties, and tested CO₂ effects on natural plankton assemblages (artificially composed communities were not considered). We then carefully checked the cited literature in these relevant studies to uncover other studies that were

missed by the initial search. Furthermore, we checked the “Ocean Acidification news stream provided by the Ocean Acidification International Coordination Centre” under the tag “phytoplankton” (<https://news-oceanacidification-icc.org/tag/phytoplankton/>) for relevant updates since December 2017 (last check on January 16, 2019).

There were two response variables of interest for the literature compilation:

1) The response of the “bulk diatom community” to high CO₂. For this we checked if the abundance of diatoms, the biomass of diatoms, or the relative portion of diatoms within the overall phytoplankton assemblage increased or decreased under high CO₂ relative to the control. We distinguished between “positive”, “negative”, and “no effect” following the statistical results provided in the individual references. When the CO₂ effect on the bulk community was derived from abundance data we also checked if there are indications for a concomitant shift in the biomass distribution among species. This is relevant because, for example, an increase in bulk abundance could coincide with a decrease in bulk biomass when the species driving the abundances is smaller. We found no indications for conflicting cases but acknowledge that not every reference provided sufficient data on morphological details to fully exclude this scenario. Furthermore, we emphasize that CO₂ can also shift the temporal occurrence of a diatom response (Bach et al., 2017). For example, a diatom bloom could occur earlier in a high CO₂ treatment than in the control but with a similar bloom amplitude (Donahue et al., 2019). In this case we assigned a “positive” response because an earlier bloom occurrence mirrors a higher net growth rate under elevated CO₂.

2) The CO₂-dependent species shifts within the diatom community with respect to taxonomic composition and/or size structure. Unfortunately, cell size of the species was not reported for all experiments. Thus, we distinguished between “no shifts”, “shifts

between species with unspecified size”, as well as “shifts towards larger or smaller species” when this information was provided. Furthermore, we noted the winners and losers within the diatom communities when these were reported (on the genus level).

In case the data was taken from factorial multiple stressor experiments (e.g. CO₂ x temperature) we only considered the control treatment at ambient conditions (e.g. at control temperature). Furthermore, we extracted various metadata from each study largely following the literature analysis of Schulz et al. (2017). All bulk diatom responses, community shifts, and metadata is compiled/described in Table 1 and most of it is self-explanatory (e.g. incubation temperature). The coordinates from where the investigated plankton communities originate are given in Table 1 and illustrated in Figure 2. Their habitats were categorized according to water depth, salinity, or life style in the case of benthic communities: “oceanic” = water depth > 200 m (unless the habitat lies within a fjord or fjord-like strait), S > 30; “coastal” = water depth < 200 m, S > 30; “estuarine” = water depth < 200 m, S < 30; “benthic” = benthic communities (diatoms growing on plates) were investigated. We reconstructed the water depth in case it was not provided in the paper using Google Earth Pro (version 7.3.2.5495). The coordinates provided in some of the experiments conducted in land-based facilities were imprecise and marked positions on land. In this case the habitats were set to coastal or estuarine depending on salinity. If salinity was not given we checked the location on Google Earth for potential fresh water sources and also checked the text for more cryptic indications (e.g. “euryhaline” in a lagoon were strong indications for an estuarine habitat). The methods with which responses of the bulk diatom communities to high OA were determined varied greatly among studies and included light microscopy (LM), pigment analyses (PA), flow cytometry (FC), genetic tools (PCR), and biogenic silica (BSi) analyses (Table 1).

2.3. Accounting for different experimental setups to balance the influence of

individual studies on the outcome of the literature analysis

The most realistic OA experiment would be one where all aspects of the natural habitat are represented correctly. Such setups are possible for benthic communities which can be sampled *in situ* along a natural CO₂ gradient at volcanic CO₂ seeps (Fabricius et al., 2011; Hall-Spencer et al., 2008; Johnson et al., 2011). However, pelagic communities are advected with currents so that it is very difficult to simulate OA in open waters. Thus, OA experiments where pelagic communities are exposed to increasing levels of CO₂ were so far always performed in closed containers even though it is well known that confinement causes experimental artefacts (Calvo-Díaz et al., 2011; Ferguson et al., 1984; Guangao, 1990; Menzel and Case, 1977). The degree by which confinement causes experimental artefacts will differ from study to study depending on factors such as the incubation volume, the length of incubation, or the selective removal of certain size classes from the incubation (Carpenter, 1996; Duarte et al., 1997; Nogueira et al., 2014). In our literature synthesis we had to deal with a large variety of experimental setups and there are very likely differences how well a given setup represents the natural environment. Therefore, we aimed to develop a metric that allows us to estimate “how well the natural system (which we are ultimately interested in) is represented by the experimental setup”. This metric – termed the “relative degree of realism (RDR)” – was used to balance the influence of individual studies on the final outcomes of the literature analysis. Most certainly, we do not mean to devalue any studies but think that the highly different scales of experiments, ranging from 0.8 L lab incubations to 75 m³ *in situ* mesocosms, should not be ignored when evaluating the literature. In the following we will first derive the equation for the RDR and introduce the underlying assumptions. Afterwards we describe aspects that were considered while conceptualizing the RDR.

The incubation volume in the studies considered herein ranged from bottle experiments

to *in situ* mesocosm studies with considerably larger incubation volumes. Smaller differences in incubation volumes (e.g. 0.5 vs. 2 L) were shown to have no, or a minor, influence on physiological rates (Fogg and Calvario-Martinez, 1989; Hammes et al., 2010; Nogueira et al., 2014; Robinson and Williams, 2005). However, they can influence food web composition (Calvo-Díaz et al., 2011; Spencer and Warren, 1996), e.g. by unrepresentatively including certain organism groups such as highly motile mesozooplankton. Larger differences of incubation volumes (e.g. 10 vs. 10000 L) are considered to have a major influence on the enclosed communities, with the larger volume being more representative of natural processes (Carpenter, 1996; Duarte et al., 1997; Sarnelle, 1997). Therefore, our first assumption to conceptualize the RDR was that larger incubation volumes represent nature generally better than smaller ones.

Plankton communities were pre-filtered in many experiments to exclude larger and often patchily distributed organisms (e.g. copepods). This is a valid procedure to reduce noise and to increase the likelihood to detect CO₂ effects but it also influences the development of plankton communities since the selective removal of certain size classes can modify trophic cascades within the food web (Ferguson et al., 1984; Nogueira et al., 2014). For example, Nogueira et al. (2014) compared plankton successions of pre-filtered (100 µm) and unfiltered communities and found that the removal of larger grazers and diatoms gave room for green algae and picophytoplankton to grow. Such manipulations make the experiment less representative for a natural food web which brought us to the second assumption for the RDR: The smaller the mesh size during the pre-filtration treatment, the less complete and thus the less realistic is the pelagic food web.

To parameterize the two abovementioned assumptions we first converted the volume information provided in each experiment into a volume-to-surface ratio (V/S). The underlying thought is that V increases with the third power to the surface area of the

incubator and is indicative for the relation of open space to hard surfaces. Therefore we first converted V into a radius (r) assuming spherical shape:

$$r = \sqrt[3]{\frac{3V}{4\pi}} \quad (1).$$

The surface (S) of the spherical volume was calculated as:

$$S = 4\pi r^2 \quad (2)$$

The assumption of spherical shape was necessary because it allowed us to calculate V/S from only knowing V which is usually the only parameter provided with respect to container characteristics. We are aware that this is a simplification because the majority of containers used in experiments will likely have had cylindrical shape. However, the conversion from volume to surface assuming cylindrical shape would have required knowledge of two dimensions (radius and height of the cylinder). Although shape can influence processes within the container (Pan et al., 2015), it is a less important factor to consider in our study because sensitivity calculations assuming reasonable cylinder dimensions showed that the V/S differences due to container shape will be small compared to the V/S differences due to the range of container volumes compared here.

The influence of pre-filtration treatments of the investigated plankton community is implemented by multiplying the V/S with the cube root of the applied mesh size (d_{mesh} in μm) so that the RDR is defined as:

$$\text{RDR} = \frac{V}{S} \sqrt[3]{d_{\text{mesh}}} \quad (3).$$

Thus, as for V/S, the influence of d_{mesh} on RDR does not increase linearly but becomes less influential with increasing d_{mesh} . The rationale for the non-linear increase is that

incubations will still have an increasing bias even if they do not have any pre-filtration treatment due to generally increasing organism motility with size. For example, when collecting a plankton community with a Niskin bottle, more motile organisms can escape from the approaching sampler so that the food web composition is still affected even without subsequent pre-filtration. For this reason we also capped the maximum d_{mesh} to 10,000 μm when there was no pre-filtration treatment applied since none of the studies included significantly larger organisms. The rationale for calculating the cube root of d_{mesh} was that in this case the influence of V/S and d_{mesh} on RDR becomes roughly similar. Figure 1 illustrates the change of RDR as a function of V and d_{mesh} . High RDRs are calculated for large-scale *in situ* mesocosm studies ($\sim 50 - 190$) while bottle experiments yield RDRs between $\sim 1 - 12$.

The key pre-requisite for an experimental parameter to be included in the RDR equation (eq. 3) was that it is reported in all studies. Many parameters that we would have liked to use for the RDR are either insufficiently reported (e.g. the light environment) or not provided quantitatively at all (e.g. turbulence). We therefore had to work with very basic properties related to the experimental setup rather than to the experimental conditions.

A particularly critical aspect of the RDR we had to deal with was the duration of the experiments (Time). Time is reliably reported in all studies and therefore principally suitable for the RDR. Our first thoughts were that a realistic community experiment should be long enough to cover relevant ecological processes such as competitive exclusion and therefore also parameterized Time in the first versions of the RDR equation. However, we decided to not account for it in the final version because the factors that define the optimal duration of an experiment are poorly constrained. For example, a 1 day experiment in a 10 L container could indeed miss important CO_2 effects caused by food web interactions. On the other hand, a 30 days experiment in the same

container could reveal such indirect effects but at the same time be associated with profound bottle effects and make the study unrepresentative for simulated natural habitat. Thus, too long and too short are both problematic and the optimum is hard to find. One such attempt to find the optimum Time was made by (Duarte et al., 1997) who analyzed the plankton ecology literature between 1990 – 1995. By correlating the experimental duration with the incubation volume of published experiments they provided an optimal length for any given volume. However, as noted by Duarte et al. (1997), their correlation is based on publication success and therefore rather reflects common practice in plankton ecology experiments and not necessarily a mechanistic understanding of bottle effects. Thus, as there is no solid ground for a parameterization of Time we ultimately decided to not consider it for the RDR.

Finally, we want to point out (and explicitly acknowledge) that the RDR approach to balance the influence of studies on the final outcome of the literature analysis is of course not the one perfect solution and most likely incomplete (see above). However, balancing a literature analysis with the RDR score may still be an improvement relative to the other case where each experiment is treated exactly equal despite huge differences in the experimental setup. Nevertheless, to account for both views (i.e. the RDR is useless vs. the RDR is useful) we will present the outcome of our literature analysis in two different ways throughout the paper: 1) by simply counting the number of outcomes and adding them to yield a cumulative score (N-based approach; left columns in Figs. 3 and 4); 2) by adding the RDR score of the experiments with a certain outcome to yield a cumulative Σ RDR score (RDR-based approach; right columns in Figs. 3 and 4).

3. Results

We found 54 relevant publications on CO₂ experiments with natural diatom assemblages.

Some publications included more than one experiment so that 69 experiments are considered hereafter (Table 1). Most were done with diatom communities from coastal environments (46 %) and oceanic (28%) environments. Estuarine and benthic communities were investigated in 16 % and 6% of the studies, respectively. 4 % of the studies did not provide coordinates where the samples were taken although the region was reported (Table 1; Fig. 2).

Among the 69 experiments, 23 (33 %, $\sum RDR = 595$) revealed a positive influence of CO₂ on the “bulk diatom community” (see section 2.2), while 13 (19 %, $\sum RDR = 266$) revealed a negative one. 5 experiments (7 %, $\sum RDR = 21$) found a CO₂ effect but did not specify whether it is a positive or negative one. 28 experiments (41 %, $\sum RDR = 706$) found no effect (Fig. 3A).

We also checked if the pCO₂ range tested in the experiments had an influence on whether the bulk diatom community responded to changing carbonate chemistry. This was done because we expected the likelihood to find an OA response to be higher when the pCO₂ difference between treatments and controls is larger. Thus, we calculated the investigated pCO₂ range (highest pCO₂ – lowest pCO₂) for each experiment and categorized the range into “small” ($\leq 300 \mu\text{atm}$), “medium (300 – 600 μatm), and “large” ($\geq 600 \mu\text{atm}$). Among the 41 experiments that found a CO₂ effect on the bulk diatom community (positive, negative, and unreported direction of change), 4 (10 %, $\sum RDR = 106$) found it within the low range, 12 (32 %, $\sum RDR = 123$) in the medium range, and 25 experiments (68 %, $\sum RDR = 653$) in the high range. Among the 28 experiments that found no CO₂ on the bulk diatom community, 3 (12 %, $\sum RDR = 12$) tested within the low range, 8 (32 %, $\sum RDR = 230$) within the medium range, and 17 experiments (68 %, $\sum RDR = 465$) within the high range. According to this analysis, the likelihood of detecting a CO₂ effect on the

316 bulk diatom community does not depend on the investigated pCO₂ range.

317 CO₂-dependent shifts in diatom species composition were investigated with light
318 microscopy except for Endo et al. (2015) who used molecular tools. Species shifts were
319 investigated in a subset of 40 of the 69 experiments (Fig. 3B). Within this subset of 40
320 studies, 12 (30 %, $\sum\text{RDR} = 265$) found a shift towards larger diatom species under high
321 CO₂, 1 (2.5 %, $\sum\text{RDR} = 10$) found a shift towards smaller diatom species, and 13 (32.5
322 %, $\sum\text{RDR} = 67$) found no CO₂ effect on diatom community composition. 14 studies (35
323 %, $\sum\text{RDR} = 141$) reported a CO₂-dependent shift but did not further specify any changes
324 in the size-class distribution (Fig. 3C).

325 We also tested if the bulk diatom response to OA in coastal, estuarine, and benthic
326 environments was different from the bulk response in oceanic environments. The
327 rationale for this comparison was that carbonate chemistry conditions in oceanic
328 environments may generally be more stable than in the often more productive coastal,
329 estuarine, and benthic environments (Duarte et al., 2013; Hofmann et al., 2011).
330 Therefore, diatoms from oceanic environments may be more sensitive to OA (Duarte et
331 al., 2013). We found 47 experiments with coastal + estuarine + benthic diatom
332 communities. Within this subset, 15 experiments (32 %, $\sum\text{RDR} = 557$) revealed a positive
333 influence of CO₂ on the “bulk diatom community” while 6 (13 %, $\sum\text{RDR} = 244$) revealed
334 a negative one. 4 experiments (9 %, $\sum\text{RDR} = 19$) found a CO₂ effect but did not specify
335 whether it is a positive or negative one. 22 experiments (47 %, $\sum\text{RDR} = 693$) found no
336 effect (Fig. 4A). In contrast, we found 19 experiments with oceanic communities. Within
337 this subset, 5 experiments (26 %, $\sum\text{RDR} = 17$) revealed a positive influence of CO₂ on
338 the “bulk diatom community” while 7 (37 %, $\sum\text{RDR} = 21$) revealed a negative one. 1
339 experiment (5 %, $\sum\text{RDR} = 2$) found a CO₂ effect but did not specify whether it is a

positive or negative one. 6 experiments (32 %, $\sum \text{RDR} = 13$) found no effect (Fig. 4B). Overall, we found a bulk diatom response to OA (positive, negative, and unreported direction of change) in 53 % of the experiments in coastal + estuarine + benthic environments as opposed to 68 % in oceanic environments. Thus, an OA response of the bulk diatom community was more frequently observed in oceanic environments which was mostly due to the higher frequency of negative OA responses (Fig. 4).

4. Discussion

Numerous physiological studies have shown that diatom growth and metabolic rates can be affected by seawater CO₂ concentrations, and that these responses vary widely among different species (Gao and Campbell, 2014). Such inter-specific differences in pCO₂ sensitivity are an important feature as this could alter the composition of diatom assemblages in a changing ocean. In this regard, it is interesting to note that paleolimnologists have long been using diatom species composition as paleo-proxy to reconstruct lake pH (Battarbee et al., 2010). Hence, there is ample evidence that high CO₂ conditions have the potential to change the diatom species composition.

Indeed, our analysis revealed that CO₂-induced changes in diatom community composition occurred in 27 out of 40 (i.e. 68 %) of community-level experiments which investigated species composition (Fig. 3C). This is certainly a conservative outcome because many studies have only looked at dominant species. In fact, one of the few experiments that investigated the diatom assemblage with higher taxonomical resolution found CO₂ effects also on subdominant species (Sommer et al., 2015) which may have been overlooked in many other experiments.

The comparison of OA effects in different environments revealed that bulk diatom communities responded more frequently to OA in oceanic than in coastal + estuarine +

benthic environments. Especially negative effects of OA were more frequent in oceanic environments (Fig. 4). This result is not particularly surprising since communities found near coasts may be adapted to larger carbonate chemistry variability and therefore be better suited to deal with OA (Duarte et al., 2013). It should be kept in mind, however, that this comparison is based on “only” 19 oceanic experiments in contrast to 47 coastal + estuarine + benthic experiments. Furthermore, our habitat characterization depends on certain criteria (mainly water depth and salinity; see section 2.2) and these may be insufficient for our habitat comparison. For example, plankton communities from near oceanic islands such as the Azores were labelled as “coastal” although they may have been moving within oceanic currents and just happened to be close to shore when they were collected. Accordingly, this type of habitat comparison would be more robust if the community had been characterized based on the prevailing carbonate chemistry they are usually exposed to. Unfortunately, information on the background carbonate chemistry is hardly ever provided.

4.1 CO₂ effects on diatom assemblages originating from (direct) physiological responses to high CO₂

Most studies that found effects of pCO₂ on diatom communities related these changes to CO₂ fertilization of photosynthesis. Concentrations of CO₂ in the surface ocean are relatively low compared to other forms of inorganic carbon, especially bicarbonate ion (HCO₃⁻) (Zeebe and Wolf-Gladrow, 2001). However, RubisCO, the primary carboxylating enzyme used in photosynthesis, is restricted to CO₂ for carbon fixation and has a relatively low affinity for CO₂ compared to O₂ (Falkowski and Raven, 2007). Therefore, diatoms (like many other phytoplankton species) operate a carbon concentrating mechanism (CCM) to enhance their CO₂ concentration at the site of fixation relative to external concentrations (e.g. by converting HCO₃⁻ to CO₂) and thereby

establish higher rates of carbon fixation than what would be possible when only depending on diffusive CO₂ uptake (Giordano et al., 2005). It is well known that the proportion of CO₂ uptake vs. HCO₃⁻ uptake for photosynthesis varies largely among diatoms (Burkhardt et al., 2001; Rost et al., 2003; Trimborn et al., 2008) and is theoretically also a function of cell size (Flynn et al., 2012; Wolf-Gladrow and Riebesell, 1997). Accordingly, increasing seawater pCO₂ may increase the proportion of diffusive carbon uptake and/or lower the energy and resource requirements for CCM operation (Raven et al., 2011). From a physiological point of view, these mechanisms could allow for increased rates of photosynthesis and cell division.

So how do these theoretical considerations align with (A) the variable and species-specific physiological responses of diatoms to increasing CO₂ (Dutkiewicz et al., 2015), and (B) the results from community-level experiments compiled in this study? Regarding the variability of physiological responses, progress has recently been made by Wu et al. (2014) who experimentally demonstrated a positive relationship between cell volume and the magnitude of the CO₂ fertilization effect on diatom growth rates. Their findings agree well with theoretical considerations, which predict that high CO₂ is particularly beneficial for carbon acquisition by larger species as they are more restricted by diffusion gradients due to lower surface-to-volume ratios than smaller cells (Flynn et al., 2012; Wolf-Gladrow and Riebesell, 1997). The outcome of our literature analysis supports this allometric concept (Fig. 3, Table 2). Twelve out of 13 experiments in which cell size was taken into account found a shift towards larger species. This is reflected in the Σ RDR score of 265 which is ~25 times higher than the opposite result (i.e. CO₂-induced shifts towards smaller diatoms, Fig. 3C). An allometric scaling of CO₂ sensitivity is particularly useful for modelling since cell size is a universal trait which is relatively easy to measure and therefore frequently available (Ward et al., 2012). Accordingly, it may lead to

significant improvements of ecological and/or biogeochemical model projections under CO₂ forcing when more than one size class for diatoms is considered.

However, although the Wu et al. (2014) allometric approach constitutes a solid starting point to help understanding the variable responses of different diatom species, it probably also still needs some further refinements. For example, central components of CCMs seem to be adapted to diatom cell sizes, thereby potentially alleviating a strict cell size dependency of CO₂ limitation (Shen and Hopkinson, 2015). Furthermore, size dependency alone cannot account for taxon-specific differences in the mode of carbon acquisition (diffusive uptake of CO₂ vs. CCM-supported uptake of HCO₃⁻) and how this will affect the competitive ability of species under increasing CO₂. OA will lead to much larger changes in dissolved CO₂ than in HCO₃⁻. Thus, species that rely to a larger extent on a resource-intensive CCM may benefit more from increasing pCO₂ on a cellular level, as they could increase the proportion of diffusive CO₂ uptake. However, it is also possible that the same species would be disadvantaged on the community-level, because their niche (i.e. being competitive at lower CO₂ due to an efficient CCM) is diminished under high CO₂ conditions. Which of the scenarios occurs in nature would also depend on how flexible species are in terms of switching carbon acquisition modes, as well as resource allocation. In this regard, it is noteworthy that only few physiological studies on OA effects have taken into account the role of changing nutrient concentrations or even a transition to nutrient limitation. The available experimental evidence suggests that increasing pCO₂ may reduce cellular nutrient requirements for CCM operations and therefore free resources for elevated maximum diatom population densities, particularly when running into nutrient limitation (Taucher et al., 2015). Unfortunately, however, the relevance of this mechanism has so far only been investigated in monoclonal laboratory experiments but not on the community-level.

These considerations illustrate that cell size is an important factor, but is not sufficient to predict physiological or even community-level of diatoms to OA. Moreover, the allometric concept as well as the additional mechanisms described above generally presume positive effects of CO₂-fertilization, thus yielding no first order explanations for observed negative responses of diatoms to changing carbonate chemistry. Obviously, increasing CO₂ concentrations are accompanied by increasing proton (H⁺) concentrations under ocean acidification. High H⁺ concentrations may reduce key metabolic rates above certain thresholds and outweigh the positive influence of CO₂ fertilization as has been observed in coccolithophores (Bach et al., 2011, 2015).

Another pathway by which ocean acidification may alter diatom communities is the pH effect on silicification and silica dissolution. Low seawater pH should theoretically facilitate silicification as the precipitation of opal occurs in a cellular compartment with low pH conditions (pH ~5) (Martin-Jézéquel et al., 2000; Vrieling et al., 1999). At the same time, a lower pH should reduce chemical dissolution rates of the SiO₂ frustule (Loucaides et al., 2012). While experimental evidence on this topic is still scarce and partly controversial (Hervé et al., 2012; Mejía et al., 2013; Milligan et al., 2004), it is not unlikely that OA-induced changes in the formation and dissolution of biogenic silica may alter the strength of the frustule and therefore the palatability of diatoms to zooplankton grazers (Friedrichs et al., 2013; Hamm et al., 2003; Liu et al., 2016; Wilken et al., 2011). As for the other physiological effects e.g. on carbon fixation, it is likely that OA impacts on silicification will vary among different diatoms species e.g. according to their species-specific intrinsic buffering capacity, thereby leading to further taxonomic shifts within diatom communities.

The response of diatoms to increasing pCO₂ in natural environments will be further modified by multiple other environmental drivers changing simultaneously. Climate

change is expected to elevate ocean temperature, as well as also irradiance and nutrient availability via changes in stratification. Physiological experiments have shown that elevated pCO₂ may have beneficial effects under low and moderate irradiance, but this effect may reverse under high light conditions due to enhanced photoinhibition (Gao 2012). Analogously, warming may have positive or negative effects on photosynthesis and metabolism in general, depending on the thermal optima of the respective species (Boyd et al., 2018). Altogether, these multiple additional drivers will also affect diatom communities, leading to shifts in their taxonomic composition and size structure, which will interact with the impacts of OA.

4.2 Indirect CO₂ effects on diatom assemblages through food web interactions

Diatom community responses can not only originate from a direct CO₂ effect on their physiology but also be caused indirectly through CO₂ responses on other components of the food web. For example, if a grazer of a diatom species is negatively affected by OA then this may benefit the prey and indirectly promote its abundance. Direct OA impacts on zooplankton communities are usually assumed to play a minor role, although there is some experimental evidence that lower pH may have physiological effects at least on some sensitive species or developmental stages (Cripps et al., 2016; Thor and Dupont, 2015; Thor and Oliva, 2015). Nevertheless, much of the currently available empirical evidence indicates that zooplankton communities are affected by OA rather via bottom-up effects, e.g. via changes in primary production or taxonomic composition of the phytoplankton community (Alvarez-Fernandez et al., 2018; Meunier et al., 2017). However, bottom-up effects on zooplankton biomass, size structure, or species composition may in turn trigger feedbacks on diatom communities, thereby leading to a feedback loop that may reinforce until a new steady state is reached. Such considerations illustrate that also second or third order effects need to be considered when assessing OA

effects on the level of ecological communities. Accounting for such indirect effects requires a holistic approach considering all key players in of the food web (something that is beyond the scope of this study). Therefore, interpretations about what the observed responses could mean for entire plankton food webs or even biogeochemical element cycles (section 4.3) should always be regarded with some healthy skepticism as they often neglect the potential for indirect effects.

4.3 Implications of changes in diatom community structure for pelagic food webs and biogeochemical cycles

The taxonomic composition and size structure of phytoplankton communities influences the transfer of energy from primary production to higher trophic levels. In theory, larger diatoms should support a more direct transfer because less trophic intermediates are needed and therefore less respiration occurs until prey items are in an appropriate size range for top predators (Azam et al., 1983; Pomeroy, 1974; Sommer et al., 2002). Such a size shift at the bottom of a food web might eventually lead to higher production in higher trophic levels such as fish. Indeed, recent experimental evidence indicated that fish (including commercially important species) could under certain constellations benefit from high CO₂ due to higher food availability, although it was not tested if this response is somehow linked to the diatom size structure (Goldenberg et al., 2018; Sswat et al., 2018).

Fluxes of elements through the oceans are (like fluxes of energy through food webs) influenced by the composition of diatom communities (Tréguer et al., 2018). This is particularly well recognized in the context of organic carbon export to the deep ocean, for which diatoms are considered to play a pivotal role (Smetacek, 1985). Given that high CO₂ favours large and perhaps more silicified diatoms over smaller ones (section 4.1),

we might expect accelerated sinking and thus a positive feedback on the vertical carbon flux. This classical hypothesis is supported by observational evidence from two consecutive years of the North Atlantic spring bloom where, despite similar primary production, particulate organic carbon sequestration into the deep ocean was much higher in the year when the larger diatom species dominated (Boyd and Newton, 1995). However, whether the positive relationship between size and carbon export holds under all circumstances is by no means clear (Tréguer et al., 2018) . It is possible that shifts towards larger sized species coincide with shifts in other traits that feed back negatively on carbon export. For example, when the size shift is associated with decreasing C:Si stoichiometry it may ultimately reduce carbon export (Assmy et al., 2013).

The abovementioned examples of trophic transfer and export fluxes illustrate the importance of the factor “diatom community structure” in the context of marine food production and biogeochemical fluxes. They also illustrate that our understanding of the feedbacks induced through changes in diatom communities is highly incomplete. Hence, with our limited understanding we can currently not go further than classifying CO₂-induced changes in diatom communities as “a potential risk” that may cause changes in key ecosystem services.

Acknowledgements

We thank Nauzet Hernández-Hernández, Ulf Riebesell, and Javier Aristegui for their comments on an earlier version of the data compilation. The research was funded by the Federal Ministry of Science and Education (Bundesministerium für Bildung und Forschung; BMBF) in the framework of the “Biological Impacts of Ocean Acidification” project (BIOACID III, FKZ 03F0728).

References

537 Alvarez-Fernandez, S., Bach, L. T., Taucher, J., Riebesell, U., Sommer, U., Aberle, N.,
538 Brussaard, C. P. D. and Boersma, M.: Plankton responses to ocean acidification: The
539 role of nutrient limitation, *Prog. Oceanogr.*, 165, 11–18,
540 doi:10.1016/j.pocean.2018.04.006, 2018.

541 Armbrust, E. V.: The life of diatoms in the world's oceans, *Nature*, 459(7244), 185–
542 192, doi:10.1038/nature08057, 2009.

543 Assmy, P., Smetacek, V., Montresor, M., Klaas, C., Henjes, J., Strass, V. H., Arrieta, J.
544 M., Bathmann, U., Berg, G. M., Breitbarth, E., Cisewski, B., Friedrichs, L., Fuchs, N.,
545 Herndl, G. J., Jansen, S., Kragefsky, S., Latasa, M., Peeken, I., Rottgers, R., Scharek,
546 R., Schuller, S. E., Steigenberger, S., Webb, a. and Wolf-Gladrow, D.: Thick-shelled,
547 grazer-protected diatoms decouple ocean carbon and silicon cycles in the iron-limited
548 Antarctic Circumpolar Current, *Proc. Natl. Acad. Sci.*, 110(51), 20633–20638,
549 doi:10.1073/pnas.1309345110, 2013.

550 Azam, F., Fenchel, T., Field, J. G., Gray, J. S., Meyer-Reil, L. A. and Thingstad, F.: The
551 Ecological Role of Water-Column Microbes in the Sea, *Mar. Ecol. Prog. Ser.*, 10, 257–
552 263, doi:10.3354/meps010257, 1983.

553 Bach, L. T., Riebesell, U. and Schulz, K. G.: Distinguishing between the effects of
554 ocean acidification and ocean carbonation in the coccolithophore *Emiliania huxleyi*,
555 *Limnol. Oceanogr.*, 56(6), 2040–2050, doi:10.4319/lo.2011.56.6.2040, 2011.

556 Bach, L. T., Riebesell, U., Gutowska, M. A., Federwisch, L. and Schulz, K. G.: A
557 unifying concept of coccolithophore sensitivity to changing carbonate chemistry
558 embedded in an ecological framework, *Prog. Oceanogr.*, 135, 125–138,
559 doi:10.1016/j.pocean.2015.04.012, 2015.

560 Bach, L. T., Alvarez-Fernandez, S., Hornick, T., Stühr, A. and Riebesell, U.: Simulated
561 ocean acidification reveals winners and losers in coastal phytoplankton, *PLoS One*,
562 12(11), e0188198, doi:10.1371/journal.pone.0188198, 2017.

563 Bach, L. T., Hernández-hernández, N., Taucher, J., Spisla, C., Sforna, C., Riebesell, U.
564 and Aristegui, J.: Effects of Elevated CO₂ on a Natural Diatom Community in the
565 Subtropical NE Atlantic, *Front. Mar. Sci.*, 6(March), 1–16,
566 doi:10.3389/fmars.2019.00075, 2019.

567 Battarbee, R. W., Charles, D. F., Bigler, C., Cumming, B. F. and Renberg, I.: Diatoms
568 as indicators as surface-water acidity, in *Diatoms: Applications for the Environmental
569 and Earth Sciences*, edited by J. P. Smol and E. F. Stoermer, pp. 98–121, Cambridge
570 University Press, Cambridge., 2010.

571 Biswas, H., Cros, A., Yadav, K., Ramana, V. V., Prasad, V. R., Acharyya, T. and Babu,
572 P. V. R.: The response of a natural phytoplankton community from the Godavari River
573 Estuary to increasing CO₂ concentration during the pre-monsoon period, *J. Exp. Mar.
574 Bio. Ecol.*, 407(2), 284–293, doi:10.1016/j.jembe.2011.06.027, 2011.

575 Biswas, H., Shaik, A. U. R., Bandyopadhyay, D. and Chowdhury, N.: CO₂induced
576 growth response in a diatom dominated phytoplankton community from SW Bay of
577 Bengal coastal water, *Estuar. Coast. Shelf Sci.*, 198, 29–42,
578 doi:10.1016/j.ecss.2017.07.022, 2017.

579 Boyd, P. and Newton, P.: Evidence of the potential influence of planktonic community
580 structure on the interannual variability of particulate organic carbon flux, *Deep Sea Res.*
581 *Part I Oceanogr. Res. Pap.*, 42(5), 619–639, 1995.

582 Boyd, P. W.: Diatom traits regulate Southern Ocean silica leakage, *Proc. Natl. Acad.*
583 *Sci.*, 110(51), 20358–20359, doi:10.1073/pnas.1320327110, 2013.

584 Boyd, P. W., Collins, S., Dupont, S., Fabricius, K., Gattuso, J. P., Havenhand, J.,
585 Hutchins, D. A., Riebesell, U., Rintoul, M. S., Vichi, M., Biswas, H., Ciotti, A., Gao,
586 K., Gehlen, M., Hurd, C. L., Kurihara, H., McGraw, C. M., Navarro, J. M., Nilsson, G.
587 E., Passow, U. and Pörtner, H. O.: Experimental strategies to assess the biological
588 ramifications of multiple drivers of global ocean change-A review, *Glob. Chang. Biol.*,
589 24, 2239–2261, doi:10.1111/gcb.14102, 2018.

590 Brzezinski, M. A. and Nelson, D. M.: Chronic substrate limitation of silicic acid uptake
591 rates in the western Sargasso Sea, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 43(2–3),
592 437–453, doi:10.1016/0967-0645(95)00099-2, 1996.

593 Burkhardt, S., Amoroso, G. and Su, D.: CO₂ and HCO₃⁻ uptake in marine diatoms
594 acclimated to different CO₂ concentrations, , 46(6), 1378–1391, 2001.

595 Calvo-Díaz, A., Díaz-Pérez, L., Suárez, L. Á., Morán, X. A. G., Teira, E. and Marañón,
596 E.: Decrease in the autotrophic-to-heterotrophic biomass ratio of picoplankton in
597 oligotrophic marine waters due to bottle enclosure, *Appl. Environ. Microbiol.*, 77(16),
598 5739–5746, doi:10.1128/AEM.00066-11, 2011.

599 Carpenter, S. R.: Microcosm Experiments Have Limited Relevance for Community and
600 Ecosystem Ecology, *Ecology*, 77(3), 667–680, 1996.

601 Cripps, G., Flynn, K. J. and Lindeque, P. K.: Ocean acidification affects the phyto-zoo
602 plankton trophic transfer efficiency, *PLoS One*, 11(4), 1–15,
603 doi:10.1371/journal.pone.0151739, 2016.

604 Davidson, A., McKinlay, J., Westwood, K., Thomson, P., van den Enden, R., de Salas,
605 M., Wright, S., Johnson, R. and Berry, K.: Enhanced CO₂ concentrations change the
606 structure of Antarctic marine microbial communities, *Mar. Ecol. Prog. Ser.*, 552, 93–
607 113, doi:10.3354/meps11742, 2016.

608 Domingues, R. B., Guerra, C. C., Galvao, H. M., Brotas, V. and Barbosa, A. B.: Short-
609 term interactive effects of ultraviolet radiation, carbon dioxide and nutrient enrichment
610 on phytoplankton in a shallow coastal lagoon, *Aquat. Ecol.*, 51(1), 91–105,
611 doi:10.1007/s10452-016-9601-4, 2017.

612 Donahue, K., Klaas, C., Dillingham, P. W. and Hoffmann, L. J.: Combined effects of
613 ocean acidification and increased light intensity on natural phytoplankton communities
614 from two Southern Ocean water masses, *J. Plankton Res.*, 41(1), 30–45,
615 doi:10.1093/plankt/fby048, 2019.

616 Duarte, C. M., Gasol, J. M. and Vaqué, D.: Role of experimental approaches in marine
617 microbial ecology, *Aquat. Microb. Ecol.*, 13(1), 101–111, doi:10.3354/ame013101,
618 1997.

619 Duarte, C. M., Hendriks, I. E., Moore, T. S., Olsen, Y. S., Steckbauer, A., Ramajo, L.,
620 Carstensen, J., Trotter, J. A. and McCulloch, M.: Is Ocean Acidification an Open-Ocean
621 Syndrome? Understanding Anthropogenic Impacts on Seawater pH, *Estuaries and*
622 *Coasts*, 36(2), 221–236, doi:10.1007/s12237-013-9594-3, 2013.

623 Dutkiewicz, S., Morris, J. J., Follows, M. J., Scott, J., Levitan, O., Dyhrman, S. T. and
624 Berman-Frank, I.: Impact of ocean acidification on the structure of future phytoplankton
625 communities, *Nat. Clim. Chang.*, 5(11), 1002–1006, doi:10.1038/nclimate2722, 2015.

626 Eggers, S. L., Lewandowska, A. M., Barcelos e Ramos, J., Blanco-Ameijeiras, S.,
627 Gallo, F. and Matthiessen, B.: Community composition has greater impact on the
628 functioning of marine phytoplankton communities than ocean acidification, *Glob.*
629 *Chang. Biol.*, 20(3), 713–723, doi:10.1111/gcb.12421, 2014.

630 Endo, H., Yoshimura, T., Kataoka, T. and Suzuki, K.: Effects of CO₂ and iron
631 availability on phytoplankton and eubacterial community compositions in the northwest
632 subarctic Pacific, *J. Exp. Mar. Bio. Ecol.*, 439, 160–175,
633 doi:10.1016/j.jembe.2012.11.003, 2013.

634 Endo, H., Sugie, K., Yoshimura, T. and Suzuki, K.: Effects of CO₂ and iron availability
635 on *rbcL* gene expression in Bering Sea diatoms, *Biogeosciences*, 12(7), 2247–2259,
636 doi:10.5194/bg-12-2247-2015, 2015.

637 Endo, H., Sugie, K., Yoshimura, T. and Suzuki, K.: Response of Spring Diatoms to
638 CO₂ Availability in the Western North Pacific as Determined by Next-Generation
639 Sequencing, *PLoS One*, 11(4), e0154291, doi:10.1371/journal.pone.0154291, 2016.

640 Fabricius, K. E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G.,
641 Okazaki, R., Muehllehner, N., Glas, M. S. and Lough, J. M.: Losers and winners in
642 coral reefs acclimatized to elevated carbon dioxide concentrations, *Nat. Clim. Chang.*,
643 1(6), 165–169, doi:10.1038/nclimate1122, 2011.

644 Falkowski, P. G. and Raven, J. A.: *Aquatic Photosynthesis*, Princeton University Press,
645 Princeton., 2007.

646 Feng, Y., Hare, C. E., Leblanc, K., Rose, J. M., Zhang, Y., DiTullio, G. R., Lee, P. A.,
647 Wilhelm, S. W., Rowe, J. M., Sun, J., Nemcek, N., Gueguen, C., Passow, U., Benner, I.,
648 Brown, C. and Hutchins, D. A.: Effects of increased pCO₂ and temperature on the north
649 atlantic spring bloom. I. The phytoplankton community and biogeochemical response,
650 *Mar. Ecol. Prog. Ser.*, 388, 13–25, doi:10.3354/meps08133, 2009.

651 Feng, Y., Hare, C. E., Rose, J. M., Handy, S. M., DiTullio, G. R., Lee, P. A., Smith, W.
652 O., Peloquin, J., Tozzi, S., Sun, J., Zhang, Y., Dunbar, R. B., Long, M. C., Sohst, B.,
653 Lohan, M. and Hutchins, D. A.: Interactive effects of iron, irradiance and CO₂ on Ross
654 Sea phytoplankton, *Deep. Res. Part I Oceanogr. Res. Pap.*, 57(3), 368–383,
655 doi:10.1016/j.dsr.2009.10.013, 2010.

656 Ferguson, R. L., Buckley, E. N. and Palumbo, A. V.: Response of marine
657 bacterioplankton to differential filtration and confinement., *Appl. Environ. Microbiol.*,
658 47(1), 49–55, doi:0099-2240/84/010049-07\$02., 1984.

659 Field, C. B., Behrenfeld, M. J., Randerson, J. T. and Falkowski, P. G.: Primary

660 Production of the Biosphere: Integrating Terrestrial and Oceanic Components, *Science*
661 (80-.), 281(5374), 237–240, doi:10.1126/science.281.5374.237, 1998.

662 Flynn, K. J., Blackford, J. C., Baird, M. E., Raven, J. A., Clark, D. R., Beardall, J.,
663 Brownlee, C., Fabian, H. and Wheeler, G. L.: Changes in pH at the exterior surface of
664 plankton with ocean acidification, *Nat. Clim. Chang.*, 2(7), 510–513,
665 doi:10.1038/nclimate1489, 2012.

666 Fogg, G. E. and Calvario-Martinez, O.: Effects of bottle size in determinations of
667 primary productivity by phytoplankton, *Hydrobiologia*, 173(2), 89–94,
668 doi:10.1007/BF00015518, 1989.

669 Friedrichs, L., Hörnig, M., Schulze, L., Bertram, A., Jansen, S. and Hamm, C.: Size and
670 biomechanic properties of diatom frustules influence food uptake by copepods, *Mar.*
671 *Ecol. Prog. Ser.*, 481(2000), 41–51, doi:10.3354/meps10227, 2013.

672 Gao, K. and Campbell, D. A.: Photophysiological responses of marine diatoms to
673 elevated CO₂ and decreased pH: A review, *Funct. Plant Biol.*, 41(5), 449–459,
674 doi:10.1071/FP13247, 2014.

675 Gao, K., Xu, J., Gao, G., Li, Y., Hutchins, D. A., Huang, B., Wang, L., Zheng, Y., Jin,
676 P., Cai, X., Häder, D. P., Li, W., Xu, K., Liu, N. and Riebesell, U.: Rising CO₂ and
677 increased light exposure synergistically reduce marine primary productivity, *Nat. Clim.*
678 *Chang.*, 2(7), 519–523, doi:10.1038/nclimate1507, 2012.

679 Gazeau, F., Sallon, A., Pitta, P., Tsiola, A., Maugendre, L., Giani, M., Celussi, M.,
680 Pedrotti, M. L., Marro, S. and Guieu, C.: Limited impact of ocean acidification on
681 phytoplankton community structure and carbon export in an oligotrophic environment:
682 Results from two short-term mesocosm studies in the Mediterranean Sea, *Estuar. Coast.*
683 *Shelf Sci.*, 186, 72–88, doi:10.1016/j.ecss.2016.11.016, 2017.

684 Giordano, M., Beardall, J. and Raven, J. A.: CO₂ concentrating mechanisms in algae:
685 mechanisms, environmental modulation, and evolution., *Annu. Rev. Plant Biol.*, 56, 99–
686 131, doi:10.1146/annurev.arplant.56.032604.144052, 2005.

687 Goldenberg, S. U., Nagelkerken, I., Marangon, E., Bonnet, A. and Camilo, M.:
688 Ecological complexity buffers the impacts of future climate on marine animals
689 Corresponding author Keywords, *Nat. Clim. Chang.*, 1–19, doi:10.1038/s41558-018-
690 0086-0, 2018.

691 Grear, J. S., Ryneerson, T. A., Montalbano, A. L., Govenar, B. and Menden-Deuer, S.:
692 pCO₂ effects on species composition and growth of an estuarine phytoplankton
693 community, *Estuar. Coast. Shelf Sci.*, 190, 40–49, doi:10.1016/j.ecss.2017.03.016,
694 2017.

695 Guangao, L.: Different types of ecosystem experiments, in *Enclosed Experimental*
696 *Marine Ecosystems: A Review and Recommendations*, edited by C. M. Lalli, pp. 7–20,
697 Springer-Verlag, New York., 1990.

698 Guiry, M. D.: How many species of algae are there?, *J. Phycol.*, 48(5), 1057–1063,
699 doi:10.1111/j.1529-8817.2012.01222.x, 2012.

700 Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S.
701 M., Rowley, S. J., Tedesco, D. and Buia, M.-C.: Volcanic carbon dioxide vents show
702 ecosystem effects of ocean acidification, *Nature*, 454, 96–99, doi:10.1038/nature07051,
703 2008.

704 Hama, T., Inoue, T., Suzuki, R., Kashiwazaki, H., Wada, S., Sasano, D., Kosugi, N. and
705 Ishii, M.: Response of a phytoplankton community to nutrient addition under different
706 CO₂ and pH conditions, *J. Oceanogr.*, 72(2), 207–223, doi:10.1007/s10872-015-0322-4,
707 2016.

708 Hamm, C. and Smetacek, V.: Armor: Why, when, and how, in *Evolution of*
709 *Phytoplankton*, edited by P. G. Falkowski and A. H. Knoll, pp. 311–332, Elsevier,
710 Boston., 2007.

711 Hamm, C. E., Merkel, R., Springer, O., Jurkojc, P., Maiert, C., Prechtelt, K. and
712 Smetacek, V.: Architecture and material properties of diatom shells provide effective
713 mechanical protection, *Nature*, 421(6925), 841–843, doi:10.1038/nature01416, 2003.

714 Hammes, F., Vital, M. and Egli, T.: Critical evaluation of the volumetric “bottle effect”
715 on microbial batch growth, *Appl. Environ. Microbiol.*, 76(4), 1278–1281,
716 doi:10.1128/AEM.01914-09, 2010.

717 Hare, C. E., Leblanc, K., DiTullio, G. R., Kudela, R. M., Zhang, Y., Lee, P. A.,
718 Riseman, S. and Hutchins, D. A.: Consequences of increased temperature and CO₂ for
719 phytoplankton community structure in the Bering Sea, *Mar. Ecol. Prog. Ser.*, 352, 9–16,
720 doi:10.3354/meps07182, 2007.

721 Hervé, V., Derr, J., Douady, S., Quinet, M., Moisan, L. and Lopez, P. J.:
722 Multiparametric Analyses Reveal the pH-Dependence of Silicon Biomineralization in
723 Diatoms, *PLoS One*, 7(10), e46722, doi:10.1371/journal.pone.0046722, 2012.

724 Hofmann, G. E., Smith, J. E., Johnson, K. S., Send, U., Levin, L. A., Micheli, F.,
725 Paytan, A., Price, N. N., Peterson, B., Takeshita, Y., Matson, P. G., de Crook, E.,
726 Kroeker, K. J., Gambi, M. C., Rivest, E. B., Frieder, C. A., Yu, P. C. and Martz, T. R.:
727 High-frequency dynamics of ocean pH: A multi-ecosystem comparison, *PLoS One*,
728 6(12), e28983, doi:10.1371/journal.pone.0028983, 2011.

729 Hopkins, F. E., Turner, S. M., Nightingale, P. D., Steinke, M., Bakker, D. and Liss, P.
730 S.: Ocean acidification and marine trace gas emissions, *Proc. Natl. Acad. Sci. U. S. A.*,
731 107, 760–765, doi:10.1073/pnas.0907163107, 2010.

732 Hoppe, C. J. M., Hassler, C. S., Payne, C. D., Tortell, P. D., Rost, B. R. and Trimbom,
733 S.: Iron limitation modulates ocean acidification effects on Southern Ocean
734 phytoplankton communities, *PLoS One*, 8(11), doi:10.1371/journal.pone.0079890,
735 2013.

736 Hoppe, C. J. M., Schuback, N., Semeniuk, D. M., Maldonado, M. T. and Rost, B.:
737 Functional Redundancy Facilitates Resilience of Subarctic Phytoplankton Assemblages
738 toward Ocean Acidification and High Irradiance, *Front. Mar. Sci.*, 4,
739 doi:10.3389/fmars.2017.00229, 2017a.

740 Hoppe, C. J. M., Schuback, N., Semeniuk, D., Giesbrecht, K., Mol, J., Thomas, H.,

741 Maldonado, M. T., Rost, B., Varela, D. E. and Tortell, P. D.: Resistance of Arctic
 742 phytoplankton to ocean acidification and enhanced irradiance, *Polar Biol.*, 41(3), 399–
 743 413, doi:10.1007/s00300-017-2186-0, 2017b.

744 Husherr, R., Levasseur, M., Lizotte, M., Tremblay, J. É., Mol, J., Thomas, H.,
 745 Gosselin, M., Starr, M., Miller, L. A., Jarníková, T., Schuback, N. and Mucci, A.:
 746 Impact of ocean acidification on Arctic phytoplankton blooms and dimethyl sulfide
 747 concentration under simulated ice-free and under-ice conditions, *Biogeosciences*, 14(9),
 748 2407–2427, doi:10.5194/bg-14-2407-2017, 2017.

749 James, R. K., Hepburn, C. D., Cornwall, C. E., McGraw, C. M. and Hurd, C. L.:
 750 Growth response of an early successional assemblage of coralline algae and benthic
 751 diatoms to ocean acidification, *Mar. Biol.*, 161(7), 1687–1696, doi:10.1007/s00227-
 752 014-2453-3, 2014.

753 Johnson, V. R., Brownlee, C., Rickaby, R. E. M., Graziano, M., Milazzo, M. and Hall-
 754 Spencer, J. M.: Responses of marine benthic microalgae to elevated CO₂, *Mar. Biol.*,
 755 160(8), 1813–1824, doi:10.1007/s00227-011-1840-2, 2011.

756 Kim, J.-M., Lee, K., Shin, K., Kang, J.-H., Lee, H.-W., Kim, M., Jang, P.-G. and Jang,
 757 M.-C.: The effect of seawater CO₂ concentration on growth of a natural phytoplankton
 758 assemblage in a controlled mesocosm experiment, *Limnol. Oceanogr.*, 51(4), 1629–
 759 1636, doi:10.4319/lo.2006.51.4.1629, 2006.

760 Kim, J. M., Lee, K., Yang, E. J., Shin, K., Noh, J. H., Park, K. T., Hyun, B., Jeong, H.
 761 J., Kim, J. H., Kim, K. Y., Kim, M., Kim, H. C., Jang, P. G. and Jang, M. C.: Enhanced
 762 production of oceanic dimethylsulfide resulting from CO₂-induced grazing activity in a
 763 high CO₂ world, *Environ. Sci. Technol.*, 44(21), 8140–8143, doi:10.1021/es102028k,
 764 2010.

765 Liu, H., Chen, M., Zhu, F. and Harrison, P. J.: Effect of Diatom Silica Content on
 766 Copepod Grazing, Growth and Reproduction, *Front. Mar. Sci.*, 3(June), 1–7,
 767 doi:10.3389/fmars.2016.00089, 2016.

768 Loucaides, S., van Cappellen, P., Roubéix, V., Moriceau, B. and Ragueneau, O.:
 769 Controls on the Recycling and Preservation of Biogenic Silica from Biomineralization
 770 to Burial, *Silicon*, 4(1), 7–22, doi:10.1007/s12633-011-9092-9, 2012.

771 Mallozzi, A. J., Errera, R. M., Bargu, S. and Herrmann, A. D.: Impacts of elevated
 772 pCO₂ on estuarine phytoplankton biomass and community structure in two
 773 biogeochemically distinct systems in Louisiana, USA, *J. Exp. Mar. Bio. Ecol.*,
 774 511(September 2018), 28–39, doi:10.1016/j.jembe.2018.09.008, 2019.

775 Malviya, S., Scalco, E., Audic, S., Vincent, F., Veluchamy, A., Poulain, J., Wincker, P.,
 776 Iudicone, D., de Vargas, C., Bittner, L., Zingone, A. and Bowler, C.: Insights into
 777 global diatom distribution and diversity in the world's ocean, *Proc. Natl. Acad. Sci.*,
 778 113(11), E1516–E1525, doi:10.1073/pnas.1509523113, 2016.

779 Mann, D. G. and Vanormelingen, P.: An Inordinate Fondness ? The Number ,
 780 Distributions , and Origins of Diatom Species, *J. Eukaryot. Microbiol.*, 60, 414–420,
 781 doi:10.1111/jeu.12047, 2013.

782 Martin-Jézéquel, V., Hildebrand, M. and Brzezinski, M. A.: Review Silicon Metabolism
783 in Diatoms: Implications for Growth, *J. Phycol.*, 36, 821–840, 2000.

784 Maugendre, L., Gattuso, J.-P., Louis, J., de Kluijver, A., Marro, S., Soetaert, K. and
785 Gazeau, F.: Effect of ocean warming and acidification on a plankton community in the
786 NW Mediterranean Sea, *ICES J. Mar. Sci.*, 72(6), 1744–1755,
787 doi:10.1093/icesjms/fsu161, 2015.

788 Mejía, L. M., Isensee, K., Méndez-Vicente, A., Pisonero, J., Shimizu, N., González, C.,
789 Monteleone, B. and Stoll, H.: B content and Si/C ratios from cultured diatoms
790 (*Thalassiosira pseudonana* and *Thalassiosira weissflogii*): Relationship to seawater pH
791 and diatom carbon acquisition, *Geochim. Cosmochim. Acta*, 123, 322–337,
792 doi:10.1016/j.gca.2013.06.011, 2013.

793 Menzel, D. W. and Case, J.: Concept and Design: Controlled Ecosystem Pollution
794 Experiment, *Bull. Mar. Sci.*, 27(1), 1–7, 1977.

795 Meunier, C. L., Algueró-Muñiz, M., Horn, H. G., Lange, J. A. F. and Boersma, M.:
796 Direct and indirect effects of near-future pCO₂ levels on zooplankton dynamics, *Mar.*
797 *Freshw. Res.*, 68(2), 373–380, doi:10.1071/MF15296, 2017.

798 Milligan, A. J., Varela, D. E., Brzezinski, M. A. and Morel, F. M. M.: Dynamics of
799 Silicon Metabolism and Silicon Isotopic Discrimination in a Marine Diatom as a
800 Function of pCO₂, *Limnol. Oceanogr.*, 49(2), 322–329, 2004.

801 Nelson, D. M., Tréguer, P., Brzezinski, M. A., Leynaert, A. and Quéguiner, B.:
802 Production and dissolution of biogenic silica in the ocean: Revised global estimates,
803 comparison with regional data and relationship to biogenic sedimentation, *Global*
804 *Biogeochem. Cycles*, 9(3), 359–372, doi:10.1029/95GB01070, 1995.

805 Nielsen, L. T., Jakobsen, H. H. and Hansen, P. J.: High resilience of two coastal
806 plankton communities to twenty-first century seawater acidification: Evidence from
807 microcosm studies, *Mar. Biol. Res.*, 6(6), 542–555, doi:10.1080/17451000903476941,
808 2010.

809 Nielsen, L. T., Hallegraeff, G. M., Wright, S. W. and Hansen, P. J.: Effects of
810 experimental seawater acidification on an estuarine plankton community, *Aquat.*
811 *Microb. Ecol.*, 65(3), 271–285, doi:10.3354/ame01554, 2012.

812 Nogueira, P., Domingues, R. B. and Barbosa, A. B.: Are microcosm volume and sample
813 pre-filtration relevant to evaluate phytoplankton growth?, *J. Exp. Mar. Bio. Ecol.*, 461,
814 323–330, doi:10.1016/j.jembe.2014.09.006, 2014.

815 Pan, Y., Zhang, Y., Peng, Y., Zhao, Q. and Sun, S.: Increases of chamber height and
816 base diameter have contrasting effects on grazing rate of two cladoceran species:
817 Implications for microcosm studies, *PLoS One*, 10(8), 1–14,
818 doi:10.1371/journal.pone.0135786, 2015.

819 Pančić, M. and Kiørboe, T.: Phytoplankton defence mechanisms: traits and trade-offs,
820 *Biol. Rev.*, 92(2), 1269–1303, doi:10.1111/brv.12395, 2018.

821 Park, K. T., Lee, K., Shin, K., Yang, E. J., Hyun, B., Kim, J. M., Noh, J. H., Kim, M.,

- 822 Kong, B., Choi, D. H., Choi, S. J., Jang, P. G. and Jeong, H. J.: Direct linkage between
823 dimethyl sulfide production and microzooplankton grazing, resulting from prey
824 composition change under high partial pressure of carbon dioxide conditions, *Environ.*
825 *Sci. Technol.*, 48(9), 4750–4756, doi:10.1021/es403351h, 2014.
- 826 Paul, A. J., Bach, L. T., Schulz, K.-G., Boxhammer, T., Czerny, J., Achterberg, E. P.,
827 Hellemann, D., Trense, Y., Nausch, M., Sswat, M. and Riebesell, U.: Effect of elevated
828 CO₂ on organic matter pools and fluxes in a summer Baltic Sea plankton community,
829 *Biogeosciences*, 12, 6181–6203, doi:10.5194/bg-12-6181/2015/, 2015.
- 830 Pomeroy, L. R.: The ocean food web - A changing paradigm, *Bioscience*, 24, 499–504,
831 1974.
- 832 Primeau, F. W., Holzer, M. and DeVries, T.: Southern Ocean nutrient trapping and the
833 efficiency of the biological pump, *J. Geophys. Res. Ocean.*, 118(5), 2547–2564,
834 doi:10.1002/jgrc.20181, 2013.
- 835 Raven, J. a, Giordano, M., Beardall, J. and Maberly, S. C.: Algal and aquatic plant
836 carbon concentrating mechanisms in relation to environmental change., *Photosynth.*
837 *Res.*, 109(1–3), 281–96, doi:10.1007/s11120-011-9632-6, 2011.
- 838 Reul, A., Muñoz, M., Bautista, B., Neale, P. J., Sobrino, C., Mercado, J. M., Segovia,
839 M., Salles, S., Kulk, G., León, P., van de Poll, W. H. D., Pérez, E., Buma, A. and
840 Blanco, J. M.: Effect of CO₂, nutrients and light on coastal plankton. III. Trophic
841 cascade, size structure and composition, *Aquat. Biol.*, 22, 59–76, doi:10.3354/ab00585,
842 2014.
- 843 Robinson, C. and Williams, P. J. le B.: Respiration and its measurement in surface
844 marine waters, in *Respiration in Aquatic Environments*, edited by P. A. Del Giorgio and
845 P. J. le B. Williams, pp. 147–180, Oxford University Press, Oxford., 2005.
- 846 Roleda, M. Y., Cornwall, C. E., Feng, Y., McGraw, C. M., Smith, A. M. and Hurd, C.
847 L.: Effect of ocean acidification and pH fluctuations on the growth and development of
848 coralline algal recruits, and an associated benthic algal assemblage, *PLoS One*, 10(10),
849 1–19, doi:10.1371/journal.pone.0140394, 2015.
- 850 Rossoll, D., Sommer, U. and Winder, M.: Community interactions dampen acidification
851 effects in a coastal plankton system, *Mar. Ecol. Prog. Ser.*, 486, 37–46,
852 doi:10.3354/meps10352, 2013.
- 853 Rost, B., Riebesell, U., Burkhardt, S. and Sültemeyer, D.: Carbon acquisition of bloom-
854 forming marine phytoplankton, *Limnol. Oceanogr.*, 48(1), 55–67, 2003.
- 855 Sala, M. M., Aparicio, F. L., Balagué, V., Boras, J. A., Borrull, E., Cardelús, C., Cros,
856 L., Gomes, A., López-Sanz, A., Malits, A., Martínez, R. A., Mestre, M., Movilla, J.,
857 Sarmiento, H., Vázquez-Domínguez, E., Vaqué, D., Pinhassi, J., Calbet, A., Calvo, E.,
858 Gasol, J. M., Pelejero, C. and Marrasé, C.: Contrasting effects of ocean acidification on
859 the microbial food web under different trophic conditions, *Ices J. Mar. Sci.*, 73(3), 670–
860 679, doi:10.1093/icesjms/fsv130, 2015.
- 861 Sarmiento, J. L., Gruber, N., Brzezinski, M. A. and Dunne, J. P.: High-latitude controls
862 of thermocline nutrients and low latitude biological productivity, *Nature*, 427(6969),

863 56–60, doi:10.1038/nature10605, 2004.

864 Sarnelle, O.: Daphnia effects on microzooplankton: comparisons of enclosure and
865 whole-lake responses, *Ecology*, 78(3), 913–928, doi:10.1016/S0010-4655(02)00300-4,
866 1997.

867 Sarthou, G., Timmermans, K. R., Blain, S. and Tréguer, P.: Growth physiology and fate
868 of diatoms in the ocean: A review, *J. Sea Res.*, 53, 25–42,
869 doi:10.1016/j.seares.2004.01.007, 2005.

870 Schulz, K. G., Riebesell, U., Bellerby, R. G. J., Biswas, H., Meyerhöfer, M., Müller, M.
871 N., Egge, J. K., Nejstgaard, J. C., Neill, C., Wohlers, J. and Zöllner, E.: Build-up and
872 decline of organic matter during PeECE III, *Biogeosciences*, 5, 707–718,
873 doi:10.5194/bg-5-707-2008, 2008.

874 Schulz, K. G., Bellerby, R. G. J., Brussaard, C. P. D., Büdenbender, J., Czerny, J.,
875 Engel, A., Fischer, M., Koch-Klavsen, S., Krug, S. A., Lischka, S., Ludwig, A.,
876 Meyerhöfer, M., Nondal, G., Silyakova, A., Stuhr, A. and Riebesell, U.: Temporal
877 biomass dynamics of an Arctic plankton bloom in response to increasing levels of
878 atmospheric carbon dioxide, *Biogeosciences*, 10, 161–180, doi:10.5194/bg-10-161-
879 2013, 2013.

880 Schulz, K. G., Bach, L. T., Bellerby, R., Bermudez, R., Boxhammer, T., Czerny, J.,
881 Engel, A., Ludwig, A., Larsen, A., Paul, A., Sswat, M. and Riebesell, U.: Phytoplankton
882 blooms at increasing levels of atmospheric carbon dioxide: experimental evidence for
883 negative effects on prymnesiophytes and positive on small picoeukaryotes, *Front. Mar.*
884 *Sci.*, 4(64), 1–18, doi:10.3389/fmars.2017.00064, 2017.

885 Segovia, M., Lorenzo, M., Maldonado, M., Larsen, A., Berger, S., Tsagaraki, T.,
886 Lázaro, F., Iñiguez, C., García-Gómez, C., Palma, A., Mausz, M., Gordillo, F.,
887 Fernández, J., Ray, J. and Egge, J.: Iron availability modulates the effects of future CO₂
888 levels within the marine planktonic food web, *Mar. Ecol. Prog. Ser.*, 565, 17–33,
889 doi:10.3354/meps12025, 2017.

890 Sett, S., Schulz, K. G., Bach, L. T. and Riebesell, U.: Shift towards larger diatoms in a
891 natural phytoplankton assemblage under combined high-CO₂ and warming conditions,
892 *J. Plankton Res.*, 40(4), 391–406, doi:10.1093/plankt/fby018, 2018.

893 Shaik, A. U. R., Biswas, H. and Pal, S.: Increased CO₂ availability promotes growth of
894 a tropical coastal diatom assemblage (Goa coast, Arabian Sea, India), *Diatom Res.*,
895 32(3), 325–339, doi:10.1080/0269249X.2017.1379443, 2017.

896 Shen, C. and Hopkinson, B. M.: Size scaling of extracellular carbonic anhydrase
897 activity in centric marine diatoms, *J. Phycol.*, 51(2), 255–263, doi:10.1111/jpy.12269,
898 2015.

899 Smetacek, V., Klaas, C., Strass, V. H., Assmy, P., Montresor, M., Cisewski, B., Savoye,
900 N., Webb, A., d’Ovidio, F., Arrieta, J. M., Bathmann, U., Bellerby, R., Berg, G. M.,
901 Croot, P., Gonzalez, S., Henjes, J., Herndl, G. J., Hoffmann, L. J., Leach, H., Losch, M.,
902 Mills, M. M., Neill, C., Peeken, I., Röttgers, R., Sachs, O., Sauter, E., Schmidt, M. M.,
903 Schwarz, J., Terbrüggen, A. and Wolf-Gladrow, D.: Deep carbon export from a
904 Southern Ocean iron-fertilized diatom bloom, *Nature*, 487(7407), 313–319,

905 doi:10.1038/nature11229, 2012.

906 Smetacek, V. S.: Role of sinking in diatom life-history: ecological, evolutionary and
 907 geological significance, *Mar. Biol.*, 84, 239–251, doi:10.1007/BF00392493, 1985.

908 Sommer, U., Stibor, H., Katchikis, A., Sommer, F. and Hansen, T.: Pelagic food web
 909 configurations at different levels of nutrient richness and their implications for the ratio
 910 fish production:primary production, *Hydrobiologia*, 484, 11–20,
 911 doi:10.1023/A:1021340601986, 2002.

912 Sommer, U., Paul, C. and Moustaka-Gouni, M.: Warming and ocean acidification
 913 effects on phytoplankton - From species shifts to size shifts within species in a
 914 mesocosm experiment, *PLoS One*, 10(5), 1–17, doi:10.1371/journal.pone.0125239,
 915 2015.

916 Sournia, A., Chrétiennot-Dinet, M. J. and Ricard, M.: Marine phytoplankton: How
 917 many species in the world ocean?, *J. Plankton Res.*, 13(5), 1093–1099,
 918 doi:10.1093/plankt/13.5.1093, 1991.

919 Spencer, M. and Warren, P. H.: The effects of habitat size and productivity on food web
 920 structure in small aquatic microcosms, *Oikos*, 75(3), 419–430 [online] Available from:
 921 <http://www.jstor.org/stable/3545882?origin=crossref>, 1996.

922 Sswat, M., Stiasny, M. H., Taucher, J., Algueró-Muñiz, M., Bach, L. T., Jutfelt, F.,
 923 Riebesell, U. and Clemmesen, C.: Food web changes under ocean acidification promote
 924 herring larvae survival, *Nat. Ecol. Evol.*, 2(May), 1–5, doi:10.1038/s41559-018-0514-6,
 925 2018.

926 Tatters, A. O., Roleda, M. Y., Schnetzer, A., Fu, F., Hurd, C. L., Boyd, P. W., Caron, D.
 927 A., Lie, A. A. Y., Hoffmann, L. J. and Hutchins, D. A.: Short- and long-term
 928 conditioning of a temperate marine diatom community to acidification and warming,
 929 *Philos. Trans. R. Soc. B Biol. Sci.*, 368, 20120437, doi:10.1098/rstb.2012.0437, 2013.

930 Tatters, A. O., Schnetzer, A., Xu, K., Walworth, N. G., Fu, F., Spackeen, J. L., Sipler,
 931 R. E., Bertrand, E. M., McQuaid, J. B., Allen, A. E., Bronk, D. A., Gao, K., Sun, J.,
 932 Caron, D. A. and Hutchins, D. A.: Interactive effects of temperature, CO₂ and nitrogen
 933 source on a coastal California diatom assemblage, *J. Plankton Res.*, 40(2), 151–164,
 934 doi:10.1093/plankt/fbx074, 2018.

935 Taucher, J., Jones, J., James, A., Brzezinski, M. A., Carlson, C. A., Riebesell, U. and
 936 Passow, U.: Combined effects of CO₂ and temperature on carbon uptake and
 937 partitioning by the marine diatoms *Thalassiosira weissflogii* and *Dactyliosolen*
 938 *fragilissimus*, *Limnol. Oceanogr.*, 60(3), 901–919, doi:10.1002/lno.10063, 2015.

939 Taucher, J., Arístegui, J., Bach, L. T., Guan, W., Montero, M. F., Nauendorf, A.,
 940 Achterberg, E. P. and Riebesell, U.: Response of Subtropical Phytoplankton
 941 Communities to Ocean Acidification Under Oligotrophic Conditions and During
 942 Nutrient Fertilization, *Front. Mar. Sci.*, 5(September), 1–14,
 943 doi:10.3389/fmars.2018.00330, 2018.

944 Thøisen, C., Riisgaard, K., Lundholm, N., Nielsen, T. G. and Hansen, P. J.: Effect of
 945 acidification on an Arctic phytoplankton community from Disko Bay, West Greenland,

- 946 Mar. Ecol. Prog. Ser., 520, 21–34, doi:10.3354/meps11123, 2015.
- 947 Thor, P. and Dupont, S.: Transgenerational effects alleviate severe fecundity loss during
948 ocean acidification in a ubiquitous planktonic copepod, *Glob. Chang. Biol.*, 21(6),
949 2261–2271, doi:10.1111/gcb.12815, 2015.
- 950 Thor, P. and Oliva, E. O.: Ocean acidification elicits different energetic responses in an
951 Arctic and a boreal population of the copepod *Pseudocalanus acuspes*, *Mar. Biol.*,
952 162(4), 799–807, doi:10.1007/s00227-015-2625-9, 2015.
- 953 Tortell, P. D., DiTullio, G. R., Sigman, D. M. and Morel, F. M. M.: CO₂ effects on
954 taxonomic composition and nutrient utilization in an Equatorial Pacific phytoplankton
955 assemblage, *Mar. Ecol. Prog. Ser.*, 236, 37–43, doi:10.3354/meps236037, 2002.
- 956 Tortell, P. D., Payne, C. D., Li, Y., Trimborn, S., Rost, B., Smith, W. O., Riesselman,
957 C., Dunbar, R. B., Sedwick, P. and DiTullio, G. R.: CO₂ sensitivity of Southern Ocean
958 phytoplankton, *Geophys. Res. Lett.*, 35, L04605, doi:10.1029/2007GL032583, 2008.
- 959 Tréguer, P., Bowler, C., Moriceau, B., Dutkiewicz, S., Gehlen, M., Aumont, O., Bittner,
960 L., Dugdale, R., Finkel, Z., Iudicone, D., Jahn, O., Guidi, L., Lasbleiz, M., Leblanc, K.,
961 Levy, M. and Pondaven, P.: Influence of diatom diversity on the ocean biological
962 carbon pump, *Nat. Geosci.*, 11(1), 27–37, doi:10.1038/s41561-017-0028-x, 2018.
- 963 Tréguer, P. J. and De La Rocha, C. L.: The World Ocean Silica Cycle, *Ann. Rev. Mar.*
964 *Sci.*, 5(1), 477–501, doi:10.1146/annurev-marine-121211-172346, 2013.
- 965 Trimborn, S., Lundholm, N., Thoms, S., Richter, K.-U., Krock, B., Hansen, P. J. and
966 Rost, B.: Inorganic carbon acquisition in potentially toxic and non-toxic diatoms: the
967 effect of pH-induced changes in seawater carbonate chemistry., *Physiol. Plant.*, 133(1),
968 92–105, doi:10.1111/j.1399-3054.2007.01038.x, 2008.
- 969 Trimborn, S., Brenneis, T., Hoppe, C. J. M., Laglera, L. M., Norman, L., Santos-
970 Echeandía, J., Völkner, C., Wolf-Gladrow, D. and Hassler, C. S.: Iron sources alter the
971 response of Southern Ocean phytoplankton to ocean acidification, *Mar. Ecol. Prog. Ser.*,
972 578, 35–50, doi:10.3354/meps12250, 2017.
- 973 Vrieling, E. G., Gieskes, W. W. C. and Beelen, T. P. M.: SILICON DEPOSITION IN
974 DIATOMS: CONTROL BY THE pH INSIDE THE SILICON DEPOSITION
975 VESICLE, *J. Phycol.*, 35(3), 548–559, doi:10.1046/j.1529-8817.1999.3530548.x, 1999.
- 976 Ward, B. A., Dutkiewicz, S., Jahn, O. and Follows, M. J.: A size-structured food-web
977 model for the global ocean, *Limnol. Oceanogr.*, 57(6), 1877–1891,
978 doi:10.4319/lo.2012.57.6.1877, 2012.
- 979 Wilken, S., Hoffmann, B., Hersch, N., Kirchgessner, N., Dieluweit, S., Rubner, W.,
980 Hoffmann, L. J., Merkel, R. and Peeken, I.: Diatom frustules show increased
981 mechanical strength and altered valve morphology under iron limitation, *Limnol.*
982 *Oceanogr.*, 56(4), 1399–1410, doi:10.4319/lo.2011.56.4.1399, 2011.
- 983 Witt, V., Wild, C., Anthony, K. R. N., Diaz-Pulido, G. and Uthicke, S.: Effects of ocean
984 acidification on microbial community composition of, and oxygen fluxes through,
985 biofilms from the Great Barrier Reef, *Environ. Microbiol.*, 13(11), 2976–2989,

- doi:10.1111/j.1462-2920.2011.02571.x, 2011.
- Wolf-Gladrow, D. and Riebesell, U.: Diffusion and reactions in the vicinity of plankton: A refined model for inorganic carbon transport, *Mar. Chem.*, 59(1–2), 17–34, doi:10.1016/S0304-4203(97)00069-8, 1997.
- Wolf, K. K. E., Hoppe, C. J. M. and Rost, B.: Resilience by diversity: Large intraspecific differences in climate change responses of an Arctic diatom, *Limnol. Oceanogr.*, 63(1), 397–411, doi:10.1002/lno.10639, 2018.
- Wu, Y., Campbell, D. A., Irwin, A. J., Suggett, D. J. and Finkel, Z. V.: Ocean acidification enhances the growth rate of larger diatoms, *Limnol. Oceanogr.*, 59(3), 1027–1034, doi:10.4319/lo.2014.59.3.1027, 2014.
- Yoshimura, T., Nishioka, J., Suzuki, K., Hattori, H., Kiyosawa, H. and Watanabe, Y. W.: Impacts of elevated CO₂ on organic carbon dynamics in nutrient depleted Okhotsk Sea surface waters, *J. Exp. Mar. Bio. Ecol.*, 395(1–2), 191–198, doi:10.1016/j.jembe.2010.09.001, 2010.
- Yoshimura, T., Suzuki, K., Kiyosawa, H., Ono, T., Hattori, H., Kuma, K. and Nishioka, J.: Impacts of elevated CO₂ on particulate and dissolved organic matter production: Microcosm experiments using iron-deficient plankton communities in open subarctic waters, *J. Oceanogr.*, 69(5), 601–618, doi:10.1007/s10872-013-0196-2, 2013.
- Young, J. N., Kranz, S. A., Goldman, J. A. L., Tortell, P. D. and Morel, F. M. M.: Antarctic phytoplankton down-regulate their carbon-concentrating mechanisms under high CO₂ with no change in growth rates, *Mar. Ecol. Prog. Ser.*, 532, 13–28, doi:10.3354/meps11336, 2015.
- Zeebe, R. E. and Wolf-Gladrow, D. A.: CO₂ in seawater: Equilibrium, kinetics, isotopes, Elsevier O., Elsevier, Amsterdam., 2001.

1010

1011 **Tables and Figures**

1012

1013 **Table 1.** Response of diatom communities to high CO₂. 69 experiments from 54
 1014 publications were considered here. Location refers to the place where diatom
 1015 communities were collected. The RDR is dimensionless (see methods). T is the average
 1016 incubation temperature in °C. DoE are days of experiment with the number of samplings
 1017 given as the second number. Pre-filt. gives the mesh size in case the collected plankton
 1018 community was pre-filtered before incubation. Setup refers to the incubation style:
 1019 undiluted volumes (batch), repeatedly diluted volumes (s.-cont.), flow-through setups

(fl.-thr.; only benthos), chemostats (chem.; only pelagic), CO₂ vent sites (seep; only benthos). Incubations can either be performed on deck (e.g. shipboards), *in situ* (e.g. *in situ* mesocosms) or under laboratory conditions. V refers to the incubation volume. Nutrient amendments were made in some but not all studies. The element indicates which nutrients were added. Asterisks indicate the presense of residual nutrients at the beginning of the study. Manipulations were done with: CO₂ saturated seawater (SWsat), acid additions (Acid), combined additions of acid and base (Comb.), CO₂ gas additions (CO₂), Aeration at target *p*CO₂ (Aer.), Passing CO₂ gas through a diffusive silicone tubing (Diff.). Meth. indicates the applied methodology to investigate diatom communities: light microscopy (LM), pigment analyses (PA), flow cytometry (FC), genetic tools (PCR), biogenic silica (BSi). The *p*CO₂ range of the experiment with the number of treatments given in brackets. The response of the bulk diatom community to CO₂: no effect (~), positive (p), negative (n), not reported (N/A). The *p*CO₂ response indicates approximately in between which treatments a CO₂ response was observed. Please note that this is based on visual inspection of the datasets and therefore involves subjectivity. Please also note that the range equals the treatment values in case only two treatments were set up. CO₂ induced shifts between diatom species can be: shift to larger species (large), shift to smaller species (small), unspecified shift (shift), no species shift detected (~), not reported (N/A). Winners or losers of the diatom community comprise: *Chaetoceros* (Chae), large *Chaetoceros* (Chae I), medium *Chaetoceros* (Chae II), small *Chaetoceros* (Chae III), *Neosyndra* (Neos), *Rhabdonema* (Rhab), *Eucampia* (Euca), *Cerataulina* (Cera), *Thalassiosira* (Thals), *Proboscia* (Prob), *Pseudo-nitzschia* (Ps-n), *Thalassionema* (Thalns), *Cylindrotheca* (Cyli), *Guinardia* (Guin), *Synedropsis* (Syned), *Dactyliosolen* (Dact), *Toxarium* (Toxa), *Leptocylindrus* (Lept), *Grammatophora* (Gram), *Bacillaria* (Baci), *Navicula* (Navi).

Reference	lat	long	RDR	S	T (°C)	Habitat	DoE/ # of sampl.	Pre-filt. (µm)	Setup	Incub.	V (L)	Nutr.	Manip.	Meth.	pCO ₂ range (µatm)	CO ₂ effect	pCO ₂ response (µatm)	Intra-taxon effect	Winners	Losers
(Bach et al., 2017)	58.264	11.479	76.2	29	7	est.	113/57	1000	batch	in situ	50000	*none	SWSat	PA, LM	(2) 380, 760	p	380 - 760	large	Cosc	
(Bach et al., 2019)	27.990	15.369	59.6	37	18.5	coastal	32/21	3000	batch	in situ	8000	N,P,Si	SWSat	LM, BSi	(7) 380 - 1120	p	380 - 1120	large	Chae, Guin, Lept	Nitz
(Biswas et al., 2011)	16.750	81.100	2.1	25	29.5	est.	5/2	200	batch	Deck	5.6	*none/N, P	Comb.	PA	(4) 230 - 1860	n	650 - 1400	N/A		
(Biswas et al., 2017)	17.000	83.000	1.5	?	?	coastal	2/1	200	batch	Deck	2	*N,P,Si,Fe, (Zn)	Comb.	LM	(2) 230, 2200	p	230 - 2200	shift	Skel	Thals
(Davidson et al., 2016)	-68.583	77.967	10.5	34	0.1	coastal	8/5	200	batch	Lab	650	*Fe	SWSat	LM	(6) 80 - 2420	n	1280 - 1850	small	Frag	Chae
(Domingues et al., 2017)	37.017	-8.500	7.4	?	23.5	est.	1/1	no	batch	Deck	4.5	N,P,Si,NH ₄	Comb.	LM, PA	(2) 420, 710	~		~		
(Donahue et al., 2019)	-45.800	171.130	2.6	34	11	oceanic	14/5	200	batch	Lab	10	*Fe	Diff.	LM, FC	(2) 350, 620	~		N/A		
(Donahue et al., 2019)	-45.830	171.540	2.6	34	11	oceanic	21/4	200	batch	Lab	10	*Fe	Diff.	LM, FC	(2) 350, 630	p	350 - 630	N/A		
(Eggers et al., 2014)	38.633	27.067	1.9	36	15	coastal	9-10/3	200	batch	Deck	4	N,P,Si	Comb.	LM	(2) 380, 910	p	380 - 910	large	Chae III	Thals
(Eggers et al., 2014)	38.650	27.250	1.9	36	15	coastal	9-10/4	200	batch	Deck	4	N,P,Si	Comb.	LM	(2) 380, 910	p	380 - 910	large	Thals, Chae II	Chae I
(Eggers et al., 2014)	38.617	27.250	1.9	36	15	oceanic	9-10/5	200	batch	Deck	4	N,P,Si	Comb.	LM	(2) 380, 910	~		N/A		
(Endo et al., 2013)	46.000	160.000	2.8	33	14	oceanic	14/3	197	batch	Deck	12	*none	Aer.	PA	(4) 230 - 1120	~		N/A		
(Endo et al., 2015)	53.083	177.000	2.8	?	8.2	oceanic	5/3	197	batch	Deck	12	*none	Aer.	PA, PCR	(2) 360, 600	n	360 - 600	~		
(Endo et al., 2016)	41.500	144.000	2.8	?	5.4	oceanic	3/3	197	batch	Deck	12	*Fe	Aer.	PA, PCR	(4) 180 - 1000	n	350 - 1000	shift		
(Feng et al., 2009)	57.580	15.320	1.7	35	12	oceanic	14/1-2	200	s.-cont.	Deck	2.7	N,P	Aer.	LM, PA	(2) 390, 690	p	390 - 690	large	Ps-n	Cyli
(Feng et al., 2010)	-74.230	179.230	1.7	34	0	oceanic	18/1-14	200	s.-cont.	Deck	2.7	none	Aer.	LM, PA	(2) 380, 750	~		large	Chae	Cyli
(Gazeau et al., 2017)	43.697	7.312	125.8	38	14	coastal	18/14	5000	batch	in situ	45000	none	SWSat	PA	(6) 350 - 1250	p	600 - 1000	N/A		
(Gazeau et al., 2017)	42.580	8.726	125.8	38	23	coastal	27/18	5000	batch	in situ	45000	none	SWSat	PA	(6) 420 - 1250	~		N/A		
(Grear et al., 2017)	41.575	71.405	9.3	?	9	est.	6/7	no	chem.	Deck	9.1	?none	Comb.	LM	(3) 220 - 720	~		~		
(Hama et al., 2016)	34.665	138.940	7.1	?	?	coastal	29/11	100	batch	Deck	400	N,P,Si	Aer.	PA	(3) 400 - 1200	~		N/A		
(Hare et al., 2007)	56.515	164.730	6.0	?	10.4	coastal	9-10/5	no	s.-cont.	Deck	2.5	Fe,N,P, Si	Aer.	LM, PA	(2) 370, 750	n	370 - 750	shift		Cyli
(Hare et al., 2007)	55.022	179.030	6.0	?	10.4	oceanic	9-10/3	no	s.-cont.	Deck	2.5	Fe	Aer.	LM, PA	(2) 370, 750	n	370 - 750	N/A		
(Hopkins et al., 2010)	60.300	5.200	99.1	?	10	coastal	21/9	no	batch	in situ	11000	N, P	Aer.	LM	(2) 300, 600	n	300 - 600	N/A		
(Hoppe et al., 2013)	-66.833	0.000	1.9	34	3	oceanic	27-30/1	200	s.-cont.	Lab	4	*none	Aer.	LM	(3) 200 - 810	N/A	400 - 810	shift	Syned	Ps-n
(Hoppe et al., 2017b)	71.406	68.601	1.9	33	9.5	oceanic	8/3	100	s.-cont.	Deck	8	N,P,Si	Aer.	PA, LM	(2) 320, 990	~		~		

(Hoppe et al., 2017a)	63.964	60.125	-	1.9	32	7.9	oceanic	13-14/3	100	s.-cont.	Deck	8	N,P,Si	Aer.	LM	(2) 300, 960	n	300 - 960	shift	Frag	Ps-n
(Hussherr et al., 2017)	71.406	70.188	-	2.6	33	4.3	oceanic	9/3-9	200	batch	Deck	10	*none	Comb.	LM, PA	(6) 510 - 3300	n	1040 - 1620	~		
(James et al., 2014)	-45.639	170.671			?	11.6	benthic	42/2		fl.-thr.	Lab	0	none	Comb.	pic	(2) 400, 1250	~		N/A		
(Johnson et al., 2011)	38.417	14.950			38	23.5	benthic	21/1		seep	in situ	0	none	NA	PA, LM	(3) 420 - 1600	p	420 - 590	large	Toxa, Gram, Baci, Navi, Cocc	Cycl, Neos, Rhab, Nitz
(Kim et al., 2006)	34.600	128.500		4.3	?	14	coastal	14/7	60	batch	in situ	150	N,P	Aer.	LM	(3) 250 - 750	N/A	400 - 750	shift	Skel	Nitz
(Kim et al., 2010)	34.600	128.500		52.1	?	12	coastal	20/22	no	batch	in situ	1600	N,P,Si	SWSat/Aer.	LM	(2) 400, 900	~		shift	Skel	Euca
(Mallozzi et al., 2019)	29.241	90.935	-	2.4	12	21	est.	112/9	80	s.-cont.	Lab	20	*none	Aer.	PA, LM	(2) 400, 1000	~		shift	Cyli	
(Mallozzi et al., 2019)	29.272	89.963	-	2.4	17	21	est.	112/9	80	s.-cont.	Lab	20	*none	Aer.	PA, LM	(2) 400, 1000	~		shift	Cyli	
(Maugendre et al., 2015)	43.667	-7.300		1.9	?	15	oceanic	12/4	200	batch	Deck	4	none	SWSat	PA	(2) 360, 630	~		N/A		
(Nielsen et al., 2010)	56.057	12.648		1.6	19	10.7	est.	14/4	175	s.-cont.	Lab	2.5	*none	Acid	LM, PA	(3) 500 - 1500	~		~		
(Nielsen et al., 2012)	-42.887	147.339		1.8	31	16	coastal	14/4	250	s.-cont.	Lab	2.5	*none	Acid	LM, PA	(3) 300 - 1200	~		~		
(Park et al., 2014)	34.600	128.500		59.6	?	17	coastal	19/17	no	batch	in situ	2400	N,P,Si	SWSat/Aer.	LM, PA	(6) 160 - 830	p	160 - 830	N/A	Cera	
(Paul et al., 2015)	59.858	23.258	112.7		6	11	est.	46/22	3000	batch	in situ	54000	none	SWSat	PA	(6) 370 - 1230	p	820 - 1000	N/A		
(Reul et al., 2014)	36.540	-4.600		3.3	?	21	coastal	7/6	200	batch	Deck	20	control/N,P	Aer.	LM, PA	(2) 500, 1000	p	500 - 1000	large		
(Roleda et al., 2015)	-45.639	170.671			34	10.8	benthic	112/7		fl.-thr.	Lab	0.65	none	Comb.	PA	(2) 430, 1170	~		N/A		
(Rossoll et al., 2013)	54.329	10.149		8.1	18	18	est.	28/7	200	batch	Lab	300	N,P,Si	Aer.	LM	(5) 390 - 4000	~		N/A		
(Sala et al., 2015)	41.667	2.800	26.1		38	14	coastal	9/2	no	batch	Lab	200	none	CO2	LM	(2) 400, 800	~		N/A		
(Sala et al., 2015)	41.667	2.800	26.1		38	22	coastal	9/2	no	batch	Lab	200	none	CO2	LM	(2) 400, 800	~		N/A		
(Schulz et al., 2008)	60.267	5.217	133.7		31	10.5	coastal	25/18-23	no	batch	in situ	27000	N,P	Aer.	PA	(3) 350 - 1050	~		N/A		
(Schulz et al., 2013)	78.937	11.893	106.1		34	3	coastal	30/26 - 30	3000	batch	in situ	45000	N,P,Si	SWSat	LM, PA	(8) 185 - 1420	~		N/A		
(Schulz et al., 2017)	60.265	5.205	125.8		32	9	coastal	38/35	3000	batch	in situ	75000	*N, P	SWSat	LM, PA	(8) 310 - 3050	n	1165 - 1425	N/A		
(Segovia et al., 2017)	60.390	5.320	99.1		?	11	coastal	22/9	no	batch	in situ	11000	control	SWSat/Aer.	FC	(2) 300, 800	~		N/A		
(Sett et al., 2018)	54.329	10.149	13.5		20	5	est.	44/26	200	batch	Lab	1400	*none	SWSat	LM, FC	(2) 540, 1020	~		~		
(Shaik et al., 2017)	15.453	43.801	5.6		35	29	coastal	2/1	no	batch	Deck	2	N,P,Si,Fe	CO2	LM	(2) 330, 1000	p	330 - 1000	~		
(Shaik et al., 2017)	15.453	43.801	5.6		36	29	coastal	9/1	no	s.-cont.	Deck	2	N,P,Si,Fe	CO2	LM	(2) 400, 1000	p	400 - 1000	~		
(Shaik et al., 2017)	15.453	43.801	5.6		35	29	coastal	2/1	no	batch	Deck	2	N,P,Si,Fe	CO2	LM	(2) 240, 780	p	240 - 780	~		

(Sommer et al., 2015)	54.329	10.149	49.8	20	9,15	est.	24/11	no	batch	Lab	1400	*none	SWsat	LM	(2) 440, 1040	~		shift		Prob, Thaln, Guin, Ps-n, Chae
(Tatters et al., 2013)	-45.752	170.81 0	0.8	35	14	coastal	14/2	80	s.-cont.	Lab	0.8	N,P,Si,Fe	Aer.	LM	(3) 230 - 570	N/A	400 - 570	shift	Cosc, Ps-n	Navi, Chae
(Tatters et al., 2018)	33.750	- 118.21 5	12.1	?	19	coastal	10/1	no	chem.	Deck	20	N/urea,P, Si	Aer.	LM	380, 800	N/A		shift		
(Taucher et al., 2018)	27.928	- 15.365	97.6	37	24-22	coastal	60/35	3000	batch	in situ	35000	N,P,Si	SWsat	LM, PA	(8) 350 - 1030	p	890 - 1030	large	Guin	Lept
(Thoisien et al., 2015)	69.217	53.367	1.4	33	3	coastal	8-17/6-9	250	s.-cont.	Lab	1.2	*none	SWsat	LM	(4) 440 - 3500	n	440 - 900	shift	Navi I	Navi II
(Tortell et al., 2002)	-6.600	- 81.017	7.1	?	?	oceanic	11/4	no	s.-cont.	Deck	4	*none	Aer.	PA, LM	(2) 150, 750	p	150 - 440	~		
(Tortell et al., 2008)	NA	NA	7.1	?	0	N/A	10-18/?	no	s.-cont.	Lab	4	*Fe	Aer.	LM, PA	(3) 100 - 800	p	100 - 400	large	Chae	Ps-n
(Tortell et al., 2008)	NA	NA	7.1	?	0	N/A	10-18/?	no	s.-cont.	Deck	4	*Fe	Aer.	LM, PA	(3) 100 - 800	p	100 - 400	large	Chae	Ps-n
(Tortell et al., 2008)	NA	NA	7.1	?	0	N/A	10-18/?	no	s.-cont.	Deck	4	*Fe	Aer.	LM, PA	(3) 100 - 800	p	100 - 400	large	Chae	Ps-n
(Trimborn et al., 2017)	-53.013	10.025	1.9	34	3	oceanic	30/4	200	s.-cont.	Lab	4	none	Aer.	LM	420, 910	n	420 - 910	shift		Ps-n
(Witt et al., 2011)	-23.450	151.91 7		?	24-25	benthic	11/4		fl.-thr.	Deck	10	none	SWsat	LM	(4) 310 - 1140	p	560 - 1140	N/A		
(Wolf et al., 2018)	78.917	11.933	1.9	?	3	coastal	10 - 13/1	200	s.-cont.	Lab	4	none	Aer.	LM	(2) 400, 1000	N/A	400 - 1000	~		
(Yoshimura et al., 2010)	49.500	148.25 0	2.7	33	13.5	oceanic	14/5	243	batch	Deck	9		Aer.	PA	(4) 150 - 590	n	150 - 280	N/A		
(Yoshimura et al., 2013)	53.390	- 177.01 0	2.8	?	8.4	oceanic	14/3	197	batch	Deck	12	*none	Aer.	PA, LM	4 (300 - 1190)	p	960 - 1190	N/A		
(Yoshimura et al., 2013)	49.020	174.02 0	2.8	?	9.2	oceanic	14/3	197	batch	Deck	12	*none	Aer.	PA, LM	(4) 230 - 1110	p	880 - 1110	N/A		
(Young et al., 2015)	-44.779	- 64.073	7.1	?	-1	coastal	21/21	no	s.-cont.	Deck	4	*none	Aer.	PA	(3) 100 - 800	~		N/A		
(Young et al., 2015)	-44.780	- 64.073	7.1	?	-0.5	coastal	16/16	no	s.-cont.	Deck	4	*none	Aer.	PA, LM	(3) 100 - 800	~		N/A		
(Young et al., 2015)	-44.780	- 64.073	7.1	?	1.5	coastal	20/20	no	s.-cont.	Deck	4	*none	Aer.	PA	(3) 100 - 800	~		N/A		

1045

1046




Figure 1. RDR as a function of incubation volume and size of the mesh that was used while filling the incubation volumes (d_{mesh}). The black and white boxes illustrate approximate ranges of the three main types of containers used in experiments. Please note that the general definition for mesocosms are volumes >1000 L (Guangao, 1990) but since most authors also use this term for open batch incubations with volumes between 150 – 1000 L we also stick to this term for the intermediate class.




Figure 2. Distribution of experiments with associated OA response of the bulk diatom

communities as listed in Table 1. Blue circles = positive effect; red triangles = negative response; grey squares = no response; orange diamonds = response not reported. Locations were slightly modified in case of geospatial overlap to ensure visibility. Please note that the three blue points in the Ross Sea at about -68, -165 are approximate locations because the reference did not provide coordinates.




Figure 3. Summary of the literature analysis. (A) Response of the bulk diatom community to ocean acidification. (B) Shifts among different diatom species due to ocean acidification. ‘Shift to large’ and ‘shift to small’ indicate that the diatom community

shifted towards the dominance of larger or smaller species, respectively. (C) Same data as in B but excluding studies where species shifts within the diatom community were not reported. This reduced the dataset from 69 to 40 studies. The left column is based on the number of studies. For example, the bulk diatom community was positively affected by OA in 29 out of 69 studies which is 33 %. The right column is based on the RDR values. For example, the \sum RDR value of all studies where the diatom community was positively affected by OA was 605 which is 36 % of the total \sum RDR.




Figure 4. Comparison of the diatom bulk response to OA in different environments. (A) coastal + estuarine + benthic environments with 47 experiments. (B) Oceanic environments with 19 experiments. The left column is based on the number of studies. For example, the bulk diatom community was positively affected by OA in 5 out of 19

1078 studies in oceanic environments which is 26 %. The right column is based on the RDR
1079 values. For example, the \sum RDR value of all studies where the oceanic diatom community
1080 was positively affected by OA was 17 which is 32 % of the total \sum RDR.

1081