

# **Marine monitoring in Europe: is it adequate to address environmental threats and pressures?**

Suzanne J. Painting<sup>1</sup>, Kate A Collingridge<sup>1</sup>, Dominique Durand<sup>4</sup>, Antoine Grémare<sup>2</sup>, Veronique Créach<sup>1</sup>, Christos Arvanitidis<sup>3</sup>, Guillaume Bernard<sup>5</sup>

Affiliations:

<sup>1</sup> Centre for Environment, Fisheries and Aquaculture Science (CEFAS), Lowestoft, NR33 0HT, UK.

<sup>2</sup> Université de Bordeaux, EPOC, UMR 5805, F-33615 Pessac, France.

<sup>3</sup> Hellenic Centre for Marine Research, Heraklion, 71500, Greece.

<sup>4</sup> COVARTEC, Bergen, 5141, Norway.

<sup>5</sup> CNRS, EPOC, UMR 5805, F-33615 Pessac, France.

*Correspondence to:* Suzanne J. Painting ([suzanne.painting@cefas.co.uk](mailto:suzanne.painting@cefas.co.uk))

## 1 Abstract

2 We provide a review of the environmental threats and gaps in monitoring programmes in  
3 European coastal waters based on previous studies, an online questionnaire, and an in-depth  
4 assessment of observation scales. Our findings underpin the JERICO-NEXT<sup>1</sup> monitoring  
5 strategy for the development and integration of coastal observatories in Europe, and support  
6 JERICO-RI<sup>2</sup> in providing high-value physical, chemical and biological datasets for addressing  
7 key challenges at a European level. This study highlights the need for improved monitoring of  
8 environmental threats in European coastal environments.

9 Participants in the online questionnaire provided new insights into gaps between environmental  
10 threats and monitoring of impacts. In total, 36 national representatives, scientists and  
11 monitoring authorities from 12 European countries (Finland, France, Germany, Greece,  
12 Ireland, Italy, Malta, Norway, Poland, Spain, Sweden, United Kingdom) completed the  
13 questionnaire, and 38 monitoring programmes were reported. The main policy drivers of  
14 monitoring were identified as the EU Water Framework Directive (WFD), Marine Strategy  
15 Framework Directive (MSFD), Regional Seas conventions (e.g. OSPAR) and local drivers.  
16 Although policy drivers change over time, their overall purposes remain similar. The most  
17 commonly identified threats to the marine environment were: marine litter, shipping,  
18 contaminants, organic enrichment, and fishing. Regime change was identified as a pressure by  
19 67% of respondents. The main impacts of these pressures or threats were identified by the  
20 majority of respondents (>70%) to be habitat loss or destruction, underwater noise, and  
21 contamination, with 60% identifying undesirable disturbance (e.g. oxygen depletion), changes  
22 in sediment/substrate composition, changes in community composition, harmful micro-  
23 organisms and invasive species as impacts.

24 Most respondents considered current monitoring of threats to be partially adequate or not  
25 adequate. The majority of responses were related to spatial and/or temporal scales at which  
26 monitoring takes place, and inadequate monitoring of particular parameters. Suggestions for  
27 improved monitoring programmes included improved design, increased monitoring effort and  
28 better linkages with research and new technologies. Improved monitoring programmes should  
29 be fit-for-purpose, underpin longer-term scientific objectives which cut across policy and other  
30 drivers, and consider cumulative effects of multiple pressures.

31 The JERICO-RI aims to fill some of the observation gaps in monitoring programmes through  
32 development of new technologies. The science strategy for JERICO-RI will pave the way to a  
33 better integration of physical, chemical and biological observations into an ecological process  
34 perspective.  
35

---

<sup>1</sup> JERICO-NEXT is the European H2020 project under grant agreement No. 654410.

<sup>2</sup> JERICO-RI is the European coastal research infrastructure (RI) community built by and through JERICO-NEXT and its predecessor JERICO (Framework 7 Grant Agreement 49 no 262584).

## 36 1. Introduction

37 Across the globe, marine monitoring networks are becoming increasingly important for the  
38 collection, dissemination and sharing of data for improved scientific understanding, assessment  
39 of the health of marine ecosystems and forecasting the likely impacts of environmental change  
40 and human activities (e.g. Schofield et al 2002; Schofield et al 2003; Proctor and Howarth  
41 2008; Duarte et al 2018; Bailey et al 2019; Bax et al 2019; Buck et al 2019; Canonico et al  
42 2019; Davidson et al 2019; Grand et al 2019; Smith et al 2019a; Smith et al 2019b). In Europe,  
43 for example, projects and infrastructures such as JERICO<sup>3</sup>, DEVOTES<sup>4</sup>, COPERNICUS<sup>5</sup>,  
44 EMODnet<sup>6</sup>, EMSO ERIC<sup>7</sup>, and AtlantOS<sup>8</sup> have played a significant role in the co-ordination  
45 and advancement of monitoring in coastal and offshore waters, from operational marine  
46 services through to delivering data products to end users. Changing pressures (e.g. due to  
47 population growth and climate change) and changing requirements to monitor, manage and  
48 mitigate the impacts of pressures require ongoing review of monitoring programmes. Over the  
49 past few decades, marine monitoring has been implemented in coastal and shelf seas around  
50 Europe in response to local/regional monitoring and oceanographic research demands.  
51 However, heterogeneity in monitoring methods and approaches has limited the integration of  
52 coastal observations. Many of the observations are driven by short-term research projects,  
53 potentially limiting the sustainability of observing systems for meeting monitoring and  
54 assessment needs.

55 The Dobris Assessment (EEA 1995) listed 56 broad environmental threats, 19 of which were  
56 relevant to the coastal domain. These include physical modifications (e.g. due to urban  
57 development, industry, energy production, military activities, fisheries, recreation),  
58 contamination and coastal pollution (e.g. due to wastewater disposal, chemical contaminants,  
59 marine litter) and loss of biodiversity and genetic resources. Recent EU policy drivers and  
60 regional sea conventions have led to improvements in water quality in many regions (notably  
61 the Baltic Sea, North Sea, Celtic Sea, Bay of Biscay). Nonetheless, the fourth assessment of  
62 the European environment (EEA 2008a; see also EEA 2015a) highlighted that some regions  
63 remain affected by eutrophication, destructive fishing practices, hazardous substances, oil  
64 pollution and invasive species. Key concerns include increasing population densities and

---

<sup>3</sup> <http://www.jerico-ri.eu/previous-project/jerico-fp7/>

<sup>4</sup> <http://www.devotes-project.eu/>

<sup>5</sup> <https://www.copernicus.eu/en>

<sup>6</sup> <http://www.emodnet.eu/>

<sup>7</sup> <http://www.emso.eu/>

<sup>8</sup> <https://www.atlantos-h2020.eu/>

65 development of built-up areas, and likely impacts of climate change on physical (e.g.  
66 temperature, currents), chemical (e.g. acidification) and biological (e.g. changes in growth,  
67 species composition and distribution, loss of organisms with carbonate shells) components.  
68 The lack of comparable data presents a major obstacle for assessments of Europe's regional  
69 seas, even for well-known problems such as eutrophication (EEA 2008b; OSPAR 2017). More  
70 and better data are needed to develop a pan-European marine protection framework that  
71 addresses environmental issues in a cost-effective way.

72 A number of studies have considered the suitability of monitoring programmes in Europe (e.g.  
73 Bean et al 2017; Borja et al 2019; DEVOTES<sup>9</sup>; Garcia-Garcia et al 2019; Tett et al 2013;  
74 Zampoukos et al 2013) for assessing good environmental status (GES) of the biodiversity suite  
75 of MSFD descriptors (D): D1 (biodiversity), D2 (non-indigenous species), D4 (food-webs) and  
76 D6 (seafloor integrity). Limitations have been identified in monitoring programmes, including  
77 limitations in spatial and temporal coverage, pressures addressed, integrated monitoring  
78 (addressing more than one descriptor and/or ecosystem component simultaneously), indicators  
79 used, and data accessibility. Differences between countries highlight budgetary constraints and  
80 differing approaches to monitoring. The Baltic region has been shown to be good at addressing  
81 multiple descriptors simultaneously, while the Mediterranean has a good history of co-  
82 ordination between countries and making good use of citizen science. Improved compatibility  
83 of datasets (for example, through standardisation of sampling methods and quality assurance  
84 of the data) and translating research activities into monitoring (e.g. for litter and noise) have  
85 been highlighted as key challenges (EEA 2008a; EU DEVOTES).

86 The EU JERICO-NEXT<sup>10</sup> project addresses the challenges of observing the complex and  
87 highly variable coastal seas at a Pan-European level, in order to improve operational marine  
88 services and meet the requirements of key policy drivers such as EU Directives. The emphasis  
89 is on providing an integrated European observing system supporting improved understanding  
90 of the coupling between physics, biogeochemistry and biology to take account of and address  
91 the complexity of the coastal environment. This requires development and application of new  
92 technologies that allow for the continuous monitoring of a larger set of parameters. It also  
93 requires an *a priori* definition of the optimal sampling strategy over very different spatial and  
94 temporal scales to develop fit-for-purpose coherent monitoring programmes. This will enable

---

<sup>9</sup> DEVOTES is an EU FP7 project

<sup>10</sup> JERICO-NEXT is a Horizon 2020 funded project, implementing the second phase of the European JERICO-RI research infrastructure aiming at multidisciplinary observations of coastal and shelf seas.

95 the JERICO community to meet the overall objective of extending the EU network of coastal  
96 observations developed during JERICO-FP7. As part of the JERICO-NEXT project, we  
97 conducted an opinion poll of experts in European countries (Figure 1) to identify current and  
98 emerging pressures or threats to the marine environment, identify gaps in monitoring these  
99 pressures, and contribute towards a forward-looking strategy for monitoring marine  
100 ecosystems.

101

102

103

104

105

106

107

108

109


110

111

112

113

114



115 *Figure 1. The countries which participated in the poll were Finland, France, Germany, Greece,*  
116 *Ireland, Italy, Malta, Norway, Poland, Spain, Sweden, United Kingdom.*

117

## 118 **2. Methodology**

119 The opinion poll was designed as an online questionnaire, which could be completed over a  
120 five-month period (29 June to 30 November 2016). The questionnaire was distributed to all  
121 partners in the JERICO-NEXT project. Partners were tasked with being national  
122 representatives and were asked to take responsibility for responding to the questionnaire  
123 and/or to collect answers from colleagues, collaborators and responsible monitoring  
124 authorities within their countries. The national representatives were also asked to forward the  
125 questionnaire to the relevant authorities in countries which were not partners within JERICO-  
126 NEXT.

127 Questionnaire development was informed by a review of existing literature on environmental  
128 pressures and threats (e.g. EEA 2008a) and the outputs of the DEVOTES project (DEVOTES  
129 2014). Threats to the marine environment were considered in terms of 'pressures' and  
130 'impacts'. Pressures were described as the human activities which have impacts on  
131 ecosystems or parts thereof (see Oosterwind et al 2016<sup>11</sup>), which is compatible with the  
132 driver-pressure-state-impact-response (DPSIR) framework (Gabrielsen and Bosch 2003;  
133 Elliott 2014).

## 134 **2.1. Format of questionnaire**

135 The questionnaire (Figure 2, for more detail see supplementary material, S1) was developed  
136 using Google Forms, and consisted of two linked forms. The first form was focussed on  
137 obtaining the views of all respondents on the environmental threats in European waters and the  
138 adequacy of current monitoring programmes. Maps were provided to ensure consistency in  
139 participant selection of 'regions of interest' (see supplementary material, S2 and S3). For  
140 questions related to pressures and impacts, respondents could select one or more responses  
141 from lists provided. They could also add free text in order to provide detail or explanations of  
142 their responses. Questions related to adequacy of existing monitoring programmes included  
143 comments boxes for free text, to allow respondents to give their views on those monitoring  
144 programmes which were not adequate or only partly adequate for addressing environmental  
145 threats, and suggestions on how to improve the monitoring of the threats identified.

146 The second form was focussed on national monitoring programmes, with the aim of obtaining  
147 a summary of sampling platforms used, variables measured, monitoring frequency and the  
148 duration of the programme. This form included a section on data accessibility.

149 An invitation to participate in the poll and complete the questionnaire was sent to all partners  
150 in JERICO-NEXT in June 2016 and subsequently forwarded to wider contact networks. It was  
151 closed to responses in November 2016.

152

---

<sup>11</sup> **Pressures** can be described as 'a result of a driver-initiated mechanism (human activity/natural process) causing an effect on any part of an ecosystem that may alter the environmental state'. **Impacts** can be defined as 'consequences of environmental state change in terms of substantial environmental and/or socio-economic effects which can be both, positive or negative'.

153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179

<p>Section 1 <a href="#">Environmental Threats and Monitoring</a></p> <ol style="list-style-type: none"><li>1. Participant Details:<ol style="list-style-type: none"><li>a. Name and contact details</li><li>b. Institute/Affiliation</li></ol></li><li>2. Region of interest (see Annex 2)<ol style="list-style-type: none"><li>a. Country</li><li>b. Region</li><li>c. Sub-Region</li></ol></li><li>3. Review of threats per region<ol style="list-style-type: none"><li>a. Pressures: What are the main pressures from human activities that are affecting the environment in this area?</li><li>b. Impacts: What are the impacts resulting from the pressures identified above?</li></ol></li><li>4. Policy Purposes: What are the main policy or other drivers behind the monitoring programme/s in each region or sub-region? These may be international conventions, EU Directives, national policies, or other requirements.</li><li>5. MSFD Descriptors: The MSFD includes 11 qualitative descriptors. Please link the threats identified to these descriptors, or any others which may be relevant in the area.</li><li>6. Names of relevant monitoring programmes:</li><li>7. Adequacy of existing monitoring programmes: are they sufficient to assess the effects of the environmental threats in the considered area?<ol style="list-style-type: none"><li>a. How are they deficient?</li><li>b. How could they be improved to better address the threats?</li></ol></li></ol> <p>Section 2 <a href="#">Monitoring programmes</a></p> <ol style="list-style-type: none"><li>1. Country</li><li>2. Monitoring programme name</li><li>3. Is the program statutory/official or unofficial?</li><li>4. Variables measured</li><li>5. Platform types</li><li>6. Number of stations</li><li>7. Is monitoring regular or ad hoc?</li><li>8. Monitoring frequency</li><li>9. Start date</li><li>10. End date</li><li>11. End reason, if not ongoing</li><li>12. Monitoring stations (in separate spreadsheet).</li><li>13. Comments</li><li>14. Data access restrictions</li><li>15. Responsible organisation</li><li>16. Responsible person and details</li><li>17. Data source institute</li><li>18. Database to which the data are submitted</li><li>19. Are data flows to central databases up to date?</li><li>20. Web links to data</li></ol>
---

*Figure 2. Format of online questionnaire.*

### **2.1.1. Data Analysis**

Once the poll was closed, responses were downloaded from Google Forms and stored in a MS Access Database. Identifying information was removed from the responses to anonymise the data. More than one response was received from some countries. Results on views or opinions on environmental threats and impacts and on monitoring programmes were analysed using responses by country, i.e. categorial responses were aggregated by country, counting each response if it appeared at least once in the individual responses for the country. Marine litter, for example (see Section 3), was identified as a threat/pressure in all of the ‘national responses’; however, it was not identified in every single individual response from each country where there were multiple responses. The aggregated responses are referred to hereafter as ‘national responses’.

180 Details of monitoring programmes and expert opinions on adequacy of monitoring  
181 programmes were analysed for all respondents. Opinions were also analysed within each  
182 country. Free-text responses from all respondents on the adequacy of monitoring programmes  
183 were extracted to summarise all opinions given, and the suggestions for improving monitoring  
184 programmes that were not adequate or partly adequate to address environmental threats.

185 To visualize the most common themes emerging from the questions on why monitoring  
186 programmes were inadequate, word clouds were created using an online software tool (Wordle  
187 2018) which emphasises the most common responses from individuals according to how many  
188 times they are mentioned.

189

### 190 **3. Results**

#### 191 **3.1. Respondents**

192 The online questionnaire was completed by representatives from 12 European countries  
193 (Finland, France, Germany, Greece, Ireland, Italy, Malta, Norway, Poland, Spain, Sweden,  
194 United Kingdom, Figure 1) representing different regional seas (Figure 3) and their sub-regions  
195 (see supplementary material, S2 and S3). From some countries, responses were received from  
196 more than one respondent, resulting in a total of 36 responses from the 12 countries. The most  
197 responses (14) were received from the UK and covered territorial waters (12 nm) as well as  
198 their Exclusive Economic Zone (EEZ) waters. Five responses were received from Greece, six  
199 from France, two from Spain, and two from Malta. Many respondents were JERICO-NEXT  
200 partners, but some were also from the wider European monitoring network. Two responses  
201 were received from people in organisations which represent multiple countries (see S3, Table  
202 S3.1). From EuroGOOS, a Swedish representative answered from a Swedish perspective. From  
203 OSPAR, a UK-based person answered for the region as a whole.

204 To reduce bias in the results due to different numbers of respondents from each country, views  
205 on threats, impacts and adequacies of monitoring programmes were aggregated to give one  
206 response per country. This was considered to represent a national response (see Section 2.1.1).  
207 Data analysis showed weak relationships between the number of pressures or impacts identified  
208 per country and the number of responses per country (data not shown).



209 **3.2. Views on environmental threats and impacts**

210 **3.2.1. Pressures from human activities**

211 Marine litter was identified as a pressure in all of the national responses (Figure 4). The next  
212 most commonly identified pressures were shipping (92%), contaminants (92%) organic  
213 enrichment (83%) and fishing (75%, Figure 4). These were followed by regime change (67%),  
214 inorganic nutrient enrichment and aquaculture (both 58%, Figure 4), dumping and aggregate  
215 extraction (50%), and atmospheric inputs, dredging of biota and construction/obstruction (all  
216 42%). Activities such as mining, water abstraction, the oil and gas industry and coastal squeeze  
217 scored considerably lower, at 10-23% of responses. Only one extra pressure was added to the  
218 list provided, unexploded ordnance (UXO).

219 Respondents noted that the pressures affecting coastal and offshore areas were not the same.  
220 Climate change related pressures (regime change and ocean acidification) were considered to  
221 have large potential for widespread harm and in all sea regions at least one respondent marked  
222 regime change as an important pressure. Thermally-driven regime change was selected in a  
223 greater proportion of responses than salinity-driven regime change.

224

225

226

227

228

229

230

231

232

233

234




235 *Figure 3. The regional seas represented by respondents to the questionnaire (see supplementary*  
236 *material for maps of regions [S2] and sub-regions of European seas [S3]).*

237


238 **3.2.1. Impacts of the pressures identified**

239 The majority of national responses (>70%) identified habitat loss or destruction, underwater  
 240 noise, and contamination as the main impacts of human activities on the marine environment  
 241 (Figure 5). Approximately 60% of national responses identified undesirable disturbance (e.g.  
 242 oxygen depletion), changes in sediment/substrate composition, changes in community  
 243 composition, harmful micro-organisms and invasive species as impacts. Fifty percent (50%)  
 244 identified changes in primary production, changes in species range, population  
 245 change/depletion of standing stocks, biofouling, physical damage, changes in suspended  
 246 sediments/turbidity and mortality of marine life.

247



248 *Figure 4. Frequency of national responses on pressures affecting the marine environment.*



250 *Figure 5. Frequency of national responses on impacts affecting the marine environment.*

251

### 252 **3.3. Views on the main drivers of marine monitoring**


#### 253 **3.3.1. Policy purposes**

254 The majority of national responses (83%) identified the main drivers of monitoring of coastal  
 255 and offshore waters as the Water Framework Directive (WFD, EU 2000) and the Marine  
 256 Strategy Framework Directive (MSFD, EU 2008, Figure 6). Other EU directives were  
 257 identified but the proportion of national responses identifying these as policy purposes for  
 258 monitoring was relatively low. Twenty five percent (25%) of national responses included the  
 259 Urban Waste Water Treatment Directive and Habitats and Birds Directive (Figure 6), and 17%  
 260 included the Bathing Waters Directive and the Nitrates Directive. Regional Seas Conventions  
 261 were also identified as drivers of marine monitoring, with OSPAR identified by 67% of  
 262 national responses and HELCOM identified by 17% of national responses. Local policy drivers  
 263 were identified by 58% of national responses, but no details were given.


264 Respondents were asked to link environmental threats in European waters to the descriptors  
 265 (D) in the MSFD (Figure 7; see EU 2008). Responses indicated that most threats (92%) affect  
 266 the biodiversity descriptor (D1, Figure 7). The next most frequent responses (83%, Figure 7)

267 were linked to descriptors for contaminants (D8), eutrophication (D5) and marine litter (D10).  
 268 Seventy five percent (75%) of threats could be linked to the energy descriptor (D11), 67% to  
 269 sea floor integrity (D6), hydrographic conditions (D7) and non-native species (D2), and 50%  
 270 to food webs (D4).

271



272 *Figure 6. Main policy or other drivers for marine monitoring.*



273 *Figure 7. MSFD Descriptors linked to environmental threats. The left axis shows the descriptor*  
 274 *number and name.*

275

276 **3.3.2. Meeting requirements of policy drivers**

277 Much of the monitoring towards older directives is now included in WFD monitoring  
278 programmes implemented under River Basin Management Plans of Member States. These  
279 results highlight that policy drivers may change over time but overall purposes may remain the  
280 same or similar. Regional Seas conventions were also identified as key policy drivers of  
281 monitoring programmes, with a greater proportion of responses for OSPAR than for  
282 HELCOM.

283 **3.4. Monitoring Programmes in each country**


284 In total, 36 responses on the monitoring section of the questionnaire were received from the 12  
285 countries who participated in the online poll. Thirty-eight (38) monitoring programmes were  
286 reported. More than half of these programmes were official or statutory programmes, and a  
287 significant proportion (28%) were project based rather than statutory. These included the  
288 Balearic Islands multi-platform observing system (SOCIB), UK BeachWatch litter project and  
289 projects in Ireland (Smartbay observatory).

290 This is not a complete inventory of monitoring in Europe, but the responses provide examples  
291 of a variety of monitoring programmes. Entries for the UK, Ireland and Greece appeared to be  
292 relatively comprehensive.

293 **3.4.1. Monitoring: variables, platforms and frequency**

294 Most monitoring programmes were reported to measure temperature and salinity. A large  
295 proportion of responses (39-45%, Figure 8) reported measurements of nutrients, chlorophyll  
296 and dissolved gases, although not all parameters are measured at all stations in a monitoring  
297 programme. Many variables, such as mammals, birds, biotoxins and marine litter are only  
298 measured in specific monitoring programmes designed for the purpose. Some variables are  
299 monitored in only a few monitoring programmes, e.g. sea level and contaminants, but this may  
300 reflect the selection of responses received. Responses to the questionnaire indicated that marine  
301 monitoring programmes provide less coverage of biological parameters (e.g. plankton 32%,  
302 fish 18%, benthos 18%, macroalgae 11%, birds 3%) than physical water column parameters  
303 (e.g. temperature, salinity, 58-61%) and chemical parameters (e.g. nutrients, dissolved gases,  
304 45% and 39%).

305



306

307

308 *Figure 8. Variables measured in marine monitoring programmes.*

309

310 Most monitoring programmes were reported to use a vessel as a monitoring platform (Figure  
 311 9), usually a research vessel or, for inshore monitoring, a small boat. Shore based monitoring  
 312 was also common (39%). The use of fixed platforms was indicated by 34% of respondents,  
 313 including those from Belgium, Greece, Ireland, Italy, Spain and the UK. The use of remote  
 314 sensing as a monitoring platform was reported by 21% of respondents (Figure 9). Other  
 315 innovative and emerging technologies, such as autonomous vehicles, FerryBoxes and ‘other’  
 316 (e.g. profiling floats) were included in  $\leq 11\%$  of the responses (Figure 9).


317

318

319

320

321



322 *Figure 9. Platform types used in marine monitoring.*

323

324 Responses to the questionnaire indicated that monitoring frequency (Figure 10) is variable. The  
325 highest proportion of responses (34%) was for continuous monitoring (e.g. from fixed  
326 platforms, moorings or gliders). Several monitoring programmes were reported to have only  
327 annual monitoring, but to be comprehensive in terms of parameters and spatial coverage.  
328 Monitoring programmes incorporating fixed platforms or gliders were more restricted in terms  
329 of spatial coverage.

330

331

332

333

334


335

336

337

338

339



340 *Figure 10. Frequency of monitoring. The main graph shows results for all options given in the*  
 341 *questionnaire. The inset combines these into three categories: continuous and intermittent are the same*  
 342 *as in the main graph, regular = all other options combined.*

343

### 344 **3.4.1. Sustainability of monitoring programmes**

345 Responses to the questionnaire showed that 68% of the monitoring programmes have been  
 346 running for longer than 10 years. The longest programme reported was the continuous plankton  
 347 recorder survey, by the Sir Alister Hardy Foundation for Ocean Science (SAHFOS), which has  
 348 been running since 1931. Several French and Scottish monitoring programmes were reported  
 349 to have been running for approximately 30 years. One respondent included a monitoring  
 350 programme which ended due to lack of funding; it is likely there were many more such cases  
 351 which were not reported.

### 352 **3.4.2. Data access**


353 The majority of respondents (71%) reported that their monitoring programmes had no  
 354 restrictions on data access. Where data access is restricted, most programmes make the data  
 355 available on request, subject to information on the intended purpose or use of the data, signing  
 356 of a licence agreement, and/or requirements to acknowledge the source of the data (e.g. through  
 357 the use of data DOIs [digital object identifiers]).



358 Respondents reported that data were submitted most commonly to local/national databases, but  
359 frequently also to ICES databases, EMODnet or Copernicus. For the majority of programmes,  
360 data flows to these central databases were considered to be not up-to-date, indicating that not  
361 all monitoring data are available centrally, or that there is a time lag in submission of data.

### 362 3.5. Gaps identified in current monitoring programmes

363 In terms of providing the information required to monitor environmental threats, 12% of all the  
364 respondents to the questionnaire considered monitoring programmes to be adequate, while 28%  
365 indicated that monitoring programmes were not adequate and 60% considered monitoring  
366 programmes to be partially adequate (Figure 11).




376 *Figure 11. Proportion of all respondents who considered their monitoring programmes to be*  
377 *adequate (Yes), inadequate (No) or partly adequate (Partly).*

378  
379 Where there was more than one respondent per country, responses were varied (Figure 12),  
380 with the majority of responses indicating inadequate monitoring. In the UK, for example, where  
381 14 responses were received, most responses (57%) were that monitoring was partly adequate,  
382 and 29% were that monitoring was not adequate. Two respondents (15%) felt that monitoring  
383 programmes were adequate. In France, where six responses were received, the majority (83%)  
384 considered monitoring was not adequate, and the remaining one felt it was adequate. In Greece,  
385 four out of five respondents (80%) felt monitoring was not adequate, and one considered it to  
386 be partly adequate. In countries with two responses (Italy, Malta and Spain), one indicated that  
387 monitoring was not adequate while one felt it was partly adequate. In countries with one

388 respondent, responses were mostly that monitoring was partly adequate (Finland, Ireland,  
 389 Norway, Sweden). In Poland, the national representative reported that monitoring was  
 390 adequate.

391



392

393 *Figure 12. Responses by country showing the proportion of respondents who considered their*  
 394 *monitoring programmes to be adequate (Yes), inadequate (No) or partly adequate (Partly). The*  
 395 *number of respondents per country ranged from 1 to 14 (see numbers in bold).*

396

### 397 **3.5.1. Where monitoring is not adequate**

398 Responses were focussed around a few key issues (see Figure 13) which appeared to be related  
 399 mostly to insufficient resolution in time and space, insufficient data or parameters measured,  
 400 and lack of integration (e.g. of monitoring programmes, indicators and descriptors).


401 A number of respondents stated that there is insufficient monitoring for some of the MSFD  
 402 descriptors. These descriptors included biodiversity, food-webs, marine litter (including micro-  
 403 plastics), underwater noise, emerging contaminants, and emerging pollutants. However, no  
 404 details were given. It was noted that coupling between physics and biology in response to  
 405 environmental pressures is typically not included in monitoring programmes focussed on  
 406 individual descriptors. One respondent indicated that methodologies and approaches were not

407 state-of-the-art, for example, the focus during benthic sampling was on taxonomy instead of  
408 ecosystem functions and services.

409

410

411



412

413

414

415

416

417

418

Figure 13. Key words used in views on partially adequate or inadequate monitoring programmes. Font sizes indicate the most common responses from individuals according to how many times they are mentioned.

419

420 Two respondents highlighted concerns about the links to policy drivers, suggesting that  
421 monitoring was reactive rather than proactive. One of these respondents commented that  
422 monitoring programmes develop to respond to pressures and impacts. The other highlighted  
423 concerns related to unexploded ordnance, for which there seems to be very little political or  
424 commercial interest in finding and making safe dumped munitions, until a person or marine  
425 life is found with injuries or abnormal growth.

426 Examples of monitoring programmes with low spatial resolution were given for point source  
427 monitoring of contaminant inputs, controls and improvements; benthic habitats for the wider  
428 environment, and deep-sea areas; and sub-regions of Mediterranean Sea. Examples of  
429 inadequate monitoring of parameters were given for the Mediterranean Sea, including  
430 zooplankton, phytoplankton compositions, marine mammals, reptiles, birds, invasive species,  
431 marine litter, and contaminants in sediment and biota.

432

#### 433 4. Improving monitoring programmes

434 The respondents highlighted key gaps between the environmental pressures or threats and the  
435 monitoring of their impacts. Suggestions were given for improving monitoring programmes

436 considered to be not adequate or partly adequate. These were focussed on improved design of  
437 monitoring programmes and increased effort, observation and research, and included:

- 438 • Improved spatial and/or temporal resolution, and assessment of emerging threats.
- 439 • Improved monitoring of biological parameters and coupling between biological and  
440 physical or chemical parameters, particularly those which provide information on  
441 ecosystem function. Examples were given for poorly covered habitats, microbes,  
442 zooplankton, marine mammals, and biodiversity components not yet monitored.
- 443 • Increased use of new technologies (e.g. remote sensing, FerryBoxes, gliders) and  
444 methodologies (e.g. molecular techniques).
- 445 • Maintaining and/or developing a limited number of long-term (fixed-point) monitoring  
446 sites to monitor changes in baseline conditions (chemistry, ecotoxicology, and ecosystem  
447 structure) in response to climate change/acidification, and diffuse inputs.
- 448 • Making better use of low-cost biochemical sensors on low-cost platforms.
- 449 • Improved data flows (submission of data to centralised and/or open-access databases).
- 450 • More-integrated cross-disciplinary approaches, e.g. through more-coordinated monitoring  
451 across descriptors.
- 452 • Improved monitoring design to create programmes which are fit for multiple purposes. e.g.  
453 to take into account regional or national specificities or requirements (e.g. sub-regions of  
454 regional seas; rigid baseline ecological assessment at local scales; increased monitoring in  
455 high-risk areas), incorporate newer threats (e.g. phosphorous-based flame-retardants,  
456 microplastics, noise), and be more proactive regarding threats likely to cause harm to or  
457 changes in biota, e.g. unexploded ordnance (UXO).
- 458 • Including flexible research/investigative monitoring to increase knowledge of specific  
459 impacts.
- 460 • Securing funding for long-term monitoring programmes.

461

## 462 **5. Discussion**

### 463 **5.1. Polling Approach**

464 The opinion poll carried out during this study had a limited number of participants, as it was  
465 targeted towards scientists and managers with the relevant expertise and experience in  
466 European countries adjoining regional and/or sub-regional seas. In order to minimise bias  
467 which might be introduced by some countries providing more individual responses than other

468 countries, project partners were expected to develop national responses, and were given  
469 approximately six months to do so. Where there was more than one response from a country,  
470 results on views or opinions were combined to represent a national view

471 Responses on monitoring programmes were not combined, as these were considered to provide  
472 useful detail on gaps in monitoring, and no monitoring programmes had duplicate responses.

473 Despite a number of limitations in the polling approach, responses provided valuable insights  
474 on the environmental pressures and their impacts, and on gaps in monitoring the impacts. A  
475 recent study in estuaries and coastal waters in the North Sea - Baltic Sea transition zone  
476 (Andersen et al. 2019) using 35 databases yielded results which are broadly similar (see below).

## 477 **5.2. Drivers of marine monitoring**

478 Most national responses were focussed on policy drivers such as EU Directives and regional  
479 conventions based on the ecosystem approach. These responses are likely to have been  
480 influenced by the overall context of the JERICO-NEXT project and its emphasis on  
481 biogeochemical processes and the coupling between physics and biology. Responses may also  
482 have been influenced by the drop-down list of options from which to select answers, although  
483 the option was given to add responses.

484 Interestingly, local drivers scored quite highly. No details or examples were given by any of  
485 the respondents but may include monitoring towards impact assessments for a variety of  
486 reasons, such as development of local fisheries or recreational activities, or to meet  
487 conservation objectives (e.g. for marine protected areas). However, such monitoring would be  
488 included under policy drivers such as the Habitats and Birds Directives or Marine Planning,  
489 and relatively few responses ( $\leq 25\%$ ) indicated these as drivers for marine monitoring. It is  
490 possible that local drivers included research projects or programmes, but this seems unlikely  
491 as the poll was focussed on monitoring rather than research. This highlights a potential  
492 weakness of the aims of this study and indeed the JERICO-NEXT project, as it did not include  
493 an objective to identify gaps in understanding, and how to provide better linkages between  
494 research and monitoring. Certainly, ongoing national monitoring programmes are focused on  
495 reporting to directives and international obligations, and not to contribute to better  
496 understanding of the possible impacts of the threats.

497 Complex linkages between pressures and impacts and the cumulative effects of multiple  
498 pressures are not currently well addressed by any of the reported monitoring programmes. The

499 MSFD was intended as a holistic approach to assessments, but descriptors are currently  
500 assessed separately. Developments are underway to move assessments towards a more  
501 integrated cross-disciplinary ecosystem approach both in Europe (e.g. OSPAR<sup>12</sup>; EEA 2011;  
502 EEA 2015b; HELCOM 2018) and globally (Schmidt et al, 2019). This will require more co-  
503 ordinated monitoring across descriptors, and a focus on acquiring long-term data sets,  
504 particularly for addressing cross-cutting issues such as climate change and ocean acidification  
505 (Tett et al 2013, Schmidt et al 2019). Responses indicating that a number of monitoring  
506 programmes have been running for more than 10 years are extremely positive, providing data  
507 to allow the detection of temporal trends on pressures and their impacts on the marine  
508 environment. Evidence that a significant proportion of monitoring is largely project-based  
509 rather than statutory, indicates some degree of risk to the sustainability of monitoring.  
510 EuroGOOS conducted a survey of sea level monitoring and found similar issues; less than half  
511 of the organisations responding considered that there were no funding issues for tide gauges  
512 and many had reduced funding or uncertain future funding (EuroGOOS 2017). Similar issues  
513 are encountered in other parts of the world where monitoring is supported by both academic  
514 and private research and hampered by lack of sustained funding from governments where  
515 grants are often short-term (Weller et al 2019).

516 With the majority of responses to the online poll indicating that the main policy drivers of  
517 current monitoring are the MSFD and WFD, rather than earlier directives such as the UWWTD,  
518 the Nitrates Directive and the Habitats Directive, it is clear that policy drivers and requirements  
519 for meeting policy needs change over time. The findings also highlight that monitoring  
520 programmes should be underpinned by high-level scientific objectives, and that research and  
521 monitoring should be well integrated. Data sharing, such as through the JERICO-NEXT  
522 research infrastructure and coastal observatories and EMODNet Data infrastructure (Miguez  
523 et al 2019), is vital to current and future integration of research and monitoring (Farcy et al  
524 2019). Furthermore, the availability of data at local and regional scales is essential for  
525 development of future monitoring and assessment approaches, particularly as new technologies  
526 and assessment tools are developed and become more readily available (e.g. Borja et al 2019;  
527 García-García et al 2019).

---

<sup>12</sup> See <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/introduction/assessment-process-and-methods/>; and <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/chapter-6-ecosystem-assessment-outlook-developing-approach-cumul/>

### 528 **5.3. Views on environmental threats and impacts**

529 Respondents were provided with comprehensive lists of key environmental threats and impacts  
530 informed by previous studies, with an option to add to the list. One item, UXO, was added to  
531 the list of pressures by one country. This pressure was considered to be outside the scope of  
532 the JERICO-NEXT project but may be useful in other contexts. No new items were added to  
533 the list of impacts in the national responses.

534 The most commonly identified pressures or threats to the marine environment due to  
535 manageable human activities were considered to be marine litter, shipping, contaminants,  
536 organic enrichment, fishing, and regime change.

537 The main impacts of threats to the marine environment (i.e. >70% of national responses) were  
538 identified to be habitat loss or destruction, underwater noise, and contamination. Sixty percent  
539 (60%) of national responses identified impacts to be undesirable disturbance (e.g. oxygen  
540 depletion), changes in sediment/substrate composition, changes in community composition,  
541 harmful micro-organisms and invasive species.

542 In a recent study, Andersen et al (2019) analysed 35 publicly available datasets from Danish  
543 marine waters and obtained broadly similar results. These authors found the main stressors  
544 (pressures) across a range of water types to be nutrients, climate anomalies, non-indigenous  
545 species, noise and contaminants. Some stressors (e.g. fisheries, contaminants, noise) were  
546 shown to have relatively higher impact in open waters, while some stressors (e.g. nutrients,  
547 shipping, physical modification) had a relatively higher impact within fjord/estuarine systems.  
548 Some of these stressors (pressures) were considered as impacts in this study, e.g. non-  
549 indigenous [invasive] species. It was recognised that it can be difficult at times to distinguish  
550 between pressures and impacts. For example, shipping is a pressure and one of its impacts can  
551 be introduction of invasive species via ballast water, but these invasive species can themselves  
552 become a pressure on the native ecosystem.

### 553 **5.4. Monitoring programmes**

554 Most respondents were of the view that current monitoring is partially adequate or not  
555 adequate. The range of views given between and within countries suggest that a broad spectrum  
556 of participants responded to the questionnaire (Figure 12). These views likely reflect different  
557 experiences of respondents in their areas of expertise and in their countries.

558 Key issues identified in responses (i.e. insufficient resolution in time and space, insufficient  
559 data or parameters measured, and lack of integration) indicate the gaps in monitoring.  
560 Suggestions for improved monitoring programmes were targeted at these gaps and need to be  
561 considered in detail to feed into science and monitoring strategies. These issues are discussed  
562 in Section 5.5.

563 Few respondents completed the second questionnaire on monitoring programmes, so a subset  
564 of European monitoring programmes was reported. Opinions may therefore reflect the views  
565 of the JERICO community, particularly on the measurement of limited parameters (with a  
566 focus on physical and biogeochemical parameters, e.g. temperature, salinity and chlorophyll).  
567 These views are supported by information available via a number of projects and infrastructures  
568 (e.g. JERICO, DEVOTES, COPERNICUS, EMODnet, EMSO ERIC, and AtlantOS; links  
569 given in Section 1), all of which indicate the need to improve the availability of datasets,  
570 especially biological components (e.g. fish, seabirds and mammals). Furthermore, limited  
571 monitoring of pressures indicates some mismatch between the pressures and impacts  
572 considered by respondents to be important and those actually monitored.

573 Several programmes were reported to be making use of alternative platforms such as remote  
574 sensing, autonomous vehicles and FerryBoxes. These technologies are likely to complement  
575 other monitoring platforms (e.g. boat-based) rather than replace them altogether. Remote  
576 sensing data, for example, are limited to surface monitoring of particular parameters, and still  
577 require *in situ* data for calibration and validation (Groom et al 2019). FerryBox monitoring can  
578 improve coverage in space and time (e.g. Grayek et al 2011) but is similarly limited in terms  
579 of depth and parameters (Petersen 2014).

580 Suggestions given for improving monitoring programmes are supported by many studies on  
581 the development of existing and new technologies. Davidson et al (2019) provide an overview  
582 of the need for operational oceanographic systems, which include a multi-platform observation  
583 network, as well as systems for data management, data assimilation and prediction, and data  
584 dissemination/accessibility. Key components of such systems include an integrated approach  
585 (She et al 2019), partnerships and shared approaches for monitoring, assessment and data (Bax  
586 et al 2019; Canonico et al 2019; Míguez et al 2019; Schmidt et al 2019; Stammer et al 2019;  
587 Tanhua et al 2019; Weller et al 2019), instrumented moorings (fixed platforms; Bailey et al  
588 2019), and new methodologies for monitoring, including *in situ* biochemical, biological and  
589 molecular sensors (reviewed by Wang et al 2019).



## 590 5.5. Resolution in time and space

591 The scale of impacts varies widely, with some activities, such as construction of a wind farm  
592 having a potentially high impact on a small area, whereas activities such as fishing are more  
593 widespread. The impact of human activities also depends on the vulnerability of the habitat in  
594 question. For example, in the southern Celtic Sea, fragile benthic habitats with cold-water  
595 corals are highly impacted by sea floor activities. Some impacts, such as noise disturbance,  
596 depend on the intensity of the activity, and will be concentrated in areas with high shipping  
597 activity, or during periods of construction.

598 Countries such as the UK adopt a risk-based monitoring approach, which was considered to  
599 result in fragmented monitoring. Examples of low spatial resolution were given for the CPR  
600 survey, one of the key plankton datasets, where spatial gaps exist throughout EU waters. Spatial  
601 resolution was also considered to be low for some habitats, as not all habitats are covered by  
602 monitoring programmes, and for monitoring of marine litter and non-native species.

603 In terms of spatial resolution, other responses indicated that not all parameters are monitored  
604 adequately. Even for parameters that were reported as monitored in many monitoring  
605 programmes, e.g. chlorophyll (47% of reported programmes), monitoring may not be adequate  
606 in space or time (see Baretta-Bekker et al 2015, Annex 1<sup>13</sup>). A more detailed analysis looking  
607 at the distribution of monitoring of different parameters in space would be required to assess  
608 this. The WFD does not require zooplankton monitoring, but some indicators under the MSFD  
609 do require information on zooplankton. Although phytoplankton is monitored inshore, the data  
610 are disparate and mainly used to report on potential health issues due to toxin producing algae.

611 For temporal resolution, examples were given for a number of threats where the monitoring  
612 period was considered to be inadequate. For example, for statutory monitoring of impacts  
613 such as those from dredging and disposal, monitoring is often over time scales which are too  
614 short (2-5 years) to properly assess the impacts on the biological communities. This also  
615 applies to seabird and cetacean monitoring, which is out of the scope of JERICO-NEXT.

616 Some monitoring programmes may be inadequate in terms of temporal frequency: 24% of  
617 monitoring programmes reported had annual monitoring, which may fail to detect impacts  
618 throughout the year. Monitoring frequency is likely to be strongly influenced by platform  
619 types, with increasing use of fixed platforms, moorings or gliders giving a high proportion

---

<sup>13</sup> See also <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/pressures-human-activities/eutrophication/chlorophyll-concentrations/>

620 (34%, Figure 10) of responses for continuous monitoring. Certainly, platforms such as  
621 moorings can provide high-frequency temporal resolution (e.g. Mills et al 2005; Greenwood  
622 et al 2010) for the parameters they measure, predominantly physical and chemical parameters  
623 (such as temperature, salinity, light, dissolved oxygen) with biological parameters limited to  
624 phytoplankton fluorescence or chlorophyll. Monitoring more complex biological parameters  
625 (such as community species composition in either the benthic and the pelagic compartment)  
626 at high frequency appears particularly challenging because of the limited degree of  
627 development of appropriate semi-automatic tools. To date, routinely using such techniques at  
628 high frequency of acquisition would still require a massive level of skilled manpower,  
629 although new developments of molecular tools would clearly help to tackle the challenge in  
630 the future. Additionally, even where low cost sensors for biological parameters exist,  
631 analysing the large volumes of data produced remains a large challenge.

632 Addressing the issue of scales is essential in establishing a future pan-European monitoring  
633 program, particularly for biological parameters. Monitoring these parameters is more limited  
634 than for physical parameters. Reasons for this include that:

635 (1) The types of biological data that can be automatically or semi-automatically acquired is  
636 low despite recent technological developments (including those achieved within FP7-  
637 JERICO and JERICO-NEXT), which limits the spatio-temporal coverage of  
638 biological/biogeochemical data sets

639 (2) Miniaturization of sensors allowing for implementation on platforms such as AUVs and  
640 floaters is more feasible for physical and chemical parameters, which results in better  
641 spatial and synoptic coverage

642 (3) Scaling-up from “point” observations to wider areas most often relies on modelling.  
643 Physical models are more advanced than biogeochemical and biological models, which  
644 also increases the importance of scales of biological observations.

#### 645 **5.5.1. Small scale threats/disturbances**

646 The majority of threats impact at relatively small spatial and temporal scales, at least initially.  
647 Examples include the accumulation of marine litter, the development of harmful algal blooms,  
648 and the invasion by non-native species, which occur locally in the first instance, as influenced  
649 by point sources and the characteristics of the abiotic and biotic components of the  
650 environment. In these examples, there is no initial discrepancy in spatial scales between

651 monitoring and threats/disturbances. However, the number of monitored habitats clearly  
652 remains too low, as indicated by responses to the questionnaire.

653 Monitoring effort should be sufficient in time and space to: (1) detect the effects of new  
654 threats/disturbances acting in different locations within the same habitat, (2) assess the  
655 consequences of an identified threat/disturbance at larger scales, and (3) assess cumulative  
656 effects of multiple threats.

### 657 **5.5.2. Large-scale threats/disturbances**

658 Some environmental threats act over large spatial scales, such as thermal regime change or  
659 ocean acidification. There is a discrepancy between the (large) spatial extent of the  
660 threat/disturbance and the (small) scale at which the monitoring is performed (station). This  
661 may be addressed to some extent by (1) the use of mobile monitoring techniques such as  
662 FerryBoxes which allow for large geographical coverage, albeit on a limited time-scale, and  
663 (2) the fact that only a small number of fixed monitored sites is required to monitor this kind  
664 of threat disturbance. Factors to consider include that:

665 (i) Different biological communities may not be affected in the same way by the same level  
666 of a given (widespread) environmental pressure. Grémare et al (1998) and Labrunne et al  
667 (2007), for example, clearly showed that in the Gulf of Lion, the composition of the two  
668 shallowest communities (i.e. littoral fine sands and littoral sandy muds) are most affected  
669 by climatic oscillations. A sound assessment of large-scale threats/disturbances at the  
670 reporting scales should therefore not rely on the sampling of a single, or even a limited  
671 number of habitats. Conversely, the monitoring strategy of large-scale  
672 threats/disturbances should ideally encompass all the habitats present in the reporting  
673 area.

674 (ii) The representativeness of monitoring data is often limited. For example, highly mobile  
675 fauna (e.g. marine mammals or birds) are often used as proxies for large scale  
676 threats/disturbances because they can be found over large spatial scales and because, as  
677 for predators, their ecophysiology and/or population dynamics tolerate a large set of  
678 ecological processes. The probability of these organisms being sampled with confidence  
679 is directly proportional to the sampling effort and to their relative accessibility. Current  
680 monitoring resources currently deployed in the UK, for example, do not have the power  
681 to detect trends in all seabird and cetacean species or identify the drivers of their  
682 population change. A similar example was given for Malta, where only the most

683 accessible marine bird nests are currently monitored as part of the seabird monitoring  
684 program.

### 685 **5.5.3. The real world: a mixture of threats/disturbances at small and larger scales**

686 At the scale of global coastal marine ecosystems, several environmental pressures act  
687 simultaneously, each having its own spatial resolution and temporal dynamics. Halpern et al  
688 (2008) and Crain et al (2009) found that no fewer than five pressures overlap anywhere in the  
689 world's oceans. Potential cumulative and/or interactive effects need to be addressed, for  
690 example by considering that:

691 (i) Monitoring should be based on the largest spatial entity within which the comparisons  
692 of community compositions are sound, e.g. habitats or ecohydrodynamic regions (van  
693 Leeuwen et al 2015).

694 (ii) The monitoring of each habitat or region should include a sample size large enough to  
695 allow for the detection and the variability in the effects of small- and large-scale  
696 threats/disturbances.

697 (iii) Within a given reporting area, a monitoring program should include the highest possible  
698 number of relevant habitats in order to facilitate the detection of new small-scale  
699 threat/disturbance and the upscaling of large-scale threat/disturbance effects.

700 Such monitoring programmes would require considerable effort, highlighting the need to  
701 define/characterize relevant environmental threats in each habitat or region.

702 The feasibility of the different suggestions for improved monitoring needs to be considered.  
703 This includes the identification of 'new technologies' and how best to incorporate them into  
704 monitoring programmes. Projects such as JERICO-NEXT work to harmonise new technologies  
705 which may be able to solve some problems of scale through high-frequency monitoring. For  
706 example, instruments such as flow cytometers and multispectral fluorometers can measure  
707 continuously on research vessels or buoys and so provide good spatial and temporal coverage.  
708 However, integrating these data types into existing monitoring presents several challenges: data  
709 may be in a very different format (continuous versus discrete samples, functional groups vs  
710 taxa), adopting new methods may affect the integrity of long time series, or there may be  
711 difficulty gaining acceptance and confidence in new methods. Similar challenges exist with  
712 using remotely sensed data instead of field measurements (e.g. for turbidity, chlorophyll), and  
713 these also still requires ongoing in situ measurements for validation (De Cauwer et al 2004).

## 714     **6.     Conclusions**

715     This study consolidates the main conclusions from the Dobris Assessment (EEA 1995) and  
716     more recent studies (e.g. EEA 2008a, b; EEA 2015a; DEVOTES; Tett et al 2013; Zampoukos  
717     et al 2013; Garcia-Garcia et al 2019), highlighting the need for improved monitoring of  
718     environmental threats in European coastal environment.

719     Most respondents to the JERICO-NEXT questionnaire considered current monitoring of threats  
720     to be partially adequate or not adequate. The majority of responses were related to spatial and/or  
721     temporal scales at which monitoring takes place, and inadequate monitoring of particular  
722     parameters. Monitoring of biological parameters was considered to be generally inadequate,  
723     with insufficient focus on coupling between biological and physical or chemical parameters

724     Suggestions for improved monitoring programmes included improved design, increased  
725     monitoring effort, better linkages with research, better use of new technologies (such as remote  
726     sensing, FerryBoxes, and gliders) and methods (such as molecular techniques), and improved  
727     data flows. Improved monitoring programmes should be fit for policy and management  
728     purposes, as well as underpin longer-term scientific objectives which cut across policy and  
729     other drivers. Improved designs of monitoring programmes need to consider cumulative effects  
730     of multiple pressures. The JERICO-RI has high potential to fill in some of the observation gaps,  
731     especially those related to physical and biogeochemical parameters, and the coupling between  
732     biology and physics across scales needed for integrated ecosystem-based understanding. The  
733     particular challenge of simultaneously observing physical, chemical and biological parameters  
734     for assessments of complex coastal processes remains an open issue in relation to the temporal  
735     scale of sampling. This will be addressed in the JERICO science strategy under development  
736     (Grémare et al 2017; Farcy et al 2019).

737     Certainly, one of the main challenges for the European marine research community is to  
738     increase the consistency and the sustainability of dispersed networks and infrastructures by  
739     integrating them within a shared pan-European framework. The long history of national  
740     monitoring programmes which have been expanded, modified and developed over time,  
741     together with methodological differences between nations, results in difficulties for the  
742     integration and holistic assessment of the data (at a regional sea level) which the JERICO-RI  
743     may contribute towards solving.

744

745 **7. Acknowledgements**

746 This project has received funding from the European Union’s Horizon 2020 research and  
747 innovation programme under grant agreement No. 654410.

748

749 **8. Author contributions**

750 SP, KC, DD, AG and GB designed the questionnaire. KC downloaded and analysed the results.

751 SP and KC prepared the manuscript with contributions from all co-authors.

752

753 **9. Competing interests**

754 Six of the authors declare that they have no conflict of interest. VC is a member of the editorial  
755 board on other topic areas in the Special Issue.

756

757 **10. References**

758 Andersen JH, Al-Hamdani Z, Harvey ET, et al Relative impacts of multiple human stressors  
759 in estuaries and coastal waters in the North Sea – Baltic Sea transition zone. *Sci Total*  
760 *Environ* 42 pp. doi: 10.1016/j.scitotenv.2019.135316, 2019.

761 Bailey K, Steinberg C, Davies C, et al: Coastal mooring observing networks and their data  
762 products: Recommendations for the next decade. *Front Mar Sci* 6:1–22. doi:  
763 10.3389/fmars.2019.00180, 2019.

764 Baretta-Bekker, Hanneke, Anne Sell, Francisco Marco-Rius, Julia Wischniewski, Pamela  
765 Walsham, Lynda, Malin Mohlin, Karin Wesslander, Hans Ruiters, Francis Gohin, Lisette En-  
766 serink: The chlorophyll case study in the JMP NS/CS project. Document produced as part of  
767 the EU project: ‘Towards joint Monitoring for the North Sea and Celtic Sea’ (Ref: ENV/PP  
768 2012/SEA). 72 pp. Annex I: Baretta-Bekker, Hanneke. Inventory of the Chlorophyll-a data in  
769 the ICES database. 23 pp, 2015.

770 Bax NJ, Miloslavich P, Muller-Karger FE, et al: A response to scientific and societal needs  
771 for marine biological observations. *Front Mar Sci* 6:1–22. doi: 10.3389/fmars.2019.00395,  
772 2019.

773 Bean, T. P., Greenwood, N., Beckett, R., Biermann, L., Bignell, J. P., Brant, J. L., Copp, G.  
774 H., Devlin, M. J., Dye, S., Feist, S. W., Fernand, L., Foden, D., Hyder, K., Jenkins, C. M.,  
775 van der Kooij, J., Kröger, S., Kupschus, S., Leech, C., Leonard, K. S., Lynam, C. P., Lyons,  
776 B. P., Maes, T., Nicolaus, E. E. M., Malcolm, S. J., McIlwaine, P., Merchant, N. D.,  
777 Paltriguera, L., Pearce, D. J., Pitois, S. G., Stebbing, P. D., Townhill, B., Ware, S., Williams,  
778 O. and Righton, D.: A review of the tools used for marine monitoring in the UK: combining  
779 historic and contemporary methods with modeling and socioeconomics to fulfill legislative  
780 needs and scientific ambitions, *Front. Mar. Sci.*, 4(263), 1–29,  
781 doi:10.3389/fmars.2017.00263, 2017.

782 Benedetti-Cecchi, L., Crowe, T., Boero, F., Christense, A., Gremare, A., Hernandez, F.,  
783 Krompamp, J., et al, Crowe, T., Boehme, L., Boero, F., Christensen, A., Grémare, A.,

784 Hernandez, F., Kromkamp, J. C., Nogueira García, E., Petihakis, G., Robidart, J., Sousa  
785 Pinto, I. and Zingone, A.: Strengthening Europe's Capability in Biological Ocean  
786 Observations, in Future Science Brief 3 of the European Marine Board, Ostend, Belgium,  
787 edited by A. Muniz Piniella, P. Kellett, K. Larkin, and J. Heymans, p. 76., 2018.

788 Borja, A., Garmendia, J. M., Menchaca, I., Uriarte, A. and Sagarmínaga, Y.: Yes, We Can!  
789 Large-Scale Integrative Assessment of European Regional Seas, using Open Access  
790 Databases, *Front. Mar. Sci.*, 6(February), 1–13, doi:10.3389/fmars.2019.00019, 2019.

791 Buck, J. J. H., Bainbridge, S. J., Burger, E. F., Kraberg, A. C., Casari, M., Casey, K. S.,  
792 Darroch, L., Rio, J. Del, Metfies, K., Delory, E., Fischer, P. F., Gardner, T., Heffernan, R.,  
793 Jirka, S., Kokkinaki, A., Loeb, M., Buttigieg, P. L., Pearlman, J. S. and Schewe, I.: Ocean  
794 Data Product Integration Through Innovation-The Next Level of Data Interoperability, *Front.*  
795 *Mar. Sci.*, 6, doi:10.3389/fmars.2019.00032, 2019.

796 Canonico G, Buttigieg PL, Montes E, et al: Global observational needs and resources for  
797 marine biodiversity. *Front Mar Sci* 6:1–20. doi: 10.3389/fmars.2019.00367, 2019.

798 Crain, C., Halpern, B., Beck, M. and Kappel, C.: Understanding and Managing Human  
799 Threats to the Coastal Marine Environment. *The Year in Ecology and Conservation Biology*,  
800 2009, *Ann. N.Y. Acad. Sci.*, 1162, 39–62, 2009.

801 Davidson FJ, Chassignet E, Vinayachandran PN, et al: Synergies in operational  
802 oceanography: The intrinsic need for sustained ocean observations. *Front Mar Sci* 6:1–18.  
803 doi: 10.3389/fmars.2019.00450, 2019.

804 DEVOTES: Catalogue of Monitoring Networks. [http://www.devotes-project.eu/devotes-  
805 release-new-version-catalogue-monitoring-networks/](http://www.devotes-project.eu/devotes-release-new-version-catalogue-monitoring-networks/), 2014.

806 De Cauwer, V., Ruddick, K., Park, Y., Nechad, B. and Kyramarios, M.: Optical remote  
807 sensing in support of eutrophication monitoring in the Southern North Sea, *EARSel*  
808 *eProceedings*, (3), 208–221 [online] Available from:  
809 <http://www.vliz.be/imis/imis.php?module=ref&refid=67132>, 2004.

810 Duarte, C. M., Poiner, I. and Gunn, J.: Perspectives on a Global Observing System to Assess  
811 Ocean Health, *Front. Mar. Sci.*, 5(August), 265, doi:10.3389/fmars.2018.00265, 2018.

812 EC: Council Directive of 12 December 1991 concerning the protection of waters against  
813 pollution caused by nitrates from agricultural sources (91/676/EEC), *Off. J. Eur.*  
814 *Communities*, L 375(1259), 1–8, 1991.

815 EC: Directive of 21 May 1991 concerning urban waste water treatment (91/271/EEC)., *Off. J.*  
816 *Eur. Communities*, L 135, 40–52 [online] Available from: [http://eur-lex.europa.eu/legal-  
817 content/EN/ALL/?uri=CELEX:31991L0271](http://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:31991L0271), 1991.

818 EEA: Europe's Environment - The Dobris Assessment. State of the environment report No  
819 1/1995. [online] Available from: <http://www.eea.europa.eu/publications/92-826-5409-5>,  
820 1995.

821 EEA: Europe's Environment - The Dobris Assessment. Chapter 35 Coastal Zone Threats and  
822 Management. State of the environment report., 6 pp [online] Available from:  
823 [www.eea.europa.eu/publications/92-826-5409-5/page035new.html](http://www.eea.europa.eu/publications/92-826-5409-5/page035new.html), 2008a.

824 EEA: Europe's Environment. The Fourth Assessment. Chapter 5 Marine and coastal  
825 environment. <http://www.eea.europa.eu/publications/92-826-5409-5>, 44 pp, 2008b.

826 EEA: Europe's environment. An Assessment of Assessments. [online] Available from:  
827 <http://www.eea.europa.eu/themes/regions/pan-european/sub-regional-assessment-of->  
828 [assessment-reports/eastern-europe-assessment-of-assessment-report](http://www.eea.europa.eu/themes/regions/pan-european/sub-regional-assessment-of-), 2011.

829 EEA: The European environment - state and outlook 2015: synthesis report, European  
830 Environment Agency, Copenhagen, 212 pp, 2015a.

831 EEA: European Ecosystem Assessment – Concept, Data and Implementation. Technical  
832 report No6/2015, 74 pp, 2015b.

833 Elliott, M.: Integrated marine science and management: wading through the morass., *Mar.*  
834 *Pollut. Bull.*, 86(1–2), 1–4, doi:10.1016/j.marpolbul.2014.07.026, 2014.

835 EU: Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000  
836 establishing a framework for community action in the field of water policy., *Off. J. Eur.*  
837 *Communities*, 327, 1–72 [online] Available from: [http://eur-lex.europa.eu/legal-](http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex:32000L0060)  
838 [content/EN/TXT/?uri=celex:32000L0060](http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex:32000L0060), 2000.

839 EU: Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008  
840 establishing a framework for community action in the field of marine environmental policy  
841 (Marine Strategy Framework Directive)., *Off. J. Eur. Union*, 164, 19–40 [online] Available  
842 from: <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32008L0056>, 2008.

843 EuroGOOS: Sea level observation networks in and around Europe Challenges. in monitoring  
844 increasing sea level hazards. Note to policymakers. [www.eurogoos.eu](http://www.eurogoos.eu), (April), 4 pp [online]  
845 Available from: [http://eurogoos.eu/download/publications/EuroGOOS-Tide-Gauge-Note-to-](http://eurogoos.eu/download/publications/EuroGOOS-Tide-Gauge-Note-to-Policy-2017-v.2.pdf)  
846 [Policy-2017-v.2.pdf](http://eurogoos.eu/download/publications/EuroGOOS-Tide-Gauge-Note-to-Policy-2017-v.2.pdf), 2017.

847 Farcy, P., Durand, D., Charria, G., Painting, S., Collingridge, K., Gremare, A., Puillat, I. and  
848 Delauney, L.: Towards a European Coastal Observing Network to provide better answer to  
849 science and to societal challenges; the JERICO Research Infrastructure, *Front Mar Sci* 6:1–  
850 13. doi: 10.3389/fmars.2019.00529., 2019.

851 Gabrielsen, P. and Bosch, P.: Environmental Indicators: Typology and Use in Reporting.  
852 EEA internal working paper., 2003.

853 García-García, L., van der Molen, J., Sivyver, D., Devlin, M., Painting, S. and Collingridge,  
854 K.: Optimizing monitoring programs: a case study based on the OSPAR Eutrophication  
855 assessment for UK waters, *Front. Mar. Sci.*, 19 pp, doi:10.3389/fmars.2018.00503, 2019.

856 Grand, M. M., Laes-Huon, A. and Al, E.: Developing autonomous observing systems for  
857 micronutrient trace metals, *Front. Mar. Sci.*, doi:10.3389/fmars.2019.00035, 2019.

858 Grayek S, Staneva J, Schulz-Stellenfleth J, et al: Use of FerryBox surface temperature and sa-  
859 linity measurements to improve model based state estimates for the German Bight. *J Mar*  
860 *Syst* 88:45–59. doi: <https://doi.org/10.1016/j.jmarsys.2011.02.020>, 2011.

861 Greenwood, N., Parker, E. R., Fernand, L., Sivyver, D. B., Weston, K., Painting, S. J., Kröger,  
862 S., Forster, R. M., Lees, H. E., Mills, D. K., Laane, R.W.P.M.: Detection of low bottom water  
863 oxygen concentrations in the North Sea; implications for monitoring and assessment of  
864 ecosystem health., *Biogeosciences*, 7(4), 1357–1373, 2010.

865 Grémare, A., Amouroux, J. and Vétion, G.: Long-term comparison of macrobenthos within  
866 the soft bottoms of the Bay of Banyuls-sur-mer (northern Mediterranean Sea), *J. Sea Res.*,  
867 281–302, 1998.

868 Grémare A, Durand D, Delauney L, et al: The JERICO-NEXT Science Strategy. JERICO-  
869 NEXT-WP1-D1.2-050717-V4.0, 2017.



870 Groom SB, Sathyendranath S, Ban Y, et al: Satellite ocean colour: Current status and future  
871 perspective. *Front Mar Sci*. doi: 10.3389/fmars.2019.00485, 2019.

872 Halpern, B., Walbridge, S., Selkoe, K., Kappel, C., Micheli, F., D'Agrosa, C., Bruno, J.,  
873 Casey, K., Ebert, C., Fox, H., Fujita, R., Heinemann, D., Lenihan, H., Madin, E., Perry, M.,  
874 Selig, E., Spalding, M., Steneck, R. and R, W.: A global map of human impact on marine  
875 ecosystems, *Science* (319 ), 948–952, 2008.

876 HELCOM: Baltic Marine Environment Protection Commission Baltic Sea Environment  
877 Proceedings 155 Baltic Sea-Second HELCOM holistic, [online] Available from:  
878 [www.helcom.fi/baltic-sea-trends/holistic-assessments/state-of-the-baltic-sea-2018/reports-](http://www.helcom.fi/baltic-sea-trends/holistic-assessments/state-of-the-baltic-sea-2018/reports-and-materials/)  
879 [and-materials/](http://www.helcom.fi/baltic-sea-trends/holistic-assessments/state-of-the-baltic-sea-2018/reports-and-materials/), 2018.

880 Labrune, C., Grémare, A., Guizien, K. and Amouroux, J.: Long-term comparison of soft  
881 bottom macrobenthos in the Bay of Banyuls-sur-Mer (north-western Mediterranean Sea): A  
882 reappraisal, *J. Sea Res.*, 58, 125–143, 2007.

883 Maximenko N, Corradi P, Law KL, et al: Towards the integrated marine debris observing  
884 system. *Front Mar Sci*. doi: 10.3389/fmars.2019.00447, 2019.

885 Míguez BM, Novellino A, Vinci M, et al: The European Marine Observation and Data  
886 Network (EMODnet): Visions and roles of the gateway to marine data in Europe. *Front Mar*  
887 *Sci* 6:1–24. doi: 10.3389/fmars.2019.00313, 2019.

888 Mills, D. K., Greenwood, N., Kröger, S., Devlin, M., Sivyer, D. B., Pearce, D. and Malcolm,  
889 S. J.: New Approaches to Improve the Detection of Eutrophication in UK Coastal Waters,  
890 2(2), 36–42, 2005.

891 Oesterwind, D., Rau, A. and Zaiko, A.: Drivers and pressures - Untangling the terms  
892 commonly used in marine science and policy, *J. Environ. Manage.*, 181, 8–15,  
893 doi:10.1016/j.jenvman.2016.05.058, 2016.

894 OSPAR: Eutrophication Status of the OSPAR Maritime Area Third Integrated Report on the  
895 Eutrophication Status of the OSPAR Maritime Area Eutrophication Series. See  
896 <https://www.ospar.org/work-areas/hasec/eutrophication/common-procedure>, 2017.

897 Palazov A, Ciliberti S, Peneva E, et al: Black sea observing system. *Front Mar Sci* 6:1–8. doi:  
898 10.3389/fmars.2019.00315, 2019.

899 Proctor, R. and Howarth, M. J.: Coastal observatories and operational oceanography: a  
900 European perspective, *Mar. Technol. Soc. J.*, 43, 10–13, doi:10.4031/002533208786842534,  
901 2008.

902 Petersen W: FerryBox systems: State-of-the-art in Europe and future development. *J Mar*  
903 *Syst* 140:4–12, 2014.

904 Schmidt JO, Bograd SJ, Arrizabalaga H, et al: Future Ocean Observations to Connect  
905 Climate, Fisheries and Marine Ecosystems. *Front Mar Sci* 6:1–18. doi:  
906 10.3389/fmars.2019.00550, 2019.

907 Schofield, O., Bergmann, T., Bissett, P., Grassle, J. F., Haidvogel, D. B., Kohut, J., Moline,  
908 M. and Glenn, S. M.: The Long-Term Ecosystem Observatory: An Integrated Coastal  
909 Observatory, *IEEE J. Ocean. Eng.*, 27(2), 146–154, 2002.

910 Schofield, O., Glenn, S., Bissett, P. W., Frazer, T. K., Iglesias-Rodriguez, D. and Moline, M.  
911 a.: Development of Regional Coastal Ocean Observatories and the Potential Benefits to  
912 Marine Sanctuaries, *Mar. Technol. Soc. J.*, 37(1), 54–67, doi:10.4031/002533203787537456,  
913 2003.

914 She, J. and Al, E.: Assessing biogeochemical monitoring networks in regional seas. Fact  
915 Sheet 2. Operational Ecology (OPEC) Marine Ecosystem Forecasting, 2 pp [online]  
916 Available from: [http://marine-opec.eu/factsheets/FS2\\_Monitoring.pdf](http://marine-opec.eu/factsheets/FS2_Monitoring.pdf), 2014.

917 She J, Piniella ÁM, Benedetti-Cecchi L, et al: An integrated approach to coastal and  
918 biological observations. *Front Mar Sci* 6:1–6. doi: 10.3389/fmars.2019.00314, 2019

919 Smith, L. M., Yarincik, K., Vaccari, L., Kaplan, M. B., Barth, J. A., Cram, G. S., Fram, J. P.,  
920 Harrington, M., Kawka, O. E., Kelley, D. S., Matthias, P., Newhall, K., Palanza, M.,  
921 Plueddemann, A. J., Vardaro, M. F., White, S. N. and Weller, R. A.: Lessons Learned From  
922 the United States Ocean Observatories Initiative, 5 (January), 1–7,  
923 doi:10.3389/fmars.2018.00494, 2019a.

924 Smith, N., Kessler, W. S. and et al: Tropical Pacific Observing Systems, *Front. Mar. Sci.*,  
925 doi:10.3389/fmars.2019.00031, 2019b.

926 Stammer D, Bracco A, AchutaRao K, et al: Ocean climate observing requirements in support  
927 of Climate Research and Climate Information. *Front Mar Sci* 6:1–18. doi:  
928 10.3389/fmars.2019.00444, 2019.

929 Tanhua T, Pouliquen S, Hausman J, et al: Ocean FAIR data services. *Front Mar Sci*. doi:  
930 10.3389/fmars.2019.00440, 2019.

931 Tett, P., Gowen, R., Painting, S., Elliott, M., Forster, R., Mills, D., Bresnan, E., Capuzzo, E.,  
932 Fernandes, T., Foden, J., Geider, R., Gilpin, L., Huxham, M., McQuatters-Gollop, A.,  
933 Malcolm, S., Saux-Picart, S., Platt, T., Racault, M.-F., Sathyendranath, S., van der Molen, J.  
934 and Wilkinson, M.: Framework for understanding marine ecosystem health, *Mar Ecol Prog*  
935 *Ser*, 494, 1–27, doi:10.3354/meps10539, 2013.

936 Tintoré J, Pinardi N, Álvarez-Fanjul E, et al: Challenges for Sustained Observing and  
937 Forecasting Systems in the Mediterranean Sea. *Front Mar Sci*. doi:  
938 10.3389/fmars.2019.00568, 2019.

939 UNEP/UNECE: GEO-6 assessment for the Pan-European region. United Nations  
940 Environment Programme, Nairobi, Kenya. 376 pp., 2016.

941 van Leeuwen, S., Tett, P., Mills, D. and van der Molen, J.: Stratified and non-stratified areas  
942 in the North Sea: Long-term variability and biological and policy implications, *J. Geophys.*  
943 *Res. Ocean.*, 120, 4670–4686, doi:10.1002/2014JC010485, 2015.

944 Wang ZA, Moustahfid H, Mueller A V., et al:Advancing Observation of Ocean  
945 Biogeochemistry, Biology, and Ecosystems With Cost-Effective in situ Sensing  
946 Technologies. *Front Mar Sci* 6:1–22. doi: 10.3389/fmars.2019.00519, 2019.

947 Weller RA, Baker DJ, Glackin MM, et al:The challenge of sustaining ocean observations.  
948 *Front Mar Sci* 6:1–18. doi: 10.3389/fmars.2019.00105, 2019.

949 Wordle: Viewed 07/06/2018. Available online at: <https://worditout.com/word-cloud/create>,  
950 2018.

951 Zampoukas, N., Piha, H., Bigagli, E., Hoepffner, N., Hanke, G. and Cardoso, A. C.: Marine  
952 monitoring in the European Union: How to fulfill the requirements for the marine strategy  
953 framework directive in an efficient and integrated way, *Mar. Policy*, 39(May 1992), 349–351,  
954 doi:10.1016/j.marpol.2012.12.004, 2013.

