

1 Seasonal variation of the sound-scattering zooplankton vertical 2 distribution in the oxygen-deficient waters of the NE Black Sea

3 Alexander G. Ostrovskii¹, Elena G. Arashkevich¹, Vladimir A. Solovyev¹, Dmitry A. Shvov¹

4 ¹ Shirshov Institute of Oceanology, Russian Academy of Sciences, 36, Nahimovskiy prospekt, Moscow, Russia, 117997

5 *Correspondence to:* Alexander G. Ostrovskii (osasha@ocean.ru)

6 **Abstract.** At the northeastern Black Sea research site, observations from 2010-2020 allowed us to study the dynamics and
7 evolution of the vertical distribution of mesozooplankton in oxygen-deficient conditions via analysis of sound-scattering
8 layers associated with dominant zooplankton aggregations. The data were obtained with profiler mooring and zooplankton
9 net sampling. The profiler was equipped with an acoustic Doppler current meter, a conductivity-temperature-depth probe,
10 and fast sensors for the concentration of dissolved oxygen [O₂]. The acoustic instrument conducted ultrasound (2 MHz)
11 backscatter measurements at 3 angles while being carried by the profiler through the oxic zone. For the lower part of the
12 oxycline and the hypoxic zone, the normalized data of 3 acoustic beams (directional acoustic backscatter ratios, *R*) indicated
13 sound-scattering mesozooplankton aggregations, which were defined by zooplankton taxonomic and quantitative
14 characteristics based on stratified net sampling at the mooring site. The time series of ~14,000 *R*-profiles as a function of
15 [O₂] at depths where [O₂] < 200 μM were analyzed to determine month-to-month variations of the sound-scattering layers.
16 From spring to early autumn, there were two sound-scattering maxima corresponding to (1) daytime aggregations mainly
17 formed by diel-vertical-migrating copepods *Calanus euxinus* and *Pseudocalanus elongatus* and chaetognaths *Parasagitta*
18 *setosa*, usually at [O₂] = 15-100 μM, and (2) persistent monospecific layer of the diapausing fifth copepodite stages of *C.*
19 *euxinus* in the suboxic zone at 3 μM < [O₂] < 10 μM. From late autumn to early winter, no persistent deep sound-scattering
20 layer was observed. At the end of winter, the acoustic backscatter was basically uniform in the lower part of the oxycline and
21 the hypoxic zone. The assessment of the seasonal variability of the sound-scattering mesozooplankton layers is important for
22 understanding biogeochemical processes in oxygen-deficient waters.

23 1 Introduction

24 The main distinguishing feature of the Black Sea environment is its oxygen stratification with an oxygenated upper layer 80-
25 200 m thick and the underlying waters containing hydrogen sulfide (Andrusov, 1890; see also review by Oguz et al., 2006).
26 Early studies of the oxic zone indicated that the vertical distribution of zooplankton hinges on oxygen stratification (Nikitin,
27 1926; Petipa et al., 1960). Later, the dives of the manned research submersible *Argus* showed that the zooplankton vertical
28 distribution was not uniform (Vinogradov et al. 1985; Flint 1989). In particular, the thin-layered structure of zooplankton
29 distribution was observed by the *Argus* research pilot in the lower part of the oxic zone. Thereafter, zooplankton sampling

30 with a vertical resolution of 3-5 m using a 150-liter sampler with an attached conductivity-temperature-depth (CTD) probe
31 indicated that the daytime deep aggregations of the zooplankton populations were associated with layers of certain water
32 density (Vinogradov and Nalbandov, 1990; Vinogradov et al., 1992). The deeper zooplankton aggregation was formed by
33 the fifth copepodite stage of *Calanus ponticus* (old name of *C. euxinus*) and its lower boundary was at the specific density
34 surface $\sigma_\theta = 15.9$, where the oxygen concentration was approximately 4 μM . The diapausing cohort of *C. ponticus* did not
35 perform vertical migrations and occupied the suboxic layer around the clock (Vinogradov et al., 1992). The accumulation of
36 a high lipid reserve, a decrease in the rate of oxygen consumption, and a delay in gonad development were defined as
37 characteristic features of diapausing *C. euxinus* (Vinogradov et al., 1992; Arashkevich et al., 1998; Svetlichny et al., 2002,
38 2006). The vertically migrating zooplankters (ctenophores *Pleurobrachia pileus*, chaetognaths *Parasagitta setosa*, and older
39 copepodites of *Pseudocalanus elongatus* and *C. euxinus*) formed daytime aggregations between isopycnals 15.7-15.5 and
40 15.4-14.9 and at an oxygen concentration of 11-40 μM . At night, the migrant zooplankters inhabited the upper layers and
41 peaked in the thermocline (Vinogradov et al., 1985). The descent of zooplankters into the hypoxic zone during the daytime
42 may give an energetic advantage to migrating specimens due to a decrease in the rate of oxygen consumption and locomotor
43 activity at low oxygen concentrations, as has been shown for females of *C. euxinus* (Svetlichny et al., 2000). This and other
44 experimental studies contributed to the development of an optimal behavioral strategy model (Morozov et al., 2019) for
45 structured populations of two species, *C. euxinus* and *P. elongatus*. The authors parameterized the model using seasonal field
46 observations in the NE Black Sea and showed that the diel vertical migrations of these species could be explained as the
47 result of a trade-off between depth-dependent metabolic costs, anoxia, available food, and predation.

48 Zooplankton aggregations result in sound-scattering layers (SSLs). Diel vertical migration was observed using ship
49 echo sounding at frequencies of 120 - 200 kHz (Erkan and Gücü, 1998; Mutlu, 2003, 2006, 2007; Stefanova and Marinova,
50 2015). The diurnal dynamics of *C. euxinus* and chaetognaths were documented from ship-borne echograms (Mutlu 2003,
51 2006). The lower boundary of the migrating *C. euxinus* was defined as $\sigma_\theta = 16.15$ -16.2 for the daytime, and the migrating
52 chaetognaths were defined as $\sigma_\theta = 15.9$ -16.0 (Mutlu 2007). In July 2013, a multifrequency (38, 120, and 200 kHz) ship-
53 borne echo-sounder survey over the southern Black Sea revealed that the daytime deep distribution of migrating *C. euxinus*
54 was bounded by σ_θ values between 15.2 and 15.9 (Sakınan and Gücü, 2016). In the above studies, the persistent layer of
55 diapausing *C. euxinus* was not detected in the echograms.

56 The 24-h rhythm in the pattern of sound scattering was a prominent feature of the 2 MHz acoustic sensing data
57 obtained by a moored profiler station (Ostrovskii and Zatsepin, 2011) in the NE Black Sea. The data obtained by a short (up
58 to 10 days) experimental deployment of a moored automatic mobile profiler equipped with an ultrasound probe operating at
59 a frequency of 2 MHz and a dissolved oxygen sensor allowed Ostrovskii and Zatsepin (2011) to define the main sound-
60 scattering zones as follows:

61 - the hydrogen sulfide zone below the specific density surface $\sigma_\theta = 15.9$ -16.0 (Yakushev et al., 2005), where sound is
62 scattered by sedimented detritus and mineral particles, whose fluxes vary temporally while being rather homogeneous at
63 different depths,

64 - above the hydrogen sulfide zone in the suboxic layer (where the concentration of dissolved oxygen [O₂] < 10 μM
65 (Murray et al., 1989, Oguz et al., 2006) and above that, in the oxycline ([O₂] increases from 10 μM to 280-300 μM with
66 decreasing depth), where sound scattering occurs from both suspended particles and mesozooplankton with characteristic
67 sizes from 200 microns to 20 mm,

68 - above the oxycline in the oxygen-rich euphotic zone, where large cell phytoplankton (Yunev et al., 2020) become an
69 additional sound-scattering agent.

70 Using a combination of ultrasound sensing and stratified zooplankton sampling was necessary to resolve the ocean
71 fine-scale vertical distribution of mesozooplankton. An analysis of both echograms and simultaneous stratified net sampling
72 showed that the SSLs at 2 MHz were associated with the zooplankton species *C. euxinus* and *P. elongatus* at $\sigma_{\theta} = 15.7-15.4$
73 and diapausing *C. euxinus* above $\sigma_{\theta} = 15.9$ (Arashkevich et al., 2013).


74 The specific theme of this study is the seasonal change in the sound-scattering zooplankton vertical distribution across
75 the oxygen gradient from the lower part of oxygenated water to the anoxic zone boundary. This theme is in line with the EU
76 Horizon 2020 BRIDGE-BS project (<http://www.unsdsn.gr/h2020-bridge-bs>), which focuses on Black Sea ecosystem
77 functioning. While the project relies on future observations and methods for understanding biogeochemical processes at
78 several pilot sites, this paper presents ongoing observations at the northeastern Black Sea Gelendzhik site. The acoustic data
79 were collected year round and analyzed to infer the SSL seasonal variability in relation to the oxygen stratification. Our
80 observational study was made possible using a moored Aqualog profiler equipped with an ultrasound probe, a CTD probe,
81 and a fast oxygen sensor. The advantage of this approach is that it provides frequent year-round measurements (with an
82 interval of up to 1 h) of collocated vertical profiles of sound scattering, temperature, salinity, and oxygen concentration in the
83 water column from the near-surface to the bottom layer with a high vertical resolution (up to 20 cm). This helps to fill in the
84 gaps due to insufficient zooplankton sampling in the winter season and resolves difficulties with sampling at precise depths,
85 thereby providing the information needed to define the displacements of the mesozooplankton aggregations.

86 The goals of the analysis are as follows: (1) to develop methods to visualize the SSLs in the lower part of the oxycline
87 and in the hypoxic zone, (2) to validate the SSLs in the oxygen-deficient waters using the taxonomic and quantitative
88 characteristics of zooplankton vertical distribution derived from stratified net sampling, and (3) to describe the seasonal
89 variations of the deep mesozooplankton SSLs, including the diapause duration of CV *C. euxinus*, in relation to oxygen
90 concentration (the oxygen bounds for the mesozooplankton SSLs).

91 **2 Measurements**

92 This study is based on the comparative analysis of the amplitude of sound backscattering data at a frequency of 2 MHz and
93 oxygen concentration data in seawater obtained in the NE Black Sea using a moored automatic mobile profiler Aqualog (Fig.
94 1) (Ostrovskii and Zatsepin, 2011, 2016; Ostrovsky et al., 2013). To obtain the depth profiles of the volume backscattering

95 strength, the Aqualog profiler was equipped with a Nortek Aquadopp acoustic Doppler current meter
96 (https://www.nortekgroup.com/assets/documents/ComprehensiveManual_Oct2017_compressed.pdf).



97

98 **Figure 1: The moored Aqualog automatic mobile profiler with a deep-water Aquadopp acoustic Doppler current meter (left).**
99 **Transducer head of the shallow-water Aquadopp acoustic Doppler current meter on the Aqualog profiler (right). Bottom right:**
100 **the acoustic beams are shown in blue and are labeled A_1 , A_2 , and A_3 .**

101 The Aquadopp is a narrow-band instrument ([https://support.nortekgroup.com/hc/en-us/articles/360029839331-The-](https://support.nortekgroup.com/hc/en-us/articles/360029839331-The-Comprehensive-Manual-ADCP)
102 [Comprehensive-Manual-ADCP](https://support.nortekgroup.com/hc/en-us/articles/360029839331-The-Comprehensive-Manual-ADCP)) that emits short sound pulses (pings) at a constant frequency and receives reflected (echo)
103 signals. Plankton and suspended matter, as well as air and gas bubbles, are the main scatterers of the sound. While sound
104 pulses are scattered in all directions when they hit particles, a small fraction of the incident sound pulse intensity is reflected.
105 The Aquadopp current meter employs a mono-static system in which 3 transducers are used to transmit and receive signals at
106 an acoustic frequency of 2 MHz. Measurements are made in the 90 dB range with a resolution of 0.45 dB. In high-accuracy
107 acoustic Doppler measurements, the acoustic beams are narrow and each has a cone angle of 1.7° . Three focused beams
108 measure the scattering strength with high sampling rates in a small volume (referred to as a single point). Two-sided acoustic
109 beams are directed horizontally with 90° spacing between the axes of the beams (Fig. 1). These beams measure the volume
110 scattering strength at the level of the transducer. The third beam is inclined at an angle of 45° to the plane formed by the axes
111 of the other two beams. The piezoelectric element of the transducer transmits sound waves when it vibrates. The vibration
112 does not stop at once but is damped over time. The speed of sound in water and the damping time of the membrane vibration
113 determine the dead zone. In our case, this distance along the acoustic beam is approximately 0.35 m from the piezoelectric

114 element of the transducer to the measurement cell (in the form of a truncated cone). The sound pulses are scattered and
115 reflected back to the transducer. In our case, the length of the cell along the axis of the acoustic beam is approximately 1.5
116 m. Therefore, the reflected sound pulse intensity obtained by the instrument is the weight average for the time during which
117 the sound wave passes the distance of 1.85 m to the far boundary of the measurement cell plus 1.85 m on the way back. The
118 received signal is processed in such a way that the greatest contribution to the average value is made by the scattering in the
119 center of the measurement cell at a distance of approximately 1.1 m from the transducer. The device can transmit up to 23
120 sound pulses every 1 s. The average value of the volume scattering strength for sound pulses transmitted and received in 1 s
121 is recorded in the device's memory.

122 The high frequency of 2 MHz allows observations of small-sized sound scatterers. Theoretically, a 2 MHz transducer
123 is most sensitive to particles with a diameter of 0.23 mm (estimates for different frequencies for standard seawater are given,
124 for example, in Hofmann and Peeters, (2013)). However, this is not entirely applicable to zooplankton due to the complex
125 shape of these organisms, their structure, their lipid composition, and the presence of gases in their bodies (Stanton et al.,
126 1994; Lavery et al., 2007, Lawson et al., 2006). However, as a simplified model, copepod species are often considered
127 cylinders, the scattering from which is defined as a function of the incident sound pressure, the acoustic wavelength, and the
128 distance between the transmitter and the animal. An approximate formula for describing sound scattering from an elongated
129 weakly scattering body of an animal also includes the angle of orientation of the body (Stanton et al., 1993, 1994).
130 Unfortunately, the manufacturer of the Aquadopp instrument does not specify information about the acoustic power of its
131 transducers. The Aquadopp measurement data for the volume scattering strength are presented in conventional units
132 (counts). Without special calibration, it is not possible to determine the amount of falling sound pressure in water at a
133 distance from the instrument transducer.

134 Since 2013, Aquadopp instruments with sideways-looking vertically mounted heads have been regularly used on the
135 Aqualog profiling carrier (Fig. 1). The carrier moves up or down at a speed of approximately 0.2 m s^{-1} , so the vertical
136 resolution of the volume scattering strength data is 0.2 m. These data are averaged every 5 s, allowing for the detection of an
137 SSL with a thickness on the order of 1 m.

138 In the context of this study, the ability to observe sound that has been reflected from zooplankton species at different
139 angles is important. In the case of settling detritus, the volume scattering strength of slanted beam A_3 and those of horizontal
140 beams A_1 or A_2 are approximately the same. If the elongated suspended particles are oriented vertically or inclined, the
141 amplitude of A_3 will significantly differ from the amplitudes of A_1 and A_2 . This was shown for copepods based on both
142 models of acoustic scattering at a frequency of 2 MHz (Stanton and Chu, 2000; Roberts and Jaffe, 2007) and laboratory
143 experiments (Roberts and Jaffe, 2008).

144 Thus, by comparing the amplitudes A_1 , A_2 , and A_3 , one can judge the predominant orientation of species in
145 zooplankton aggregations. It is assumed that aggregation's characteristic size is greater than the length of the acoustic
146 measurement cell, that is, not less than $\sim 2 \text{ m}$, and its lifetime is longer than 10 s. Therefore, during the Aqualog carrier
147 movement at a speed of 0.2 m s^{-1} , the slanted and horizontal acoustic beams scan the same zooplankton aggregation. The

148 complexity and variability of the acoustic backscatter makes it difficult to compare the acoustic signals obtained for different
149 observational periods. Proper normalization of the signals is needed to evaluate the seasonal change in the vertical
150 distribution of the mesozooplankton SSLs from many profiles despite the variability of the amplitude of the acoustic
151 backscatter. For the Aquadopp instrument, such normalization is the ratio of the volume scattering strength of the horizontal
152 beams to the volume scattering strength of the slanted beam


$$R = (A_1 + A_2)/2A_3. \quad (1)$$

154 It allows for a drastic reduction in the noise associated with clouds of sinking particles, which have an approximately equal
155 area in the horizontal projection to the projection with a 45° angle of inclination. In some cases, the suspended particles can
156 completely obscure the signal associated with the aggregation of mesozooplankton. However, in this study, there were
157 usually only a few such cases. As will be shown below in Section 3, typically at depths from 60 to 120 m during the day, the
158 directional acoustic backscatter ratio $R = 1.05$ - 1.2 , and at night, $R < 1.05$. In Appendix, we will consider whether the
159 mesozooplankton specimens' vertical orientation is tilted in the deep aggregations. The analysis will be based on calculation
160 of the ratio of the volume scattering strength of the horizontal beams A_1/A_2 assuming that due to the tilt the standard
161 deviation of A_1/A_2 should be greater than 0.

162 In addition to the Aquadopp instrument, a SeaBird 52MP CTD probe and Aanderaa 4330F and SBE 43F dissolved
163 oxygen fast sensors were incorporated into the Aqualog profiler aerobic zone (Ostrovskii and Zatsepin, 2016). The SeaBird
164 52MP CTD was specially designed for a moored profiling application in which the instrument makes vertical profile
165 measurements from a carrier that travels vertically beneath a subsurface floatation ([https://www.seabird.com/moving-
166 platform/sbe-52-mp-moored-profiler-ctd-optional-do-sensor/family?productCategoryId=54627473795](https://www.seabird.com/moving-platform/sbe-52-mp-moored-profiler-ctd-optional-do-sensor/family?productCategoryId=54627473795)). The CTD is
167 equipped with a pump that controls a flow at a constant speed through a single small diameter opening to ensure the
168 minimization of salinity spiking in the measurement data by the temperature and conductivity cell. On the Aqualog profiling
169 carrier slowly moving at $\sim 0.2 \text{ m s}^{-1}$, the CTD sampling rate of once per second provides sufficient data to resolve ocean fine-
170 scale thermohaline structure. The accuracy of the CTD probe is $0.002 \text{ }^\circ\text{C}$ for the temperature, $\pm 0.0003 \text{ S/m}$ for the
171 conductivity and $\pm 0.1\%$ of the full scale range for the pressure. The SBE 43F accuracy should be no worse than $\pm 2\%$
172 saturation, which can be compared with 5% for Aanderaa 4330F with a resolution better than $1 \text{ } \mu\text{M}$ or 0.4%
173 (https://www.aanderaa.com/media/pdfs/d378_aanderaa_oxygen_sensor_4330_4330f.pdf). In practice, in the Black Sea, SBE
174 43F showed very robust results in detecting the lower boundary of the oxic zone, consistent with observations of the sigma-
175 density structure and definition of the oxic zone boundary for the northeastern region of the Sea (Ostrovskii and Zatsepin,
176 2016). The SeaBird 52MP CTD with SBE 43F was regularly calibrated at the facility of the Southern Branch of Shirshov
177 Institute of Oceanology, Gelendzhik. The dissolved oxygen measurements by using the Aanderaa 4330F and SBE 43F
178 sensors at the profiler were described in (Ostrovskii and Zatsepin, 2016) and later in a companion paper (Ostrovskii et al.,
179 2018). The fast response sensing foils of the Aanderaa 4330F sensor were replaced by new foils two times in the past four
180 years. The CTD and dissolved oxygen sensors were mounted at the leading edge of the Aqualog profiler pointing into
181 horizontal oncoming flow, while hydrodynamic cowling (vertically oriented, wing-like) helped to stabilize the profiler

182 orientation with respect to the flow direction. It should be noted that the Black Sea environment is particularly suitable for
183 profiling measurements since there is no biological fouling on the sensors of the profiler, which is usually submerged into
184 the hydrogen sulfide zone for ~10 min every 1-2 h. Finally, the dissolved oxygen sensor data were verified with the water
185 samples at standard depths for determination of dissolved oxygen by Winkler method (not shown here).

186 The profiler mooring station was deployed approximately four nautical miles from the coast at the uppermost part
187 of the continental slope at 44°29.3'N and 37°58.7'E (Fig. 2). From June 2010 to April 2021, 16 surveys lasting from a few
188 days to 3 months were carried out (Table 1) (Solovyev et al., 2021). During the surveys, the device automatically performed a
189 profiling cycle usually every 1-2 h, descending to the near-bottom depth of 200-220 m and ascending to the upper layer
190 while remaining submerged at a depth of 20-40 m. In particular, in 2016-2020, more than 14,000 multiparameter sets of
191 vertical profiles were collected year-round (except March).



192
193 **Figure 2: The Black Sea coastline (<http://openstreetmapdata.com/data/coastlines>). The observational site off Gelendzhik is shown**
194 **by a red dot. © OpenStreetMap contributors 2021. Distributed under the Open Data Commons Open Database License (ODbL)**
195 **v1.0.**

196 To acquire taxonomic and quantitative features of zooplankton vertical distribution, stratified net samples were
197 taken from R/V Ashamba (Table 1) near the moored profiler Aqualog with a Juday net (mouth area 0.1 m², mesh size 180
198 µm) equipped with a closing device. The towing speed was 0.9-1.0 m s⁻¹ and the net was closed without stopping the upward
199 movement. The sampling was carried out in calm weather so that the wire angle was not higher than 10 degrees. The
200 sampling was carried out at earlier stages of this project in June 2010, October 2013, and July 2014, as well as later in
201 October 2016.

202 **Table 1. Deployments of the profiler Aqualog-6 with Nortek Aquadopp current meter in the NE Black Sea and the dates**
203 **of the zooplankton sampling near the profiler mooring site in 2010-2021. Since 2013, the profiler Aqualog was equipped with SBE**
204 **52MP CTD probe with SBE 43F DO sensor. Additional sensors used at the profiler were as follows Oxygen Aanderaa 4330F,**
205 **Seapoint Turbidimeter, and Seapoint Fluorometer.**

Survey	Start	End	Profile cycle interval, h	Profile depth range, m	Number of profiles	Additional sensors at the profiler	Stratified net sampling for zooplankton / Sampling for determination of dissolved oxygen by Winkler method
1	21.06.2010 16:03	22.06.2010 16:50	1	19-245	25	*No dissolved oxygen sensor	Zooplankton: 21.06.2010 18:05-19:00 21.06.2010 21:10-21:55 22.06.2010 00:05-00:50 22.06.2010 05:30-06:20 22.06.2010 09:00-09:50
2	02.10.2013 12:42	07.10.2013 9:14	1	30-220	234	--	Zooplankton: 06.10.2013 12:30-13:20
3	28.06.2014 10:46	02.07.2014 13:24	1	20-240	198	4330F	Zooplankton: 01.07.2014 13:30-14:30 02.07.2014 02:30-03:30 Dissolved oxygen: 12.07.2020, 14.07.2020 16.07.2020
4	06.10.2014 5:50	17.12.2014 12:02	6	30-220	860	4330F	--
5	01.01.2016 18:00	06.03.2016 6:00	2	28-208	1490	4330F	--
6	06.10.2016 5:47	10.10.2016 10:21	2	25-220	98	4330F, Fluorometer, Turbidimeter	Zooplankton: 04.10.2016 22:00-23:00 05.10.2016 11:05-11:50
7	10.10.2016 13:24	12.11.2016 12:45	2	30-220	790	4330F, Fluorometer, Turbidimeter	--
8	10.02.2019 12:00	24.02.2019 4:08	2	25-206	328	Turbidimeter	--
9	16.04.2019 11:34	28.05.2019 9:24	1	46-206	2016	4330F, Turbidimeter,	--
10	01.06.19 10:32	27.08.19 12:02	1-2	22-200	2784	Turbidimeter	Dissolved oxygen: 06.07.2019, 08.07.2019 12.07.2019
11	30.08.19 16:00	15.10.19 20:26	1-2 (16 cpd)	22-200	1482	Turbidimeter	--
12	28.10.19 14:00	24.12.2019 20:36	1-2 (16 cpd)	21-204	1491	4330F	--
13	28.03.2020 11:30	24.05.2020 2:03	1-2 (16 cpd)	20-200	1584	4330F, Fluorometer	--
14	16.07.2020 5:00	26.07.2020 23:13	1-2 (16 cpd)	23-201	444	4330F, Fluorometer	Dissolved oxygen: 17.07.2020, 20.07.2020
15	03.10.2020 5:00	27.11.2020 9:37	2	20-203	1320	4330F, Fluorometer	--
16	11.12.2020 9:06	07.04.2021 1:04	4	21-203	1399	4330F, Fluorometer **Nortek Aquadopp broken	--

208

209 The net hauls targeted the backscattering aggregation considering that their locations were associated with specific
210 isopycnal layers (Ostrovskii and Zatsepin, 2011, Fig. 9). Vertical profiles of temperature, salinity, and density were obtained
211 with a ship-borne SeaBird 19plus CTD probe prior to mesozooplankton sampling. Depth strata were chosen based on the
212 CTD profiles to sample the upper mixed layer (UML), the thermocline layer, the layer from the oxycline upper boundary (σ_θ
213 = 14.25) to the lower boundary of the thermocline, and two layers in the oxygen-deficient zone: the layer from depths of σ_θ
214 15.7 to σ_θ 15.4 and the layer from 2-3 m below σ_θ 15.9 to σ_θ 15.7.


215 The time of sampling corresponded to the day/night vertical distribution and upward/downward migration of
216 zooplankton (June 2010), the daytime distribution (October 2013), and the day/night distribution (July 2014 and October
217 2016). The samples were immediately fixed with buffered formaldehyde (4% final concentration of seawater–formaldehyde
218 solution). The volume of filtered sea water was estimated from the area of the net mouth and the length of the released wire.
219 Organisms were identified and counted under a stereomicroscope equipped with an ocular micrometer. Zooplankters were
220 identified at the level of species and age stages of copepods and size classes (with an interval of 2 mm) of chaetognaths and
221 ctenophores. The smallest organisms (meroplankton, appendicularians, copepod nauplii and ova) considered in the analysis
222 were 180 μm in size. Mesozooplankton biomass in terms of dry weight (DW) was estimated based on the published length-
223 DW regressions for different species summarized in (Arashkevich et al., 2014, Table 2). Biomass values were standardized
224 to mg DW m^{-3} or mg DW m^{-2} . The intensity of the echo signal strongly depends on the material properties of the organism's
225 tissue (Stanton et al., 1994); therefore, when comparing the pattern of the scattering signal intensity with the pattern of
226 zooplankton distribution in a community containing different taxa, it was reasonable to express zooplankton biomass as DW
227 or carbon (Flagg and Smith, 1989; Heywood et al., 1991; Ashjian et al., 1998). For a graphical presentation of the results, six
228 components of zooplankton were considered: copepods *Calanus euxinus* and *Pseudocalanus elongatus*, small crustaceans
229 (*Acartia clausi*, *Paracalanus parvus*, *Oithona similis* and cladocerans), heterotrophic dinoflagellate *Noctiluca scintillans*,
230 chaetognaths *Parasagitta setosa*, and varia (ctenophores *Pleurobrachia pileus*, appendicularians, meroplankton, decapod
231 larvae, Pisces ova).

232 One method for calculating vertical migration speed of zooplankton from the sound backscatter data of the acoustic
233 current meter at the profiler Aqualog was described in (Pezacki et al., 2017). However the vertical migration speed of
234 mesozooplankton is beyond the focus of this study. Only once when discussing the pattern of the diel vertical migration, the
235 slope of the migration track on the echogram (see Figure 9 below) is considered to give rough idea about the dive and the
236 ascend of mesozooplankton. Much more effort would certainly be needed to visualize the specimens' vertical swimming.

237 3 Results

238 3.1 Acoustic scattering by mesozooplankton aggregations

239 The first validation data for the Aquadopp observations were obtained on 21-22 June 2010. The sound scattering layers were
240 identified at the raw echogram (Fig. 3) as mesozooplankton aggregations by comparison with the net sampling data (Fig. 4).
241 The zooplankton net sampling data were consistent with the acoustic backscatter, indicating short-term variations in biomass
242 and diel vertical migration of zooplankton.




243

244 **Figure 3: Diurnal motions of the sound scattering layers in the oxic zone. The depth-time scatterplot for profiles of acoustic**
245 **backscatter amplitude at 2 MHz was obtained using the Aquadopp instrument at the moored profiler during verification study**
246 **involving net sampling of zooplankton on 21-22 June 2010.**


247 The total mesozooplankton biomass in the entire water column varied from 0.99 to 3.57 g DW m⁻². Zooplankton
248 was dominated by the copepod *Calanus euxinus*, which made up the mean 58% with the standard deviation (SD) ±14% of
249 the total biomass (below for the sake of brevity, such estimates are denoted as mean±SD%). The contribution of
250 chaetognaths *Parasagitta setosa* was 21±11%, followed by copepods *Pseudocalanus elongatus* (13±7% of the total
251 biomass). The sum share of other groups of mesozooplankton did not exceed 7% of the total biomass.

252 The pattern of the vertical distribution of mesozooplankton biomass reveals a relatively uniform distribution over
253 depth in the evening twilight (18:05-19:00) and at dawn (05:30-06:20) (Fig. 4, left column). At night (21:10-21:55 and
254 00:05-00:50), the highest concentration of zooplankton was observed in the thermocline layer, while in the daytime (09:00-
255 09:50), the zooplankton maximum was in the layer between the density surfaces σ_{θ} 15.7 and 15.4 (Fig. 4, left), in accordance
256 with the diurnal changes in the volume backscatter strength (Fig. 3). The deepest layer bounded by isopycnals σ_{θ} 15.9 and
257 15.7 was inhabited by nonmigrating copepods, the fifth copepodite stage (CV) of *C. euxinus* (median prosome length 2.3

258 mm), persistently staying at this depth throughout the day (Figs. 3 and 4, middle column). Visual inspection of live samples
 259 revealed quiescent behavior of these specimens and large oil sac volume inside their body, suggesting a diapausing state in
 260 *C. euxinus* CV collected from the deepest layer (Vinogradov et al., 1992).



261
 262 **Figure 4: The diel changes in vertical distributions of (left) total mesozooplankton biomass, (middle) zooplankton composition,**
 263 **(right) temperature (*T*) and density (σ_{θ}) near the mooring site on 21-22 June 2010. The temperature and density profiles were used**
 264 **for the selection of sampling strata. CE – *Calanus euxinus*; PE – *Pseudocalanus elongatus*; SC – small crustaceans; NS – *Noctiluca***
 265 ***scintillans*; PE – *Parasagitta setosa*; Var – varia.**



266


267 **Figure 4: Continued.**

268 Three migrating species, copepodites CIV-CVI of *C. euxinus* (median prosome length 2.6 mm), CV-CVI *P.*
 269 *elongatus* (median prosome length 0.92 mm), and chaetognaths *P. setosa* (median length 19 mm), formed daytime
 270 zooplankton aggregations in the oxygen-deficient zone (Fig. 4, middle panel). Ctenophore *Pleurobrachia pileus*, also
 271 inhabiting the deep layers in the daytime, contributed negligibly to the total biomass due to the low dry matter content in
 272 their gelatinous bodies and their low abundance (shown as Var in Fig. 4). At night, most of the migrating zooplankters were
 273 concentrated in the thermocline and did not ascend to the warm UML, which was inhabited by small copepods, cladocerans,
 274 and small (<6 mm) chaetognaths.

275 3.2 Zooplankton aggregations visualized using the directional acoustic backscatter ratio

276 The echograms based on the data from horizontal-beam transducers A_1 and A_2 often reveal aggregations of zooplankton at
 277 depths of 80-120 m in the daytime (Fig. 5). The aggregations begin to rise around sunset and descend before dawn. Thin,
 278 nearly vertical lines on the echogram indicate acoustic traces of the migrating mesozooplankton species. The echogram also
 279 shows patches that occupy the entire water column, from the upper to the lower measurement depth, penetrating below the
 280 surface σ_θ 15.9 and then deeper into the hydrogen sulfide zone. These are clouds of suspended particles (see, for example,


281 Klyuvitkin et al., 2016). Acoustic scattering by clouds of particles sinking through the water column can obscure
 282 zooplankton aggregations.



283
 284 **Figure 5: a -** The time-depth graph of the Aquadopp horizontal-beam $(A_1 + A_2)/2$ echogram showing acoustic backscatter intensity
 285 (counts) on 5-7 October 2013. The Aqualog profiler with the Aquadopp instrument performed ascending/descending cycles every 1
 286 h. The upper, middle and lower white lines are for isopycnals σ_θ 15.4, 15.7 and 15.9, respectively. **b -** Time-depth scatterplot of the
 287 Aquadopp directional acoustic backscatter ratio $R = (A_1 + A_2)/2A_3$. Colored lines show $[\text{O}_2] = 50 \mu\text{M}$ (upper black line); $[\text{O}_2] = 10$
 288 μM (lower black line); and $\sigma_\theta = 15.9 \text{ kg m}^{-3}$ (white line), which can be taken as a proxy for the boundary of the oxygen zone in the
 289 NE Black Sea (Glazer et al., 2006, Ostrovskii and Zatsepin, 2016). Notice that due to upwelling the oxycline was moved upward.
 290 Vertical dotted white lines indicate a 17.3 h time interval, which is equal to the period of inertial oscillations at the latitude of the
 291 observation. They approximately coincide with troughs of inertial waves.

292
 293 The layers of elevated acoustic backscatter amplitude due to deep zooplankton aggregations are accounted for using
 294 the R graphs (Fig. 5b) that were validated by net sampling on 6 October 2013 (Fig. 6), although sampling was not performed
 295 at night due to stormy weather. Since the depths of the isopycnals of 15.9 and 15.7 differed by only 3 m, the integrated

296 zooplankton sample was taken in the layer between $\sigma_\theta = 15.9$ and $\sigma_\theta = 15.4$. In this layer, the contributions of *Calanus*
 297 *euxinus*, *Parasagitta setosa*, and *Pseudocalanus elongatus* to the total biomass were 60%, 26%, and 12%, respectively (Fig.
 298 6 b). The extremely low zooplankton biomass ($< 2 \text{ mg DW m}^{-3}$) in the upper 50 m layer (Fig. 6 a, b) is consistent with data
 299 on a fourfold decrease in the annual average biomass of upper-dwelling zooplankton in 2013 compared to previous years
 300 (Arashkevich et al., 2015).




301

302 **Figure 6: The daytime vertical distributions of (a) total mesozooplankton biomass, (b) zooplankton composition, (c) temperature**
 303 **(T) and density (σ_θ) near the mooring site at 12:30-13:20 on 6 October 2013. Temperature and density profiles (c) indicate the**
 304 **selection of sampling strata. CE – *Calanus euxinus*; PE – *Pseudocalanus elongatus*; SC – small crustaceans; NS – *Noctiluca***
 305 ***scintillans*; PE – *Parasagitta setosa*; Var – *varia*.**

306 Zooplankton diel vertical migration trajectories in the R graph are noticeably clear below 40 m (Fig. 5b). The
 307 explanation of these phenomena could be that the scattering area of the elongated bodies of the zooplankton species is larger
 308 in the horizontal projection than in the inclined projection at an angle of 45° . Therefore, the orientation of the bodies of
 309 mesozooplankton species appears to be mainly vertical during migration. At night, these specimens are randomly oriented in
 310 the upper layer, where $R \approx 1$. In addition to the diel vertical migrations, intraday vertical fluctuations of zooplankton occur
 311 with an inertial period (Fig. 5b). The vertical displacements of the daytime deep mesozooplankton aggregations are coherent
 312 with the vertical displacements of both isopycnals and isooxylines. The displacements of isopycnals with amplitudes up to
 313 20 m are mainly due to near-inertial waves.


314 In October 2016, persistent aggregation of diapausing *C. euxinus* was detected in the acoustic backscatter signal (Fig.
 315 7), unlike October 2013 (Fig. 5). Zooplankton sampling was performed at midnight and midday on 4-5 October 2016 (Fig.
 316 8). The pattern of zooplankton distribution was similar to that in June 2010 (Fig. 4), both in terms of the total biomass and
 317 composition of zooplankton and in terms of the day/night vertical distribution. The total mesozooplankton biomass of 1.8-
 318 2.3 g DW m^{-2} was dominated by three species: *C. euxinus* (59-76%), *P. setosa* (9-22%), and *P. elongatus* (5-10%). At night,
 319 the maximum aggregation of migrating zooplankters was in the thermocline layer, and at midday, it was in the layer between
 320 isopycnals σ_θ 15.7 and 15.4 (Fig. 8a). Daytime zooplankton aggregation consisted mainly of *C. euxinus* (92% of total
 321 biomass) with a small contribution from chaetognaths (7% of total biomass) (Fig. 5b). The layer between isopycnals σ_θ 15.9
 322 and 15.7 was persistently occupied by diapausing *C. euxinus* CVs. The chaetognaths found in this layer were represented by

323 spent specimens and corpses (Fig. 8b). UML was inhabited by nonmigrating small copepods, cladocerans, and small
324 chaetognaths.



325

326 **Figure 7: a - The time-depth scatterplot of the Aquadopp horizontal-beam echo $(A1 + A2)/2$ on 6-10 October 2016. b - The time-**
327 **depth graph of the Aquadopp directional acoustic backscatter ratio R . The isopycnals and isooxylines are superimposed near the**
328 **SSLs.**




329

330 **Figure 8: Day/night vertical distribution of (a) total mesozooplankton biomass, (b) zooplankton composition, (c) temperature (T , °C) and density (σ_θ) near the mooring site at 22:00-23:00 of 4 October (night) and 11:05-11:50 of 5 October (day) in 2016.**
 331 **Temperature and density profiles (c) indicate the selection of sampling strata. CE – *Calanus euxinus*; PE – *Pseudocalanus elongatus*; SC – small crustaceans; NS – *Noctiluca scintillans*; PE – *Parasagitta setosa*; Var – varia.**
 332
 333

334 The net sampling data on the day/night vertical distribution of mesozooplankton agreed broadly with the acoustic
 335 backscatter observations obtained during the next few days (Fig. 7). On the echogram, one can see a persistently existing
 336 backscattering layer associated with the isopycnal layer near $\sigma_\theta = 15.9$ as well as patches of the high-volume backscattering
 337 strength at depth during the daytime and their movement into shallower layers at night.


338 The two-layered structure was also observed at the end of June – early July 2014 (Fig. 9) and validated by day/night
 339 zooplankton sampling on 1-2 July (Fig. 10). Deeper zooplankton aggregation was monospecific, consisting only of
 340 diapausing *C. euxinus* CVs (Fig. 10b) and formed a thin layer (5 - 10 m thick). This layer was visible all day and night and
 341 was usually located above the isopycnal surface of 15.9. It is clearly distinguished by the value $R > 1.1$ (Fig. 9b). The
 342 daytime zooplankton aggregation consisted of three migrating species and their different developmental stages, *C. euxinus*,
 343 CIV-CVIs, *P. elongatus*, CV-CVIs, and *P. setosa*, 14-22 mm in size (Fig. 10b). Since the amplitude of vertical migration is
 344 different for different components of this assembly, the daytime deep aggregation reached 35 m in thickness. Before sunset,
 345 migrating zooplankters began to move upward and at night formed aggregations at depths above 40 m (Fig. 9a) and peaked
 346 in the thermocline at 17-25 m (Fig. 10a).



347
 348 **Figure 9: a** — The time-depth graph of the Aquadopp horizontal-beam echo $(A1 + A2)/2$ during the moored profiler survey on 28
 349 **June - 2 July 2014**. The isopycnals are superimposed near the SSLs. **b** - The graph of time-depth variation in R based on the
 350 measurements of sound backscattering. The upper and lower black lines are iso-oxylines of 50 and 10 μM , respectively. The white
 351 line indicates isopycnal $\sigma_\theta = 15.9$. There is a persistent SSL under isooxylene $[\text{O}_2] = 10 \mu\text{M}$. Thin white arrows schematically show
 352 the diel migration of mesozooplankton. The maximum depth of the diel vertical migration is 120-150 m, although some specimens
 353 dive to depths of only 80-100 m. The slope of the straight arrow pointing downwards corresponds to a diving speed of $\sim 1.5 \text{ cm s}^{-1}$.
 354 The ascent is accelerated and reaches values of approximately 2.5 cm s^{-1} in the upper 60 m depth.

355


356



357

358 **Figure 10: The day/night vertical distribution of (a) total mesozooplankton biomass, (b) zooplankton composition, (c) temperature**
 359 **(T) and density (σ_θ) near the mooring site on 1 July (13:30-14:30) and 2 July (02:30-03:30) of 2014. Temperature and density**
 360 **profiles (c) indicate the selection of sampling strata. CE – *Calanus euxinus*; PE – *Pseudocalanus elongatus*; SC – small crustaceans;**
 361 **NS – *Noctiluca scintillans*; PE – *Parasagitta setosa*; Var – varia.**

362 Since migrating zooplankton aggregations were observed in the deep layers only during the daytime, it is worth
 363 comparing the daytime average R profile with that for the nighttime (Fig. 11). Such a comparison clearly reveals the deep
 364 maximum of R at the daytime migration depths of mesozooplankton at 90-120 m as well as the persistent maximum of the
 365 diapause layer within the deeper layer at 125-140 m. Notably, the depths of the persistent SSL change by approximately 5 m
 366 from night to the daytime, while they completely overlap when considered versus the density. Such variations in the depth of
 367 the SSL might be linked to inertial oscillations (Ostrovskii et al., 2018).



368


369 **Figure 11: The time averages of R and [O2] for the daytime of 1 July 2014 (red), and the nighttime of 2 July 2014 (blue), when net**
370 **sampling (Fig. 10) took place. Left - The depth profiles of the time averages R . Middle – the daytime and nighttime averages R**
371 **versus the specific density, σ_θ . Right – Distribution of the daytime and nighttime averages of the dissolved oxygen concentration**
372 **versus σ_θ .**

373 **3.3 The seasonal variation in mesozooplankton dynamics in relation to dissolved oxygen concentration**

374 In subsection 3.2, it was shown that the mesozooplankton species float on isopycnals in the lower part of the oxycline and in
375 the hypoxic zone. Both the diapausing aggregations and the daytime aggregations are displaced coherently by near-inertial
376 waves. The deep aggregations of mesozooplankton are bounded by certain isopycnal surfaces and iso-oxyines.

377 Since the oxygen stratification strongly depends on the density stratification in the pycnocline (e.g., Vinogradov and
378 Nalbandov, 1990, Codispoti, et al., 1991, Konovalov et al., 2005, see also example at Fig. 12), it becomes possible to switch
379 from the depth profiles of the directional acoustic backscatter ratio $R(z)$, where z is the depth, to the $R([O_2])$ profiles to
380 investigate the seasonal changes of the sound-scattering mesozooplankton layers in terms of R versus [O2].


381



382

383 **Figure 12: a** – Example of the monthly long time series of the vertical profiles of the directional acoustic backscatter ratio R for
 384 **November 2019** (in total 960 profiles from the moored profiler survey). The upper and lower black lines are iso-oxylines of 50 and
 385 **10 μM** , respectively. The white line indicates isopycnal $\sigma_\theta = 15.9$. **b** – Evolution of the dissolved oxygen at the profiler mooring site
 386 **in November 2019**. The colored lines indicate the following: the depths of the isopycnals $\sigma_\theta = 14$ and 15.9 (top and bottom white
 387 **lines**); the depth of the temperature minimum (green line); and $[\text{O}_2] = 50$ and $10 \mu\text{M}$ (blue lines). The inset shows the diagram of
 388 **the concentration of dissolved oxygen versus the potential density, $[\text{O}_2]-\sigma_\theta$** , plotted from the moored profiler data of November
 389 **2019**. In this example, as well as for other observational periods, the concentration of dissolved oxygen deviates very little from
 390 **isopycnal surfaces in the lower part of the oxycline where $[\text{O}_2] < 200 \mu\text{M}$** .

391 The average monthly profiles of $R([O_2])$ were constructed from $R(z)$ and $[O_2](z)$ data for every month when the data
 392 were available. To compute the averages, the daytime was defined as a period beginning 2 h after the local time of sunrise (at
 393 a given date) and ending 2 h before sunset. The nighttime was defined as a period beginning 1 h after sunset and ending 1 h
 394 before sunrise. Example plots of the average profiles $\langle R([O_2]) \rangle$ computed as arithmetic and bootstrap mean values along
 395 with 95% bootstrap confidence intervals are shown for November 2019 in Fig. 13. In the hypoxic zone, the average values
 396 $\langle R([O_2]) \rangle$ for the daytime are significantly higher than those for the nighttime. The daytime averages $\langle R([O_2]) \rangle >$
 397 1.06 were in the range of $[O_2] = 9-40 \mu\text{M}$ in November 2019.




398
 399 **Figure 13: Example profiles of the daytime and nighttime averages $\langle R([O_2]) \rangle$ in November 2019. The inset shows the same plots**
 400 **with the Y-axis drawn logarithmically to reflect the lower parts of the profiles (the hypoxic zone) in more detail.**


401 The average monthly $\langle R([O_2]) \rangle$ profiles show the seasonal evolution of the mesozooplankton distribution (Fig. 14).
 402 The SSLs are barely discernible in January. One can note some activity in the upper part of the oxycline in February.
 403 Although we unfortunately do not have data for March, in April, two peaks appear in the $\langle R([O_2]) \rangle$ profiles in the layers
 404 where the concentration of dissolved oxygen is 25-60 μM and 4-9 μM . These maxima correspond to the daytime

405 mesozooplankton aggregations and the diapause layer, respectively. The upper maximum of $\langle R([O_2]) \rangle$, which corresponds
 406 to the daytime aggregations of mesozooplankton, may weaken in June-July. However, it becomes stronger again at the end
 407 of summer and in autumn. The largest value for this maximum over the entire observation period $\langle R([O_2]) \rangle = 1.18$ is
 408 observed in October. At that time, the maximum shifts into the layer where $[O_2]$ is 10-25 μM . In December 2019, this peak
 409 was between the 10 μM and 30 μM iso-oxylines.

410 The maximum of diapause mesozooplankton was strongest in May and July 2020, reaching almost 1.2 at $[O_2] = 5-8$
 411 μM . In August, the diapause mesozooplankton layer shifts in the lower part of the suboxic zone where $[O_2] = 3-7$ μM . It
 412 becomes substantially weaker in September. In October, this layer degrades further. In November, it tends to disappear.



413



414
 415 **Figure 14: Top - The monthly averaged profiles of $\langle R([O_2]) \rangle$ for the daytime over the upper part of the continental slope near**
 416 **Gelendzhik in the NE Black Sea. Bottom – The same for the nighttime.**

417 **4 Discussion**

418 **4.1 Visualization of the sound-scattering mesozooplankton aggregations**

419 Previously, acoustic measurements at a frequency of 2 MHz were not considered a tool for observations of the
 420 mesozooplankton SSLs in the sea due to the limited range of soundings. However, with the advent of ocean profilers with
 421 acoustic Doppler current meters, such as the Nortek Aquadopp, it has become possible to obtain the depth profiles of the
 422 volume scattering strength at 2 MHz frequency in the entire water column and to study the vertical distribution of

423 zooplankton, such as those in the Black Sea (Ostrovskii and Zatsepin, 2011; Pezacki et al., 2017). Acoustic sounding of
424 mesozooplankton at two angles is made possible by using the side-looking head of the Nortek Aquadopp instrument. The
425 combination of horizontal and tilted beam signals allows, on the one hand, eliminating the patches of particles and equalizing
426 the background scattering level of the echogram and, on the other hand, determining the preferred orientation of
427 mesozooplankton species migrating through the oxycline. Earlier, Stanton and Chu (2000) reproduced the influence of the
428 orientation of a 3-mm calanoid copepod (modeled as a high-resolution approximation of an animal profile) on the acoustic
429 target strength at 2 MHz with respect to an incident sonar beam. The reduction was found to be 5-15% when copepod
430 orientation was shifted from 0° (broadside incidence) to 30-60°. Benfield et al. (2000) carried out field observations using
431 the Video Plankton Recorder on George Bank and showed that most *Calanus finmarchicus* (75%) in the depth range of 10-
432 70 m were within $\pm 30^\circ$ of the prosome-up or prosome-down orientation. It was suggested that one reason for the behavior
433 underlying the head-up orientation pattern might be due to the predator avoidance strategy aimed at reducing the
434 conspicuousness of *C. finmarchicus* when viewed from above. Such individuals would present a significantly reduced cross-
435 sectional area to an echo-sounder's transducer with correspondingly diminished target strength. It was concluded that it is
436 necessary to know how the orientation of individuals changes with depth to correctly account for the biomass of
437 mesozooplankton. Experiments using a multiple-angle acoustic receiver array on live copepods and mysids in a laboratory
438 tank showed that it is possible to use the scattered acoustic signal to distinguish among zooplankton taxa (Roberts and Jaffe,
439 2008). Reflections in the frequency range from 1.5 to 2.5 MHz were recorded from untethered 1 – 4 mm calanoid copepods
440 and 8 – 12 mm mysids over an angular range of 0–47°. That study demonstrated the utility of a multiple-angle acoustic array
441 for zooplankton identification.

442 To distinguish the SSLs against the background patterns of vertical flow of settling particles and to study the
443 orientation of zooplankton species, we propose a simple method for the processing of ultrasound sensing data at three angles.
444 This acoustic 3-beam geometry provides a partial pragmatic solution for the quest towards the multiple-angle scatter
445 measurements suggested by models (Stanton and Chu, 2000; Roberts and Jaffe, 2007) and laboratory experiments (Roberts
446 and Jaffe, 2008). Since the late 1990s, researchers' efforts have been focused on creating multichannel instruments to
447 measure acoustic backscatter (volume scattering strength) at several frequencies, which contain information about the size
448 composition of the scatterers, since different frequencies bounce off objects of different sizes (Wiebe et al., 2002, Smeti et
449 al., 2015). Multichannel instruments in conjunction with video cameras are fairly expensive systems that are used for the
450 identification of mesozooplankton in its natural habitat. Plausibly, a multichannel 3-angle system featuring several relatively
451 cheap short-range 3-beam acoustic units each operating at an individual frequency when installed on a vertically profiling
452 carrier would be a very effective tool for visualizing of zooplankton aggregations.

453 **4.2 The SSLs validated from the stratified net sampling**

454 Comparison of the Nortek Aquadopp acoustic backscatter observations with the data obtained by stratified zooplankton
455 sampling showed good agreement of the features of the diel vertical distribution of zooplankton. This was made possible by

456 sampling narrow depth strata (10-15 m layers) targeting deep-water aggregations visualized on the echograms. The two-
457 layered structure of the aggregations seen on echograms and *R*-graphs in the daytime (Figs. 3, 7, 9) reflected the species
458 composition of zooplankton in these layers (Figs. 4, 8, 10). The deepest layer bounded by isopycnals 15.9 and 15.7 was
459 visible in the suboxic zone all day and night and was formed by diapausing CV *Calanus euxinus*. To some extent, this
460 monospecific layer was contaminated by crustacean exuviae and carcasses, spent females, and zooplankters' remains sinking
461 from the upper layers and apparently retained on the density gradient. The existence of a nonmigrating diapausing stock
462 located in the suboxic layer from mid-spring to mid-autumn is confirmed by observations from submersible Argus
463 (Vinogradov et al. 1985; Flint 1989), by high vertical resolution sampling with 150 l water bottles (Vinogradov et al., 1992),
464 and by zooplankton net sampling (Arashkevich et al., 1998; Besiktepe, 2001; Svetlichny et al., 2009). However, for some
465 unknown reasons, this nonmigrating layer was not detected by ship-borne echo sounders at frequencies of 38 - 200 kHz
466 (Erkan and Gücü, 1998; Mutlu, 2003, 2007; Stefanova and Marinova, 2015, Sakınan and Gücü, 2016), unlike our data
467 obtained by Aquadopp at a frequency of 2 MHz.

468 The inclusion of diapause (or dormant stage) in the life cycle of all Calanidae species living in high-latitude and
469 temperate environments is well known (e.g., see review Baumgartner and Tarrant, 2017). Having accumulated a large
470 amount of lipids, diapausing copepods descend into deeper ocean layers where they can exist for several months at the
471 expense of energy reserves. Decreased metabolic rate and developmental delay are characteristic features of diapausing
472 copepods. In the Black Sea, a decrease in the metabolic rate in diapausing *C. euxinus* is caused not only by internal
473 physiological reasons but also by hypoxia in their dormant layer. The oxygen consumption rate in diapausing CV *C. euxinus*
474 in hypoxia decreases by almost an order of magnitude, and the rate of ammonia excretion decreases six times compared with
475 those in their active counterparts in normoxia (Svetlichny et al., 1998).

476 During the daytime, the upper SSL mostly located above $\sigma_{\theta} = 15.7$ consisted of four species, copepods *C. euxinus* and
477 *Pseudocalanus elongatus*, chaetognaths *Parasagitta setosa*, and ctenophores *Pleurobrachia pileus*; the latter had a negligible
478 contribution to dry biomass. This assembly had a wide range of body lengths from approximately 1 mm in *P. elongatus* to 22
479 mm in *P. setosa*. The different species had different swimming speeds. It was also possible that these species had different
480 physiological tolerances to oxygen deficiency. This confirms earlier observations from the manned submersible, which
481 showed that the daytime aggregation of migrating zooplankton had a layered structure: the lower layer was formed by
482 chaetognaths, whereas the older stages of *C. euxinus* were located above, and ctenophores inhabited the upper part of the
483 aggregation (Vinogradov et al., 1985; Flint, 1989). Furthermore, the different developmental stages of copepods *C. euxinus*
484 and *P. elongatus* occupied different depths, deepening as their size increased (Morozov et al., 2019).

485 In the evening approximately two hours before sunset, zooplankters begin to ascend to the upper layers, where they
486 spend all the dark hours concentrating in the thermocline layer and below it. In this layer, while feeding, they move in
487 different directions and are oriented randomly (Kiørboe et al., 2009), so they cannot be discernible in the *R*-graphs.
488 According to our data, cold-water herbivorous *C. euxinus* and *P. elongatus* only occasionally ascend into the warm UML,
489 mainly inhabiting colder layers rich in phytoplankton (see also the supplement to the paper by Morozov et al., 2019).

490 Predator chaetognaths *P. setosa* move upward following copepods, their main prey (Drits and Utkina, 1988). The time of
491 zooplankton migration clearly visible on the echograms is confirmed by the results of net sampling and is consistent with
492 other published data (see for references Morozov et al., 2019).

493 The vertical migration of zooplankton can increase the vertical flow of carbon and thus contribute to the functioning
494 of the biological pump in the ocean (Tutasi et al., 2020). The mesozooplankton that feed at the surface but metabolize and
495 excrete at depth contribute to the transport of organic matter; more quantitatively, this contribution is estimated to be
496 between approximately 10–50% of the local sinking flux of organic particles (Bianchi et al. (2013) and citation therein).

497 In the lower part of the oxic zone, the vertical displacements of SSLs coincide with the oscillations of isopycnal
498 surfaces (Figs. 5 and 10). The dissolved oxygen concentration profile tightly hinges on the density stratification in the Black
499 Sea since both are basically due to vertical mixing processes (e.g., Ostrovskii et al., 2018). Hence, displacements of the SSLs
500 with regard to the oxy-isolines are much smaller than those versus the depths. The vertical oscillations with a period of
501 approximately 17 h near the mooring site are due to near-inertial waves (Ostrovskii et al., 2018). Irregular changes in
502 isopycnal depths occur due to hydrodynamic events, such as individual internal waves, oceanic fronts, and jets.
503 Occasionally, the isopycnal depth may change by 30–40 m within a day (Ostrovskii and Zastsepin, 2016).

504 It is unlikely that copepods maintain their positions on certain isopycnal surfaces by swimming, as displacements of
505 such large amplitudes as tens of meters require an additional depletion of energy reserves. A more beneficial strategy would
506 be to adjust their buoyancy to neutral. Having neutral buoyancy in the hypoxic zone, the copepods would not need to spend
507 much additional energy floating up and down following crests and troughs of internal waves while avoiding entrainment into
508 the suboxic layer. Indeed, direct observations from manned submersibles revealed a quiescent behavior of diapausing
509 copepods and their slow response to light and noise produced by underwater vehicles, both in the Santa Barbara basin
510 (Alldredge et al., 1984) and in the Black Sea (M.V. Flint, personal communication). Neutral buoyancy has been
511 hypothesized to be regulated by changes in lipid composition (Visser and Jónasdóttir, 1999); however, Campbell and Dower
512 (2003) argued that this buoyancy regulation mechanism is inherently unstable because wax esters are more compressible
513 than seawater. An alternative mechanism for buoyancy regulation in diapausing copepods that involves the replacement of
514 heavy ions with lighter ammonium ions in hemolymph has been proposed by Sartoris et al. (2010) by analogy with other
515 invertebrates. Later, Schründer et al. (2013) found high concentrations of ammonium ions in the hemolymph of a diapausing
516 species, *Calanoides acutus*, and suggested that these copepods could achieve neutral buoyancy through their biochemical
517 body composition without swimming movements. This mechanism obviously would better explain the observed
518 phenomenon of diapausing copepod movement synchronized with the displacements of the isopycnal surfaces in the Black
519 Sea.

520 **4.3 Seasonal variations of the deep mesozooplankton SSLs**


521 For most of the year, from April to October, the sound scattering profiles in the deeper part of the oxic zone were bimodal
522 during the day (Fig. 14), reflecting the vertical distributions of two different zooplankton cohorts, migrating and diapausing.

523 The oxygen concentration at which the maximum backscattering signal from migrating zooplankton was observed decreased
524 throughout the year, from ca. 70-90 μM in January-February to 30-40 μM in April-August and to 15-20 μM in September-
525 December. Two explanations can be considered for the seasonal shift in the preferred oxygen concentration in these
526 zooplankters. On the one hand, this shift can be attributed to the deepening of the suboxic layer in January-February,
527 followed by gradual shallowing from April onwards (Fig. 15). If we assume that the depth of daytime zooplankton
528 aggregation depends to a large extent on the species-specific migration amplitude then with a deepened suboxic layer ($[\text{O}_2]$
529 $< 10 \mu\text{M}$), the migrating zooplankton will reach shallower depths, and its aggregation will be at a higher oxygen
530 concentration. Conversely, with a shoaling suboxic layer, zooplankton localize at lower oxygen levels.

531 This assumption is consistent with the data of Vinogradov et al. (1992), who found the daytime aggregation
532 maximum of migrating CV and female *C. euxinus* at an oxygen concentration of 18 μM when the suboxic layer was at a
533 depth of 110 m and at $[\text{O}_2] = 36 \mu\text{M}$ when the suboxic layer was at a depth below 170 m. This suggests that the trade-off
534 between the additional metabolic cost for extended swimming and metabolism reduction caused by low oxygen is in favor of
535 a decrease in the diel migration amplitude of *C. euxinus*. This differs from observations (Wishner et al., 2020) on the
536 migration of *Lucicutia hulsemannae* in the eastern tropical North Pacific, where this species changes its daytime location in
537 response to changes in the depth of the oxygen minimum zones (OMZ). For example, at the lower oxycline, the depth of
538 maximum abundance for *L. hulsemannae* shifted from ~ 600 to ~ 800 m in an expanded OMZ compared to a thinner OMZ
539 but remained at similar low oxygen levels in both situations. *L. hulsemannae* is an example of a “hypoxiphilic” species
540 (Wishner et al., 2020). However, unlike *Calanus* spp., *L. hulsemannae* is a strong swimmer capable of diel vertical migration
541 with amplitudes as large as approximately 1000 m.

542 Another explanation for the seasonal shift in the depth of the daytime aggregation in the Black Sea is the change in
543 taxonomic and age composition of the migrating cohort. A decreasing/increasing share of strong/weak swimmers with
544 different tolerances to oxygen deficiency may lead to a shift in the depth of daytime aggregation. The oxygen concentration
545 was in the range of 15-60 μM in the layer of daytime aggregation of migrating species in the Black Sea. Similarly, the
546 vertical distribution of migrating CV and adult *Calanus chilensis* off northern Peru was characterized by high abundance in
547 hypoxic waters at oxygen concentrations between 5 and 50 μM (Hirche et al., 2014).

548
549
550



551
 552 **Figure 15: The suboxic boundary depth ($[O_2] = 10 \mu M$) as observed by the moored profiler in 2016, 2019, 2020, and 2021.**

553 Based on our data, a diapausing cohort of *C. euxinus* appeared in April and terminated in November (Fig. 14).
 554 According to Vinogradov et al. (1985) and Svetlichny et al. (2009), the diapausing stage of *C. euxinus* was not found in the
 555 Black Sea in March. Hence, it is assumed that the diapausing stock is formed in April, when the offspring of the first
 556 generation of *C. euxinus* develop into the CV and accumulate sufficient lipid reserves. The energy reserve and a decrease in
 557 metabolic rate allow diapausing CV to exist without food for seven months in the suboxic zone (Vinogradov et al., 1992),
 558 which agrees with our observation of the diapause duration. The suboxic zone provides a refuge from large visual predators,
 559 which generally need higher oxygen concentrations (e.g., Bianchi et al., 2013).

560 The diapausing layer is bound by $3 \mu M$ and $10 \mu M$ and peaks at $5-7 \mu M$ oxygen. Earlier, the oxygen survival
 561 threshold for diapausing *C. euxinus* was determined experimentally at $[O_2] = 1.8 \mu M$ (Vinogradov et al., 1992).
 562 Physiological tolerance for hypoxia in diapausing *C. euxinus* resembles that reported for diapausing stage CV *Calanus*
 563 *pacificus*, which formed narrow dense aggregations at an oxygen concentration of $6.25 \mu M$ in the Santa Barbara Basin
 564 (Alldredge et al., 1984). The diapausing *C. pacificus* at a depth of 450 m was characterized by quiescent behavior, low
 565 laminarinase activity (as a proxy of feeding), and high lipid storage.

566 Similar tolerance for hypoxia was reported for diapausing *Eucalanus inermis*. Based on lactate dehydrogenase
 567 activity in the deep-dwelling CV and female *E. inermis*, their oxygen tolerance threshold was defined at the level of $4.47 \mu M$
 568 in the Pacific Ocean (Flint et al., 1991). Wishner et al. (2020) found a monospecific aggregation of *E. inermis* diapausing at
 569 extremely low oxygen, $1.0-5.7 \mu M$, in the eastern tropical North Pacific. In the Black Sea, the diapause depth is directly
 570 associated with certain density surfaces and consequently with a specific concentration of oxygen. From April to November,
 571 the isopycnal surfaces bounded by the suboxic layer move upward from depths of 130-160 m to 110-140 m. On this seasonal
 572 trend, the superimposed surfaces exhibit strong variations up to 60 m in amplitude at time scales from 17 h to several days

573 (Fig. 15). Diapause layers varied in depth along with isopycnal oscillations, allowing the copepods to remain in a constant-
574 low-oxygen habitat.

575 **5 Conclusions**

576 The key to using high-frequency sound in this study is to deploy the acoustic transducer in a manner that gets it sufficiently
577 close to the animal aggregations of interest. To visualize the mesozooplankton SSLs over the echogram with background
578 vertical flows of settling particles, we take advantage of the differences in acoustic scattering that is isotropic on the settling
579 particles and anisotropic on zooplankton species due to the elongated shape of the animals because their side view area is
580 larger than the head-view area or the tail-view area. The calculations of the ratio R of the volume scattering strength of the
581 horizontal acoustic beams to the volume scattering strength of the slanted beam allow visualizations of the mesozooplankton
582 aggregations of the specimen to be oriented vertically. This three-beam approach enhances the capability of underwater
583 ultrasound sensing to observe the mesozooplankton layers.

584 Linking the values of R to oxygen concentration enables us to derive the monthly averages from many profiles despite
585 the fluctuations in vertical distribution. The analysis of the oxygen-deficient zone allows us to describe the seasonal
586 evolution of diel zooplankton migrations, to determine the preferred oxygen regime for migrating and nonmigrating
587 zooplankters and to define the timing of formation, termination, and duration of diapause in CV *Calanus euxinus* in the
588 Black Sea.

589 Aggregations of vertically migrating zooplankton, consisting mainly of the older copepodite stages of *C. euxinus* and
590 *Pseudocalanus elongatus* and large-sized chaetognaths *Parasagitta setosa*, are observed within the hypoxic zone during the
591 daytime and mostly in the thermocline layer at night. The volume scattering strength in migrating SSL in the hypoxic layer
592 varies seasonally, with a minimum in winter and a maximum in late summer - early autumn. The location of this SSL also
593 changes in relation to the oxygen concentration in the range $[O_2]$ between 10 and 100 μM . Roughly, the deeper the suboxic
594 zone is located, the higher the oxygen concentration at the layers where the migrating species are aggregated. These
595 variations are hypothesized to address seasonal changes in the taxonomic and age composition of migratory zooplankton.
596 The maximum depth of zooplankton vertical diel migration is limited by the upper boundary of the suboxic zone ($[O_2] = 10$
597 μM).

598 The nonmigrating diapause SSL is observed at low $[O_2] = 3-10 \mu\text{M}$ from the beginning of April to the end of
599 October, suggesting a seven-month duration of diapause in *C. euxinus*. This persistent layer does not exceed 5-10 m in
600 thickness. The volume scattering strength in this monospecific layer may exceed that in the overlying daytime SSL,
601 apparently indicating the tighter aggregation of diapausing copepods compared to the aggregations of multispecies migrating
602 zooplankton. Diapause layers vary in depth along with isopycnal oscillations, allowing copepods to remain in a constant-
603 low-oxygen habitat.


604 Fluctuations of the SSLs are subject to interannual changes. It is necessary to maintain moored profiling acoustic
605 observatories in the Black Sea for a detailed analysis of year-to-year variability.

606

607 Appendix

608 Is the mesozooplankton specimens' vertical orientation tilted in the deep aggregations?

609 In the deep aggregations, the mesozooplankton species, while being oriented vertically in general, might be tilted with
610 respect to the vertical axis. Furthermore the specimens are probably occasionally tilted i.e., their azimuth angles are
611 distributed randomly so that the broadside incident angles of the horizontal acoustic beams would dominate the acoustic
612 backscattering data. Since the horizontal beams of the Nortek Aquadopp instrument are orthogonal, one can calculate the
613 standard deviation of the ensemble of the ratio of the acoustic backscatter of horizontal beams A_1/A_2 to check for the
614 possibility of a tilt. There is a high probability that $\langle A_1/A_2 \rangle = 1$ for the aggregations of tilted species, while the standard
615 deviation should be greater than 0. The persistent SSL caused by the diapausing mesozooplankton appears on the spring and
616 summer profiles of $R([O_2])$ as the maximum in the layer where $[O_2] < 10 \mu\text{M}$, i.e., in the suboxic layer. The orientation of
617 the mesozooplankton species in this layer is mostly vertical but tends to be slightly more tilted than in the daytime
618 aggregations of migrating mesozooplankton, as indicated by the A_1/A_2 ratio (see example for July 2019 at Fig. A1).



619

620 **Figure A1: Monthly averages of the depth profiles of the acoustic backscattering amplitude ratio A_1/A_2 for the time series of the**
621 **daytime (solid red line) and nighttime (blue solid line) data in July 2019. Dotted lines indicate the values of the standard deviations**
622 **from the means.**

623

624 *Data availability.* Underlying research data can be accessed via [HTTPS://DOI.ORG/10.13140/RG.2.2.28470.73285](https://doi.org/10.13140/RG.2.2.28470.73285),
625 License CC BY-NC 4.0 and [HTTPS://DOI.ORG/10.13140/RG.2.2.27548.62084](https://doi.org/10.13140/RG.2.2.27548.62084), License CC BY 4.0.

626

627 *Author contributions.* AO analyzed the moored profiler Aqualog data and wrote the parts of the paper related to the profiler
628 mooring measurements and data analysis. EA analyzed the net zooplankton data and wrote the parts of the paper related to
629 the zooplankton distribution analysis. VS deployed the mooring and handled the profiler sensors. DS is a design engineer of
630 the profiler who also maintained the profiler.

631

632 *Competing interests.* The authors declare that they have no conflict of interest

633

634 *Acknowledgements.* We thank A. Zatsepin for promoting of the moored profiler measurement program in the Black Sea.
635 The Black Sea field study is carried out in the framework of Russian Ministry of Science and High Education Assignment
636 No. 0128-2021-0016. The net sampling data analysis is supported by Russian Science Foundation grant No.20-17-00167.
637 The Aqualog profiler data processing and analysis are supported via grants No. 19-05-00459 and 19-45-230012 by Russian
638 Fund for Basic Research and Krasnodarsky Kray Ministry of Science, Education, and Youth Policy. We are grateful to editor
639 and three anonymous referees for helpful comments on this paper.

640

641 **References**

642 Alldredge, A. L., Robison, B. H., Fleminger, A., Torres, J. J., King, J. M., and Hamner, W. M.: Direct sampling and in situ
643 observation of a persistent copepod aggregation in the mesopelagic zone of the Santa Barbara Basin, *Mar. Biol.*, 80, 75–81,
644 1984.

645 Andrusov, N. I.: Preliminary report on the Black Sea cruise. *Izvestiya Imperatorskogo Russkogo Geographicheskogo*
646 *Obschestva [Bulletin of the Imperial Russian Geographic Society]*, 26, 398-409, 1890 (in Russian).

647 Arashkevich, E., Svetlichny, L., Gubareva, E., Besiktepe, S., Gücü, A. C., and Kideys, A. E.: Physiological and ecological
648 studies of *Calanus euxinus* (Hulsemann) from the Black Sea with comments on its life cycle, in: *Ecosystem modelling as a*
649 *management tool for the Black Sea*, edited by: Ivanov L. I., and Oguz, N., Kluwer Academic Publishers, Dordrecht, 351-
650 365, 1998.

651 Arashkevich, E. G., Louppova, N. E., Nikishina, A. B., Pautova, L. A., Chasovnikov, V. K., Drits, A. V., Podymov, O. I.,
652 Romanova, N. D., Stanichnaya, R. R., Zatsepin, A. G., Kuklev, S. B., and Flint, M. V.: Marine environmental monitoring in
653 the shelf zone of the Black Sea: Assessment of the current state of the pelagic ecosystem. *Oceanology*, 55, 871–876,
654 <https://doi.org/10.1134/S0001437015060016>, 2015.

655 Arashkevich, E., Ostrovskii, A., and Solovyev, V.: Observations of water column habitats by combining acoustic backscatter
656 data and zooplankton sampling in the NE Black Sea, 40th CIESM Congress Proceedings, Marseille, France, 28 October - 1
657 November 2013, 40, 722, (<http://www.ciesm.org/online/archives/abstracts/pdf/40/index.php>), 2013.

658 Arashkevich, E. G., Stefanova, K., Bandelj, V., Siokou, I., Kurt, T. T., Örek, Y. A., Timofte, F., Timonin, A., and Solidoro,
659 C.: Mesozooplankton in the open Black Sea: Regional and seasonal characteristics, *J. Mar. Syst.*, 135, 81–96,
660 <https://doi.org/10.1016/j.jmarsys.2013.07.011>, 2014.

661 Ashjian, C. J., Smith, S. L., Flagg, C. N., and Wilson, C.: Patterns and occurrence of diel vertical migration of zooplankton
662 biomass in the Mid-Atlantic Bight described by an acoustic Doppler current profiler. *Cont. Shelf Res.*, 18, 831-858, 1998.

663 Baumgartner, M.F. and Tarrant, A.M.: the physiology and ecology of diapause in marine copepods, *Annu. Rev. Mar. Sci.*, 9,
664 387-411, <https://doi.org/10.1146/annurev-marine-010816-060505>, 2017.

665 Benfield, M. C., Davis, C. S., and Gallagher, S. M.: Estimating the in situ orientation of *Calanus finmarchicus* on Georges
666 Bank using the Video Plankton Recorder, *Plankton Biol. Ecol.*, 47, 69–72, 2000.

667 Besiktepe, S.: Diel vertical distribution, and herbivory of copepods in the south-western part of the Black Sea, *J. Mar. Syst.*
668 28, 281–301, [https://doi.org/10.1016/S0924-7963\(01\)00029-X](https://doi.org/10.1016/S0924-7963(01)00029-X), 2001,

669 Bianchi, D., Galbraith, E., Carozza, D., Mislán, K. A. S., and Stock, C. A.: Intensification of open-ocean oxygen depletion
670 by vertically migrating animals, *Nature Geosci.*, 6, 545–548, <https://doi.org/10.1038/ngeo1837>, 2013.

671 Campbell, R. W. and Dower J. F.: Role of lipids in the maintenance of neutral buoyancy by zooplankton, *Mar. Ecol. Prog.*
672 *Ser.*, 263, 93–99, <https://doi.org/10.3354/meps263093>, 2003.

673 Codispoti, L. A., Friederich, G. E., Murray, J. W., and Sakamoto, C. M.: Chemical variability in the Black Sea:
674 implications of continuous vertical profiles that penetrated the oxic/anoxic interface. *Deep-Sea Res. I*, 38, Suppl. 2,
675 S691-S710, 1991.

676 Drits, A. V. and Utkina, S. V.: *Sagitta setosa* feeding in the deep layers of high plankton concentration during daytime in the
677 Black Sea, *Oceanology*, 28, 1014-1018, 1988.

678 Erkan, F. and Gücü, A. C.: Analyzing shipborne ADCP measurements to estimate distribution of southern Black Sea zoo
679 plankton. *Proceedings of the 4th European Conference on Underwater Acoustics, Rome*, 1, 267–27, 1998.

680 Flagg, C. N. and Smith, S. L.: On the use of the acoustic Doppler current profiler to measure zooplankton abundance. *Deep*
681 *Sea Res.*, 36, 455-474, 1989.

682 Flint, M. V.: Vertical distribution of mass zooplankton species in lower layers of aerobic zone in relation to the structure of
683 oxygen field, in: *Structure and production characteristics of planktonic populations in the Black Sea*, edited by
684 Vinogradov, M. E. and Flint, M. V., Nauka, Moscow, 187–213, 1989 (in Russian).

685 Flint, M. V., Drits, A. V., and Pasternak, A. F.: Characteristic features of body composition and metabolism in some in some
686 interzonal copepods, *Mar. Biol.*, 111, 199-205, 1991.

687 Glazer, B. T., Luther, G. W., Konovalov, S. K., Friederich, G. E., Trouwborst, R. E., and Romanov, A. S.: Spatial and
688 temporal variability of the Black Sea suboxic zone, *Deep-Sea Res. II* 53, 1756–1768, [doi:10.1016/j.dsr2.2006.03.022](https://doi.org/10.1016/j.dsr2.2006.03.022), 2006.

689 Heywood, K. J., Scrope-Howe, S., and Barton, E.D.: Estimation of zooplankton abundance from shipborne ADCP
690 backscatter, *Deep Sea Res.*, 38, 667-691, 1991.

691 Hirche, H., Barz, K., Ayon, P., and Schulz, J.: High resolution vertical distribution of the copepod *Calanus chilensis* in
692 relation to the shallow oxygen minimum zone off northern Peru using LOKI, a new plankton imaging system, *Deep Sea Res.*
693 Part I, 88, 63–73, <https://doi.org/10.1016/j.dsr.2014.03.001>, 2014.

694 Hofmann, H. and Peeters, F.: *In-Situ* optical and acoustical measurements of the buoyant cyanobacterium *P. Rubescens*:
695 Spatial and temporal distribution patterns. *PLoS ONE* 8(11):e80913, <https://doi.org/10.1371/journal.pone.0080913>, 2013.

696 Kiørboe, T., Andersen, A., Langlois, V. J., Jakobsen, H. H., and Bohr, T.: Mechanisms and feasibility of prey capture in
697 ambush-feeding zooplankton. *P. Natl. Acad. Sci. USA*, pnas.0903350106, <https://doi.org/10.1073/pnas.0903350106>, 2009.

698 Klyuvitkin, A. A., Ostrovskii, A. G., Novigatskii, A. N., and Lisitzin, A. P.: Multidisciplinary experiment on studying short-
699 period variability of the sedimentary process in the northeastern part of the Black Sea. *Doklady Earth Sciences*, 469, 771–
700 775, <https://doi.org/10.1134/S1028334X16070230>, 2016.

701 Konovalov, S. K., Murray, J. W., and Luther, G. W.: Basic processes of Black Sea biogeochemistry, *Oceanography*, 18, 24–
702 35, <https://doi.org/10.5670/oceanog.2005.39>, 2005.

703 Lavery, A. C., Wiebe, P. H., Stanton, T. K., Lawson, G. L., Benfield, M. C., and Copley, N.: Determining dominant
704 scatterers of sound in mixed zooplankton populations, *J. Acoust. Soc. Am.*, 122, 3304–3326.
705 <https://doi.org/10.1121/1.2793613>, 2007.

706 Lawson, G. L., Wiebe, P. H., Ashjian, C. J., Chu, D., and Stanton, T. K.: Improved parameterization of Antarctic krill target
707 strength models, *J. Acoust. Soc. Am.* 119, 232–242. <https://doi.org/10.1121/1.2141229>, 2006.

708 Morozov, A., Kuzenkov, O. A., Arashkevich, E. G.: Modelling optimal behavioural strategies in structured populations
709 using a novel theoretical framework, *Sci. Rep.*, 9:15020. <https://doi.org/10.1038/s41598-019-51310-w>, 2019.

710 Murray, J. W., Jannash, H. W., Honjo, S., Anderson, R. F., Reeburgh, W. S., Top, Z., Friederich, G.,E., Codispoti, L.,A., and
711 Izdar, E.: Unexpected changes in the oxic/anoxic interface in the Black Sea, *Nature*, 338, 411–413, 1989.

712 Mutlu, E.: Acoustical identification of the concentration layer of a copepod species, *Calanus euxinus*, *Mar. Biol.*, 142, 517–
713 523, <https://doi.org/10.1007/s00227-002-0986-3>, 2003.

714 Mutlu, E.: Diel vertical migration of *Sagitta setosa* as inferred acoustically in the Black Sea, *Mar. Biol.*, 149, 573–584,
715 <https://doi.org/10.1007/s00227-005-0221-0>, 2006.

716 Mutlu, E.: Compared studies on recognition of marine underwater biological scattering layers, *J. Appl. Biol. Sci.*, 1 (3), 113-
717 119, 2007.

718 Nikitin, V. N.: La distribution verticale du plancton dans la mer Noire. I. Copepoda et Cladocera. *Trudy Osoboi*
719 *Zoologicheskoi Laboratorii i Sevastopol'skoi Biologicheskoi Stantsii*, Ser. 2, No. 5–10, 93–140, 1926. (in Russian; abstract
720 in French).

721 Oguz, T., Turgul, S., Kideys, A.E., Ediger, V., and Kubilay, N.: Physical and biogeochemical characteristics of the Black
722 Sea, in: *The Sea, Volume 14B: The Global Coastal Ocean Interdisciplinary Regional Studies and Syntheses*, edited by
723 Robinson, A. R. and Brink, K. H. 1331-1369, ISBN 9780674021174, 2006.

724 Ostrovskii, A. G. and Zatsepin, A. G.: Short-term hydrophysical and biological variability over the northeastern Black Sea
725 continental slope as inferred from multiparametric tethered profiler surveys, *Ocean Dyn.*, 61, 797-806,
726 <https://doi.org/10.1007/s10236-011-0400-0>, 2011.

727 Ostrovskii, A. G. and Zatsepin, A. G.: Intense ventilation of the Black Sea pycnocline due to vertical turbulent exchange in
728 the Rim Current area, *Deep Sea Res. I* 116, 1–13, <https://doi.org/10.1016/j.dsr.2016.07.011>, 2016.

729 Ostrovskii, A. G., Zatsepin, A. G., Solovyev, V. A., and Soloviev, D. M.: The short timescale variability of the oxygen
730 inventory in the NE Black Sea slope water, *Ocean Sci.*, 14, 1567–1579, <https://doi.org/10.5194/os-14-1567-2018>, 2018.

731 Ostrovskii, A. G., Zatsepin, A. G., Soloviev, V. A., Tsibulsky, A. L., and Shvoev, D. A.: Autonomous system for vertical
732 profiling of the marine environment at a moored station, *Oceanology*, 53, 233–242, <https://doi.org/10.1134/S0001437013020124>, 2013.

734 Petipa, T. S., Sazhina, L. I., and Delalo, E. P.: Vertical distribution of zooplankton in the Black Sea in relation to the
735 hydrological conditions, *Dokl. Akad. Nauk SSSR*, 133, 964–967, 1960 (in Russian).

736 Pezacki, P. D., Gorska, N., and Soloviev, V.: An acoustic study of zooplankton diel vertical migration in the Black Sea,
737 *Hydroacoustics*, 20, 139-148, 2017.

738 Roberts, P. L. D. and Jaffe, J. S.: Multiple angle acoustic classification of zooplankton, *J. Acoust. Soc. Am.*, 121, 2060–
739 2070, <https://doi.org/10.1121/1.2697471>, 2007.

740 Roberts, P. L. D. and Jaffe, J. S.: Classification of live, untethered zooplankton from observations of multiple-angle acoustic
741 scatter, *J. Acoust. Soc. Am.*, 124, 796–802. <https://doi.org/10.1121/1.2945114>, 2008.

742 Sakinan, S. and Gücü, A. C.: Spatial distribution of the Black Sea copepod, *Calanus euxinus*, estimated using multi-
743 frequency acoustic backscatter, *ICES J. Mar. Sci.*, 74, 832–846, <https://doi.org/10.1093/icesjms/fsw183>, 2016.

744 Sartoris, F. J., Thomas, D. N., Cornils, A., and Schnack-Schiel, S. B.: Buoyancy and diapause in Antarctic copepods: the role
745 of ammonium accumulation. *Limnol. Oceanogr.* 55, 1860–64, <https://doi.org/10.4319/lo.2010.55.5.1860>, 2010.

746 Schründer, S., Schnack-Schiel, S. B., Auel, H., Sartoris, F. J.: Control of diapause by acidic pH and ammonium
747 accumulation in the haemolymph of Antarctic copepods. *PLOS ONE*, 8:e77498,
748 <https://doi.org/10.1371/journal.pone.0077498>, 2013.

749 Smeti, H., Pagano, M., Menkes, C., Boissieu, F., Lebourges-Dhaussy, A., Hunt, B. P. V., Allain, V., Rodier, M., Kestenare,
750 E., and Sammari, C.: Spatial and temporal variability of zooplankton off New Caledonia (Southwestern Pacific) from
751 acoustics and net measurements, *J. Geophys. Res. C: Oceans*, 120, 2676–2700, <https://doi.org/10.1002/2014JC010441>, 2015.

752 Solovyev, V. A., Shvoev D. A., and Ostrovskii, A. G.: Metadata of Aqualog profiler surveys in 2013-2021, Research Gate,
753 DOI: <https://doi.org/10.13140/RG.2.2.27548.62084>, last access: 12 June 2021.

754 Stanton, T. K., Chu, D., Wiebe, P. H., and Clay, C. S.: Average echoes from randomly oriented random-length finite

755 cylinders: Zooplankton models. J. Acoust. Soc. Am., 94, 3463–3472, <https://doi.org/10.1121/1.407200>, 1993.

756 Stanton, T. K. and Chu, D.: Review and recommendations for the modelling of acoustic scattering by fluid-like elongated
757 zooplankton: euphausiids and copepods, ICES J. Mar. Sci., 57, 793–807, <https://doi.org/jmsc.1999.0517>, 2000.

758 Stanton, T. K., Wiebe, P. H., Chu, D., Benfield, M., Scanlon, L., Martin, L., and Eastwood, R. L.: On acoustic estimates of
759 zooplankton biomass, ICES J. Mar. Sci., 51, 505–512. 1994.

760 Stefanova, K. and Marinova, V.: Zooplankton distribution and sound scattering layers in the Bulgarian Black Sea area – A
761 case study Cape Galata Transect, Intl. J. Appl. Eng. Res., 10(15), 35998-36003, 2015.

762 Svetlichny, L. S., Hubareva, E. S., and Arashkevich, E. G.: Physiological and behavioural response to hypoxia in active and
763 diapausing stage V copepodites of *Calanus euxinus*, in: Arch. Hydrobiol. Spec. Issues, Advanc. Limnol., Evolutionary and
764 ecological aspects of crustacean diapause edited by Brendonck, L., de Meester, L., and Hairston, 52, 507-519, 1998.

765 Svetlichny, L. S., Hubareva, E. S., Erkan, F., and Gucu, A. C.: Physiological and behavioral aspects of *Calanus euxinus*
766 females (Copepoda: Calanoida) during vertical migration across temperature and oxygen gradients, Mar. Biol., 137,
767 963–971, 2000.

768 Svetlichny, L., Gubareva, E., and Arashkevich, E.: Effect of oxygen concentration on energy metabolism in the migrating
769 and diapausing copepods *Calanus euxinus* in the Black Sea, Oceanology, 42, 670–676, 2002.

770 Svetlichny, L. S., Kideys, A. E., Hubareva, E. S., Besiktepe, S., and Isinibilir, M.: Development and lipid storage in *Calanus*
771 *euxinus* from the Black and Marmara seas: variabilities due to habitat conditions, J. Mar. Syst., 59, 52–62,
772 <https://doi.org/10.1016/j.jmarsys.2005.09.003>, 2006.

773 Svetlichny, L. S., Yuneva, T. V., Hubareva, E. S., Schepkina, A. M., Besiktepe, S., Kideys, A. E., Bat, L., and Sahin F.:
774 Development of *Calanus euxinus* during spring cold homothermy in the Black Sea, Mar. Ecol. Prog. Ser., 374, 199–213,
775 <https://doi.org/10.3354/meps07740>, 2009.

776 Tutasi, P. and Escribano, R.: Zooplankton diel vertical migration and downward C flux into the oxygen minimum zone in the
777 highly productive upwelling region off northern Chile, Biogeosciences, 17, 455–473, <https://doi.org/10.5194/bg-17-455->
778 2020, 2020.

779 Vinogradov, M. E., Flint, M. V., and Shushkina, E. A. Vertical distribution of mesozooplankton in the open Black Sea, Mar.
780 Biol., 89, 95–107, 1985.

781 Vinogradov, M. E. and Nalbandov, Yu. R.: Influence of water density on the distribution of physical, chemical, and
782 biological characteristics of the Black Sea pelagic ecosystem, Oceanology, 30, 769–777, 1990.

783 Vinogradov, M. E., Arashkevich, E. G., and Ilchenko, S. V.: The ecology of the *Calanus ponticus* population in the deeper
784 layer of its concentration in the Black Sea, J. Plankton Res., 14, 447-458, 1992.

785 Visser, A. W. and Jónasdóttir S. H.: Lipids, buoyancy, and the seasonal vertical migration of *Calanus finmarchicus*, Fish.
786 Oceanogr., 8 (Suppl 1), 100–106, 1999.

787 Wiebe, P. H., Stanton, T. K., Greene, C. H., Benfield, M. C., Sosik, H. M., Austin, T. C., Warren, J. D., and Hammar, T.:
788 BIOMAPER-II: an integrated instrument platform for coupled biological and physical measurements in coastal and oceanic
789 regimes, *IEEE J. Oceanic Eng.*, 27, 3, 700-716, 2002, <https://doi.org/10.1109/JOE.2002.1040951>, 2002.

790 Wishner, K. F., Seibel, B., and Outram, D.: Ocean deoxygenation and copepods: coping with oxygen minimum zone
791 variability, *Biogeosciences*, 17, 2315–2339, doi: 10.5194/bg-17-2315-2020, 2020.

792 Yakushev, E. V., Podymov, O. I., and Chasovnikov, V. K.: Seasonal changes in the hydrochemical structure of the Black
793 Sea redox zone, *Oceanography*, 18, 48–55, <https://doi.org/10.5670/oceanog.2005.41>, 2005.

794 Yunev, O. A., Carstensen, J., Stelmakh, L. V., Belokopytov, V. N., and Suslin, V. V.: Reconsideration of the
795 phytoplankton seasonality in the open Black Sea, *Limnology Oceanogr.: Lett.*, 6, 51-59, doi: 10.1002/lo12.10178, 2020.