

Freshwater in the Arctic Ocean 2010-2019

Amy Solomon^{1,2}, Céline Heuzé³, Benjamin Rabe⁴, Sheldon Bacon⁵, Laurent Bertino⁶, Patrick Heimbach⁷, Jun Inoue⁸, Doroteaciro Iovino⁹, Ruth Mottram¹⁰, Xiangdong Zhang¹¹, Yevgeny Aksenov⁵, Ronan McAdam⁹, An Nguyen⁷, Roshin P. Raj⁶, Han Tang¹¹

5 ¹Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, USA

²Physical Sciences Laboratory, NOAA Earth System Research Laboratory, Boulder, Colorado, USA

³Department of Earth Sciences, University of Gothenburg, Gothenburg, Sweden

⁴Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany

⁵National Oceanography Centre, Southampton, UK

10 ⁶Nansen Environmental and Remote Sensing Center and Bjerknes Center for Climate Research, Bergen, Norway

⁷University of Texas, Austin, Texas, USA

⁸National Institute of Polar Research, Tachikawa, Japan

⁹Centro Euro-Mediterraneo per i Cambiamenti Climatici, Bologna, Italy

¹⁰Danish Meteorological Institute, Copenhagen, Denmark

15 ¹¹University of Alaska, Fairbanks, Alaska, USA

Correspondence to: Amy Solomon (amy.solomon@noaa.gov)

Abstract. The Arctic climate system is rapidly transitioning into a new regime with a reduction in the extent of sea ice, enhanced mixing in the ocean and atmosphere, and thus enhanced coupling within the ocean-ice-atmosphere system; these physical changes are leading to ecosystem changes in the Arctic Ocean. In this review paper, we assess one of the critically important aspects of this new regime, the variability of Arctic freshwater, which plays a fundamental role in the Arctic climate system by impacting ocean stratification and sea ice formation. Liquid and solid freshwater exports also affect the global climate system, notably by impacting the global ocean overturning circulation. We assess how this budget has changed relative to the 2000-2010 period. We include discussions of processes not included in all previous assessments, such as runoff from the Greenland Ice Sheet, the role of snow on sea ice, and vertical redistribution. Notably, the sea ice cover has become more seasonal and more mobile, the mass loss of the Greenland Ice Sheet has increased in the 2010s (particularly in the west, north, and south regions), and imported warm, salty Atlantic waters has shoaled. We show that the trend in Arctic freshwater content in the 2010s has stabilized relative to the 2000s, potentially due to an increased compensation between a freshening of the Beaufort Gyre and a reduction in freshwater in the rest of the Arctic Ocean. However, large inter-model spread across the ocean reanalyses and uncertainty in the observations used in this study prevent a definitive conclusion about the degree of this compensation.

1 Freshwater in the Arctic Ocean

Rapid changes in the Arctic climate system are impacting marine resources and industries, coastal Arctic environments, and large-scale ocean and atmosphere circulations. The Arctic climate system is rapidly transitioning into a new regime with a reduction in the extent of sea ice (Stroeve and Notz, 2018), a thinning of the ice cover (Kwok, 2018), a warming and freshening

35 of the Arctic Ocean (Timmermans and Marshall, 2020), enhanced mixing in the ocean and atmosphere and enhanced coupling
within the ocean-ice-atmosphere system (Timmermans and Marshall, 2020); these physical processes are leading to cascading
changes in the Arctic Ocean ecosystems (Bluhm et al., 2015; Polyakov et al., 2020). The emergent properties of this new
regime, termed the “New Arctic” (Jeffries et al., 2013), are yet to be determined since altered feedback processes are expected
to further impact upper ocean heat and freshwater content, atmospheric and oceanic stratification, the interactions between
40 subsurface/intermediate warm waters and surface cold and fresh layer, among other properties. In this review we assess one
of the critically important aspects of this new regime, the variability of Arctic freshwater.

Freshwater in the Arctic Ocean plays a critical role in the global climate system by impacting large-scale overturning ocean
circulations (Sévellec et al., (2017), see Figure 1 showing basins and upper circulation), ocean stratification that determines
sea ice growth, biological primary productivity (Ardyna and Arrigo, 2020; Lewis et al., 2020), and ocean mixing (Aagaard
45 and Carmack, 1989); and emerging freshwater regimes that couple variability in land, atmosphere, and ocean systems (e.g.,
Jeffries et al., 2013; Wood et al., 2013), among other impacts. Arctic Ocean freshwater is a balance between:

- sources (Pacific and Atlantic oceanic inflow, precipitation, river runoff, ice sheet discharge, sea ice melt) (Aagaard and
Woodgate, 2001; Serreze et al., 2006; Bamber et al., 2012),
- sinks (sea ice growth, evaporation, liquid and solid transport through oceanic gateways) (Aagaard and Carmack, 1989;
50 Rudels et al., 1994; Serreze et al., 2006; Haine et al., 2015),
- redistribution between Arctic basins (e.g., Timmermans et al., 2011; Morison et al., 2012; Proshutinsky et al., 2015).

These processes are not necessarily independent and are largely driven by atmospheric variability both within the Arctic and
from lower-latitudes.


Oceanographers have long been accustomed to the use of "freshwater" as an identifiable and separable component of seawater,
55 either as a freshwater volume or a freshwater flux component of a seawater volume or flux. It usually manifests as a small
fraction of the seawater volume or flux, where the fraction takes the form $(\delta S/S_{ref})$, and where $\delta S = S - S_{ref}$ is the deviation of
the seawater salinity S from a reference value S_{ref} . However, scientists' familiarity with this usage perhaps disguises the fact
that it is an arbitrary construct: the existence of such a concept as "reference salinity" and values attributed to it are not
rigorously mathematically and physically defined. Indeed, this state of affairs prompted Schauer & Losch (2019) to write a
60 paper entitled "*Freshwater' in the ocean is not a useful parameter in climate research*", in which they argue their preference
for the uniquely-defined salt budget as an absolute and well-posed physical quantity. The significant freshwater flux
differences that can arise from use of different reference salinities are illustrated and quantified by Tsubouchi et al. (2012), as
well as by Schauer and Losch (2019).

In contrast, Bacon et al. (2015) observed that there is, in fact, one place in the ocean where a true freshwater flux occurs without
65 ambiguity, and that is at the surface, where freshwater is exchanged between ocean and atmosphere (via precipitation and
evaporation) and where the ocean receives freshwater input from the land (as river or other runoff). They recognize that a
surface flux requires definition of a surface area. They then use a time-varying ice and ocean control volume (or "budget")
approach, combined with mass and salinity conservation, to generate a closed mathematical expression where the surface
freshwater flux is given by the sum of three terms: (i) the divergence of the (scaled) salinity flux around the boundary of the
70 control volume, (ii) the change in total (ice and ocean) seawater mass within the control volume (or change in mass storage),
and (iii) the (scaled) change in mass of salt within the control volume (the change in salinity storage). The "scaling" term that
emerges from the mathematics performs the same function as the traditional reference salinity, but in its place is the control
volume's ice and ocean boundary mean salinity, which has uncomfortable implications in that it can vary in time and with
boundary geography. This is a consequence of the nature of the calculation, which quantifies surface freshwater fluxes.
75 Carnack et al. (2016) interpret the Arctic case thus: the surface freshwater flux is what is needed to dilute all the ocean inflows
to become the outflows, allowing for interior storage changes. An exactly equivalent interpretation is that surface freshwater
fluxes and the relatively fresh Bering Strait sea water inflow combine to dilute the relatively saline Atlantic water inflow,
which then become the outflows (allowing for storage) – where "relatively" means relative to the boundary mean salinity.

Is "ocean freshwater flux" purely a mirage, therefore? Forryan et al. (2019) pursue the surface freshwater flux approach, noting
80 that (as is well known, e.g., Östlund and Hut, 1984) evaporation and freezing are distillation processes that leave behind a
geochemical imprint via oxygen isotope anomalies on the affected freshwater in the sea ice and seawater. In the case of
evaporation, distillation (here, isotopic fractionation) preferentially removes lighter oxygen isotopes from seawater, leaving
behind in the seawater a proportion of heavier isotopes. The lighter isotopes that are now in the atmosphere return to the land
or sea surface as precipitation. Those falling on land can (eventually) transfer from land to sea by river runoff or by other
85 glacial processes, or by further cycles of evapo-transpiration and precipitation. For sea ice, the ice contains the lighter isotopes
while heavier isotopes are contained in the brine that drains out of the ice during freezing, to re-enter the ocean. The
isotopically-lighter meteoric fractions are used to quantify freshwater that originates from the atmosphere (directly or
indirectly), and the isotopically-heavier fractions similarly quantify the signal of brine rejected from sea ice, and thereby the
amount of ice formed from that seawater.

90 Forryan et al. (2019) show that, within uncertainties, the geochemical approach returns the same surface freshwater flux as the
budget approach. However, we are still left with a conundrum, as per the argument of Schauer and Losch (2019), in that the
formal definition of a fixed ice and ocean "reference salinity" remains elusive. Tsubouchi et al. (2018) find that in practice,
time variability in ice and ocean boundary mean salinity (for a fixed boundary) has no significant impact on surface freshwater
flux calculation. Since this is a review of existing literature and in light of established practice, we continue to employ here the
95 "traditional" approach to freshwater flux calculation by use of a fixed reference salinity.

Freshwater input to the Arctic Ocean is almost entirely confined to the upper water column and comes in the form of continental runoff, waters of Pacific origin, various coastal currents and precipitation. Freshwater input from the Greenland ice sheet has two subsurface contributions: (i) melting from calved icebergs (Moon et al., 2017), and (ii) submarine melt rates which may produce a freshwater plume that becomes neutrally buoyant below the surface (Straneo et al., 2011). The upper Arctic Ocean hence is characterized by salinity values lower than that of the inflow of waters of largely Atlantic origin through the Fram Strait and the Barents Sea opening. The result is an extremely stratified Arctic Ocean, with a shallow seasonal mixed layer on average less than 100 m thick and a very fresh halocline that is the mixture result of all the inflows (McLaughlin et al., 1996; Rudels et al., 2004). As noted by Rudels et al. (2004), the term “halocline” is misleading yet common practice. In the rest of the world ocean, halocline denotes the depth range where salinities abruptly change as two water masses mix; in the Arctic, the halocline is a cold and fresh water mass. From the bottom of the halocline (ca 300 m depth) to ca 800 m sits the so-called Atlantic layer, which is comparatively warm and salty, and below this is the Arctic Ocean deep waters (Aagaard et al., 1985; Rudels, 2012). Vertical fluxes of freshwater are generally low due to this strong stratification and very low vertical turbulent mixing / diffusion (e.g., Fer, 2009). The reviews of Carmack et al. (2016) and Haine et al. (2015) confirm the picture above; hence, they mainly considered the Arctic freshwater budget in the near-surface layers. This current study expands on their work and describes the processes impacting the vertical (re)distribution of freshwater throughout the entire water column.




115 **Figure 1: Map of the Arctic Ocean with names of major basins and shelf seas, and ocean circulation features: major river and Pacific inflow (cyan and turquoise) and surface outflows (purple), 2020 minimum sea ice edge (yellow), cold and fresh upper ocean circulations (Polar Surface Water and halocline, blues), and warm and salty Atlantic water circulation (red). Areas shallower than 1000 m are referred to as shelf areas in the text. BG stands for Beaufort Gyre; TPD, Trans-polar drift; BC, Barrow Canyon; CAA, Canadian archipelago; SAT, St Anna Trough.**

120 Assessments of Arctic freshwater for the 2000-2010 period relative to 1980-2000 were completed as part of the WCRP/IASC/AMAP Arctic Freshwater Synthesis (Prowse et al., 2015; Carmack et al., 2016; Vihma et al., 2015) and the Arctic-Subarctic Ocean Fluxes program (Haine et al., 2015). These projects found that liquid freshwater increased by 25% (5000 km³) in the Beaufort Gyre; the Beaufort High was stronger than normal with higher sea level, a deeper halocline, stronger anticyclonic flow, and stronger Trans-polar Drift (Proshutinsky et al., 2009; McPhee et al., 2009; Rabe et al., 2011; Haine et al., 2015). However, estimates of fluxes through the ocean gateways were either too uncertain or insignificantly different, leading to speculation that freshwater had accumulated in the Arctic Ocean, which, if released through the Fram Strait could

125 significantly impact the global climate system through changes in the global ocean overturning circulation. In these studies,
processes such as the redistribution of freshwater between basins, vertical redistribution due to turbulent mixing, and discharge
from the Greenland Ice Sheet (among other processes) were not taken into account, leading to uncertainty in this speculation.
For example, Rabe et al. (2011) and Morison et al. (2012) found that from the early to late 2000s, the increased deep basin
freshwater content in the Beaufort Gyre was largely balanced by a decrease in the rest of the Arctic Ocean.

130 The observed Beaufort Gyre freshening is illustrated in Figure 2, which shows 1993-2019 annual mean Arctic Ocean
freshwater from seven state-of-the-art global ocean reanalyses (ORAs, see Table 1 for a description of the models used in this
study). Significant freshening in the Beaufort Gyre is seen in 2010-2017 means minus 2000-2010 means in six ORAs (Figure
2b, not including ASTE_R1). However, this freshening is partly compensated by a reduction in freshwater in the rest of the
Arctic Ocean (Figure 2b,c). This compensation increases in 2010-2018 compared to 2000-2010, which flattens the total Arctic
Ocean freshwater trend when extended to 2019 (Figure 2a). However, there is a significant spread in estimates of freshwater
135 content in the Beaufort Gyre and the rest of the Arctic Ocean (Figure 2d), which prevents a definitive estimate of the degree
of this compensation. This highlights the need to be able to estimate the redistribution of freshwater when assessing changes
in Arctic Ocean freshwater, as well as the recent reduction in total Arctic Ocean freshening relative to the 2000-2010 period.

In this review we assess to what extent the 2010-2019 freshwater budget has changed relative to the 2000-2010 period. This
study is not meant to be a comprehensive assessment of all processes that contribute to Arctic freshwater. Instead, we focus
140 on specific aspects that provide insight into how the variability has changed since 2010 and the role of processes not considered
in previous assessments.



145 **Figure 2: Ordered counter-clockwise: A) Time series of annual freshwater content integrated from 70-90°N and down to the 34.8 isohaline for the period 1993-2019 from 7 ORAs (in 10^3 km^3). Multi-model mean shown in red, darker red indicates all 7 ORAs included. B) Difference (in m) between 2010-2017 and 2000-2010 means in 6 ORAs (not including ASTE_R1). C) Multi-model mean of differences (in m) shown in (B). D) Annual Freshwater content separated into contribution from Beaufort Gyre (blue) and the rest of the Arctic Ocean (red). Lower right figure shows the multi-model mean with ± 1 standard deviation shown with shading. The Beaufort Gyre is defined as 70-80°N, 128-180°W to be consistent with the satellite estimates below.**

	Institution	Horiz Resol.	Vert. Resol.	Fluxes	Atmo. Forcing	Ocean-Sea Ice Model	Observations assimilated	Reference
C-GLORSv7	CMCC	0.25°	L75	CORE	ERA-I	NEMO3.2-LIM2	EN3v2a, SIC, PIOMASSIT, T, S, SLA, SST, SSS, MDT	Storto & Masina (2016)
FOAM	UK MetOffice	0.25°	L75	CORE	ERA-I	NEMO3.2-CICE	SIC, T, S, SLA, SST	Blockley et al. (2014)
GLORYS2V4	CMEMS	0.25°	L75	CORE	ERA-I	NEMO3.1-LIM2	CORA4v1, SST, SLA, SIC, runoff (Dai and Trenberth)	Garric et al. (2017)
ORAS5	ECMWF	0.25°	L75	CORE + wave forcing	ERA-I until 2014 ECMWF NWP after 2014	NEMO3.4-LIM2	EN4, XBTs, CTDs, SLA, SIC, SST	Zuo et al. (2019)
ASTE_R1	U. Texas Austin	0.3°	L50	CORE	Adjusted JRA55	MITgcm	SST, SLA, MDT, SIC, insitu Argo, ITP, ICES, XBT, CTD, T/S at mooring arrays at Arctic gateways	Nguyen et al. (2021)
ECDA CM2.1	NOAA - GFDL	1.0°	L50	Bulk O-M	Coupled DA using atmos. model AM2	MOM4	WOD T/S, XBT, CTD, SST analysis, moorings, ARGO	Chang et al. (2013)
SODA3.3.2	U. Maryland	0.25°	L50	COARE4	MERRA2	MOM5-SIS1	WOD T/S, ICOADS and satellite SST, river runoff (Dai), Greenland discharge (Bamber)	Carton et al. (2018)

Table 1: Global ocean reanalyses used in this study.

1.1 Arctic freshwater estimates from in-situ and satellite measurements

1.1.1 Satellite measurements

A major challenge in the retrieval of freshwater fluxes in the Arctic Ocean is associated with the lack of availability of in-situ observations. Direct measurements are non-homogenous in both time and space and rely on spatial as well as temporal interpolation resulting in large uncertainties. The ability to estimate freshwater content of the Arctic region indirectly from satellite observations is a major breakthrough. The methodology which exploits the satellite derived ocean mass change and satellite altimeter data is detailed in Giles et al. (2012), Morison et al. (2012), and Armitage et al. (2016). Our understanding of the Earth's gravity field has improved considerably during the recent decade, thanks to the Gravity Recovery and Climate Experiment (GRACE) mission launched in 2002. GRACE is the only satellite mission designed to be directly sensitive to mass changes by means of gravity. The variability in spatiotemporal characteristics of the Earth's gravitational field resulting in very small deviations in the separation between the two satellites of the GRACE mission are measured with micrometer precision and are used to infer the Earth's gravity field, which can then be used to estimate changes in ocean mass (Peralta-Ferriz et al., 2014; Armitage et al., 2016). Here we use the latest Release-06 gridded GRACE ocean mass products from the Jet Propulsion Laboratory (Watkins et al., 2015). Satellite radar altimeters on the other hand can retrieve sea surface heights in the open ocean with variable precision depending on the number of flying altimeters and has been uninterrupted since 1993. CryoSat-2, launched in 2010, is a satellite altimeter that provides coverage up to 88°N with much better spatial resolution than before. However, constructing precise altimeter derived sea level data in the Arctic Ocean is a challenge, mainly due to the changing sea-ice cover which affects the range correction.


Satellites can monitor some important pieces of the Arctic freshwater puzzle. Here, we use the state-of-the-art sea level product produced as part of the recently concluded climate change initiative (CCI) project (Sea level budget closure; Horwath et al., 2020) of the European Space Agency. This Arctic sea level product (DTU/TUM SLA record; Rose et al., 2019) is the first one which includes a physical retracker (ALES+) for retrieving the specular waveforms from open leads in the sea cover. The sea state bias corrected using ALES+ improves the sea level estimates of the region (Passaro et al., 2018). The latest version (v3.1) of the DTU/TUM SLA record is a complete reprocessing of the former DTU Arctic sea level product (Andersen et al., 2016) by dedicated Arctic retracking. The current study thus takes advantage of the state-of-the-art satellite datasets to study the freshwater content of the region following Giles et al. (2012) and Armitage et al. (2016).

Time series from 2002 to 2018 using GRACE-derived ocean bottom pressure (OBP) anomalies (<https://podaac.jpl.nasa.gov/GRACE>) and satellite altimeter data provide insights into the redistribution of freshwater in the Arctic Ocean (Figure 3). While initial results from GRACE suggest an overall OBP decrease caused by a fresher Arctic surface (Morison et al., 2007), results on the now-longer time series show more complex interannual variability, in agreement with modelling data (e.g., de Boer et al., 2018). Figure 3a (red line) shows that freshwater content increases in the Beaufort Gyre during the time-period 2002-2010, followed by a stabilizing phase where the increase flattens out. However, including

freshwater content outside of the Beaufort Gyre (blue line in Figure 3a, defined as the region contoured in Figure 3b) results
185 in a reduction in freshwater content during the time-period 2010-2016, indicating increased compensation between freshwater
content in the Beaufort Gyre and outside the Beaufort Gyre after 2009. Raj et al. (2020) noted a similar signature in the
altimeter derived sea surface height anomaly and the halosteric component of the sea surface height anomaly and attributed it
to the change in the dominant atmospheric forcing over the Arctic, which changed from the Arctic dipole pattern to the Arctic
Oscillation respectively during the time-periods prior-to and after 2010. These results are qualitatively consistent with
190 estimates in Figure 2 using the ocean reanalyses. In addition, Figure 2c shows that the regions not included in Figure 3 make
only small contributions to the time series In Figure 2.

1.1.2 In-situ measurements

Figure 3a includes estimates of Arctic freshwater content from in-situ hydrographic observations (black line). The timeseries
of freshwater content for the whole basin to the 34 isohaline is extended from Rabe et al. (2014). Details of the mapping
195 procedure and the distribution of hydrographic stations until 2012 is given in Rabe et al. (2014). Further data is based on the
data sources listed in Table 2. Interestingly, the Arctic satellite and in-situ time series in Figure 3a are relatively consistent
before 2009, but do not show the same variability after 2009. This difference may stem from the lack of data coverage in the
in-situ measurements, the different regions used in the time series and the choice of time period for the mean used to obtain
anomalies. The satellite time series uses the region contoured in Figure 3b and the in-situ time series uses observations within
200 the basin excluding the shelves, indicating a good part of the difference after 2009 may be due to the contribution by the rest
of the basin outside the Beaufort Gyre. In addition, the in-situ time series is not well constrained after 2013. Due to the method
the annual values are biased towards prior years near the end of the timeseries, as the mapping analysis only includes data up
to 2015.



205




Figure 3: Anomalies of freshwater content (in 10^3 km^3) from satellite sea-surface height data analysis and GRACE OBP data and from objectively mapped in-situ hydrographic observations. A) Annual mean time series of freshwater content from satellite measurements in the Beaufort Gyre (red) and Arctic region (blue), and Arctic basin using in-situ hydrographic observations (black). B) Difference between 2010-2017 and 2002-2010 freshwater content means from satellite measurements, in units of 10^3 km^3 . Anomalies in (A) are relative to the corresponding mean of the period 2003-2006 in each time series using a reference practical salinity of 35 and a layer from the surface to the 34 isohaline. The Beaufort Gyre region is defined as 70-80°N, 120-180°W. The time series are calculated using observations from the Arctic Ocean with a water depth deeper than 500 m and a cut-off at 82°N north of the Fram Strait for the in-situ estimates, and the contoured region shown in (B) for the satellite estimates. The black line in (A) is an update of the time series in Rabe et al. (2014), the additional data used is listed in Table 2.

Expedition, Project	Year(s)	Platform	Source URL or contact
Beaufort Gyre Project	2012-2013	various ships	http://www.whoi.edu/beaufortgyre/
NPEO	2012-2014	Airborne and ice-based	ftp://psc.apl.washington.edu/NPEO Data Archive/NPEO Aerial CTDs/
WHOI	2012-2015	ITP	http://www.whoi.edu/itp
PS86	2014	RV Polarstern	http://doi.pangaea.de/10.1594/PANGAEA.853768 (Vogt et al., 2015)
PS87	2014	RV Polarstern	http://doi.pangaea.de/10.1594/PANGAEA.853770 (Roloff et al., 2015)
PS94	2015	RV Polarstern	https://doi.org/10.1594/PANGAEA.859558 (Rabe et al., 2015)

Table 2: Sources of salinity data used in the objective analysis to derive the black curve in Figure 3. The listed data sources are for the data used in addition to the data described in Rabe et al. (2014) and published in Rabe et al. (2014b). Abbreviations: ITP -- Ice-Tethered Profiler, NPEO -- North Pole Environmental Observatory, WHOI -- Woods Hole Oceanographic Institution.

2 Changes in Arctic Freshwater Sources and Sinks

The most recent estimates of Arctic freshwater sources and sinks have been developed by Østerhus et al. (2019), Haine et al. (2015), Prowse et al. (2015), Carmack et al. (2016), and Vihma et al. (2016). Only Østerhus et al. (2019) covers a more recent period through 2015. One issue is that not all these estimates use the same reference salinity; a discussion of freshwater versus salt transports and reference salinities is provided in Bacon et al. (2015), Schauer and Losch (2019), and Tsubouchi et al. (2018). Another more recent development over the last decade is the inclusion of freshwater fluxes from the Greenland Ice Sheet (GIS) and smaller Arctic glaciers and ice caps (GICs) into these basins. GIS FW fluxes were estimated by Bamber et al. (2012) and updated by Bamber et al. (2018) to include GIC FW fluxes (see also Dukhovskoy et al., 2019).

2.1 River Discharge

Observations suggest a linkage between the Arctic Oscillation (AO) and the North American (mainly Mackenzie River) runoff pathways (Yamamoto-Kawai et al., 2009; Fichot et al., 2013). There has been a shift from a rather direct outflow via the Canadian Arctic Archipelago (CAA) in early 2000s to a northward pathway into the Beaufort Gyre around 2006, coinciding with a change to a strongly positive AO. In addition, for high AO indices, river runoff entering the Eurasian shelves is mainly transported into the Canada Basin, while for low AO indices, the transport is mainly towards the Fram Strait by a strengthened transpolar drift (Morison et al., 2012; Alkire et al., 2015).

Observations of runoff rates for Eurasian rivers are available since 1936, and for North American rivers since 1964 (Shiklomanov et al., 2021). There has been a decline since about 1990 in the total gauged area, by ~10%, in Siberia and Canada (Shiklomanov et al., 2021), due to the closure or mothballing of gauging stations. Regardless, only the most important rivers are gauged: knowledge of net (continent-scale) river discharge rates require estimation of the substantial ungauged runoff fraction, typically one third of the total. The long-term, multi-decadal, gauged annual mean runoff rates are given by Shiklomanov et al. (2021) as $1800 \text{ km}^3 \text{ yr}^{-1}$ (Eurasia, 1936-2015) and $1150 \text{ km}^3 \text{ yr}^{-1}$ (North America, 1964-2015), for a total of $2950 \text{ km}^3 \text{ yr}^{-1}$. Shiklomanov et al. (2021) also note the increase (with uncertainties) in these records as 2.9 ± 0.4 (Eurasia, using 1935-2015 period) and 0.7 ± 0.3 (North America, using 1964-2015 period) $\text{km}^3 \text{ yr}^{-2}$. The significant Eurasian trend is of order 15% per century. However, the weakly-significant North American trend over the shorter period disguises an apparent signal of multi-decadal variability similar to that observed by Florindo-Lopez et al. (2020), who suggest it to be part of the evidence for much wider-area atmospheric and oceanic teleconnections.


2.2 Precipitation and Atmospheric Moisture Transport

Precipitation over the Arctic is the main source of freshwater into the Arctic Ocean, including that from river discharge from the large continental drainage basins. Precipitation is largely driven by atmospheric moisture transport. Based on a mass-corrected atmospheric moisture transport dataset, Zhang et al. (2013) found that the observed increase in the Eurasian Arctic river discharge was decisively driven by an enhanced poleward atmospheric moisture transport into the river basins. Using

the same dataset, Villamil-Otero et al. (2017) also found a continual enhancement of the poleward atmospheric moisture transport across 60°N into the Arctic Ocean from the 1950s to the mid 2010s. An update of the transport using ERA5 reanalysis shows a continuation of the enhancement across 60°N (Figure 4). Nygard et al. (2020) also found an increase in poleward moisture transport from 1979-2018 using the ERA5 data. Interestingly, they also found that evaporation shows a negative trend due to suppression by the horizontal moisture transport.

Corresponding to the enhanced atmospheric moisture transport, the large-scale atmospheric circulation may play a dynamic driving role. A statistical analysis indicates a temporally-varying relationship between the annual moisture transport and the annual mean Arctic Oscillation (AO, Thompson and Wallace, 1998; Figure 4), showing a negative and a positive correlation before and after 2000. In the 1990s, the variability of and changes in the atmospheric circulation was mainly characterized by the AO. The positive phase of the AO indicates a strengthening of the westerlies, transporting the atmospheric moisture to the Eurasian continent and leading to an increase in precipitation over the landmass (e.g., Kryzhov and Gorelits, 2015). However, although a positive correlation occurs between the transport and AO after 2000, the AO phase transition lagged the transport variability and may not show main driving or modulating role. In fact, during this time period, the atmospheric circulation spatial pattern has experienced a radical change in particular during winter seasons, as revealed in Zhang et al. (2008). This changed spatial pattern, named the Arctic Rapid change Pattern (ARP), exhibits a predominant role in driving the poleward moisture transport (Zhang et al., 2013). This driving role can also be manifested by a poleward extension and intensification of the Icelandic Low in the negative ARP phase. Considering that temporal-varying features of AO and the seasonal preference of the emergence of the spatially transformed ARP, the dynamic driving role of the atmospheric circulation needs to be further investigated. In addition, synoptic-scale analysis also suggested the propagation of intense storms into the Arctic played an important role in the enhanced poleward moisture transport and resulting increase in precipitation (e.g., Villamil-Otero et al., 2017; Webster et al., 2019).

Much of the precipitation in the Arctic falls as snow but projections show an increasing amount of rain as the climate warms. This appears to have been tentatively observed in Greenland (Doyle et al., 2015; Haine et al., 2015; Boisvert et al., 2018; Oltsmanns et al., 2019), where consequences for surface melt, surface runoff, and ice dynamics from increased rainfall over the ice sheet have been observed (e.g., Lenaerts et al., 2019). Similarly, Webster et al. (2019) note an increased frequency of rain on sea ice. Unfortunately, precipitation is notoriously difficult to measure, particularly in the solid phase, and as with other observations in the Arctic, reliable observations of precipitation are few and far between. Estimates of the precipitation flux are therefore forced to rely on model reanalysis, which have large uncertainties (e.g., Bromwich et al., 2018), on indirect measures such as river runoff, which may also be affected by glacier melt or on GNSS data analysis of solid earth movements in response to localized precipitation (e.g., Bevis et al., 2019).



285 **Figure 4: Time series of annual poleward atmospheric moisture transport (in $\text{km}^3 \text{yr}^{-1}$) across 60°N updated using ERA5 reanalysis dataset following Zhang et al. (2013) and the annual mean Arctic Oscillation (AO) Index constructed by NOAA Climate Prediction Center from 1979-2019. The transport was integrated from surface to the top of the atmosphere and along 60°N .**

2.3 Sea Ice

290 Freshwater stored in sea ice, i.e., sea ice volume, decreased by up to 50% over 2000-2010, but remained stable at approx. 12000 km^3 over 2010-2020 both in ORAs and in CryoSat-2 estimates (Figure 5). Kwok (2018) explained the flattening by the predominance of seasonal ice. Using a different approach, Liu et al. (2020) converted sea ice age into volume and also found a decrease in sea ice volume over the entire Arctic of $-411 \text{ km}^3 \text{yr}^{-1}$ over 1984-2018, most pronounced until 2010; their monthly trend ranges between $-537 \text{ km}^3 \text{yr}^{-1}$ in May and $-251 \text{ km}^3 \text{yr}^{-1}$ in September. The decrease in sea ice thickness is

295 responsible for 80% of this trend in winter and 50% in summer.




Figure 5: Time series of annual freshwater volume stored as sea ice from 7 ORAs and CRYOSAT2 (red stars), (in 10^3 km^3). The sea ice volume is calculated as the product of sea ice area and thickness. Annual volume maxima are shown by bold lines, while annual minima are shown by dashed lines.

300 Sea ice in the Arctic forms predominantly over the continental shelf. Estimates based on satellite imagery puts the cumulative sea ice formation of all Arctic coastal polynyas to 3000 km^3 per year (Tamura and Oshima, 2011), i.e., about a quarter of the total mean Arctic sea ice volume. Consequently, although the shelves receive large amounts of freshwater from rivers, their largest contribution to freshwater exchanges comes from sea ice export (e.g., Volkov et al., 2020), as the sea ice that forms on the shelves does not stay there. Sea ice is instead slowly transported across the Arctic by the Trans-polar Drift (Serreze et al., 305 1989), taking one to three years to travel from the Laptev Sea to Fram Strait (Pfirman et al., 1997; Steele et al., 2004). The Trans-polar Drift and ice deformation rates have been observed to be accelerating since the early 2000's (Rampal et al., 2011; Spreen et al., 2011); just recently, the MOSAiC drift expedition (<https://mosaic-expedition.org/>; Krumpfen et al., 2020) has shown that the Trans-polar Drift can, indeed, be unusually fast.

Fram Strait sea ice export is the largest dynamic sink of the Arctic freshwater cycle. The increase in Fram Strait sea ice export 310 detected from long-term monitoring of sea ice area has been suspected as the cause of Arctic sea ice volume loss, in particular for the multiyear thick sea ice within the Arctic Ocean (Smedsrud et al., 2017; Ricker et al., 2018). Using the more recent sea ice thickness retrievals, Spreen et al. (2020) actually showed that in volume, the Fram Strait export has in fact been decreasing at 27% per decade over 1992-2014, on par with the Fram Strait and Arctic ice thickness. In addition to the changes caused by thinned sea ice, changes in the atmospheric circulation pattern has also significantly contributed to the decrease in Fram Strait 315 sea ice export since the mid 1990s (Wei et al., 2019). Sea ice export from the Siberian shelf has increased by 46% over 2000-2014 compared to 1988-1999; from Amerasia to Europe, by 37% (Newton et al., 2017). But the summer survival rate of sea

ice on the Siberian shelves is decreasing by 15% per decade (Krumpen et al., 2019). That is, in the 1990s, 50% of first year ice entered the Trans-polar Drift; now, it is less than 20%, as the rest melts before reaching the Trans-polar Drift (Spall, 2019; Krumpen et al., 2019).

320 Snow on sea ice is crucial for surface heat budgets through its high albedo, and sea-ice growth through its thermal insulating effect. Therefore, snow on sea ice plays a significant role in determining where and when sea ice melts (Bigdeli et al., 2020). Although the delay of freeze up during early winter, partly depending on the anomalies of oceanic and atmospheric circulations (e.g., Kodaira et al., 2020), would cause a delay of snow accumulation on sea ice, the increase in precipitation and snow depth associated with the increase in storm activities in the Pacific Arctic contributes to a rapid build-up of snow cover on first year
325 ice (and a potential delay in spring/summer sea ice melt). These feedbacks were reported by Sato and Inoue (2018) based on the analysis of Ice Mass Balance buoys and CFSR reanalysis data sets. In the Atlantic sector, precipitation associated with six major storm events in 2014/2015 during the N-ICE2015 field campaign (Merkouriadi et al., 2017) caused the snow depth to be substantially greater than climatology.

2.4 Greenland Ice Sheet Discharge

330 The Greenland Ice Sheet has shown an increasing tendency for net ice sheet loss since the early 2000s (Shepherd et al., 2020), though with wide spatial and large temporal variability from year to year. This ice loss takes three forms: 1) liquid meltwater runs off from surface or basal melting, 2) submarine melt at outlet glaciers in contact with the ocean, 3) a solid component of ice loss driven by the calving of icebergs. As all components of ice loss have seen recent increases (Shepherd et al., 2020), the Greenland ice sheet is thus potentially a major source of change in freshwater fluxes in the Arctic Basin compared to mean
335 conditions established in earlier climatological periods.

Calculation of liquid runoff from Greenland needs to take into account both meltwater production, based either on surface energy budget considerations or using temperature index scaling, as well as the refreezing or storage of meltwater in the snowpack. Recent model intercomparisons of modelled Surface Mass Budget (SMB) (Fettweis et al., 2020) and refreezing in firn (Vandecrux et al., 2020) show that the primary source of variability in model estimates is still the amount of melt. This is
340 primarily modulated by surface albedo, but is also determined by the amount and spatial variability in the distribution of snowfall from models as the difference in surface properties between fresh snow and bare glacier ice lead to a melt-albedo feedback that is triggered when bare glacier ice is exposed (e.g., Hermann et al., 2018).

Ice loss also occurs from calving discharge and submarine melting where calving termini meet the ocean in fjords (Bamber et al., 2012, 2018). The Ice sheet Mass Balance Inter-comparison Exercise (IMBIE) (Shepherd et al., 2012) and IMBIE2
345 (Shepherd et al., 2020) results show a steady increase in net mass loss from around $-119 \pm 16 \text{ Gt yr}^{-1}$ for the period 1992–2011, followed by a reduction to -244 ± 28 in the 2012 to 2017 period with a peak in 2012 of $345 \pm 66 \text{ Gt yr}^{-1}$ (see also Helm

et al., 2014). Their analysis emphasizes that the increase in ice loss is due to both enhanced calving and submarine melting at outlet glaciers and increased surface melt and runoff through the period. In the mid-2010s a series of cooler summers, wetter winters and a slowing in calving rates from Sermeq Kujalleq (Jakobshavn Isbræ) led to a short-lived slowing in the rate of mass loss. In fact, Simonsen et al. (2021) found that 2017 is the first year in the 21st century with a neutral annual mass budget. However, they and others also further note the resumption of high ice loss in 2018 and particularly 2019, which although outside the IMBIE2 period of mass change has led to further decreases in the decadal mass balance of the ice sheet (Tedesco and Fettweis, 2020; Sasgen et al., 2020).

It is important to note that net mass loss is not the same as net freshwater flux from the ice sheet. The total flux is much larger than the former, however it is much harder to measure and modelled SMB is often used in association with remote sensing and field observations to assess the different components of the ice budget. The ice sheet only accumulates ice via precipitation. The GIS SMBMIP (Fettweis et al., 2020) compared results from 13 different models over Greenland and these figures are helpful to define both the net mass loss but also the components, including the freshwater flux. Typical values for the mean annual snowfall from regional climate models, and statistical downscaling of reanalysis are in the range of 500 to 800 Gt per year for the mean annual snowfall. The modelled liquid runoff by comparison is in the range of 200 to 500 Gt though note that many of the highest snowfall models also have runoff so the models converge to a smaller range of SMB values. To assess the calving and submarine melting components of freshwater flux from Greenland, remote sensing observations have focused on two separate techniques, a discharge estimate based on the observed velocity of outlet glaciers through flux gates and a gravimetry method where the total ice sheet change in mass is computed from gravimetric observations from the GRACE and GRACE-Follow On satellites. Modelled SMB is subtracted from the total mass change to give an estimate of the total discharge. The discharge component of the freshwater fluxes thus encompasses both submarine melting and iceberg calving.

Mankoff et al. (2019) produced the most recent assessment of the freshwater flux from Greenland based on ice dynamical discharge. This study measured ice velocity from satellite observations and determined the shape and size of flux gates on every outlet around the ice sheet to calculate how much ice leaves the ice sheet every year. Their estimate of $488 \pm 49 \text{ Gt yr}^{-1}$ is consistent with that produced by King et al. (2018) of $484 \pm 9 \text{ Gt yr}^{-1}$ and Kjeldsen et al. (2015) of -465.2 ± 65.5 , both for the 2003-2010 period. All three studies note that while the amount of discharge over the whole ice sheet has steadily increased through the 20th century (based on comparison with aerial photos and mapped glacier extents (Kjeldsen et al., 2015) to the 2010s, the rate of increase has largely stabilized at a high level in the last few years. However, the spatial pattern of discharge varies through time and space. Initial high discharge numbers in the 2000s were driven by accelerations in outlet glaciers in especially western Greenland (e.g., Jakobshavn/ Sermeq Kujalleq in the west). More recently, deceleration at Sermeq Kujalleq but additional accelerations in ice flow speeds at other outlets (e.g., Helheim glacier in south east Greenland) have been observed and these are sufficient to compensate and keep the overall discharge numbers high.

Taken together, the modelled runoff and ice discharge figures given in this section indicate that Greenland adds on average between 680 to 1000 Gt of fresh water to the oceans each year. However, the spatial variability in ice discharge complicates the interpretation of implications for the Arctic freshwater balance. The main regions of accelerating ice loss in Greenland drain out to the North Atlantic particularly in the high melt and high calving regions of western and south east Greenland. There has also been an observable increase in both calving and runoff from the outlet glaciers of northern Greenland (Hill et al., 2018; Solgaard et al., 2018; Shepherd et al., 2019, Extended Figure 4), which directly drains to the Arctic Ocean. The northern coast of Greenland as well as the Canadian Arctic Archipelago draining into the same region have seen a succession of ice shelf collapses and associated changes in the fjords most likely related to sub-shelf melting and increased atmospheric air temperatures in the region since the 1950s (e.g. Copland et al, 2007) indicating an increase in freshwater contribution from both Greenland as well as some of the smaller Arctic glaciers in the region directly into the Arctic Ocean basin. Mankoff et al. (2019) show a relatively stable 26 gigatonnes of ice discharge per year in the northern Greenland drainage basin that drains directly to the Arctic Ocean basin. This figure does not include surface melt and runoff but analysis by Fettweis et al. (2020) indicates a similar annual gain by SMB processes in the same basin up until 2013 but declining thereafter as mass loss has increased in this region. The analysis of Arctic freshwater flux from land ice presented by Bamber et al. (2018) reaches a similar conclusion. They estimate that including land ice from other parts of the Arctic as well as the Greenland ice sheet, the total freshwater flux is around 1300 Gt per year in the period since 2010 where they also identify a marked increase in runoff and discharge compared to a climatology period of 1960 to 1990. They also note, as we do, that the distribution of the freshwater flux is not even around Greenland spatially, but also temporally, with both runoff and iceberg discharge peaking in summer but being rather low (though not zero) in winter.

2.5 Ocean Transport Through Gateways

The latest reviews of the Arctic freshwater budget and fluxes (e.g., Haine et al., 2015; Carmack et al., 2016; Østerhus et al., 2019) conclude that observations of liquid freshwater transport through the Bering, Davis and Fram straits do not show significant trends between 1980-1990 and the 2000s. A recent study by Woodgate (2018) has shown that the Bering Strait exhibited a significant increase in volume and freshwater import to the Arctic between 2001 and 2014. Florindo-Lopez et al. (2020) analysed several decades of summertime hydrographic data at the eastern side of the Labrador Sea to find that freshwater transports in the boundary current were generally lower in the period mid-1990s to 2015 than the pre-1990s transports. The long-term variability was of the order of 30 mSv.

Polyakov et al. (2020) have described the contrasting changes in the Eurasian and Amerasian basins, where the latter has shown increasing stratification in recent years. They relate this to an increased import of low-salinity waters through the Bering Strait (see Woodgate, 2018). In the Eurasian Basin, Polyakov et al. (2020) relate the weakening stratification and enhanced sea ice melt, a process referred to as the Atlantification of the Arctic (Polyakov et al., 2017), to injection of (warmer) relatively salty water from the Barents Sea into the Eurasian Basin halocline, flowing at shallower depths. Although they do not show

410 any clear link to the Fram Strait imports, they find a small but statistically significant correlation between observed salinity in the Eastern Eurasian Basin halocline and the northern Barents Sea upper water column. Thus, in agreement with the box model estimates of Tsubouchi et al. (2021), there appears to be no trend in volume fluxes at the boundaries, and no evidence for a dominant link between changes in the freshwater fluxes at the boundaries and changes in the upper Arctic Ocean, this also true for the Atlantic water volume inflow.

415 3 Redistribution of Arctic Freshwater

The freshwater surface circulation in the Arctic has two, non-independent components; a density-driven circulation, linked mostly to river runoff and sea ice processes; and a wind-driven circulation, consisting mostly of the anticyclonic/convergent Beaufort Gyre and the cyclonic/divergent Trans-polar Drift. The wind-driven circulation produces local accumulation or thinning of the surface layer (Timmermans and Marshall, 2020). Although, the exchanges with the Atlantic and Pacific
420 influence the large scale salinity gradients across the Arctic Ocean (Polyakov et al., 2020), the combined effects of the density-driven and wind-driven circulations primarily drive a strong freshwater gradient through the Arctic, of up to 25 m freshwater equivalent (Rabe et al., 2011), with a maximum freshwater content in the Beaufort Gyre and a minimum in the Nansen Basin towards the Barents Sea. Morison et al. (2012) and Alkire et al. (2007) in particular have shown the regional changes in steric height and sea level pressure, respectively, can redistribute relatively fresh water near the surface along the boundaries of the
425 deep basin and the shelves. Recent studies suggest that the Beaufort Gyre has stabilised or reached a new normal high-freshwater content state. Dewey et al. (2018) attributes this to a switch from a system driven by surface ice- and wind-stress that affects a passive ocean, to one where it is the ocean that drives the ice (often in the absence of wind). Zhong et al. (2019) in contrast attribute it to higher energy input to the ocean, and suggest that the transition is not complete, i.e., the Beaufort Gyre is not "saturated" yet. Zhong et al. (2019) further concludes that the recent increase in cyclonic activity reduces this
430 energy input, and hence should result in future decrease of freshwater stored in the Beaufort Gyre. This surface circulation transports meteoric water (and hence nutrients, e.g., Bluhm et al., 2015) throughout the Arctic. On average, 10% of the Arctic surface waters are made up of meteoric waters (shallower than ~200 m depth; see Forryan et al. 2019, their figure 5b) and this number has so far been constant since the early 2000s (Alkire et al., 2017; Proshutinsky et al., 2019).

The wind also contributes to vertical redistribution via wind-driven coastal up and downwelling. On average, only the Laptev
435 and Kara are dominated with downwelling; the rest of the Arctic, especially the Amerasian basin, is upwelling dominated (Williams and Carmack, 2015). On average, only the Laptev and Kara are dominated with downwelling; the rest of the Arctic, especially the Amerasian basin, is upwelling dominated (Williams and Carmack, 2015). However, bathymetric features can reverse the sign of this Ekman transport (Randelhoff and Sundfjord, 2018; Danielson et al., 2020). Even more relevant for freshwater, at locations where upwelling occurs, river plumes are pushed offshore (Williams and Carmarck, 2015; Våge et al.,
440 2016). Downward flows of water can be generated by the wind or by an increase in density that destabilizes the water column.

On the Arctic shelf, dense water can form as a result of cooling or brine rejection during sea ice formation, especially in polynyas (Ivanov et al., 2004). Cascading plumes entrain waters during their descent, explaining how cascading of cold and saline surface waters can result in warmer (if entraining Atlantic water) or fresher (if entraining halocline) deeper levels (Backhaus et al., 1997). Preconditioning of the shelf waters due to the mixing with the upwelled Atlantic water also can result
445 in the cold and saline cascading plumes (Luneva et al., 2020). Furthermore, cascading is becoming more common in the Arctic; it is more effective in mixing and ventilating upper and low intermediate Arctic waters than open ocean deep convection and can reach deep into the water column (e.g., Luneva et al., 2020). Cascading and entrainment in the Beaufort Sea during upwelling events re-injects cold and fresh water into the halocline (Ivanov et al., 2004). Janout et al. (2017) observed shelf processes and the modification of warm Atlantic Water leading to flux of the modified water, and hence an effective freshwater
450 flux, toward the basin. From two expeditions in 2013 and 2014 and one year of mooring deployment in between, Janout et al. (2017) found a dual behavior in Vilkitsky Trough, between the Kara and Laptev Sea: strong winds can cause an upward diversion of the along-slope freshwater transport onto the shelf; the addition of sea ice formation results in the formation of water with a higher density than that found at 3000 m, suggesting possible sinking of these waters to the Nansen basin.

The wind also impacts the depth of the mixed layer. The Arctic surface mixed layer varies both seasonally and geographically,
455 as reviewed by Peralta-Ferriz and Woodgate (2015). Using all available observations from 1979 to 2012, Peralta-Ferriz and Woodgate (2015) find a shoaling trend in the whole Arctic in winter; in summer, the mixed layer trend is of a deepening in ice free parts of Barents and Beaufort, but also a shoaling in the Eurasian basin. Polyakov et al. (2017) found an opposite trend using moorings and ice-tethered profilers: an increased winter convection caused by sea ice formation over a weakened stratification in the eastern Eurasian basin. They argue that the entire Eurasian basin is becoming similar to the Atlantic sector
460 of the Nansen basin and hence dubbed this phenomenon “the Atlantification of the Arctic” (or more recently, “the Borealisation of the Arctic”).

The Beaufort Gyre is a retainer of liquid freshwater in the Arctic Ocean, governed by three factors: wind stress, the dynamic feedback between ice motion and upper ocean currents (ice-ocean governor) and lateral eddy fluxes (Doddridge et al., 2019). Observations and an idealised two-layer model study indicate that the “ice-ocean governor”, controlling Ekman pumping, is
465 five times more important than eddy dynamics in regulating the retention and release of freshwater from the Beaufort Gyre (Meneghello et al., 2020). Regan et al. (2020) have shown that the mean kinetic energy dominated over eddy kinetic energy (isopycnal slope / potential for baroclinic instability) in governing Beaufort Gyre dynamics during the spin-up in the past one to two decades. Armitage et al. (2020) predict that eddies will become more important in stabilizing the Beaufort Gyre. In addition, idealised simulations with and without a continental slope by Manucharyan and Isachsen (2019) demonstrate that
470 eddy dynamics prevail in the Beaufort Gyre only in the presence of the slope. Further, Liang and Losch (2018) show not only that the positive feedback loop “enhanced vertical mixing = less sea ice” reduces the halocline-to-Atlantic Water (AW) stratification, but also leads to a colder AW and hence mixing down of salt from AW to the deep Arctic.

4 Summary

Our review of recent work suggests that Arctic freshwater content in the 2010s has stabilized relative to the 2000s. This
475 stabilization is due in part to the compensation between an increase in the Beaufort Gyre and a decrease in the rest of the Arctic
Ocean. However, large inter-model spread in the ocean reanalyses and uncertainty in the observations used in this study
prevents a definitive estimate of the degree of this compensation. The most notable differences between the 2010s and the
2000s are the switch to an increasingly seasonal and mobile sea ice cover, whose impacts on the Arctic ocean and atmosphere
are still being debated; an increase in mass loss from the Greenland ice sheet including in the northern region that drains
480 directly into the Arctic ocean; and the import of warm Atlantic waters that has shoaled. The review also suggests that large
uncertainties remain in quantifying regional patterns, changes, and individual contributors to freshwater content variability,
motivating the need for long term monitoring, in-situ in rivers and ocean, and from space, to distinguish trends from low
frequency variability.

5 References

485 Aagaard, K. and Carmack, E. C.: The role of sea ice and other fresh water in the Arctic circulation. *J. Geophys. Res.*, 94,
14,485–14,497, 1989.

Aagaard, K. and R. A. Woodgate, R. A.: Some thoughts on the freezing and melting of sea ice and their effects on the ocean,
Ocean Modell., 3, 127– 135, 2001.

Aagaard, K., Swift, J. H., and Carmack, E. C.: Thermohaline circulation in the Arctic mediterranean seas. *J. Geophys. Res.*,
490 90: 4833-4846, 1985.

Alkire, M. B., Morison, J., and Andersen, R.: Variability in the meteoric water, sea-ice melt, and Pacific water contributions
to the central Arctic Ocean, 2000–2014. *J. Geophys. Res.*, 120, 1573–1598. doi:10.1002/2014JC010023, 2015.

Alkire, M. B., Falkner, K. K., Rigor, I., Steele, M., and Morison, J.: The return of Pacific waters to the upper layers of the
central Arctic Ocean, *Deep-Sea Res. I*, 54, 1509-1529, 2007.

495 Andersen, O. B. and Piccioni, G.: Recent Arctic Sea Level Variations from Satellites. *Front. Mar. Sci.*, 3, 76, 2016.

Ardyna, M. and Arrigo, K. R.: Phytoplankton dynamics in a changing Arctic Ocean. *Nature Climate Change*, 10, 892-903,
doi:10.1038/s41558-020-0905-y, 2020.

Armitage, T. W. K., Manucharyan, G. E., Petty, A. A., Kwok, R., and Thompson, A. F.: Enhanced eddy activity in the Beaufort
Gyre in response to sea ice loss, *Nature Comm.*, 11, 761, doi:10.1038/s41467-020-14449-z, 2020.

- 500 Armitage, T. W. K., Bacon, S., Ridout, A. L., Thomas, S. F., Aksenov, Y., and Wingham, D. J.: Arctic sea surface height variability and change from satellite radar altimetry and GRACE, 2003–2014. *J. Geophys. Res. Oceans*, 121, 4303–4322, 2016.
- Backhaus, J. O., Fohrmann, H., Kämpf, J., and Rubino, A.: Formation and export of water masses produced in Arctic shelf polynyas—Process studies of oceanic convection. *ICES J. Marine Sci.*, 54, 3, 366–382, 1997.
- 505 Bacon, S., Aksenov, Y., Fawcett, S., and Madec, G.: Arctic mass, freshwater and heat fluxes: methods and modelled seasonal variability. *Philos. T. Roy. Soc. A.*, 373(2052), 20140169–16. doi:10.1098/rsta.2014.0169, 2015.
- Bamber, J. L., Tedstone, A. J., King, M. D., Howat, I. M., Enderlin, E. M., van den Broeke, M. R., and Noël, B.: Land Ice Freshwater Budget of the Arctic and North Atlantic Oceans: 1. Data, Methods, and Results. *J. Geophys. Res. Oceans*, 44(4), 3728–11. doi:10.1002/2017JC013605, 2018.
- 510 Bamber, J., van den Broeke, M., Ettema, J., Lenaerts, J., and Rignot, E.: Recent large increases in freshwater fluxes from Greenland into the North Atlantic. *Geophys. Res. Lett.*, 39(19), L19501. doi:10.1029/2012GL052552, 2012.
- Bevis, M., Harig, C., Khan, S. A., Brown, A., Simons, F. J., Willis, M., Fettweis, X., van den Broeke, M. R., Madsen, F. B., Kendrick, E., Caccamise II, D. J., van Dam, T., Knudsen, P., and Nylén, T.: Accelerating changes in ice mass within Greenland, and the ice sheet’s sensitivity to atmospheric forcing. *Proc. Natl. Acad. Sci. U.S.A.* 116, 1934–1939.
- 515 doi:10.1073/pnas.1806562116, 2019.
- Bigdeli, A., Nguyen, A. T., Pillar, H. R., Ocana, V., and Heimbach, P.: Atmospheric Warming Drives Growth in Arctic Sea-Ice: A Key Role for Snow. *Geophys. Res. Lett.*, 47, doi:10.1029/2020GL090236, 2020.
- Blockley, E. W., Martin, M. J., McLaren, A. J., Ryan, A. G., Waters, J., Lea, D. J., Mirouze, I., Peterson, K. A., Sellar, A., and Storkey, D.: Recent development of the Met Office operational ocean forecasting system: an overview and assessment of
- 520 the new Global FOAM forecasts, *Geosci. Model Dev.*, 7, 2613–2638, doi:10.5194/gmd-7-2613-2014, 2014.
- Bluhm, B. A., Kosobokova, K. N., and Carmack, E. C.: A tale of two basins: An integrated physical and biological perspective of the deep Arctic Ocean. *Prog. Oceanog.*, 139, 89–121, 2015.
- Boisvert, L. N., Webster, M. A., Petty, A. A., Markus, T., Bromwich, D. H., and Cullather, R. I.: Intercomparison of precipitation estimates over the Arctic Ocean and its peripheral seas from reanalyses. *J. Clim.*, 31(20), 8441–8462,
- 525 doi:10.1175/JCLI-D-18-0125.1, 2018.

- Bromwich, D. H., Wilson, A. B., Bai, L., Liu, Z., Barlage, M., Shih, C.-F., Maldonado, S., Hines, K. M., Wang, S.-H., Woollen, J., Kuo, B., Lin, H.-C., Wee, T.-K., Serreze, M. C., and Walsh, J. E.: The Arctic System Reanalysis, Version 2. *Bull. Amer. Meteor. Soc.*, 99(4), 805–828, doi:10.1175/BAMS-D-16-0215.1, 2018.
- 530 Carmack, E. C., Yamamoto Kawai, M., Haine, T. W. N., Bacon, S., Bluhm, B. A., Lique, C., Melling, H., Polyakov, I. V., Straneo, F., Timmermans, M.-L., and Williams, W. J.: Freshwater and its role in the Arctic Marine System: Sources, disposition, storage, export, and physical and biogeochemical consequences in the Arctic and global oceans. *Journal of Geophys. Res. Biogeosci.*, 121(3), 675–717. doi:10.1002/2015JG003140, 2016.
- Carton, J.A., Chepurin, G. A., and Chen, L.: SODA3: a new ocean climate reanalysis, *J. Clim.*, 31, 6967-6983, doi:10.1175/JCLI-D-18-0149.1, 2018.
- 535 Chang, Y.-S., Zhang, S., Rosati, A., Delworth, T. L., and Stern, W.F.: An assessment of oceanic variability for 1960–2010 from the GFDL ensemble coupled data assimilation. *Clim. Dyn.*, 40(3-4), doi:10.1007/s00382-012-1412-2, 2013.
- Copland, L., Mueller, D. R., and Weir, L.: Rapid loss of the Ayles Ice Shelf, Ellesmere Island, Canada, *Geophys. Res. Lett.*, 34, L21501, doi:10.1029/2007GL031809, 2007.
- 540 Danielson, S. L., Hennon, T. D., Hedstrom, K. S., Pnyushkov, A. V., Polyakov, I. V., Carmack, E., Filchuk, K., Janout, M., Makhotin, M., Williams, W. J., and Padman, L.: Oceanic Routing of Wind-Sourced Energy Along the Arctic Continental Shelves. *Front. Mar. Sci.* 7:509. doi:10.3389/fmars.2020.00509, 2020.
- Dewey, S., Morison, J., Kwok, R., Dickinson, S., Morison, D., and Andersen, R.: Arctic ice-ocean coupling and gyre equilibration observed with remote sensing. *Geophysical Research Letters*, 45. doi: 10.1002/2017GL076229, 2018.
- 545 Dickson, R. R., Meincke, J., Malmberg, S. A., and Lee, A. J.: The “Great Salinity Anomaly” in the northern North Atlantic 1968–1982. *Prog. Oceanogr.*, 20:103–151. doi:10.1016/0079- 6611(88)90049-3, 1988.
- Doddridge, E. W., Meneghello, G., Marshall, J., Scott, J., and Lique, C.: A Three-Way Balance in the Beaufort Gyre: The Ice-Ocean Governor, Wind Stress, and Eddy Diffusivity, *J. Geophys. Res. Oceans*, doi:10.1029/2018JC014897, 124, 5, 3107-3124, 2019.
- 550 Doyle, S. H., Hubbard, A., van de Wal, R. S. W., Box, J. E., van As, D., Scharrer, K., Meierbachtol, T. W., Smeets, P. C. J. P., Harper, J. T., Johansson, E., Mottram, R. H., Mikkelsen, A. B., Wilhelms, F., Patton, H., Christoffersen, P., and Hubbard, B.: Amplified Melt and Flow of the Greenland Ice Sheet Driven by Late-Summer Cyclonic Rainfall. *Nature Geosci.*, 8 (8), 647–53. doi:10.1038/ngeo2482, 2015.

- Dukhovskoy, D. S., Yashayaev, I., Proshutinsky, A., Bamber, J. L., Bashmachnikov, I. L., Chassignet, E. P., Lee, C. M., and Tedstone, A. J.: Role of Greenland Freshwater Anomaly in the Recent Freshening of the Subpolar North Atlantic. *J. Geophys. Res. Oceans*, 21(1), 181–28. doi:10.1029/2018JC014686, 2019.
- 555
- Fer, I.: Weak vertical diffusion allows maintenance of cold halocline in the central Arctic. *Atmos. Oceanic Sci. Lett.*, 2, 148–152, 2009.
- Fettweis, X., Hofer, S., Krebs-Kanzow, U., Amory, C., Aoki, T., Berends, C. J., Born, A., Box, J. E., Delhasse, A., Fujita, K., Gierz, P., Goelzer, H., Hanna, E., Hashimoto, A., Huybrechts, P., Kapsch, M.-L., King, M. D., Kittel, C., Lang, C., Langen, P. L., Lenaerts, J. T. M., Liston, G. E., Lohmann, G., Mernild, S. H., Mikolajewicz, U., Modali, K., Mottram, R. H., Niwano, M., Noël, B., Ryan, J. C., Smith, A., Streffing, J., Tedesco, M., van de Berg, W. J., van den Broeke, M., van de Wal, R. S. W., van Kampenhout, L., Wilton, D., Wouters, B., Ziemen, F., and Zolles, T.: GrSMBMIP Intercomparison of the modelled 1980–2012 surface mass balance over the Greenland Ice sheet, *The Cryosphere*, 14, 3935–3958, doi:10.5194/tc-2019-321, 2020.
- 560
- Fichot, G. G., Kaise, K., Hooker, S. B., Amon, R. M. W., Babin, M., Bélanger, S., Walker, S. A., and R. Benner, R.: Pan-Arctic distributions of continental runoff in the Arctic Ocean, *Sci. Rep.*, 3, 1053, doi:10.1038/srep01053, 2013.
- 565
- Florindo-Lopez, C., Bacon, S., Aksenov, Y., Chafik, L., Colbourne, E., and Holliday, N. P.: Arctic Ocean and Hudson Bay freshwater exports: new estimates from 7 decades of hydrographic surveys on the Labrador shelf. *J. Clim.*, doi:10.1175/JCLI-D-19-0083.1, 2020.
- Forryan, A., Bacon, S., Tsubouchi, T., Torres-Valdes, S., and Naveira Garabato, A. C.: Arctic freshwater fluxes: sources, tracer budgets and inconsistencies. *The Cryosphere* 13 2111–2131, doi:10.5194/tc-13-2111-2019, 2019.
- 570
- Garric, G., Parent, L., Greiner, E., Dréville, M., Hamon, M., Lellouche, J.M., Régnier, C., Desportes, C., Le Galloudec, O., Bricaud, C., Drillet, Y., Hernandez, F., Dubois, C., and Le Traon, P.-Y.: Performance and quality assessment of the global ocean eddy-permitting physical reanalysis glorys2v4. In: Buch E, Fernandez V, Nolan G, Eparkhina D (eds) Operational oceanography serving sustainable marine development. Proceedings of the eight EuroGOOS international conference, 2018, 3–5 October 2017, Bergen, Norway (ISBN 978-2-9601883-3-2), 2017.
- 575
- Giles, K. A., Laxon, S. W., Ridout, A. L., Wingham, D. J., and Bacon, S.: Western Arctic Ocean freshwater storage increased by wind-driven spin-up of the Beaufort Gyre, *Nat. Geosci.*, 5, 194–197, 2012.
- Haine, T. W. N., Curry, B., Gerdes, R., Hansen, E., Karcher, M., Lee, C., Rudels, B., Spreen, G., de Steur, L., Stewart, K. D., and Woodgate, R.: Arctic freshwater export: Status, mechanisms, and prospects. *Global Planet. Change*, 125(C), 13–35. doi:10.1016/j.gloplacha.2014.11.013, 2015.
- 580

- Helm, V., Humbert, A., and Miller, H.: Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2, *The Cryosphere*, 8(4), 1539–1559, 2014.
- Hermann, M., Box, J. E., Fausto, R. S., Colgan, W. T., Langen, P. L., Mottram, R., Wuite, J., Noël, B., van den Broeke, M. R., and van As, D.: Application of PROMICE Q-Transect in Situ Accumulation and Ablation Measurements (2000–2017) to
585 Constrain Mass Balance at the Southern Tip of the Greenland Ice Sheet. *J. Geophys. Res. Earth Surf.*, 123, 1235–1256, 2018.
- Holland, M., Finniss, J., and Serreze, M. C.: Simulated Arctic Ocean freshwater budgets in the twentieth and twenty-first centuries. *J. Clim.*, 19:6221–6242. doi:10.1175/JCLI3967.1, 2006.
- Horwath, M., Novotny, K., Cazenave, A., Palanisamy, H., Marzeion, B., Paul, F., Döll, P., Cáceres, D., Hogg, A., Shepherd, A., et al.: ESA Climate Change Initiative (CCI) Sea Level Budget Closure (SLBC_cci) Executive Summary Report D4.4;
590 Version 1.0; ESA: Rome, Italy, 2020.
- Ivanov, V.V., Shapiro, G.I., Huthnance, J.M., Aleynik, D.L., and Golovin, P.N.: Cascades of dense water around the world ocean. *Prog. Oceanog.*, 60(1), 47-98, 2004.
- Janout, M. A., Hölemann, J., Timokhov, L., utjahr, O., and Heinemann, G.: Circulation in the northwest Laptev Sea in the eastern Arctic Ocean: Crossroads between Siberian River water, Atlantic water and polynya-formed dense water, *J. Geophys. Res. Oceans*, 122, 6630–6647, doi:10.1002/2017JC013159, 2017.
595
- Jeffries, M. O., Overland, J. E., and Perovich, D. K.: The Arctic shifts to a new normal. *Phys. Today*, 66(10):35–40. doi:10.1063/PT.3.2147, 2013.
- King, M. D., Howat, I. M., Jeong, S., Noh, M. J., Wouters, B., Noël, B., and van den Broeke, M. R.: Seasonal to decadal variability in ice discharge from the Greenland Ice Sheet, *The Cryosphere*, 12, 3813–3825, doi:0.5194/tc-12-3813-2018, 2018.
- 600 Kjeldsen, K., Korsgaard, N., Bjørk, A. et al.: Spatial and temporal distribution of mass loss from the Greenland Ice Sheet since AD 1900. *Nature* 528, 396–400, doi:10.1038/nature16183, 2015.
- Kodaira, T., Waseda, T., Nose, T., and Inoue, J.: Record high Pacific Arctic seawater temperatures and delayed sea ice advance in response to episodic atmospheric blocking. *Sci. Rep.*, 10, 20830, doi:10.1038/s41598-020-77488-y, 2020.
- Kruppen, T., Belter, H. J., Boetius, A., Damm, E., Haas, C., Hendricks, S., Nicolaus, M., Nöthig, E.-M., Paul, S., Peekem, I.,
605 Ricker, R., and Stein, Rudiger: Arctic warming interrupts the Transpolar Drift and affects long-range transport of sea ice and ice-rafted matter. *Sci. Rep.*, 9, 5459, doi:0.1038/s41598-019-41456-y, 2019.

- Kruppen, T., Birrien, F., Kauker, F., Rackow, T., von Albedyll, L., Angelopoulos, M., Belter, H. J., Bessonov, V., Damm, E., Dethloff, K., Haapala, J., Haas, C., Harris, C., Hendricks, S., Hoelemann, J., Hoppmann, M., Kaleschke, L., Karcher, M., Kolabutin, N., Lei, R., Lenz, J., Morgenstern, A., Nicolaus, M., Nixdorf, U., Petrovsky, T., Rabe, B., Rabenstein, L., Rex, M.,
610 Ricker, R., Rohde, J., Shimanchuk, E., Singha, S., Smolyanitsky, V., Sokolov, V., Stanton, T., Timofeeva, A., Tsamados, M.,
and Watkins, D.: The MOSAiC ice floe: sediment-laden survivor from the Siberian shelf, *The Cryosphere*, 14, 2173–2187,
doi:10.5194/tc-14-2173-2020, 2020.
- Kryzhov, V.N., and Gorelits, O.V.: The Arctic Oscillation and its impact on temperature and precipitation in Northern Eurasia
in the 20th Century. *Russ. Meteorol. Hydrol.* 40, 711–721, doi:10.3103/S1068373915110011, 2015.
- 615 Kwok, R.: Arctic sea ice thickness, volume, and multiyear ice coverage: losses and coupled variability (1958-2018). *Environ.
Res. Lett.*, 13(10), 105005. doi:10.1088/1748-9326/aae3ec, 2018.
- Lenaerts, J. T. M., Medley, B., Broeke, M. R., and Wouters, B.: Observing and Modeling Ice Sheet Surface Mass Balance. *Rev. Geophys.*, 41(1), 1–45, doi:10.1029/2018RG000622, 2019.
- Lewis, K. M., van Dijken, G. L., and Arrigo, K. R.: Changes in phytoplankton concentration now drive increased Arctic Ocean
620 primary production. *Science*, 369(6500), 198–202. doi:10.1126/science.aay8380, 2020.
- Liang, X. and Losch, M.: On the effects of increased vertical mixing on the Arctic Ocean and sea ice. *J. Geophys. Res. Oceans*,
123(12), pp.9266-9282, 2018.
- Liu, Y., Key, J. R., Wang, X., and Tschudi, M.: Multidecadal Arctic sea ice thickness and volume derived from ice age, *The
Cryosphere*, 14, 1325–1345, <https://doi.org/10.5194/tc-14-1325-2020>, 2020.
- 625 Luneva, M. V., Ivanov, V. V., Tuzov, F., Aksenov, Y., Harle, J.D., Kelly, S., and Holt, J.T.: Hotspots of dense water cascading
in the Arctic Ocean: Implications for the Pacific water pathways. *J. Geophys. Res.*, 125 (10), e2020JC016044.
doi:10.1029/2020JC016044, 2020.
- Mankoff, K. D., Colgan, W., Solgaard, A., Karlsson, N. B., Ahlstrøm, A. P., van As, D., Box, J. E., Khan, S. A., Kjeldsen, K.
K., Mouginot, J., and Fausto, R. S.: Greenland Ice Sheet solid ice discharge from 1986 through 2017, *Earth Syst. Sci. Data*,
630 11, 769–786, doi:10.5194/essd-11-769-2019, 2019.
- Manucharyan, G.E. and Isachsen, P.E.: Critical role of continental slopes in halocline and eddy dynamics of the Ekman-driven
Beaufort Gyre. *J. Geophys. Res. Oceans*, 124(4), pp.2679-2696, 2019.

- McLaughlin, F. A., Carmack, E. C., Macdonald, R. W., and Bishop, J. K. B.: Physical and geochemical properties across the Atlantic/Pacific water mass front in the southern Canadian Basin, *J. Geophys. Res.*, 101(C1), 1183–1197, doi:10.1029/95JC02634, 1996.
- 635
- McPhee, M. G., Proshutinsky, A., Morison, J. H., Steele, M., and Alkire, M. B.: Rapid change in freshwater content of the Arctic Ocean. *Geophys. Res. Lett.*, 36(10), L10602. doi:10.1029/2009GL037525, 2009.
- Meneghello G., Doddridge, E. W., Marshall, J., Scott, J., and Campin, J.-M.: Exploring the role of the “Ice-Ocean governor” and mesoscale eddies in the equilibration of the Beaufort Gyre: lessons from observations. *J. Phys. Oceanogr.*, 50(1), doi:10.1175/JPO-D-18-0223.1, 2020.
- 640
- Merkouriadi, I., Gallet, J.-C., Graham, R. M., Liston, G. E., Polashenski, C., Hudson, S., Rösel, A., and Gerland, S.: Winter snow conditions on Arctic sea ice north of Svalbard during the Norwegian young sea ICE (N-ICE2015) expedition. *J. Geophys. Res. Atmos.*, 122, 10,837- 10,854, 2017.
- Moon, T., Sutherland, D. A., Carroll, D., Felikson, D., Kehrl, L., & Straneo, F.: Subsurface iceberg melt key to Greenland fjord freshwater budget. *Nature Geoscience*, 11(1), 1–8. doi:10.1038/s41561-017-0018-z, 2017.
- 645
- Morison, J., Wahr, J., Kwok, R., and Peralta-Ferriz, C.: Recent trends in Arctic Ocean mass distribution revealed by GRACE. *Geophys. Res. Lett.*, 34, L07602, doi:10.1029/2006GL029016, 2007.
- Morison, J., Kwok, R., Peralta-Ferriz, C., Alkire, M., Rigor, I., Andersen, R., and Steele, M.: Changing arctic ocean freshwater pathways. *Nature*, 481(7379), 66-70, 2012.
- 650
- Newton, R., Pfirman, S., Tremblay, B., and DeRepentigny, P.: Increasing transnational sea-ice exchange in a changing Arctic Ocean. *Earth's Future*, 5(6), 633-647, 2017.
- Nguyen, A. T., Pillar, H., Ocaña, V., Bigdeli, B., Smith, T. A., and Heimbach, P.: The Arctic Subpolar gyre sTate Estimate (ASTE): Description and assessment of a data-constrained, dynamically consistent ocean-sea ice estimate for 2002--2017, doi:10.1002/essoar.10504669.1, 2021.
- 655
- Nygaard, T., Naakka, T., and Vihma, T.: Horizontal moisture transport dominates the regional moistening patterns in the Arctic. *J. Clim.*, 33, 6793-6807, 2020.
- Oltmanns, M., Straneo, F., and Tedesco, M.: Increased Greenland melt triggered by large-scale, year-round cyclonic moisture intrusions, *The Cryosphere*, 13, 815–825, doi:10.5194/tc-13-815-2019, 2019.

- Østerhus, S., Woodgate, R., Valdimarsson, H., Turrell, B., de Steur, L., Quadfasel, D., Olsen, S. M., Moritz, M., Lee, C. M.,
660 Larsen, K. M. H., Jónsson, S., Johnson, C., Jochumsen, K., Hansen, B., Curry, B., Cunningham, S., and Berx, B.: Arctic
Mediterranean exchanges: a consistent volume budget and trends in transports from two decades of observations, *Ocean Sci.*,
15, 379–399, doi:10.5194/os-15-379-2019, 2019.
- Östlund, G. H. and Hut, G.: Arctic Ocean water mass balance from isotope data. *J. Geophys. Res.*, 89, 6373–6381,
doi:10.1029/JC089iC04p06373, 1984.
- 665 Passaro, M., Rose, S., Andersen, O., Boergens, E., Calafat, F., Dettmering, D., and Benveniste, J.: ALES+: Adapting a
homogenous ocean retracker for satellite altimetry to sea ice leads, coastal and inland waters. *Remote Sens. Environ.* 211,
456–471, 2018.
- Peralta-Ferriz, C. and Woodgate, R. A.: Seasonal and interannual variability of pan-Arctic surface mixed layer properties from
1979 to 2012 from hydrographic data, and the dominance of stratification for multiyear mixed layer depth shoaling. *Prog.*
670 *Oceanogr.*, 134, 19–53, doi:10.1016/j.pocean.2014.12.005, 2015.
- Peralta-Ferriz, C., Morison, J. H., Wallace, J. M., Bonin, J. A., and Zhang, J.: Arctic Ocean Circulation Patterns Revealed by
GRACE. *Journal of Climate*, 27(4), 1445–1468. doi:10.1175/JCLI-D-13-00013.1, 2014.
- Pfirman, S. L., Colony, R., Nürnberg, D., Eicken, H., and Rigor, I.: Reconstructing the origin and trajectory of drifting Arctic
sea ice. *J. Geophys. Res. Oceans*, 102(C6), 12575-12586, 1997.
- 675 Polyakov, I. V., Pnyushkov, A., Alkire, M., Ashik, I., Baumann, T., Carmack, E., Goszczko, I., Guthrie, J., Ivanov, V.,
Kanzow, T., Krishfield, R., Kwok, R., Sundfjord, A., Morison, J., Rember, R., and Yulin, A.: Greater role for Atlantic inflows
on sea-ice loss in the Eurasian Basin of the Arctic Ocean, *Science*, 356, 285–291, doi:10.1126/science.aai8204, 2017.
- Polyakov, I. V., Alkire, M. B., Bluhm, B. A., Brown, K. A., Carmack, E. C., Chierici, M., Danielson, S. L., Ellingsen, I.,
Ershova, E. A., Gårdfeldt, K., Ingvaldsen, R. B., Pnyushkov, A. V., Slagstad, D., and Wassmann, P.: Borealization of the
680 Arctic Ocean in Response to Anomalous Advection from Sub-Arctic Seas. *Front. Mar. Sci.* 7:491.
doi:10.3389/fmars.2020.00491, 2020.
- Proshutinsky, A., Dukhovskoy, D., Timmermans, M.-L., Krishfield, R., and J. Bamber, J.: Arctic circulation regimes, *Philos.*
Trans. R. Soc., A, 373, doi:10.1098/rsta.2014.0160, 2015.
- Proshutinsky, A., Krishfield, R., Timmermans, M.-L., Toole, J., Carmack, E., McLaughlin, F., Williams, W. J., Zimmermann,
685 S., Itoh, M., and Shimada, K.: The Beaufort Gyre Fresh Water Reservoir: State and variability from observations, *J. Geophys.*
Res., 114, C00A10, doi:10.1029/2008JC0055104, 2009.

- Proshutinsky, A., Krishfield, R., Toole, J. M., Timmermans, M.-L., Williams, W., Zimmermann, S., Yamamoto-Kawai, M., Armitage, T. W. K., Dukhovskoy, D., Golubeva, E., Manucharyan, G. E., Platov, G., Watanabe, E., Kikuchi, T., Nishino, S., Itoh, M., S.-H. Kang, S.-H., Cho, K.-H., Tateyama, K., and Zhao, J.: Analysis of the Beaufort Gyre freshwater content in 690 2003–2018. *J. Geophys. Res. Oceans*, 124, doi:10.1029/2019JC015281, 2019.
- Prowse, T., Bring, A., Mård, J., Carmack, E., Holland, M., Instanes, A., and Wrona, F. J.: Arctic Freshwater Synthesis: Summary of key emerging issues. *J. Geophys. Res. Biogeosci.*, 120(10), 1887–1893. doi:10.1002/2015JG003128, 2015.
- Rabe, B., Karcher, M., Schauer, U., Toole, J., Krishfield, R., Pisarev, S., Kauker, F., Gerdes, R., and Kikuchi, T.: An assessment of Arctic Ocean freshwater content changes from the 1990s to 2006–2008, *Deep-Sea Res. I*, 58, 173–185, 695 doi:10.1016/j.dsr.2010.12.002, 2011.
- Rabe, B., Karcher, M., Kauker, F., Schauer, U., Toole, J. M., Krishfield, R. A., Pisarev, S., Kikuchi, T., and Su, J.: Arctic Ocean liquid freshwater storage trend 1992–2012, *Geophys. Res. Lett.*, 41(3), 961–968, doi:10.1002/2013GL058121, 2014.
- Rabe, B., Karcher, M. J., Kauker, F., Schauer, U., Toole, J. M., Krishfield, R. A., Pisarev, S., Kikuchi, T., and Su, J.: (Table 1) Physical oceanography from several POLARSTERN cruises to the Arctic Ocean, 1993 - 2012. *PANGAEA*, 700 doi:10.1594/PANGAEA.815756, 2014b.
- Raj, R. P., Andersen, O. B., Johannessen, J. A., Benjamin, G., Chatterjee, S., Rose, S. K., Bonaduce, A., Horwath, M., Ranndal, H., Richter, K., Palanisamy, H., Ludwigsen, C. A., Bertino, L., Nilsen, J. E. O., Knudsen, P., Hogg, A., Cazenave, A., and Benveniste, J.: Arctic Sea level Budget Assessment During the GRACE/Argo Time Period. *Remote Sens.*, 125(2837). doi:0.3390/rs12172837, 2020.
- 705 Rampal, P., Weiss, J., Dubois, C., and Campin, J.-M.: IPCC climate models do not capture Arctic sea ice drift acceleration: Consequences in terms of projected sea ice thinning and decline. *J. Geophys. Res.*, 116, C00D07. doi:10.1029/2011JC007110, 2011.
- Randelhoff, A. and Sundfjord, A.: Short commentary on marine productivity at Arctic shelf breaks: upwelling, advection and vertical mixing. *Ocean Sci.*, 14(2), 293. doi:10.5194/os-14-293-2018, 2018.
- 710 Regan, H., Lique, C., Talandier, C., and Meneghello, G.: Response of Total and Eddy Kinetic Energy to the Recent Spinup of the Beaufort Gyre. *J. Phys. Oceanogr.*, 50, 575–594, doi:10.1175/JPO-D-19-0234.1, 2020.
- Ricker, R., Girard-Arduin, F., Krumpen, T., and Lique, C.: Satellite- derived sea ice export and its impact on Arctic ice mass balance. *Cryosphere* 12(9):3017–3032. doi:10.5194/tc-12-3017-2018, 2018.

- Roloff, A., Rabe, B., Kikuchi, T., Wisotzki, A.: Physical oceanography from 49 XCTD stations during POLARSTERN cruise PS87 (ARK-XXVIII/4). Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, PANGAEA, doi:10.1594/PANGAEA.853770, 2015.
- Rose, S. K., Andersen, O. B., Passaro, M., Ludwigsen, C. A., and Schwatke, C.: Arctic Ocean Sea Level Record from the Complete Radar Altimetry Era: 1991–2018. *Remote Sens.*, 11, 1672, 2019.
- Rudels, B.: Arctic Ocean circulation and variability—Advection and external forcing encounter constraints and local processes, *Ocean Sci.*, 8, 261–286, 2012.
- Rudels, B., Jones, E. P., Anderson, L. G., and Kattner, G.: On the intermediate depth waters of the Arctic Ocean, in *The Polar Oceans and Their Role in Shaping the Global Environment*, Geophys. Monogr. Ser., vol. 85, edited by O. M. Johannessen, R. D. Muench, and J. E. Overland, pp. 33–46, AGU, Washington, D. C., 1994.
- Rudels, B., Jones, E. P., Schauer, U., and Eriksson, P.: Atlantic sources of the Arctic Ocean surface and halocline waters. *Polar Res.*, 23, 181–208, doi:10.3402/polar.v23i2.6278, 2004.
- Sasgen, I., Wouters, B., Gardner, A.S., King, M. D., Tedesco, M., Landerer, F. W., Dahle, C., Save, H., and Fettweis, X.: Return to rapid ice loss in Greenland and record loss in 2019 detected by the GRACE-FO satellites. *Commun. Earth Environ.*, 1, 8, doi:10.1038/s43247-020-0010-1, 2020.
- Sato, K. and Inoue, J.: Comparison of Arctic sea ice thickness and snow depth estimates from CFSR with in situ observations. *Climate Dyn.*, 50, 289-301, 2018:
- Schauer, U. and Losch, M.: “Freshwater” in the Ocean is Not a Useful Parameter in Climate Research. *J. Phys. Oceanog.*, 49(9), 2309–2321. doi:10.1175/JPO-D-19-0102.1, 2019.
- Serreze, M. C., McLaren, A. S., and Barry, R. G.: Seasonal variations of sea ice motion in the Transpolar Drift Stream. *Geophys. Res. Lett.*, 16(8), 811-814, 1989.
- Serreze, M. C., Barrett, A. P., Slater, A. G., Woodgate, R. A., Aagaard, K., Lammers, R. B., Steele, M., Moritz, R., Meredith, M., and Lee, C. M.: The large-scale freshwater cycle of the Arctic, *J. Geophys. Res.*, 111, C11010, doi:10.1029/2005JC003424, 2006.
- Sévellec, F., Fedorov, A. V., and Liu, W.: Arctic sea-ice decline weakens the Atlantic meridional overturning circulation. *Nat. Climate Change*, 7, 604–610, doi:10.1038/nclimate3353, 2017.

- 740 Shepherd, A. and The IMBIE Team: Mass balance of the Greenland Ice Sheet from 1992 to 2018. *Nature*, 579(7798), 1–21. doi:10.1038/s41586-019-1855-2, 2020.
- Shepherd, A., Ivins, E. R., Geruo, A., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Joughin, I., King, M. A., Lenaerts, J. T., Li, J., Ligtenberg, S. R., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R., Milne, G., Mouginot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, 745 E., Rott, H., Sørensen, L. S., Scambos, T. A., Scheuchl, B., Schrama, E. J., Smith, B., Sundal, A. V., van Angelen, J. H., van de Berg, W. J., van den Broeke, M. R., Vaughan, D. G., Velicogna, I., Wahr, J., Whitehouse, P. L., Wingham, D. J., Yi, D., Young, D., and Zwally H. J.: A reconciled estimate of ice-sheet mass balance. *Science*, 338(6111), 1183-1189, doi:10.1126/science.1228102, 2012.
- Shiklomanov, A., Déry, S., Tretiakov, M., Yang, D., Magritsky, D., Georgiadi, A., and Tang, W.: River Freshwater Flux to 750 the Arctic Ocean. In: D. Yang and D. Kane (eds) *Arctic Hydrology, Permafrost and Ecosystems*. Springer, Cham., doi:10.1007/978-3-030-50930-9_24, 2021.
- Simonsen, S. B., Barletta, V. R., Colgan, W. T., and Sørensen, L. S.: Greenland Ice Sheet mass balance (1992–2020) from calibrated radar altimetry. *Geophysical Research Letters*, 48, e2020GL091216. doi:10.1029/2020GL091216, 2021.
- Smedsrud, L. H., Halvorsen, M. H., Stroeve, J. C., Zhang, R., and Kloster, K.: Fram Strait sea ice export variability and 755 September Arctic sea ice extent over the last 80 years. *The Cryosphere*, 11 (1) pp. 65-79. 10.5194/tc-11-65-2017, 2017.
- Solgaard, A. M., Simonsen, S. B., Grinsted, A., Mottram, R., Karlsson, N. B., Hansen, K. et al.: Hagen Bræ: A surging glacier in North Greenland—35 years of observations. *Geophysical Research Letters*, 47, e2019GL085802. doi:10.1029/2019GL085802, 2020.
- Spall, M. A.: Dynamics and thermodynamics of the mean transpolar drift and ice thickness in the Arctic Ocean. *J. 760 Clim.*, 32, 8449–8463, doi:10.1175/JCLI-D-19-0252.1, 2019.
- Spreen, G., Kwok, R., and Menemenlis, D.: Trends in Arctic sea ice drift and role of wind forcing: 1992–2009. *Geophys. Res. Lett.*, 38(19), 2011.
- Spreen, G., de Steur, L., Divine, D., Gerland, S., Hansen, E., and Kwok, R.: Arctic sea ice volume export through Fram Strait from 1992 to 2014. *J. Geophys. Res. Oceans*, 125(6), doi:10.1029/2019JC016039, 2020.
- 765 Steele, M., Morison, J., Ermold, W., Rigor, I., Ortmeier, M., and Shimada, K.: Circulation of summer Pacific halocline water in the Arctic Ocean. *J. Geophys. Res. Oceans*, 109(C2), 2004.

- Storto, A. and Masina, S.: C-GLORSv5: An improved multipurpose global ocean eddy-permitting physical reanalysis. *Earth Syst. Sci. Data*, 8(2), 679, 2016.
- 770 Straneo, F., Curry, R. G., Sutherland, D. A., Hamilton, G. S., Cenedese, C., Våge, K., and Stearns, L. A.: Impact of fjord dynamics and glacial runoff on the circulation near Helheim Glacier. *Nature Geoscience*, 4(5), 322–327. doi:10.1038/ngeo1109, 2011.
- Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons. *Environ. Res. Lett.*, 13(10), 103001, 2018.
- Tamura, T. and Ohshima, K. I.: Mapping of sea ice production in the Arctic coastal polynyas. *J. Geophys. Res. Oceans*, 116(C7), 2011.
- 775 Tedesco, M. and Fettweis, X.: Unprecedented atmospheric conditions (1948-2019) drive the 2019 exceptional melting season over the Greenland ice sheet. *The Cryosphere*, 14, 1209-1223, doi:10.5194/tc-14-1209-2020, 2020.
- Thompson, D. W. J. and Wallace, J. M.: The Arctic Oscillation Signature in Wintertime Geopotential Height and Temperature Fields, *Geophys. Res. Lett.*, 25, 1998.
- 780 Timmermans, M.-L. and Marshall, J.: Understanding Arctic Ocean Circulation: A Review of Ocean Dynamics in a Changing Climate. *J. Geophys. Res. Oceans*, 125(4), C04S02. doi:10.1029/2018JC014378, 2020.
- Timmermans, M. L., Proshutinsky, A., Krishfield, R. A., Perovich, D. K., Richter-Menge, J. A., Stanton, T. P., and Toole, J. M.: Surface freshening in the Arctic Ocean’s Eurasian Basin: An apparent consequence of recent change in the wind-driven circulation. *J. Geophys. Res.*, 116, C00D03, doi:10.1029/2011JC006975, 2011.
- 785 Tsubouchi, T., Våge, K., Hansen, B., Larsen, K. M. H., Østerhus, S., Johnson, C., Jónsson, S., and Valdimarsson, H.: Increased ocean heat transport into the Nordic Seas and Arctic Ocean over the period 1993–2016. *Nature Climate Change* 11, 21-26, 2021.
- Tsubouchi, T., Bacon, S., Naveira Garabato, A. C., Aksenov, Y., Laxon, S. W., Fahrbach, E., Beszczynska-Möller, A., Hansen, E., Lee, C. M., and Ingvaldsen, R. B.: The Arctic Ocean in summer: a quasi-synoptic inverse estimate of boundary fluxes and water mass transformation. *J. Geophys. Res.*, 117, C01024, doi:10.1029/2011JC007174, 2012.
- 790 Tsubouchi, T., Bacon, S., Aksenov, Y., Naveira Garabato, A. C., Beszczynska-Möller, A., Hansen, E., de Steur, L., Curry B., and Lee, C. M.: The Arctic Ocean seasonal cycles of heat and freshwater fluxes: observation-based inverse estimates. *J. Phys. Oceanogr.*, 48, 2029-2055, doi:10.1175/JPO-D-17-0239.1, 2018.

- 795 Vandecrux, B., Fausto, R., Van As, D., Colgan, W., Langen, P. L., Haubner, K., Ingeman-Nielsen, T., Heilig, A., Stevens, C. M., Macferrin, M., Niwano, M., Steffen, K., and Box, J. E.: Firn cold content evolution at nine sites on the Greenland ice sheet between 1998 and 2017. *J. Glaciology*, 66(258), 591-602. doi:10.1017/jog.2020.30, 2020.
- Vihma, T. J., Screen, J., Tjernström, M., Newton, B., Zhang, X., Popova, V., Deser, C., Holland, M., and Prowse, T.: The atmospheric role in the Arctic water cycle: A review on processes, past and future changes, and their impacts. *J. Geophys. Res. Biogeosci.*, 121, 586–620, doi:10.1002/2015JG003132, 2016.
- 800 Villamil-Otero, G. A., Zhang, J., He, J., and Zhang, X.: Role of extratropical cyclones in the recently observed increase in poleward moisture transport into the Arctic Ocean. *Adv. Atmos. Sci.*, 35, 85–94, doi:10.1007/s00376-017-7116-0, 2018.
- Vinje, T., Nordlund, N., and Kvambekk, A.: Monitoring ice thickness in Fram Strait. *J. Geophys. Res. Oceans* 103:10437–10449. doi:10.1029/97jc03360, 1998.
- Volkov, V. A., Mushta, A., and Demchev, D.: Sea Ice Drift in the Arctic. In *Sea Ice in the Arctic*. 301-313. Springer, Cham, 2020.
- 805 Våge, K., Pickart, R. S., Pavlov, V., Lin, P., Torres, D. J., Ingvaldsen, R., Sundfjord, A., and Proshutinsky, A.: The Atlantic Water boundary current in the Nansen Basin: Transport and mechanisms of lateral exchange, *J. Geophys. Res. Oceans*, 121, 6946–6960, doi:10.1002/2016JC011715, 2016.
- 810 Vogt, M., Rabe, B., Kikuchi, T., and Wisotzki, A.: Physical oceanography from 15 XCTD stations during POLARSTERN cruise PS86 (ARK-XXVIII/3 AURORA). Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, PANGAEA, doi:10.1594/PANGAEA.853768, 2015.
- Watkins, M. M., Wiese, D. N., Yuan, D.-N., Boening, C., and Landerer, F. W.: Improved methods for observing Earth's time variable mass distribution with GRACE using spherical cap mascons. *Journal of Geophysical Research: Solid Earth*, 120(4), 2648–2671. doi:10.1002/2014JB011547, 2015.
- 815 Wei, J, Zhang, X., and Wang, Z.: Impacts of extratropical storm tracks on Arctic sea ice export through Fram Strait. *Clim. Dyn.*, 52, 2235-2246, doi:10.1007/s00382-018-4254-8, 2019.
- Webster, M. A., Parker, C., Boisvert, L., and Kwok, R.: The role of cyclone activity in snow accumulation on Arctic sea ice, *Nature Commun.*, 10, 2019.
- Williams, W. and Carmack, C.E.: The ‘interior’ shelves of the Arctic Ocean: Physical oceanographic setting, climatology and effects of sea-ice retreat on cross-shelf exchange. *Prog. Oceanogr.* 139, 24–41. doi:10.1016/j.pocean.2015.07.008, 2015.

820 Wood, K.-R., Overland, J. E., Salo, S. A., Bond, N. A., Williams, W. J., and Dong, X.: Is there a “new normal” climate in the Beaufort Sea? *Polar Res.*, 32, 19552. doi:10.3402/polar.v32i0.19552, 2013.

Woodgate, R. A.: Increases in the Pacific inflow to the Arctic from 1990 to 2015, and insights into seasonal trends and driving mechanisms from year-round Bering Strait mooring data. *Prog. Oceanog.*, 160, 124–154. doi:10.1016/j.pocean.2017.12.007, 2018.

825 Yamamoto-Kawai, M., McLaughlin, F. A., Carmack, E. C., Nishino, S., Shimada, K., and Kurita, N.: Surface freshening of the Canada Basin, 2003–2007: River runoff versus sea ice meltwater, *J. Geophys. Res.*, 114, C00A05, doi:10.1029/2008JC005000., 2009.

Zhang, X., Ikeda, M., and Walsh, J.: Arctic sea ice and freshwater changes driven by the atmospheric leading mode in a coupled sea ice–ocean model. *J. Clim.*, 16:2159–2177, doi:10.1175/2758.1, 2003.

830 Zhang, X., Sorteberg, A., Zhang, J., Gerdes, R. and Comiso, J. C.: Recent radical shifts of atmospheric circulations and rapid changes in Arctic climate system. *Geophys. Res. Lett.* 35, L22701, 2008.

Zhang, X., He, J. Zhang, J., Polyakov, I., Gerdes, R., Inoue, J., and Wu, P.: Enhanced poleward moisture transport and amplified northern high-latitude wetting trend. *Nat. Climate Change*, 3, 47–51, doi:10.1038/nclimate1631, 2013.

835 Zhong, W., Zhang, J., Steele, M., Zhao, J., and Wang, T.: Episodic extrema of surface stress energy input to the western Arctic Ocean contributed to step changes of freshwater content in the Beaufort Gyre. *Geophys. Res. Lett.*, 46, 12173– 12182, doi:10.1029/2019GL084652, 2019.

Zuo, H., Balmaseda, M. A., Tietsche, S., Mogensen, K., and Mayer, M.: The ECMWF operational ensemble reanalysis-analysis system for ocean and sea-ice: a description of the system and assessment, *Ocean Sci.*, 15, 779–808, doi:10.5194/os-15-779-2019, 2019.