

# Laboratory experiments on the influence of stratification and a bottom sill on seiche damping

Karim Medjdoub<sup>1</sup>, Imre M. Jánosi<sup>2,3</sup>, and Miklós Vincze<sup>1,4</sup>

<sup>1</sup>von Kármán Laboratory of Environmental Flows; Eötvös Loránd University, Pázmány P. sétány 1/A, Budapest H-1117, Hungary

<sup>2</sup>Max Planck Institute for the Physics of Complex Systems, Nöthnitzer Str. 38, 01187 Dresden, Germany

<sup>3</sup>University of Public Service, Faculty of Water Sciences, Ludovika sqr. 1, H-1083 Budapest, Hungary

<sup>4</sup>MTA-ELTE Theoretical Physics Research Group, Pázmány P. sétány 1/A, Budapest H-1117, Hungary

**Correspondence:** Miklós Vincze (mvincze@general.elte.hu)

**Abstract.** The damping of water surface standing waves (seiche modes) and the associated excitation of baroclinic internal waves are studied experimentally in a quasi-two-layer laboratory setting with a topographic obstacle at the bottom, representing a seabed sill. We find that topography-induced baroclinic wave drag indeed contributes markedly to seiche damping in such systems. Two major pathways of barotropic-baroclinic energy conversions were observed: the stronger one – involving short-  
5 wavelength internal modes of large amplitudes – may occur when the node of the surface seiche is situated above the close vicinity of the sill. The weaker, less significant other pathway is the excitation of long waves, internal seiches along the pycnocline that may resonate with the low frequency components of the decaying surface forcing.

## 1 Introduction

Energy conversion over bathymetric formations is a key component of the global ocean dynamics and mixing (Wunsch and  
10 Ferrari, 2004). The coupling between barotropic tidal waves at the sea surface and internal gravity waves facilitates heat and material exchange between the uppermost and deeper layers (Garrett, 2003; Lelong and Kunze, 2013; Morozov, 2018; Vic et al., 2019; Rippeth and Green, 2020; Stanev and Ricker, 2020). This largely interconnected dynamics is particularly pertinent in semi-enclosed basins, bays, and fjords with density profiles characterized by sharp gradients (Rattray Jr., 1960; Niiler, 1968; Bell Jr., 1975; Stigebrandt, 1980; Stigebrandt and Aure, 1989; Chapman and Giese, 1990; Münnich, 1996; Parsmar and  
15 Stigebrandt, 1997; Stigebrandt, 1999; Antenucci and Imberger, 2001, 2003; Inall et al., 2004; Cushman-Roisin et al., 2005; Johnsson et al., 2007; Boegman and Ivey, 2012; Park et al., 2016; Staalstrøm and Røed, 2016; Castillo et al., 2017; Roget et al., 2017; Stanev and Ricker, 2020; Xue et al., 2020).

The exchange between waves at the surface and at the pycnocline can be especially well studied in the situation where standing waves develop at the water surface, known as seiches (Chapman and Giese, 2001; de Carvalho Bueno et al., 2020).  
20 Such oscillatory motions of a water body are typically initiated by temporally changing wind stress, especially pulse-like wind bursts, or storms. Strong wind shear gets balanced by a certain tilt of the water surface throughout the basin, but when the storm subsides, the restoring forces, predominantly gravity, yield a damped “sloshing” of the free surface, until it settles in its

equilibrium (horizontal) position. Tidal barotropic waves, even tsunamis and other seismic disturbances are known to generate large inflows into coastal harbors and may also yield strong seiche activity (Chapman and Giese, 2001).

25 The energy dissipation rate (or decay rate) of surface seiches in natural enclosed water bodies, e.g. lakes or fjords is determined by the physical properties of the fluid body (e.g. its stratification profile) and the geometry of the basin. In the presence of bottom topography, surface gravity waves generate internal waves, a process also referred to as barotropic to baroclinic energy conversion, or – in the specific case of tidal surface waves – tidal conversion. The first demonstration of such a wave excitation mechanism over a tilted bottom topography dates back to the work of Rattray Jr. (1960) in the case of a two-layer stratification. Niiler (1968) explained the observed current oscillation at the Florida Straits with the interaction of barotropic tides and the continental slope that yields a generation of baroclinic tidal waves, periodically modulating the flow. Although these pioneering studies did not consider the dissipative effect of the excited baroclinic modes on the surface waves, it is obvious that freely propagating internal waves use up a significant fraction of the energy stored in the surface oscillations, and hence speed up the decay of the latter.

35 There are, however, substantial differences between the seiche modes excited in enclosed and semi-enclosed basins. In the former configurations the boundary conditions “select” such standing waves that have antinodes at all boundaries (at least in the idealized case of vertical sidewalls), whereas, in the semi-enclosed systems the open boundary (e.g. the mouth of the bay) facilitates maximum horizontal flux, leading to the formation of a wave node at the surface. Obviously, the internal waves generated by the waves over topographic obstacles may also exhibit different behaviour in the two settings. In the semi-infinite domain travelling internal waves are excited which “radiate” away the kinetic energy of the surface waves when the topographic obstacles at the bottom reach up to the pycnocline (Davies et al., 2009). If, however, the basin is closed, resonant mode selection can also occur as internal seiches are generated at the pycnocline.

The effect of damping on the frequencies of the seiche modes was addressed by the paper of Cushman-Roisin et al. (2005) who concluded that in enclosed water bodies the oscillation periods of surface seiche modes do not depend on the stratification significantly. However, in semi-enclosed systems the surface seiche period was found to be more sensitive to the stratification. This effect of internal wave excitation on the decay of surface seiches was further investigated by Wynne et al. (2019). When the characteristic (e.g. diurnal) frequency of the wind forcing matches that of a seiche eigenmode, a resonant amplification of the latter may occur. However, as the seasonal changes of the freshwater inflow (e.g. from glacier runoff) influence the depth of the pycnocline, the natural frequencies of the internal waves also change, and – as the resonant frequencies shift – so does the decay coefficient of the dominant surface seiche mode, as shown by Antenucci and Imberger (2001, 2003).

55 The basin geometry of certain fjords, e.g. the Gullmar fjord of Sweden is especially interesting, as their bottom topography involves a sill reaching up to the pycnocline between the saline seawater and the upper freshwater layer. Parsmar and Stigebrandt (1997) investigated the damping of tidally excited surface seiches in the Gullmar fjord and concluded that the drag caused by baroclinic internal wave excitation at the sill contributes far more to the damping of the seiche than bottom friction or other phenomena.

An earlier work in our laboratory (Vincze and Bozóki, 2017) investigated interfacial internal wave generation experimentally in a quasi-two-layer wave-tank with a bottom obstacle at its center, reaching up to the pycnocline, motivated by the fjord

60 geometry. We found that shear instability developing at the tip of the obstacle and the emerging billows press down the pycnocline at the lee side of the obstacle (with respect to the direction of the oscillating horizontal flow in the upper layer) and generate the observed internal waves. However, the energetics of the damping of surface seiches was not addressed there.

Internal wave excitation by surface seiche modes has also been investigated in laboratory experiments by Boegman and Ivey (2012). They modeled resonant barotropic-baroclinic energy conversion of a periodically forced shallow water system in a rectangular tank without topographic obstacles in order to quantify the energy flux pathways between the applied forcing and the internal wave field. A similar setting (i.e. a rectangular tank with a flat bottom) has also been studied recently by Xue et al. (2020), with a special focus on internal wave excitation and seiche damping. **Similar laboratory models can provide illuminating results for the interpretation of direct oceanic observations, because they make possible to focus on specific phenomena that are inevitably loaded with disturbing effects in natural environments.**

Here we report on laboratory experiments in a narrow water tank, filled up with quasi two-layer salinity stratified water, with a vertical obstacle installed in the middle of the domain in the bottom layer. We analyze the damping of various surface and internal wave modes that develop after pulse-like initial seiche excitation at the water surface. To the best of our knowledge, these are the first experiments in the literature on surface seiche damping in the presence of a bottom sill yielding considerable baroclinic wave drag.


The paper is organized as follows. Section 2 describes the experimental set-up and the applied data acquisition methods. In Section 3 we present our results on the relationship between surface seiche damping and the physical and geometrical properties of the experimental settings. Finally, we briefly discuss and summarize our findings in Section 4.

## 2 Set-up and methods


Our experiments were conducted in a transparent rectangular acrylic (Plexiglas) tank of length  $L = 80$  cm and width  $w = 5$  cm, filled up with quasi-two-layer stratified water, as shown in Fig. 1a. The bottom layer of thickness  $H_2 = 4$  cm consisted of saline water colored with blue food dye, whereas the upper layer of thickness  $H_1$  was freshwater of the same temperature, colored red. To inhibit mixing at the interface (i.e. diapycnal mixing) during the filling-up procedure the freshwater of the upper layer was entering the surface slowly, through a sponge.

With each prepared stratification profile two configurations were studied. One with an upside-down U-shaped small plastic obstacle (Fig. 1a) that was placed in the middle of the tank to the bottom. The obstacle's height  $h$  matched the thickness of the bottom (saline) layer:  $h = H_2 = 4$  cm. In the other series of experiments the obstacle was removed from the tank (slowly, in the vertical direction, to minimize turbulent mixing) and the dynamics was studied in the absence of the topographic obstacle. **The experimental parameters and the list of other notations used throughout the paper are given in Table 1.**

Each experiment series consisted of five runs with and five runs without the obstacle. A run was initiated as follows. A suspended wave maker consisting of a metallic rod and six rectangular rubber foam "bumpers" (Fig. 1b) was placed above the water surface. The bumpers could be placed in various configurations along the rod to facilitate the excitation of surface seiche (standing wave) modes, as sketched in Fig. 1b. The standing wave pattern with a dominant mode of  $m = 5$  – where  $m$



**Figure 1.** (a) The schematics of the set-up. The geometrical parameters of the tank are  $L = 80$  cm,  $w = 5$  cm, and  $H_2 = h = 4$  cm. The shape of the obstacle is sketched in the bottom right corner ( $h = 4$  cm,  $r = 1.75$  cm). (b) The configurations of the six rubber foam bumpers of the wave maker for the excitation of the various surface seiche modes and the sketch of the corresponding waveform. (c) The vertical motion of the bottom of the rubber foam bumpers at the initiation. The position is given with respect to the unperturbed water surface. The width of the line represents the error of reproducibility.



**Figure 2.** Density  $\rho$  as a function of vertical position  $z$  from the bottom, for all experiments series (cf. Table 2). The sketch in the inset demonstrates the meaning of the parameters of the two-layer approximation.

denotes the number of nodes of the standing waveform – was found to be very hard to excite with this device, therefore our analysis was restricted to experiments with dominant surface seiche modes  $m = 1, 2, 3, 4,$  and  $6$ . In the beginning of each



**Table 1.** List of parameters and notations.

notation	meaning	value	unit
$L$	length of the tank	80	cm
$w$	width of the tank	5	cm
$H$	total fluid depth	15;17;19	cm
$H_1$	upper layer depth	11;13;15	cm
$H_2$	bottom layer depth	4	cm
$h(= H_1)$	obstacle height	4	cm
$\rho_1$	upper layer density	0.998	kg/l
$\rho_2$	bottom layer average density	1.001–1.114	kg/l
$\Delta\rho(= \rho_2 - \rho_1)$	density contrast	0.003–0.116	kg/l
$k$	wavenumber		rad/cm
$m$	dominant surface mode number	1;2;3;4;6	–
$\omega$	angular frequency		rad/s
$f(= \omega/(2\pi))$	frequency		Hz
$\lambda(= 2\pi/k)$	wavelength		cm
$\eta$	surface displacement		cm
$\chi$	pycnocline displacement		cm
$A_i$	amplitude of surface wave component $i$		cm
$C_i$	decay coefficient of surface wave component $i$		1/s
$\varphi_i$	phase of surface wave component $i$		rad

**Table 2.** Geometrical and physical characteristics of the experiment series.

Experiment series	#1	#2	#3	#4	#5	#6	#7	#8	#9
$H_1$ (cm)	11	11	15	15	13	13	13	13	13
$\rho_2$ (kg/l)	1.0592	1.0076	1.0872	1.0019	1.1114	1.1136	1.1114	1.0437	1.0084

run, the suspended rod was pushed down (only once) to the water surface, reaching a maximum depth of  $1.65 \pm 0.25$  cm in a standardized manner and was then instantaneously pulled up, out of the water. The vertical motion of the bottom of the bumper at initiation is shown in Fig. 1c, as acquired from video tracking. Apparently, the characteristic timescale of the impact, i.e. the duration that the bumper spent below the unperturbed water height (indicated with dashed line) was approximately  $(0.4 \pm 0.06)$  s, i.e. shorter than the basin-crossing timescale of a surface perturbation.

The prepared density profiles of the experiment series are presented in Fig. 2, as measured by a conductivity probe. The physical parameters of the two-layer approximation, namely layer thicknesses  $H_1$  and  $H_2$  and the respective densities  $\rho_1$  and


100  $\rho_2$  were obtained as follows.  $H_1$  and  $H_2$  were measured directly (with a ruler) based on their color (and indeed, the intended thickness  $H_2 = h = 4$  cm matching the obstacle height could be achieved within  $\pm 0.25$  cm in all cases). The characteristic density  $\rho_2$  of the saline layer was taken as an average of the measured density profile in the  $z \leq H_2$  domain, as sketched in the inset of Fig. 2. The relevant adjustable parameters of the stratification are summarized in Table 2 for all nine experiment settings.

105 The damping of the initiated surface sloshing and the internal dynamics were recorded by an HD camera (at frame rate 30 fps and frame size 1080 px  $\times$  1920 px) facing the long sidewall of the tank perpendicularly, close to the middle of the tank. Afterwards, the open source correlation based feature tracking software Tracker (<https://physlets.org/tracker/>) was used to acquire the time series of surface and pycnocline motion.

### 3 Results

#### 110 3.1 Qualitative description

The flow dynamics in the set-up is driven by the decaying surface waves initiated by the aforementioned standardized instantaneous push of the wave maker. In the qualitative sense, the decay of the resulting surface standing waves is not affected by the presence or absence of the bottom obstacle (although the actual values of the decay rates are substantially different in the two configurations as will be discussed later). However, the dynamics at the pycnocline can be vastly different in the two cases: if the obstacle is installed, velocity shear can develop between the bottom and top layers that yields the excitation of baroclinic (internal) wave modes along the pycnocline (Vincze and Bozóki, 2017). Whereas in the control runs – i.e. without obstacle – the vertical displacement of the surface and the pycnocline is found to be co-aligned at each time instant, as if in a homogeneous fluid (barotropic wave modes).




**Figure 3.** (a) Snapshots from experiments with  $m = 1$  (top) and  $m = 2$  (bottom) surface seiche modes (configuration #3, cf. Table 2).

Yet, even if the obstacle is present, large amplitude internal waves do not necessarily get excited, as demonstrated in Figs. 3 and 4. Fig. 3 shows snapshots from two experiments, characterized by the leading mode of the surface seiche that is  $m = 1$  (base mode) and  $m = 2$  in the top and bottom images, respectively. One can notice that in the former case, where a node of

the surface wave form is located at the center of the tank, indicating purely horizontal flow right above the obstacle, slowly propagating internal waves of short wavelength appear “radiating away” from the obstacle with fairly large amplitudes. In the latter experiment, however, where the surface displacement has an antinode – and hence vertical oscillation of the fluid parcels – at the obstacle location, no progressive internal waves are visible.

125 – at the obstacle location, no progressive internal waves are visible.



**Figure 4.** Space-time plots of the two experiments of Fig. 3, showing the vertical displacement of the surface  $\eta(x,t)$  (a) and the pycnocline  $\chi(x,t)$  (b) in the whole tank for dominant surface wave modes  $m = 1$  (left) and  $m = 2$  right (obstacle location is at  $x = 40$  cm). Vertical dashed lines in the top panels denote the expected locations of nodes.

The corresponding space-time plots showing the displacement of the surface  $\eta(x,t)$  and of the pycnocline  $\chi(x,t)$  with respect to their initial levels are presented in Figs. 4a and 4b, respectively, as obtained via video analysis. The dashed vertical lines in Fig. 4a represent the locations of the nodes of the dominant mode ( $m = 1$  in the left and  $m = 2$  in the right panels). After the initiation of the surface seiche at  $t = 0$ , propagating baroclinic waves dominate the picture in the  $m = 1$  run (Fig. 4b left); later the waves get reflected at the sidewall and thus evolve into internal standing wave-like interference patterns. In the

130

$m = 2$  case, the large-scale decaying barotropic oscillation characterizes the entire domain with hardly any noticeable slowly propagating structures (Fig. 4b right). This duality of internal wave excitation is apparent throughout the higher seiche modes as well: antisymmetric surface wave forms associated with odd values of  $m$  (see Fig. 1b) tend to excite larger baroclinic wave activity via the obstacle placed in the geometrical center than the symmetric waves (even values of  $m$ ).

### 135 3.2 Surface waves

Vertical displacement time series of the water surface  $\eta(x, t)$  and the pycnocline  $\chi(x, t)$  were logged at the vicinity of the sidewall (i.e. at  $x \approx 0$  or  $L$ ) in each experiment to ensure that all standing wave modes have antinode at the measurement location. **Note, that due to the boundary conditions only here it is guaranteed that all standing wave components exhibit maximum amplitude.** Examples of such records are shown in the panels of Fig. 5a and b for the  $\eta$  and  $\chi$  signals (red and blue curves, respectively) from runs with dominant modes  $m = 1, 2$ , and 4 (from the left).

The surface time series acquired at location  $x = 0$  can be approximated as a sum of decaying sinusoidal oscillations in the following form:


$$\eta(0, t) \approx \sum_{i=1}^N A_i \exp(-C_i t) \sin(\omega_i t + \varphi_i) , \quad (1)$$

where  $\omega_i$ ,  $\varphi_i$  and  $A_i$  denote the frequency, phase shift and initial amplitude of the  $i$ -th component, respectively, whereas  $C_i(\omega_i)$  is the decay coefficient of the given mode. **Eq. (1) was fitted to the time series with  $N = 2$  as the limit. The standard deviations of the residuals indicated that such two-term sums were sufficient to account for over 90% of the observed displacements in all cases and in many cases even the single-mode fit ( $N = 1$ ) was enough to reach the same precision. It is to be noted here that by using Eq. (1), the frequency-dependant coefficients of decay  $C_i(\omega_i)$  could theoretically be extracted from the “sloshing” of the water surface with any arbitrary initial condition. We chose our initiation method with the different bumper configurations (shown in Fig. 1b) representing “quasi pure” modes only for the practical reason of making the regressions to the time series as simple as reasonably achievable.**

The Fourier amplitude spectra  $A_\eta(f)$  and  $A_\chi(f)$  of the  $\eta(t)$  and  $\chi(t)$  signals are presented in Fig. 5c with a color coding identical to the corresponding time series themselves. The frequencies of the largest spectral peaks of  $A_\eta(f)$  from each experiment were found to be in good agreement (i.e. **within 5%**) with the linear dispersion relation of homogeneous (non-stratified) surface gravity waves, which reads as

$$\omega = \sqrt{gk \tanh(kH)} , \quad (2)$$

where  $H$  is the total water depth,  $g$  is the gravitational acceleration, and the wave number  $k$  is to be taken at  $k(m) = \pi m/L$  for the dominant standing wave mode excited in the given run (cf. Fig. 1). The match is demonstrated in Fig. 6. In terms of the  $\omega(m)$  relation, no systematic bias was found from this simple (linear, undamped) theoretical formula regardless of whether the experiment included the obstacle or not, or even whether the water was stratified or homogeneous.




**Figure 5.** Time series and spectra from three exemplary runs (configuration #1, cf. Table 2) of dominant modes  $m = 1$ ,  $m = 2$ , and  $m = 4$  from the left. The panels of row a) show the water surface displacement as a function of time  $t$  from the initiation, as tracked in the close vicinity of the sidewall. The displacement of the pycnocline  $\chi$  at the sidewall is presented in row b) (note, that here, unlike in the  $\eta(t)$  plots, the vertical range is different in all three panels). The Fourier spectra of the  $\eta(t)$  (red) and  $\chi(t)$  (blue) records are visible in row c), whereas the panels in d) show the transfer function  $T(f) = A_\chi(f)/A_\eta(f)$ , i.e. the ratio of the spectra of the corresponding two spectra above.

### 3.3 Source-filter dynamics

The surface waves exhibit faster damping than their internal counterparts and the dominant frequency components of the two may also largely differ. It is clearly visible from Fig. 5 that the “internal”  $\chi(t)$  signals (panels in row b) possess more pronounced low-frequency variability than the  $\eta(t)$  oscillations (row a), even in the  $m = 2$  and 4 modes, where the classic, topography-induced baroclinic wave generation is practically inhibited – due to the lack of considerable velocity shear at the obstacle – as discussed earlier.

To understand this mechanism it is important to note that even if the surface seiche was a perfectly “monochromatic”, single-frequency source signal, its exponential decay would still unavoidably introduce nonzero amplitudes into the low-frequency



**Figure 6.** (a) The dispersion relation of water surface waves Eq. (2) expressed as a function of dominant mode index  $m$ . The solid curves represent the theoretical relationship for the shallowest ( $H = 15$  cm) and deepest ( $H = 19$  cm) configurations, and the data points indicate the measured values of angular frequency  $\omega$ . The coloring is based on the total water height  $H$ . (b) Contrasting the calculated angular frequency  $\omega$  from Eq. (2) to its measured value. The various experiment configurations are indicated by the shapes of the symbols in both panels (see legend).

range of its spectrum (see, e.g. French (1971)), making it suitable for the excitation of slow internal oscillations. The resulting  
 170 signal at the pycnocline can thus be understood as the outcome of a resonance-like amplification of certain characteristic  
 frequency bands of the surface source signal.

The transfer function  $T(f)$  of this frequency “filtering”, a widely used tool, e.g. in acoustics (Stevens, 2001), can be defined  
 as  $T(f) = A_\chi(f)/A_\eta(f)$ . Such empirical transfer functions for the experiments of Fig. 5 are shown in the panels of row d).  
 In all cases, the maxima of  $T(f)$  appear well below the fundamental frequencies  $f_i = \omega_i/(2\pi)$  of the source signal Eq. (1),  
 175 consistently with the rule of thumb that interfacial internal propagation is typically around  $\sqrt{\Delta\rho/\rho_0}$  times slower than that  
 of surface waves. The question arises of what kind of process determines the observed low-frequency amplification bands of  
 $T(f)$ .

In the two-layer approximation, the dispersion relation of small amplitude interfacial internal waves (Massel, 2015) takes  
 the form

$$180 \quad \omega(k) = \sqrt{\frac{gk\Delta\rho}{\rho_1 \coth(H_1 k) + \rho_2 \coth(H_2 k)}} \quad , \quad (3)$$

where  $k$  denotes the wavenumber (in this case, along the pycnocline),  $H_1$  and  $H_2$  are the thicknesses and  $\rho_1$  and  $\rho_2$  are the densities of the top and bottom fluid layers, respectively (cf. Table 2), and  $\Delta\rho = \rho_2 - \rho_1$  is their difference.

185 A straightforward way to contrast the theoretical dispersion relation with the observed wave propagation is taking the two-dimensional Fourier transform of space-time plots like the ones shown in Fig. 4. Then, the time axis is mapped onto the frequency domain  $\omega$  and the spatial axis is transformed to the wavenumber ( $k$ ) space. Marked spectral amplitudes are expected to occur along the  $\pm\omega(k)$  dispersion relation curves. Examples of such spectra are presented in Fig. 7 for the cases of dominant surface seiche modes  $m = 4$  and  $m = 6$  in experiment series #3 (see Table 2) for the surface (panels a) and the pycnocline (panels b). In the surface spectra of panels a) the “quasi pure” initiation yielded sharp peaks (cf. Fig. 5) corresponding to the dominant modes (white circles) that indeed lie along the surface dispersion relation Eq. (2), plotted with black lines (cf. 190 Fig. 6a). The spectra of the pycnocline motion exhibit much more disperse responses, nevertheless we also find that the formula of Eq. (3) shown again with black lines match the observed distribution of large amplitudes in the low wave-number domain. (Towards higher values of  $k$ , three-layer corrections may be necessary as shown e.g. in Vincze and Bozóki (2017).)


Combining the transfer functions  $T(f)$  of all five experimental runs (corresponding to different dominant  $m$  values) in a given stratification setting, one can define a cumulative transfer function  $\bar{T}(f)$  by assigning the largest value of  $T(f)$  obtained 195 throughout all five runs to each frequency  $f$ , as represented by the gray shaded area of the combined spectrum in Fig. 8a. The dispersion relation Eq. (3) enables us to transform the  $T(f)$  functions to the wave-number domain, as shown in Fig. 8b for four selected stratification settings. Here the frequency  $f = \omega/(2\pi)$  is given on the horizontal axis and the wave-number on the vertical one is expressed in the nondimensional units of  $L/\lambda = Lk/(2\pi)$ , with  $\lambda$  being the internal wavelength and  $L$  the length of the tank. (This is a rescaled inverse of the positive branches of the solid black curves in Fig. 7b.) The curves therefore 200 represent the inverse of the dispersion relation Eq. (3), and their color scale marks the corresponding normalized cumulative transfer function  $\bar{T}(f)$  (the normalization was carried out by setting the maximum of the  $T(f)$  function as unity for each stratification).

By reorganizing the cumulative transfer functions this way, one can notice that the amplification peaks (high values of  $\bar{T}$ , colored yellow) tend to appear in the close vicinity of integer values of the nondimensional wave number, e.g. at  $L/\lambda =$  205 1; 2; 3; 5 and 8, as marked by dotted horizontal lines in Fig. 8b. These low frequency resonance peaks are thus seem to be related to standing wave generation at the pycnocline – internal seiche modes – whose wavelengths can fit integer times into the tank. Despite having much lower oscillation frequencies than the main peak of the “source” signal, these modes can still get excited. It is due to the fact that the damping of the surface seiche excites considerable amplitudes in the low-frequency spectral domain with which the internal standing waves can resonate. It is to be emphasized that combining the 210  $L/\lambda = n/2$  (where  $n = 1, 2, \dots$ ) criterion with the dispersion relation Eq. (3) provides the same information as calculating the corresponding internal seiche frequencies directly from the geometrical parameters of the setting.

### 3.4 Topographic energy conversion

The damping of surface waves in a stratified system is caused by the combination of factors such as the friction with the basin boundaries, diapycnal mixing, and barotropic-to-baroclinic energy conversion due to topographic effects. To separate






**Figure 7.** Two-dimensional Fourier transforms of surface displacement  $\eta(x,t)$  (a) and of the pycnocline displacement  $\chi(x,t)$  (b) in the tank for dominant surface seiche mode  $m = 4$  (left) and  $m = 6$  (right), experimental configuration #3. The white circles mark the dominant surface wave peak, whereas the black solid lines represent the respective dispersion relations Eqs. (2) and (3).

215 the contribution of the latter we conducted control experiments with each studied stratification profile where the obstacle was removed from the tank. We then compared the observed decay rates with the ones acquired with the obstacle in place.

The scatter plot in Fig. 9a contrasts the the decay coefficient of the dominant surface seiche mode  $C_o$  – from fitting formula (1) – in each experiment with obstacle (hence the index  $o$ ), with the corresponding no-obstacle value  $C_{no}$ . It is clearly visible that almost all data points spread in the  $C_o > C_{no}$  domain, but no systematic relationship could be found with the density ratio  
 220 of the two layers (indicated by the coloring), or total water height  $H$  (marked by the different symbol shapes, see legend).

There is, however, a marked difference between the two configurations in terms of the  $m$ -dependence of the decay coefficients, as shown in Fig. 9b, where the  $C_o/C_{no}$  ratios are plotted for each experiment pair as a function of mode index  $m$ .






**Figure 8.** (a) The transfer functions  $T(f)$  of all dominant excitation modes  $m$  in experiment series #1, and their cumulative transfer function (grey shaded area). (b) The internal wave (inverse) dispersion relations as a function of frequency  $f$  for four exemplary stratification settings (representing series #1, #2, #3, and #4). The coloring marks the normalized cumulative transfer functions  $\bar{T}(f)$  as acquired from the corresponding experiment series.

Apparently, the relative increase of barotropic energy dissipation (damping) compared to the no-obstacle setting is the largest – up to a factor of 2.1 – at the odd wave-numbers  $m = 1$  and 3 (note that the  $m = 5$  mode could not be excited due to technical reasons, as mentioned earlier). This is in concert with the qualitative description provided in subsection 3.1, where we demonstrated that large amplitude internal waves having the same frequency as the surface wave are excited when the obstacle is situated below a surface seiche node.

#### 4 Conclusions

In this experimental study, we analyzed the coupling between surface seiche modes and internal wave dynamics in a quasi-two-layer stratified system, and the effect of a topographic obstacle on the damping of the surface seiche. The decaying surface oscillations were initiated by a wave maker in a pulse-like manner, and the adjustable shape of this device enabled us to excite surface standing waves of rarely observable larger wave-numbers.



**Figure 9.** (a) Comparison of dominant mode surface seiche decay coefficients in paired experiments without (horizontal axis) and with (vertical axis) bottom obstacle. The solid line represents  $y = x$ . (b) The ratio of the damping coefficients  $C_o/C_{no}$  of paired experiments with and without obstacle as a function of dominant surface seiche mode index  $m$ . Symbol shapes and coloring mark the total water depth and the density ratio of the two layers, respectively, in both panels (see legend).

Two pathways of barotropic-to-baroclinic conversion were uncovered. Firstly, the “direct” excitation of short-wavelength propagating internal waves via the horizontal velocity shear emerging in the vicinity of the obstacle. These waves were easily noticeable along the pycnocline due to their relatively large amplitudes. Their periods matched the dominant sloshing timescale associated with the damping surface seiche. It is to be noted that the most pronounced wave excitation, and thus the most prominent increase in the damping coefficient of the surface seiche was detected in the situations where the surface (standing) waveform was such that a node of the surface displacement was located above the bottom obstacle, that – in the geometry investigated here – means seiches of odd wave-number indices.

Secondly, we found evidence for the excitation of slow, long wavelength internal seiche modes, whose half wavelength can fit roughly integer times onto the pycnocline between one sidewall and the obstacle (i.e. onto the half of the total basin length). The occurrence of such internal oscillations in this system is interesting as they have much lower characteristic frequencies than the surface seiche, i.e. the source signal. It appears, indeed, that a nontrivial source-filter dynamics can describe this phenomenon: the spectral structure of the decaying source signal includes **non-zero amplitudes in the low-frequency range** that can resonate with certain internal standing wave modes whose wavelengths are such that they fulfil the geometrical boundary conditions, representing a “band-pass filtering”.

Comparing the dominant seiche decay rates with those from control runs without bottom topography, we demonstrated that in almost all cases internal wave activity yields a detectable increase of the damping, also for even seiche mode indices, where the “short wave” excitation is inhibited, as discussed above. In these cases internal wave dynamics can be attributed to the aforementioned slow wave (internal seiche) excitation pathway, but even this lesser effect appeared to be enough to yield substantially larger damping than the flat-bottom control runs.

Future research is planned to address the source-filter feedback effects in the system to better understand the relationship between the decay rates of the waves at the surface and the ones along the pycnocline. and to explore the dynamics of a more realistic, three-dimensional configuration, possibly involving the effect of Earth’s rotation.

*Author contributions.* K.M. developed the set-up and performed the experiments. M.V., I.M.J. and K.M. conducted the evaluation of the data. All authors have contributed to the interpretation of results and the writing of the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We are thankful for the fruitful discussions with T. Tél and for the highly useful comments of the three anonymous reviewers. This paper was supported by the National Research, Development and Innovation Office (NKFIH) under Grant FK125024. The work of K. M. is supported by the Stipendium Hungaricum Scholarship of the Tempus Public Foundation. This paper is also supported by the ÚNKP-18-4 New National Excellence Program (M.V.) of the Ministry of Human Capacities of Hungary.

## References

- Antenucci, J. P. and Imberger, J.: Energetics of long internal gravity waves in large lakes, *Limnol. Oceanogr.*, 46, 1760–1773, <https://doi.org/https://doi.org/10.4319/lo.2001.46.7.1760>, 2001.
- 265 Antenucci, J. P. and Imberger, J.: The seasonal evolution of wind/internal wave resonance in Lake Kinneret, *Limnol. Oceanogr.*, 48, 2055–2061, <https://doi.org/https://doi.org/10.4319/lo.2003.48.5.2055>, 2003.
- Bell Jr., T. H.: Topographically generated internal waves in the open ocean, *J. Geophys. Res.* (1896-1977), 80, 320–327, <https://doi.org/https://doi.org/10.1029/JC080i003p00320>, 1975.
- Boegman, L. and Ivey, G. N.: The dynamics of internal wave resonance in periodically forced narrow basins, *J. Geophys. Res.: Oceans*, 117, <https://doi.org/https://doi.org/10.1029/2012JC008134>, 2012.
- 270 Castillo, M. I., Pizarro, O., Ramírez, N., and Cáceres, M.: Seiche excitation in a highly stratified fjord of southern Chile: the Reloncaví fjord, *Ocean Sci.*, 13, 145–160, <https://doi.org/https://doi.org/10.5194/os-13-145-2017>, <https://os.copernicus.org/articles/13/145/2017/>, 2017.
- Chapman, D. and Giese, G.: Seiches, in: *Encyclopedia of Ocean Sciences* (2nd Edition), edited by Steele, J. H., pp. 5:344–350, Academic Press, Oxford, <https://doi.org/https://doi.org/10.1016/B978-012374473-9.00128-4>, 2001.
- 275 Chapman, D. C. and Giese, G. S.: A model for the generation of coastal seiches by deep-sea internal waves, *J. Phys. Oceanogr.*, 20, 1459–1467, [https://doi.org/https://doi.org/10.1175/1520-0485\(1990\)020<1459:AMFTGO>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0485(1990)020<1459:AMFTGO>2.0.CO;2), 1990.
- Cushman-Roisin, B., Willmott, A. J., and Biggs, N. R.: Influence of stratification on decaying surface seiche modes, *Cont. Shelf Res.*, 25, 227–242, <https://doi.org/https://doi.org/10.1016/j.csr.2004.09.008>, 2005.
- Davies, A. M., Xing, J., and Willmott, A. J.: Influence of open boundary conditions and sill height upon seiche motion in a gulf, *Ocean dynamics*, 59, 863–879, 2009.
- 280 de Carvalho Bueno, R., Bleninger, T., Yao, H., and Rusak, J. A.: An empirical parametrization of internal seiche amplitude including secondary effects, *Env. Fluid Mech.*, Published online: 18 September, <https://doi.org/https://doi.org/10.1007/s10652-020-09767-1>, 2020.
- French, A.: *Vibrations and Waves*, M.I.T. introductory physics series, Taylor & Francis, <https://books.google.hu/books?id=RqE26vDmd5wC>, 1971.
- 285 Garrett, C.: Internal tides and ocean mixing, *Science*, 301, 1858–1859, <https://doi.org/https://doi.org/10.1126/science.1090002>, 2003.
- Inall, M., Cottier, F., Griffiths, C., and Rippeth, T.: Sill dynamics and energy transformation in a jet fjord, *Ocean Dyn.*, 54, 307–314, <https://doi.org/https://doi.org/10.1007/s10236-003-0059-2>, 2004.
- Johnsson, M., Green, J. A. M., and Stigebrandt, A.: Baroclinic wave drag from two closely spaced sills in a narrow fjord as inferred from basin water mixing, *J. Geophys. Res.: Oceans*, 112, <https://doi.org/https://doi.org/10.1029/2006JC003694>, 2007.
- 290 Lelong, M.-P. and Kunze, E.: Can barotropic tide — eddy interactions excite internal waves?, *J. Fluid Mech.*, 721, 1—27, <https://doi.org/https://doi.org/10.1017/jfm.2013.1>, 2013.
- Massel, S. R.: *Internal gravity waves in the shallow seas*, Springer, 2015.
- Morozov, E. G.: *Oceanic Internal Tides: Observations, Analysis and Modeling: A Global View*, Springer International Publishing, Cham, <https://doi.org/https://doi.org/10.1007/978-3-319-73159-9>, 2018.
- 295 Münnich, M.: The influence of bottom topography on internal seiches in stratified media, *Dyn. Atmos. Oceans*, 23, 257–266, [https://doi.org/https://doi.org/10.1016/0377-0265\(95\)00439-4](https://doi.org/https://doi.org/10.1016/0377-0265(95)00439-4), 1996.
- Niiler, P. P.: On the internal tidal motions in the Florida Straits, *Deep Sea Res. Oceanogr. Abst.*, 15, 113–123, [https://doi.org/https://doi.org/10.1016/0011-7471\(68\)90031-4](https://doi.org/https://doi.org/10.1016/0011-7471(68)90031-4), 1968.

- Park, J., MacMahan, J., Sweet, W. V., and Kotun, K.: Continuous seiche in bays and harbors, *Ocean Sci.*, 12, 355–368, <https://doi.org/https://doi.org/10.5194/os-12-355-2016>, <https://os.copernicus.org/articles/12/355/2016/>, 2016.
- 300 Parsmar, R. and Stigebrandt, A.: Observed damping of barotropic seiches through baroclinic wave drag in the Gullmar Fjord, *J. Phys. Oceanogr.*, 27, 849–857, [https://doi.org/https://doi.org/10.1175/1520-0485\(1997\)027<0849:ODOBST>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0485(1997)027<0849:ODOBST>2.0.CO;2), 1997.
- Rattray Jr., M.: On the coastal generation of internal tides, *Tellus*, 12, 54–62, <https://doi.org/https://doi.org/10.1111/j.2153-3490.1960.tb01283.x>, 1960.
- 305 Rippeth, T. and Green, J. A. M.: Tides, the Moon and the kaleidoscope of ocean mixing, in: *Oceanography and Marine Biology - An Annual Review*, Vol. 58, edited by Hawkins, S., Allcock, A., Bates, A., Evans, A., Firth, L., McQuaid, C., Russell, B., Smith, I., Swearer, S., and Todd, P., pp. 319–350, CRC Press, <https://books.google.hu/books?id=Mp4FEAAAQBAJ>, 2020.
- Roget, E., Khimchenko, E., Forcat, F., and Zavialov, P.: The internal seiche field in the changing South Aral Sea (2006–2013), *Hydrol. Earth Syst. Sci.*, 21, 1093–1105, <https://doi.org/https://hess.copernicus.org/articles/21/1093/2017/>, 2017.
- 310 Staalstrøm, A. and Røed, L. P.: Vertical mixing and internal wave energy fluxes in a sill fjord, *J. Marine Syst.*, 159, 15–32, <https://doi.org/https://doi.org/10.1016/j.jmarsys.2016.02.005>, 2016.
- Stanev, E. V. and Ricker, M.: Interactions between barotropic tides and mesoscale processes in deep ocean and shelf regions, *Ocean Dyn.*, 70, 713–728, <https://doi.org/https://doi.org/10.1007/s10236-020-01348-6>, 2020.
- Stevens, K. N.: *Acoustic phonetics*, 2001.
- 315 Stigebrandt, A.: Barotropic and baroclinic response of a semi-enclosed basin to barotropic forcing from the sea., in: *Fjord Oceanography. NATO Conference Series (IV Marine Sciences)*, vol 4., edited by Freeland, H., Farmer, D., and Levings, C., pp. 141–164, Springer, Boston, MA., [https://doi.org/https://doi.org/10.1007/978-1-4613-3105-6\\_5](https://doi.org/https://doi.org/10.1007/978-1-4613-3105-6_5), 1980.
- Stigebrandt, A.: Resistance to barotropic tidal flow in straits by baroclinic wave drag, *J. Phys. Oceanogr.*, 29, 191–197, [https://doi.org/https://doi.org/10.1175/1520-0485\(1999\)029<0191:RTBTFI>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0485(1999)029<0191:RTBTFI>2.0.CO;2), 1999.
- 320 Stigebrandt, A. and Aure, J.: Vertical mixing in basin waters of fjords, *J. Phys. Oceanogr.*, 19, 917–926, [https://doi.org/https://doi.org/10.1175/1520-0485\(1989\)019<0917:VMIBWO>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0485(1989)019<0917:VMIBWO>2.0.CO;2), 1989.
- Vic, C., Naveira Garabato, A. C., Green, J. A. M., Waterhouse, A. F., Zhao, Z., Melet, A., de Lavergne, C., Buijsman, M. C., and Stephenson, G. R.: Deep-ocean mixing driven by small-scale internal tides, *Nat. Commun.*, 10, 2099, <https://doi.org/https://doi.org/10.1038/s41467-019-10149-5>, 2019.
- 325 Vincze, M. and Bozóki, T.: Experiments on barotropic–baroclinic conversion and the applicability of linear n-layer internal wave theories, *Experiments in Fluids*, 58, 1–12, 2017.
- Wunsch, C. and Ferrari, R.: Vertical mixing, energy, and the general circulation of the oceans, *Annu. Rev. Fluid Mech.*, 36, 281–314, <https://doi.org/https://doi.org/10.1146/annurev.fluid.36.050802.122121>, 2004.
- Wynne, Z., Reynolds, T., Bouffard, D., Schladow, G., and Wain, D.: A novel technique for experimental modal analysis of barotropic seiches for assessing lake energetics, *Environ. Fluid Mech.*, 19, 1527–1556, <https://doi.org/https://doi.org/10.1007/s10652-019-09677-x>, 2019.
- 330 Xue, M.-A., Kargbo, O., and Zheng, J.: Seiche oscillations of layered fluids in a closed rectangular tank with wave damping mechanism, *Ocean Eng.*, 196, 106842, <https://doi.org/https://doi.org/10.1016/j.oceaneng.2019.106842>, 2020.