



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

Preface: Developments in the Science and History of Tides

Philip L. Woodworth¹, J.A. Mattias Green², Richard D. Ray³ and John M. Huthnance¹

1. National Oceanography Centre, Joseph Proudman Building, 6 Brownlow Street, Liverpool, L3 5DA, United Kingdom

2. School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, LL59 5AB, United Kingdom

3. Geodesy and Geophysics Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, USA

Correspondence to: Philip L. Woodworth (plw@noc.ac.uk)

Abstract

This special issue marks the 100th anniversary of the founding of the Liverpool Tidal Institute (LTI), one of a number of important scientific developments in 1919. The preface gives a brief history of how the LTI came about and the roles of its first two Directors, Joseph Proudman and Arthur Doodson. It also gives a short overview of the research on tides at the LTI through the years. Summaries are given of the 26 papers in the special issue. It will be seen that the topics of many of them could be thought of as providing a continuation of the research first undertaken at the LTI. Altogether, they provide an interesting snapshot of work on tides now being made by groups around the world.

Keywords: ocean and earth tides; history of tidal science

1. Introduction

The idea for this special issue on tides came about when we realised that the 100th anniversary of the founding of the Liverpool Tidal Institute (LTI) was coming up in 2019, and we thought that a special issue of a journal would be one good way of celebrating it. The year following the end of the First World War saw the establishment of a number of organisations which have had a lasting importance for geophysical research. Notably, the International Union of Geodesy and Geophysics (IUGG) was founded in that year (Ismail-Zadeh and Joselyn, 2019). In addition, as examples at a national level, the American Geophysical Union (AGU) and American Meteorological Society (AMS) were established in the same year (McEntee, 2018; McFarquhar, 2020).

The suggestion for a special institute in the UK dedicated to research on tides can be said to have arisen from a report written in 1916 for the British Association for the Advancement of Science (BAAS) by Sir Horace Lamb, Professor of Applied Mathematics at Manchester, and Joseph Proudman, a lecturer in Applied Mathematics at Liverpool (Figure 1). The proposal was accepted by Liverpool University and funding was obtained from Sir Alfred and Charles Booth of the Booth Shipping Line in order to ‘prosecute continuously scientific research into all aspects of knowledge of the tides’ (Doodson, 1924; Cartwright, 1980, 1999). These are the basic facts, although Carlsson-Hyslop (2020) has described how the founding of the LTI came about only after a considerable amount of prior discussion and argument between the various stakeholders in academia (including Liverpool University and the BAAS), industry (including the Mersey Docks and Harbour Board) and the Navy (Admiralty).



51 The first LTI office was set up in March 1919 in the George Holt Building (then the Physics building) on
52 the Liverpool University campus.¹ Proudman became its Honorary (unpaid) Director while Arthur
53 Doodson was recruited as its Secretary. Proudman continued with his other university responsibilities
54 and was promoted to Professor of Applied Mathematics. Meanwhile, in that same year, Liverpool
55 University established the first university Oceanography department in the UK.

56
57 However, space was short at the university after the war, and during the 1920s the LTI relocated across
58 the River Mersey to Bidston Observatory in the Wirral where there was more room for research and
59 where Doodson took up residence in 1929, now as Associate Director (Nature, 1928). Proudman
60 transferred from Applied Mathematics to be Professor of Oceanography in 1933, a position which he
61 held until retirement in 1954, having handed over as Director of the LTI to Doodson in 1945. By then,
62 the LTI had become an acknowledged centre of expertise for research into ocean and earth tides, and
63 it was to further extend its reputation in the following years into research on storm surges, sea level
64 changes (Permanent Service for Mean Sea Level), and the measurement and modelling of coastal and
65 shelf processes. The LTI underwent many name changes through the years, most notably being
66 renamed as the Proudman Oceanographic Laboratory twice in 1987 and 2000, and became a
67 component of the present National Oceanography Centre in 2010.

68
69 Proudman and Doodson both became Fellows of the Royal Society. Proudman was particularly expert
70 on the dynamical theories of tides, while Doodson was superb at simplifying large numerical
71 calculations in the days before digital computers. One topic in which Doodson excelled was tidal
72 prediction, from both a theoretical foundation (his development of the tidal potential, Doodson, 1921)
73 and in practical application (the construction and use of tidal prediction machines (TPMs), see below).
74 These topics and other historical aspects of work at the LTI will be mentioned briefly. Excellent
75 biographies of Proudman and Doodson can be found in Cartwright and Ursell (1976) and Proudman
76 (1968) respectively.
77

¹ This is another link between Liverpool University and the shipping companies, George Holt being the founder of the Holt Line and the brother of Alfred Holt, co-founder of the Blue Funnel Line. Alfred Holt's daughter was Jane Herdman after whom the Geology building on the campus is named. She was married to Sir William Herdman, Professor of Natural History, who became the first Professor of Oceanography.



78

79

80 Figure 1. Joseph Proudman CBE, FRS (1888-1975) photographed in 1931 by Walter Stoneman. ©
81 National Portrait Gallery, London.

82

83 2. The Special Issue

84

85 A special issue such as this was ideally suited to the European Geosciences Union (EGU) family of
86 journals published by Copernicus. Tides occur in the ocean, solid earth and atmosphere, with each of
87 those areas of research being served by one or more journals. However, Copernicus provides a good
88 mechanism for papers to appear in their individual journals and be linked together eventually to
89 provide an overall special issue.

90

91 Initial enquiries with the various Executive Editors suggested that five journals might be involved, the
92 largest number which would have taken part in a special issue so far. However, in the event only three
93 journals were included: History of Geo- and Space Sciences, Solid Earth and, in particular, Ocean
94 Science. However, these three journals cover more than adequately most of the research into tides
95 which was undertaken at the LTI, and also a great deal of the tidal research which is being conducted
96 today. Potential authors were invited to address virtually any aspect of research into ocean or earth
97 tides including the history of that research. As examples, the former could possibly include the present
98 accuracy of coastal, regional and global tide models, tidal dissipation and its role in geophysics,
99 internal tides and their role in mixing the ocean and in the global ocean circulation, secular changes in
100 tides, and new techniques for measuring tides and analysing the data. The special issue opened for
101 submissions on 1 January 2018, and stayed open through 2019 to take advantage of new findings



102 presented at the AGU, EGU and IUGG conferences in that year. It closed finally to new submissions in
103 the middle of 2020 having received and accepted 26 papers across the three journals.

104
105 To complement the special issue, a number of special sessions on tides were held at international
106 conferences. These included presentations on tides at the EGU General Assembly in Vienna in April
107 2019 and the IUGG General Assembly in Montreal in July 2019. Events were also held locally to mark
108 the LTI anniversary, including a large meeting for the public at the Merseyside Maritime Museum in
109 May 2019 (Woodworth, 2020b) and in talks to Liverpool history groups.

110 111 2.1 History of Tides in the Special Issue 112

113 One of the first papers in the special issue was concerned with the tidal data obtained by James Cook
114 during his remarkable voyage to Tahiti to observe the Transit of Venus followed by the first landings
115 by Europeans in New Zealand and Australia (Woodworth and Rowe, 2018). This paper marked yet
116 another anniversary in 2019, in this case being 250 years since Cook's arrival at Tahiti. The paper
117 discusses how puzzled Cook was by diurnal inequality in the tide, a factor which led to the near-sinking
118 of the *Endeavour*, and a topic that was later to be an important aspect of tidal research in the 19th
119 century.

120
121 Agnew (2020) discusses how tidal fluctuations in gravity played an important role in the accuracy of
122 timing measurements using pendulum clocks, which were up until the 1940s the best timekeepers
123 available. The paper shows that the best pendulum clocks were able to detect tides long before the
124 advent of measurements by gravimeters or Global Navigation Satellite System (GNSS) technology. This
125 particular aspect of pendulum clocks does not appear to have been addressed at the LTI. However,
126 good timing was certainly one of the LTI's general interests, as it had been in Liverpool since the middle
127 of the 19th century, first by the astronomer John Hartnup of the Liverpool Observatory at Waterloo
128 Dock and then at Bidston Observatory.² Recent papers (not in the special issue) discuss how timing
129 formed an important aspect of local scientific research and services over many years. These include
130 the famous time transfer experiments between Liverpool and Boston; the provision of a chronometer
131 calibration service for seafarers; and through the time control of the Liverpool One O'Clock Gun and
132 time balls (Schmidt and Dearden, 2019, 2020; Thomas and Thomas, 2020a,b).

133
134 Other historical topics are addressed by Carlsson-Hyslop (2020) who, as mentioned above, provides
135 many details of the complicated discussions which resulted in the founding of the LTI. One topic
136 mentioned in that paper concerns the importance of the provision of tidal predictions to its overall
137 finances. The use of TPMs became essential to such work, as discussed by Woodworth (2020a), and
138 the LTI, under the direction of Doodson, became a world leader in the design and operation of TPMs.

139
140 Finally, a reminder that tidal measurements have a long history in many other countries is provided
141 by Raicich (2020) who discusses recording at Trieste since 1782. This paper is an example of what has
142 become known as 'data archaeology', whereby the information contained in sometimes vulnerable
143 paper records is being converted into computer form so that the important data can be used in studies
144 of long-term climate change.

145 146 2.2 Tidal Science at the LTI 147

148 The history of a branch of science can often be marked by the introduction of new technologies which
149 have revolutionised and reinvigorated the research. Tidal science, and sea-level science in general,
150 provides a good example. In the following, we mention some of the research areas at the LTI which

² A good summary of the history of research at first the Liverpool Observatory and then at Bidston Observatory, marking the centenary of the founding of the former, was provided by Doodson in LOTI (1945).



151 involved tides and sea levels and which benefitted from new technology. This provides one way of
152 relating the LTI's history to the research papers contained in the special issue. The areas include:

153

- 154 • Tidal Prediction

155

156 Woodworth (2020a) explains how it became possible to determine tidal harmonic constants from a
157 set of tide gauge data. In principle, these constants could be used to provide tidal predictions by means
158 of tedious hand computations. However, the technological leap provided by Kelvin's invention of the
159 TPM, a type of analogue computer, speeded up the determination of predictions considerably.
160 Between the 1870s and 1960s, over 30 TPMs were constructed around the world, of which the
161 majority were made in the UK. Only three of them were used for tidal prediction at the LTI itself.
162 However, there were more for which Doodson played a major role in their design or supervised closely
163 their manufacture. Of course, TPMs were superseded by another technological leap, the advent of
164 digital computers in the 1960s.

165

- 166 • Storm Surge Modelling

167

168 Proudman (1968) remarked that towards the end of Doodson's tenure as LTI Director he strenuously
169 opposed the use of modern digital computers, claiming that they would increase the cost of providing
170 tidal predictions to harbour authorities. That reservation is understandable given Doodson's lifetime
171 involvement with the TPMs and the fact that they were still a source of income. It was left to his
172 successor Jack Rossiter to introduce modern computers to the LTI.

173

174 The study of 'meteorological effects on the tides' (i.e. storm surges) had been included in the LTI's
175 terms of reference since its founding. However, the study was given impetus by the major floods of
176 1953 (Wolf and Flather, 2005). Attempts were made by Shizuro Ishiguro at the National Institute of
177 Oceanography (NIO) in the UK (in the south of England, not then associated with the LTI) to predict
178 surges using electronic analogue computers (Kennard, 2016; Wolf, 2017). However, the work became
179 much easier once advances in technology had led to the availability of powerful digital computers. In
180 turn, this enabled the development of numerical storm surge models. Work at the LTI was led by
181 Norman Heaps and Roger Flather (e.g. Heaps, 1983), with their models adopted for operational use
182 by the Meteorological Office and Environment Agency for flood warning around the coast and control
183 of the Thames Barrier.

184

- 185 • Sea Level Measurements

186

187 The measurement of the tide, and sea level variations in general, using tide gauges has always been
188 an LTI interest. In 1933, Doodson together with Chadburns of Liverpool installed instrumentation to
189 the tide gauge in Birkenhead docks, two miles away, so that data could be transmitted to Bidston
190 Observatory, an early example of near-real-time data reporting. He also designed a current meter for
191 use on annual Irish Sea cruises (Proudman, 1968) (Figure 2). Eventually, the LTI became a centre of
192 expertise for tide gauge technology and was responsible for on-going sea level measurements in both
193 the UK and abroad. It also became a data centre for tide gauge information, the UK data being quality
194 controlled by the British Oceanographic Data Centre, and mean sea level information from around the
195 world being archived by the international Permanent Service for Mean Sea Level (PSMSL,
196 <https://www.psmsl.org>). The latter had been initiated at Bidston in 1933 with Proudman as its first
197 Secretary. The PSMSL has become one of the main services of the International Association for the
198 Physical Sciences of the Oceans (Smythe-Wright et al., 2020), and its data set is used within a wide
199 range of geophysical research, including the regular scientific assessments of the Intergovernmental
200 Panel on Climate Change (IPCC).

201



202 Doodson also experimented in the 1930s with a number of Favé pressure gauges with which
203 measurements of the tide offshore were obtained. However, it was not until the 1960s that the serious
204 measurement of deep-sea tides using bottom pressure records (BPRs) began in the UK, led by the
205 group of David Cartwright at the NIO. That small team transferred to Bidston Observatory in 1974
206 when Cartwright became Assistant Director of the Institute of Oceanographic Sciences (Bidston), as
207 the LTI was then called. The team became the world leaders in the measurement of tides at depths
208 down to 5000m (Figure 3). The same instruments were also used for ocean transport measurements
209 during the World Ocean Circulation Experiment (WOCE) (Spencer and Vassie, 1997). Cartwright's own
210 interests in sea level measurements were to be extended later into the use of satellite altimeter data,
211 mostly in studies of tides, in collaboration with US and UK researchers. Cartwright became the third
212 Fellow of the Royal Society associated with the LTI. A detailed biography may be found in Webb (2017).

213

214 • Earth Tides

215

216 Bidston Observatory has two levels of basements cut into the sandstone of Bidston Hill. In 1909, a
217 horizontal pendulum seismometer was installed by John Milne, specifically to study the tidal loading
218 of the solid Earth. This showed clearly the tilt in the north-south direction due to loading by the tide
219 in the adjacent Irish Sea (Milne, 1910).

220

221 After measurements using different instruments over many years by different groups, it was
222 concluded that tilt measurements were overly sensitive to local geology (Baker, 1980), and gravity
223 measurements took their place, the Bidston research group being equipped with LaCoste and
224 Romberg Earth Tide meters. These were used at many locations in the UK and abroad, and the
225 measurements by the Bidston group were shown to be particularly accurate due to their unique
226 electrostatic feedback mechanism and the careful recalibration of the gravimeters on the Hannover
227 gravity baseline in Germany. From the mid-1980s onwards, superconducting gravimeters were
228 deployed at a number of sites around the world (but not in the UK), with calibrations mainly provided
229 by new Micro-g LaCoste FG5 absolute gravity meters (see below). The results from these
230 superconducting gravimeters are in close agreement with the earlier results of the Bidston group's
231 LaCoste and Romberg Earth Tide gravimeters (Baker and Bos, 2003).

232

233 The development of GNSS technology in the 1990s, of which the Global Positioning System (GPS) is
234 the most well-known, enabled the vertical variations of the loading tide to be measured in an global
235 reference frame for the first time (and also its smaller horizontal components). The LTI group was
236 among the first to demonstrate the capabilities of GNSS in this way and it is now the main technique
237 for tidal loading studies. The result of all this body of work has been to demonstrate which ocean tide
238 models are more accurate than others, through computation of their corresponding loading tide
239 distributions, followed by comparison of those distributions to the gravimeter or GNSS data. In
240 addition, one learns a considerable amount concerning the physical properties of the solid Earth (Bos
241 et al., 2015). For an authoritative history of this part of the LTI's research, the reader is referred to the
242 unpublished article by Baker (2016).

243

244 • Geodetic Measurements

245

246 Alongside the GNSS measurements for tidal loading studies, the LTI group embarked in the 1990s on
247 collaborative research, especially with Nottingham University, on using the technique to measure
248 long-term rates of vertical land movements in the UK. Such GNSS data sets are now used by many
249 groups around the world in order to remove land movements from time series of relative sea level
250 measurements provided by tide gauges. This is becoming a well-established technique, although
251 problems remain, such as the stability of the reference frame in which measurements are made
252 (Wöppelmann and Marcos, 2016). Eventually, the group also acquired two Micro-g LaCoste FG5



253 absolute gravity meters, which measure small changes in the local acceleration due to gravity that can
254 be interpreted as equivalent to changes in vertical land movement (e.g. Teferle et al., 2006). However,
255 although absolute gravity continues to be used by other groups, especially where land movements are
256 particularly large (e.g. due to Glacial Isostatic Adjustment in Canada), it is no longer used in this role
257 in the UK.

258

259 • Other Research

260

261 The above sub-headings inevitably omit those research topics at the LTI which were not particularly
262 dependent on technology but which nevertheless had an important connection to tidal science. The
263 most obvious ones are the many dynamical studies of Proudman, a good summary of which can be
264 found in Cartwright (1980). They also omit many other aspects of the LTI's work which are less
265 applicable to the special issue. These include studies of coastal and shelf processes, modelling of water
266 quality and ecosystems, and investigations into the use of wave and tidal energy. For a more complete
267 overview of work at the LTI, the reader might consult publications such as Scofield (2006) or its regular
268 reports through the years.

269



270
271
272
273
274
275
276

Figure 2. Joseph Proudman (right), Arthur Doodson (centre) and R.J. Daniel, a marine biologist from Liverpool University (left), during an Irish Sea cruise in the late 1930s aboard the *Zephyr*. Photograph courtesy of Valerie Gane. Other photographs of Doodson may be found in Carlsson-Hyslop (2020) and Woodworth (2020a).



277
278
279

280 Figure 3. David Cartwright, FRS (1926-2015) (left) and colleagues around a 'Mark IV' bottom pressure
281 recorder deployed to measure the tide in over 3000 m of water NE of the Azores in 1980. Others
282 appearing the photograph (left to right): Ken Parry, Ian Vassie, Bev Hughes and Bob Spencer.
283 Photograph courtesy of the National Oceanography Centre.

284
285
286

287 2.3 Tidal Science in the Special Issue

288

289 It is possible to make connections in many cases between papers in the special issue and the areas of
290 work at the LTI mentioned above.

291

292 For example, Woodworth and Hibbert (2018) discuss long period ocean tides (Mf, Mm and Mt) in the
293 Drake Passage, and how their variation over a nodal cycle compares to expectations from the
294 equilibrium tide. They were indeed found to vary as expected. However, such a study would have been
295 impossible without the many years of BPR measurements in that 'choke point' area undertaken for
296 the WOCE and in the following years.

297

298 The term 'radiational potential' was introduced by Walter Munk to account for motions of a tidal
299 nature, which are caused, directly or indirectly, by the Sun's radiation, instead of being of astronomical
300 tidal origin due to the Moon or Sun. The largest oceanic radiational tides are due to annual and



301 semiannual variations in air pressure, and the ocean's response via the inverse barometer effect, and
302 also due to diurnal meteorological variability. The magnitude of such radiational tidal contributions
303 was estimated by Cartwright and Tayler (1971), work which came just before Cartwright's move to
304 the LTI. Williams et al. (2018) take a fresh look at these quasi-tidal variations and consider how they
305 may be double-counted in storm surge forecasts and also how estimates of Highest Astronomical Tide
306 might be affected.

307
308 Third-degree tides represent a much under-studied aspect of tidal research, probably because they
309 are very small (usually millimetric) at most locations and so are of less interest to people primarily
310 interested in tidal predictions. Nevertheless, the different spatial pattern of their forcing in the tidal
311 potential to those of the more familiar second-degree semidiurnal and diurnal tides provides another
312 way of testing our understanding of the ocean's response to astronomical forcing. Woodworth (2019)
313 returned to this topic using the global set of tide gauge data in the GESLA-2 data set and a global
314 numerical model. M1, the largest third-degree tide, was confirmed to have a geographical variation
315 consistent with the suggestions of Platzman and Cartwright, that it is generated in the ocean as a
316 consequence of the spatial and temporal overlap of M1 in the tidal potential and one (or at least a
317 small number of) diurnal ocean normal mode(s). It is remarkable that several of the larger (but still
318 tiny) third-degree tides have also begun to be mapped reliably by making use of the many years of
319 precise satellite altimetry (Ray, 2020).

320
321 One result of Cartwright's sabbatical at the Scripps Institution of Oceanography with Walter Munk
322 during 1963-1965 was the development of the response method of tidal analysis in which tides are
323 treated as the overall oceanic response to the astronomical forcing by the Moon and Sun (Munk and
324 Cartwright, 1966). In this special issue, Byun and Hart (2020b) further develop their own response-
325 type method of analysis with application to a mixed tidal regime around Antarctica. In an earlier paper
326 in the issue (Byun and Hart, 2020a), the same authors explained their classification scheme for tides
327 around New Zealand in terms of the relative proportions of the S2, N2, and M2 constituents, N2 being
328 relatively more important than S2 along some parts of the coast. Their new index provides a useful
329 addition to the usual Form Factor classification which simply describes the relative importance of
330 diurnal and semidiurnal components. Tidal analysis is also the topic of the paper by Boesch and Müller-
331 Navarra (2019) who make use of a technique called the Harmonic Representation of Inequalities
332 (HROI) method, that combines aspects of harmonic and non-harmonic methods, to reassess long-
333 period constituents for tidal predictions along the German North Sea coast and its tidally influenced
334 rivers. The usefulness of tidal predictions arising from tidal analysis in providing information on actual
335 high tides for use in the international Witness King Tides project (<http://kingtides.net/>) is assessed in
336 a paper by Hunter (2020).

337 Global tide modelling is represented in the special issue by Lyard et al. (2020) who discuss the
338 performance of the most recent global tide model of the Toulouse group (FES2014). This model
339 provides a significant improvement on earlier versions and is one of the most obvious demonstrations
340 of the value of many years of satellite altimetry. Regional modelling is represented by Medvedev et
341 al. (2020), who provide a model of the tides of the Caspian Sea, making use of available tide gauge
342 data around its coast. The importance of bathymetry in the modelling of tides in extensive shallow
343 water regions is demonstrated by Rasquin et al. (2020), who consider the uncertainties in how tides
344 might change in the German Bight following a rise in sea level. The importance of bathymetry also
345 enters into the study of Fofonova et al. (2019) who consider the non-linear aspects of tidal dynamics
346 in an even smaller part of German-Danish coastal waters.

347
348 The need for good bathymetric information in tide modelling of coastal waters is also discussed by
349 Green and Pugh (2020). The authors make use of a large number of short-term tidal measurements
350 around Bardsey Island off the coast of North Wales to investigate how well the dynamics of tidal
351 streams around the island compare to present knowledge derived from satellite altimetry or regional



352 tide models, both having their acknowledged spatial resolution limitations. The set of tide gauge
353 measurements infers much larger tidal currents than anticipated with important consequences for the
354 computation of local tidal energetics.

355
356 There is an agreed need to understand better the reasons for variability in tides on longer timescales
357 for the whole global coastline (Haigh et al., 2020). A possible connection between long-term changes
358 in the tide in the North Atlantic and the North Atlantic Oscillation climate index is discussed in this
359 issue by Pineau-Guillou et al. (2020), while Harker et al. (2019) discuss possible changes to tides
360 around Australia due to a rise in mean sea level.

361
362 Tidal loading investigations such as those described above at the LTI are represented in the special
363 issue by two papers (Wang et al., 2020; Matviichuk et al., 2020). The former paper considers
364 asthenospheric anelasticity effects on ocean tide loading around the East China Sea using GNSS data
365 and employing a number of ocean tide models from which loading is computed. The latter paper
366 investigates the potential improvements in GNSS loading measurements around the UK and western
367 Europe using GLONASS data in combination with GPS. These two papers demonstrate the present
368 maturity of using GNSS in tidal loading studies.

369
370 Internal tides were primarily a regional numerical modelling activity at the LTI (e.g. Xing and Davies,
371 1999). They are now being modelled worldwide by several groups (e.g. Zaron, 2019b). This topic is
372 represented in the special issue by a regional study of the predictability of Caribbean internal tides by
373 Zaron (2019a), and an accuracy assessment of various global internal tide models by Carrere et al.
374 (2020). The benefits of technology for the local measurement of internal tides using gliders and
375 moored acoustic Doppler current profilers is described by Hall et al. (2019).

376
377 The only paper in the issue concerned with the role of the tide in marine biology is that of Petrusevich
378 et al. (2020) who consider the impact of tidal dynamics on diel vertical migration of zooplankton in
379 Hudson Bay.

380
381 In possibly the most charming paper in the special issue, Cooper et al. (2018) discuss the ability to
382 learn about tidal currents by observing the positions of sea birds (razorbills) resting on the water
383 surface and thereby functioning as ‘drifters of opportunity’. This paper provides a good demonstration
384 of how a new technology (in this case GNSS tracking) can complement existing techniques for
385 measuring tidal currents.

386 387 3. A Short Forward Look on Tides 388

389 The availability of precise satellite altimeter data in the early 1990s revolutionised the development
390 of regional and global tide models (Stammer et al., 2014). These models provide accurate maps of
391 amplitude and phase lag for the largest constituents of the barotropic tide throughout the global
392 ocean, and enable a reliable determination of tidal energy budgets (Ray and Egbert, 2017). However,
393 this encouraging situation is far from complete. For example, there is still a need to know more about
394 seasonal variability in the tides, including those at high latitudes under ice shelves. Measurements in
395 polar regions require the use of several different types of *in situ* tide gauge or GNSS instrumentation
396 as well as altimetry.

397
398 A particular high priority is to have a better understanding of tides in coastal waters, comparable in
399 accuracy and completeness to our knowledge of deep-ocean tides. For example, one of the tests of
400 the Stammer et al. (2014) review was to calculate differences in tidal heights from the eight major
401 constituents in the models and the heights derived using the same constituents obtained from
402 analyses of *in situ* (tide gauge) data. The best models showed root-sum-square agreement of



403 approximately 0.9, 5.0, and 6.5 cm for pelagic (deep ocean), shelf and coastal regions, respectively.
404 This demonstrated the centimetric accuracy for the best models over most of the deep ocean.
405 However, the modelled tides were found to be considerably less accurate close to the coast, where
406 problems remain due to the inherent limitations of spatial and temporal sampling by altimeters,
407 technical issues to do with land contamination in altimeter and radiometer footprints, and the fact
408 that coastal tides are more complex than those of the deep ocean with a multiplicity of shallow-water
409 constituents (Ray et al., 2011; Woodworth et al., 2019).

410
411 The need to do better near the coast has resulted in a whole new field of research being created called
412 ‘coastal altimetry’ (Vignudelli et al., 2019). It has spurred more sophisticated processing (retracking)
413 of radar returns from existing and past missions, in order to remove as much as possible of the land
414 contamination from the data. In addition, the altimeters of later missions such as Cryosat-2, Sentinel-
415 3 or Sentinel-6 are capable of providing measurements closer to the coast (for the global coastline for
416 the latter, only certain sections of coast for the former). It remains to be seen how well data from
417 these missions will benefit global tide models in general. Attention is now turning to the next
418 generation of altimeters for tidal research (e.g. SWOT, Morrow et al., 2019) which will provide
419 considerably greater information on tidal variations on short spatial scales including in coastal waters.

420
421 Many opportunities for future tidal studies exist for interested researchers, for example in a better
422 estimation of the global tide through geological history. Such studies are important for a more
423 complete understanding of topics such as species evolution. The same modelling techniques can be
424 applied to the study of putative tides on other planets and even exoplanets. In addition, there is much
425 to be learned how tides contribute to the variability in the climate of our own planet throughout
426 modulation in the strength of the over-turning circulation. There have been recent papers on all these
427 subjects (too many to list here) but it is clear that much interesting research remains.

428
429 4. Conclusions

430
431 We hope that this special issue has been useful in marking an anniversary of an important institution
432 in the history of tides. The work started at the LTI is far from over and continues as part of the present-
433 day National Oceanography Centre. In addition, we hope that the special issue will also have been of
434 interest in providing a snapshot of present-day tidal research.

435
436 Competing interests. The authors declare that they have no conflicts of interest.

437 Special issue statement. This article is part of the special issue “Developments in the science and
438 history of tides (OS/ACP/HGSS/NPG/SE inter-journal SI)”. It is not associated with a conference.

439 Acknowledgements

440
441 We thank all authors, reviewers and topic editors for their involvement in the special issue. All papers
442 were edited by one of the present four authors or, on occasions when one of us was an author, by Jo
443 Williams or Kristian Schlegel.

444
445



- 446 Papers in the Special Issue in Chronological Order
- 447 1. Woodworth, P. L. and Rowe, G. H.: The tidal measurements of James Cook during the voyage of the
448 Endeavour, *Hist. Geo Space. Sci.*, 9, 85-103, <https://doi.org/10.5194/hgss-9-85-2018>, 2018.
- 449 2. Woodworth, P. L. and Hibbert, A.: The nodal dependence of long-period ocean tides in the Drake
450 Passage, *Ocean Sci.*, 14, 711-730, <https://doi.org/10.5194/os-14-711-2018>, 2018.
- 451 3. Williams, J., Irazoqui Apecechea, M., Saulter, A., and Horsburgh, K. J.: Radiational tides: their double-
452 counting in storm surge forecasts and contribution to the Highest Astronomical Tide, *Ocean Sci.*, 14,
453 1057-1068, <https://doi.org/10.5194/os-14-1057-2018>, 2018.
- 454 4. Cooper, M., Bishop, C., Lewis, M., Bowers, D., Bolton, M., Owen, E., and Dodd, S.: What can seabirds
455 tell us about the tide?, *Ocean Sci.*, 14, 1483-1490, <https://doi.org/10.5194/os-14-1483-2018>, 2018.
- 456 5. Harker, A., Green, J. A. M., Schindelegger, M., and Wilmes, S.-B.: The impact of sea-level rise on tidal
457 characteristics around Australia, *Ocean Sci.*, 15, 147-159, <https://doi.org/10.5194/os-15-147-2019>,
458 2019.
- 459 6. Hall, R. A., Berx, B., and Damerell, G. M.: Internal tide energy flux over a ridge measured by a co-
460 located ocean glider and moored acoustic Doppler current profiler, *Ocean Sci.*, 15, 1439-1453,
461 <https://doi.org/10.5194/os-15-1439-2019>, 2019.
- 462 7. Woodworth, P. L.: The global distribution of the M1 ocean tide, *Ocean Sci.*, 15, 431-442,
463 <https://doi.org/10.5194/os-15-431-2019>, 2019.
- 464 8. Zaron, E. D.: Predictability of non-phase-locked baroclinic tides in the Caribbean Sea, *Ocean Sci.*, 15,
465 1287-1305, <https://doi.org/10.5194/os-15-1287-2019>, 2019a.
- 466 9. Boesch, A. and Müller-Navarra, S.: Reassessment of long-period constituents for tidal predictions
467 along the German North Sea coast and its tidally influenced rivers, *Ocean Sci.*, 15, 1363-1379,
468 <https://doi.org/10.5194/os-15-1363-2019>, 2019.
- 469 10. Fofonova, V., Androsov, A., Sander, L., Kuznetsov, I., Amorim, F., Hass, H. C., and Wiltshire, K. H.:
470 Non-linear aspects of the tidal dynamics in the Sylt-Rømø Bight, south-eastern North Sea, *Ocean Sci.*,
471 15, 1761-1782, <https://doi.org/10.5194/os-15-1761-2019>, 2019.
- 472 11. Rasquin, C., Seiffert, R., Wachler, B., and Winkel, N.: The significance of coastal bathymetry
473 representation for modelling the tidal response to mean sea level rise in the German Bight, *Ocean Sci.*,
474 16, 31-44, <https://doi.org/10.5194/os-16-31-2020>, 2020.
- 475 12. Medvedev, I. P., Kulikov, E. A., and Fine, I. V.: Numerical modelling of the Caspian Sea tides, *Ocean*
476 *Sci.*, 16, 209-219, <https://doi.org/10.5194/os-16-209-2020>, 2020.
- 477 13. Wang, J., Penna, N. T., Clarke, P. J., and Bos, M. S.: Asthenospheric anelasticity effects on ocean
478 tide loading around the East China Sea observed with GPS, *Solid Earth*, 11, 185-197,
479 <https://doi.org/10.5194/se-11-185-2020>, 2020.
- 480 14. Raicich, F.: A 1782-1794 sea level record at Trieste (northern Adriatic), *Hist. Geo Space. Sci.*, 11, 1-
481 14, <https://doi.org/10.5194/hgss-11-1-2020>, 2020.
- 482 15. Woodworth, P. L.: Tide prediction machines at the Liverpool Tidal Institute, *Hist. Geo Space. Sci.*,
483 11, 15-29, <https://doi.org/10.5194/hgss-11-15-2020>, 2020a.



- 484 16. Petrusevich, V. Y., Dmitrenko, I. A., Niemi, A., Kirillov, S. A., Kamula, C. M., Kuzyk, Z. Z. A., Barber,
485 D. G., and Ehn, J. K.: Impact of tidal dynamics on diel vertical migration of zooplankton in Hudson Bay,
486 Ocean Sci., 16, 337-353, <https://doi.org/10.5194/os-16-337-2020>, 2020.
- 487 17. Hunter, J.: Are tidal predictions a good guide to future extremes? - a critique of the Witness King
488 Tides project, Ocean Sci., 16, 703-714, <https://doi.org/10.5194/os-16-703-2020>, 2020.
- 489 18. Carlsson-Hyslop, A.: How the Liverpool Tidal Institute was established: industry, navy and
490 academia, Hist. Geo Space. Sci., 11, 139-156, <https://doi.org/10.5194/hgss-11-139-2020>, 2020.
- 491 19. Byun, D.-S. and Hart, D. E.: A monthly tidal envelope classification for semidiurnal regimes in terms
492 of the relative proportions of the S₂, N₂, and M₂ constituents, Ocean Sci., 16, 965-977,
493 <https://doi.org/10.5194/os-16-965-2020>, 2020a.
- 494 20. Agnew, D. C.: Time and tide: pendulum clocks and gravity tides, Hist. Geo Space. Sci., 11, 215-224,
495 <https://doi.org/10.5194/hgss-11-215-2020>, 2020.
- 496 21. Byun, D.-S. and Hart, D. E.: Predicting tidal heights for extreme environments: from 25h
497 observations to accurate predictions at Jang Bogo Antarctic Research Station, Ross Sea, Antarctica,
498 Ocean Sci., 16, 1111-1124, <https://doi.org/10.5194/os-16-1111-2020>, 2020b.
- 499 22. Matviichuk, B., King, M., and Watson, C.: Estimating ocean tide loading displacements with GPS
500 and GLONASS, Solid Earth, 11, 1849-1863, <https://doi.org/10.5194/se-11-1849-2020>, 2020.
- 501 23. Green, J. A. M. and Pugh, D. T.: Bardsey – an island in a strong tidal stream: underestimating
502 coastal tides due to unresolved topography, Ocean Sci., 16, 1337–1345, <https://doi.org/10.5194/os-16-1337-2020>, 2020.
- 504 24. Pineau-Guillou, L., Lazare, P., and Wöppelmann, G.: Large-scale changes of the semidiurnal tide
505 along North Atlantic coasts from 1846 to 2018, Ocean Sci., 17, 17–34, <https://doi.org/10.5194/os-17-17-2021>, 2021.
- 507 25. Carrere, L., Arbic, B. K., Dushaw, B., Egbert, G. D., Erofeeva, S. Y., Lyard, F., Ray, R. D., Ubelmann,
508 C., Zaron, E., Zhao, Z., Shriver, J. F., Buijsman, M. C., and Picot, N.: Accuracy assessment of global
509 internal tide models using satellite altimetry, Ocean Sci. Discuss., <https://doi.org/10.5194/os-2020-57>,
510 in review, 2020.
- 511 26. Lyard, F. H., Allain, D. J., Cancet, M., Carrère, L., and Picot, N.: FES2014 global ocean tides atlas:
512 design and performances, Ocean Sci. Discuss., <https://doi.org/10.5194/os-2020-96>, in review, 2020.
- 513
514
515
516
517



518 Other References

519

520 Baker, T.F.: Tidal tilt at Llanrwst, North Wales: tidal loading and Earth structure, *Geophys. J. Roy. Astr.*
521 *S.*, 62, 269–290, <https://doi.org/10.1111/j.1365-246X.1980.tb04855.x>, 1980.

522

523 Baker, T.F. Earth tides and ocean tide loading, <http://www.bidstonobservatory.org.uk/earth-tides/>,
524 2016.

525

526 Baker, T.F. and Bos, M.S.: Validating Earth and ocean tide models using tidal gravity measurements,
527 *Geophys. J. Int.*, 152, 468-485, <https://doi.org/10.1046/j.1365-246X.2003.01863.x>, 2003.

528

529 Bos, M.S., Penna, N.T., Baker, T.F., and Clarke, P.J.: Ocean tide loading displacements in western
530 Europe: 2. GPS-observed anelastic dispersion in the asthenosphere, *J. Geophys. Res. Solid Earth*, 120,
531 6540–6557, <https://doi.org/10.1002/2015JB011884>, 2015.

532

533 Cartwright, D.E. and Tayler, R.J.: New computations of the tide-generating potential, *Geophys. J. Roy.*
534 *Astr. S.*, 23, 45-74, <https://doi.org/10.1111/j.1365-246X.1971.tb01803.x>, 1971.

535

536 Cartwright, D.E. and Ursell, F.J.: Joseph Proudman, 30 December 1888 - 26 June 1975, *Biographical*
537 *Memoirs of Fellows of the Royal Society*, 22, 319-333, <https://doi.org/10.1098/rsbm.1976.0014>, 1976.

538

539 Cartwright, D. E.: The historical development of tidal science, and the Liverpool Tidal Institute, in:
540 *Oceanography of the Past*, edited by: Sears, M. and Merriman, D., New York, Springer-Verlag, 812 pp.,
541 240–251, 1980.

542

543 Cartwright, D. E.: *Tides: a scientific history*, Cambridge, Cambridge University Press, 292 pp., 1999.

544

545 Doodson, A. T.: The harmonic development of the tide-generating potential, *Proc. R. Soc. London*, 100,
546 305-329, <https://doi.org/10.1098/rspa.1921.0088>, 1921.

547

548 Doodson, A. T.: The tides and the work of the Tidal Institute, Liverpool, *Geogr. J.*, 63, 134–144,
549 available at: <http://www.jstor.org/stable/1781626> (last access: 16 March 2020), 1924.

550

551 Haigh, I.D., Pickering, M.D., Green, J.A.M., Arbic, B.K., Arns, A., Dangendorf, S., Hill, D., Horsburgh, K.,
552 Howard, T., Idier, D., Jay, D.A., Lee, S.B., Müller, M., Schindelegger, M., Talke, S.A., Wilmes, S-B., and
553 Woodworth, P.L.: The tides they are a-changin', *Rev. Geophys.*, 57, e2018RG000636,
554 <https://doi.org/10.1029/2018RG000636>, 2020.

555

556 Heaps, N.S.: Storm surges 1967-1982, *Geophys. J. Roy. Astr. S.*, 74, 331-376,
557 <https://doi.org/10.1111/j.1365-246X.1983.tb01883.x>, 1983.

558

559 Ismail-Zadeh, A. and Joselyn, J. A.: IUGG: beginning, establishment, and early development (1919–
560 1939), *Hist. Geo Space. Sci.*, 10, 25–44, <https://doi.org/10.5194/hgss-10-25-2019>, 2019.

561

562 Kennard, C.: Understanding storm surges in the North Sea: Ishiguro's electronic modelling machine,
563 *Science Museum Group Journal*, Issue 6 (Autumn 2016), <http://dx.doi.org/10.15180/160603>, 2016.

564

565 LOTI: Liverpool Observatory and Tidal Institute Centenary Report and Annual Reports (1940-1945).
566 University of Liverpool, Liverpool, 1945.

567



- 568 McEntee, C.: AGU launches its centennial celebration, *Eos*, 99,
569 <https://doi.org/10.1029/2018EO101321>, 2018. Published on 18 June 2018.
570
- 571 McFarquhar, G. (Ed.): A century of progress in atmospheric and related sciences: celebrating the
572 American Meteorological Society centennial, 2 vols., Amer. Met. Soc., Boston, 1428pp., 2020.
573
- 574 Milne, J.: Surface deformation and the tides, *Nature*, 82, 427-427, <https://doi.org/10.1038/082427a0>,
575 1910.
576
- 577 Morrow, R., Fu, L.-L., Ardhuin, F., Benkiran, M., Chapron, B., Cosme, E., et al.: Global observations of
578 fine-scale ocean surface topography with the Surface Water and Ocean Topography (SWOT)
579 mission, *Front. Mar. Sci.*, 6, 232, <https://doi.org/10.3389/fmars.2019.00232>, 2019.
580
- 581 Munk W.H. and Cartwright, D.E.: Tidal spectroscopy and prediction, *Philos. T. R. Soc. London*, A259,
582 533-581, <https://doi.org/10.1098/rsta.1966.0024>, 1966.
583
- 584 *Nature: Liverpool Observatory and Tidal Institute (Nature News Item)*, *Nature*, 122, 979–
585 980, <https://doi.org/10.1038/122979b0>, 1928.
586
- 587 Proudman, J.: Arthur Thomas Doodson, 1890–1968, *Biographical Memoirs of Fellows of the Royal*
588 *Society*, 14, 189–205, <https://doi.org/10.1098/rsbm.1968.0008>, 1968.
589
- 590 Ray, R.D.: First global observations of third-degree ocean tides. *Sci. Adv.*, 6, eabd4744,
591 <https://doi.org/10.1126/sciadv.abd4744>, 2020.
592
- 593 Ray, R.D. and Egbert, G.D.: Tides and satellite altimetry. pp. 427-458 (Chapter 13) in, *Satellite Altimetry*
594 *Over Oceans and Land Surfaces* (eds. D. Stammer and A. Cazenave). CRC Press: Boca Raton. 617pp.,
595 2017.
596
- 597 Ray, R.D., Egbert, G.D., and Erofeeva, S.Y.: Tide predictions in shelf and coastal waters: status and
598 prospects. pp.191-216 (Chapter 8) in, *Coastal Altimetry* (S. Vignudelli et al., eds.), Berlin: Springer-
599 Verlag, 2011.
600
- 601 Schmidt, R.E. and Dearden, P.: The Liverpool Observatory at Waterloo Dock. Part 1. Origins and
602 controversy, *The Antiquarian Astronomer*, Issue 13, 2-22, 2019.
603
- 604 Schmidt, R.E. and Dearden, P.: The Liverpool Observatory at Waterloo Dock. Part 2. Greenwich on the
605 Mersey, *The Antiquarian Astronomer*, Issue 14, 77-108, 2020.
606
- 607 Scofield, J.: *Bidston Observatory: the place and the people*, Merseyside, Countrywise Ltd., 344 pp.,
608 2006.
609
- 610 Smythe-Wright, D., Gould, W. J., McDougall, T. J., Sparnocchia, S., and Woodworth, P. L.: IAPSO: tales
611 from the ocean frontier, *Hist. Geo Space. Sci.*, 10, 137–150, [https://doi.org/10.5194/hgss-10-137-](https://doi.org/10.5194/hgss-10-137-2019)
612 2019, 2019.
613
- 614 Spencer, R. and Vassie, J.M.: The evolution of deep ocean pressure measurements in the U.K., *Prog.*
615 *Oceanogr.*, 40, 423-435, [https://doi.org/10.1016/S0079-6611\(98\)00011-1](https://doi.org/10.1016/S0079-6611(98)00011-1), 1997.
616
- 617 Stammer, D., Ray, R.D., Andersen, O.B., Arbic, B.K., Bosch, W., Carrère, L., Cheng, Y., Chinn, D.S.,
618 Dushaw, B.D., Egbert, G.D., Erofeeva, S.Y., Fok, H.S., Green, J.A.M., Griffiths, S., King, M.A., Lapin, V.,



- 619 Lemoine, F.G., Luthcke, S.B., Lyard, F., Morison, J., Müller, M., Padman, L., Richman, J.G., Shriver, J.F.,
620 Shum, C.K., Taguchi, E., and Yi, Y.: Accuracy assessment of global barotropic ocean tide models. *Rev.*
621 *Geophys.*, 52, 243-282, <https://doi.org/10.1002/2014RG000450>, 2014.
622
- 623 Teferle, F.N., Bingley, R.M., Williams, S.D.P., Baker, T.F., and Dodson, A.H.: Using continuous GPS and
624 absolute gravity to separate vertical land movements and changes in sea-level at tide-gauges in the
625 UK, *Philos. T. R. Soc. London*, A364, 917-930, <https://doi.org/10.1098/rsta.2006.1746>, 2006.
626
- 627 Thomas, S. and Thomas, D.: Getting the right time. Liverpool's contribution in the mid-nineteenth
628 century. Part 1, *Antiquarian Horology*, 41(1), 20–43, 2020a.
629
- 630 Thomas, S. and Thomas, D.: Getting the right time. Liverpool's contribution in the mid-nineteenth
631 century. Part 2, *Antiquarian Horology*, 41(2), 214–234, 2020b.
632
- 633 Vignudelli, S., Birol, F., Benveniste, J., Fu, L-L., Picot, N., Raynal, M., and Roinard, H.: Satellite altimetry
634 measurements of sea level in the coastal zone, *Surv. Geophys.*, 40, 1319-1349.
635 <https://doi.org/10.1007/s10712-019-09569-1>, 2019.
636
- 637 Webb, D.J.: David Edgar Cartwright, 21 October 1926 — 2 December 2015, *Biographical Memoirs of*
638 *Fellows of the Royal Society*, 63, 99-115, <https://doi.org/10.1098/rsbm.2017.0001>, 2017.
639
- 640 Wöppelmann, G and Marcos, M.: Vertical land motion as a key to understanding sea level change and
641 variability, *Rev. Geophys.*, 54, 64-92, <https://doi.org/10.1002/2015RG000502>, 2016.
642
- 643 Wolf, J. and Flather, R.A.: Modelling waves and surges during the 1953 storm, *Philos. T. R. Soc. London*,
644 A363, 1359-1375, <https://doi.org/10.1098/rsta.2005.1572>, 2005.
645
- 646 Wolf, J.: The connection between storm surges in the North Sea and the new British Nobel Laureate,
647 Kazuo Ishiguro, <http://www.bidstonobservatory.org.uk/ishiguro/>, 2017.
648
- 649 Woodworth, P.L.: Celebrating 100 years of tidal science on Merseyside, *Ocean Challenge*, 24(1), 10-
650 11, 2020b.
651
- 652 Woodworth, P.L., Melet, A., Marcos, M., Ray, R.D., Wöppelmann, G., Sasaki, Y.N., Cirano, M., Hibbert,
653 A., Huthnance, J.M., Montserrat, S., and Merrifield, M.A.: Forcing factors affecting sea level changes
654 at the coast, *Surv. Geophys.*, 40(6), 1351-1397, <https://doi.org/10.1007/s10712-019-09531-1>, 2019.
655
- 656 Xing, J.X. and Davies, A.M.: The influence of topographic features and density variations upon
657 the internal tides in shelf edge regions, *Int. J. Numer. Meth. Fl.*, 31, 535-577,
658 [https://doi.org/10.1002/\(SICI\)1097-0363\(19991015\)31:3<535::AID-FLD887>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1097-0363(19991015)31:3<535::AID-FLD887>3.0.CO;2-9), 1999.
659
- 660 Zaron, E.D.: Baroclinic tidal sea level from exact-repeat mission altimetry, *J. Phys. Oceanogr.*, 49, 193–
661 210, <https://doi.org/10.1175/JPO-D-18-0127.1>, 2019b.
662
663