



Oxygen and nutrient trends in the Tropical Oceans

Lothar Stramma and Sunke Schmidtke

GEOMAR Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20, 24105 Kiel, Germany

Correspondence to: Lothar Stramma (lstramma@geomar.de)

Abstract. A vertical expansion of the intermediate-depth low-oxygen zones (300 to 700 m) is seen in time series for selected tropical areas for the period 1960 to 2008, in the eastern tropical Atlantic, the equatorial Pacific and the eastern tropical Indian Ocean. These nearly five decade-long time series were extended to 68 years by including rare historic data starting in 1950 and more recent data. For the extended time series between 1950 and 2018 the deoxygenation trend for the layer 300 to 700 m is similar to the deoxygenation trend seen in the shorter time series. Additionally, temperature, salinity and nutrient time series in the upper ocean layer (50 to 300 m) of these areas were investigated since this layer provides critical pelagic habitat for biological communities. Generally, oxygen is decreasing in the 50 to 300 m layer except for an area in the eastern tropical South Atlantic. Nutrients also showed long-term trends in the 50 to 300 m layer in all ocean basins and indicates overlying variability related to climate modes. Nitrate increased in all areas. Phosphate also increased in the Atlantic and Indian Ocean areas, while it decreased in the two areas of the equatorial Pacific Ocean. Silicate decreased in the Atlantic and Pacific areas but increased in the eastern Indian Ocean. Hence oxygen and nutrients show trends in the tropical oceans, though nutrients trends are more variable between ocean areas than the oxygen trends, therefore we conclude that those trends are more dependent on local drivers in addition to a global trend. Different positive and negative trends in temperature, salinity, oxygen and nutrients indicate that oxygen and nutrient trends cannot be completely explained by local warming.

1 Introduction

Temperature, oxygen and nutrient changes in the ocean have various impacts on the ecosystem. These impacts span from habitat compression in the open ocean (Stramma et. al., 2012) and affect all marine organisms through multiple direct and indirect mechanisms (Gilly et al., 2013) to affect the ecophysiology of marine water-breathing organisms with regard to distribution, phenology and productivity (Cheung et al., 2013). Despite its far-reaching consequences for humanity, the focus on climate change impacts on the ocean lags behind the concern for impacts on the atmosphere and land (Allison and Bassett, 2015). An oceanic increase in stratification, thus reduction in ventilation as well as decrease of oceanic dissolved oxygen are two of the less obvious but important expected indirect consequences of climate change on the ocean (Shepherd et al., 2017). Warming leads to lighter water in the surface layer and increased stratification reducing the mixing and deep ventilation of oxygen-rich surface water to the subsurface layers. Increasing ocean stratification over the last half century of



about 5% is observed in the upper 200 m (Li et al. 2020). The subsequent previously observed deoxygenation (e.g. Stramma et al, 2008, Schmidtko et al 2017) of the open ocean is one of the major manifestations of global change. This temperature oxygen relation can also be seen for the 0-1000 m layer of the global ocean, as the oxygen inventory is negatively correlated with the ocean heat content ($r=-0.86$; 0-1000 m) (Ito et al., 2017). Oxygen-poor waters often referred to as oxygen minimum zones (OMZ) occupy large volumes of the intermediate-depth eastern tropical oceans. In an investigation of six selected areas for the 300 to 700 m layer in the tropical oceans for the time period 1960 to 2008 Stramma et al. (2008) observed declining oxygen concentrations of -0.09 to $-0.34 \mu\text{mol kg}^{-1} \text{ year}^{-1}$ and a vertical expansion of the intermediate depth low oxygen zone. Such a vertical expansion of the OMZ that is entered and passed by diel vertical migrators and sinking particles could have widespread effects on species distribution, the biological pump and benthic-pelagic coupling (Wishner et al., 2013). The areas of the world ocean investigated for oxygen changes can be extended and in a quantitative assessment of the entire world ocean oxygen inventory by analysing dissolved oxygen and supporting data for the complete oceanic water column over the past 50 years since 1960. Schmidtko et al. (2017) reported that the global oceanic oxygen content of 227.4 ± 1.1 petamoles (10^{15}mol) has decreased by more than two percent (4.8 ± 2.1 petamoles). However, these oxygen changes vary by region with some areas showing increasing oxygen values on time scales related to climate modes.

The nutrient distribution is in addition to oxygen a key parameter controlling the marine ecosystems. However, very little is known about long term nutrient changes in the ocean. The transformation of carbon and nutrients into organic carbon, its sinking into the in the deep ocean, and its decomposition at depth, is known as the biological carbon pump. As a consequence, nutrients are consumed and thus lower in the surface ocean and released and thus higher in the deep ocean. The oceanic distribution of nutrients and patterns of biological production are controlled by the interplay of biogeochemical and physical processes, and external sources (Williams and Follows, 2003). In the upper 500 to 1000 m of the tropical oceans the nutrient concentration is higher than in the subtropics and is decreasing westwards (Levitus et al., 1993). In the subarctic North Pacific surface nutrient concentration decreased during 1975 to 2005, and is strongly correlated with a multidecadal increasing trend of sea surface temperature (SST) (Ono et al., 2008). Below the surface, however, oxygen decreased and nutrients increased in the subarctic Pacific pycnocline from the mid-1980s to around 2010 (Whitney et al., 2013). Nutrients would be expected to vary inversely with oxygen, if the dominant process was the remineralization of marine detritus (Whitney et al., 2013). In a recent study the trends of nutrients in the open Pacific Ocean were investigated (Stramma et al., 2020) and in the open Pacific Ocean nutrient trends were observed and seemed to be related to oxygen trends. The supply of nutrients to the sunlit surface layer of the ocean has traditionally been attributed solely to vertical processes. However, horizontal advection may also be important in establishing the availability of nutrients in some regions. Palter et al. (2005) showed that the production and advection of North Atlantic Subtropical Mode Water introduces spatial and temporal variability in the subsurface nutrient reservoir beneath the North Atlantic subtropical gyre. By means of a coupled ecosystem circulation model Oschlies (2001) described for the North Atlantic that the long term change in the North Atlantic Oscillation (NAO; e.g. Hurrell and Deser, 2010) between the 1960s and 1990s may have induced significant regional changes in the upper ocean's nutrient supply. These include a decrease of nitrate supply to the surface waters of by about 30% near



Bermuda and in mid latitudes, and a simultaneous 60% increased nitrate flux in the upwelling region off West Africa. On the other side of the globe the Indonesian throughflow (ITF) is a chokepoint in the upper ocean thermohaline circulation, carrying Pacific waters through the strongly mixed Indonesian Seas and into the Indian Ocean (Ayers et al., 2014). Ayers et al. (2014) determined the depth- and time-resolved nitrate, phosphate, and silicate fluxes at the three main exit passages of the ITF: Lombok Strait, Ombai Strait, and Timor Passage. Nutrient flux as well as its variability with depth and time differed greatly between the passages. They estimated the effective flux of nutrients into the Indian Ocean and found that the majority of ITF nutrient supply to the Indian Ocean is to thermocline waters, where it is likely to support new production and significantly impact Indian Ocean biogeochemical cycling.

Here we investigate the extend of changes in oxygen, temperature and salinity trends for the six tropical areas with longer time series compared to the previously about one third shorter timeseries. Additionally, trends in the biological active near surface layer 50 to 300 m are investigated. As the upper ocean provides critical pelagic habitat for biological communities, nutrient time series of the six tropical areas since 1950 are investigated at 50 to 300 m depth, as nutrient changes in combination with hydrographic changes will influence the biological productivity of the ocean (Sigman and Hain, 2012). The upper boundary of 50 m was chosen to reduce the influence of the seasonal cycle in the upper 50 m although the seasonal cycle in the tropics is weaker than in most subtropical and subpolar regions (Louanchi and Najjar, 2000). As there are indications that climate modes and the El Niño-Southern Oscillation (ENSO) events have an influence on the trends, we check whether these signals are apparent in the data in the near surface layer.

2 Data and methods

Stramma et al. (2008) investigated the temperature and oxygen trends for the period 1960 to 2008 in the 300 to 700 m layer of six tropical ocean areas. There were three areas in the tropical Atlantic (A: 10°–14°N, 20°–30°W; B: 3°S–3°N, 18°–28°W; C: 14°S–8°S, 4°–12°E), two areas in the eastern and central tropical Pacific (D: 5°S–5°N, 105°–115°W; E: 5°S–5°N, 165°–175°W) and one in the eastern Indian Ocean (F: 5°S–0°N, 90°–98°E) (Figure 1). Here these time series were extended with more recent data as well as back in time to 1950 for the regions with available data (Table 1 and Figure 2).




Figure 1: Climatological mean dissolved oxygen concentration ($\mu\text{mol kg}^{-1}$ shown in color) at 400 m depth contoured at 20 $\mu\text{mol kg}^{-1}$ intervals from 10 to 230 $\mu\text{mol kg}^{-1}$ (black lines). Analysed areas A to F (Table 1) are enclosed by black boxes (Stramma et al., 2008).

Despite long-term trends in ocean oxygen also climate signal related influence on the trends was observed in recent years. More recently also long-term trends and climate signal related influence was observed for nutrients. The areas D and E were also used for the layer 50 to 300 m for oxygen changes in Stramma et al. (2020), but not for nutrient trends due to the low amount of available nutrients data. However, here we list also the nutrients trends for these two areas, despite the low amount of data does not make these calculations statistically significant (Table 2).

The main hydrographic data set is similar to the one used and described in Schmidtke et al. (2017), relying on Hydrobase and World Ocean Database bottle data for nutrient data. Quality control and handling is described in Schmidtke et al. (2017) for oxygen and is used here similarly for nutrients. The only divergence to the described procedure was that bottle data with missing temperature and/or salinity were assigned the temporal and spatial interpolated temperature and salinity derived from MIMOC (Schmidtke et al., 2013). This was done to ensure all data were in $\mu\text{mol kg}^{-1}$ and not requiring the discarding



of already sparse data due to missing water density values. This enables us to use data in mol l⁻¹ or ml l⁻¹ which otherwise could not be used.

In the Atlantic the hydrographic and nutrient data were extended with some *RV Meteor*, *RV Merian* and *RV Poseidon* cruises. For the area A data from Meteor cruises M68/2 (2006), M83/1 (2008), M97 (2010), M119 (2015) and M145 (2018) and Merian MSM10/1 (2008) were added. For area B Meteor cruises M106 (2014), M130 (2016) and M145 (2018) were added. For area C cruise data from Poseidon P250 (1999), Merian MSM07 (2008), Meteor M120 (2015), Meteor M131 (2016) and Meteor M148 (2018) were included.

The Pacific the region at 5°N–5°S, 165–175°W (area E) which had data until 2009 was supplemented with data from a *RV Investigator* cruise at 170°W from June 2016. The region 5°N–5°S, 105–110°W (area D), which had data up to 2008, was supplemented with data from a *RV Ron Brown* cruise at 110°W in December 2016.

Climate indices considered include the NAO, the AMO, the PDO, ENSO, as well as the Indian Ocean Dipole Mode (IOD). The NAO is an extratropical climate signal of the North Atlantic. As our areas are tropical regions the three Atlantic areas were investigated relative to the Atlantic Multidecadal Oscillation (AMO) index (Montes et al., 2016) before and after 1995. The AMO was high before 1963, low until 1995 and high since 1995. In the Pacific the central equatorial area at 5°N–5°S, 165°–175°W (area E in Stramma et al., 2008) which had hydrodata until 2009, was supplemented with data from a *RV Investigator* cruise at 170°W from June 2016. The eastern equatorial area 5°N–5°S, 105°–115°W (area D in Stramma et al. 2008), which had hydrodata until 2008, was supplemented with data from a *RV Ron Brown* cruise at 110°W in December 2016. The data were investigated in relation to the Pacific Decadal Oscillation (PDO; e.g. Deser et al., 2010) before and after 1977. The PDO was negative from 1944 to 1976, positive from 1977 to 1998, variable from 1998 to 2013 and positive after 2013. In the Indian Ocean the available data covered the area F only after 1960 but until 2016. The area F (0° to 5°S, 90° to 98°E) is shown in relation to the IOD (Saji et al., 1999), which slightly increased after 1990.

Linear trends and their 95% confidence interval were computed using annual averages (all measurements from one year were attributed to that year) of the profiles linearly to standard vertical depth levels. A computation routine was used which used the effective number of degrees of freedom for the computation of the confidence interval. The data used for the oxygen time series were interpolated with an objective mapping scheme (Bretherton et al., 1976) with Gaussian weighting. In the 50 to 300 m layer and the 300 to 700 m a temporal half folding range of 0.5 year and a vertical half folding range of 50 m with maximum ranges of 1 year and 100 m were applied. The covariance matrix was computed from 100 local data points and 50 random data points within the maximum range, for the diagonal of the covariance matrix a signal to noise ratio of 0.7 was set (see Schmidt et al. 2013, for details). A more improved mapping scheme was used compared to the one used in Stramma et al. (2008) where larger temporal ranges were used (1-year half folding and a maximum temporal range of 2 years).

Nutrients nitrite (NO₂⁻), nitrate (NO₃⁻), phosphate (PO₄³⁻) and silicic acid (Si(OH)₄ referred to as silicate hereafter) on the recent cruises were measured on-board with a QuAatro auto-analyzer (Seal Analytical). For recent autoanalyzer measurements precisions are 0.01 μmol kg⁻¹ for phosphate, 0.1 μmol kg⁻¹ for nitrate, and 0.5 μmol kg⁻¹ for silicate and 0.02 mL L⁻¹ (~ 0.9 μmol kg⁻¹) for oxygen from Winkler titration (Bograd et al., 2015). For older uncorrected nutrient data, offsets



are estimated to be 3.5% for nitrate, 6.2% for silicate and 5.1% for phosphate (Tanhua et al., 2010). One problem with nutrient data is that certified reference material (CRM) was applied to some measurements while for other measurements only a bias was applied.

The ENSO cycle of alternating warm El Niño and cold La Niña events is the climate system's dominant year-to year signal. ENSO originates in the tropical Pacific through interaction between the ocean and the atmosphere, but its environmental and socioeconomic impacts are felt worldwide (McPhaden et al., 2006). Three month running mean SST anomalies (ERSST.v5 SST anomalies) in the Niño 3.4 region (equatorial Pacific: 5°N to 5°S, 120°W to 170°W) of at least +0.5°C and lasting for at least 5 consecutive three months periods are defined as El Niño events and 5 consecutive three months periods of at least -0.5°C are defined as La Niña events (http://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v5.php). In case of measurements in ENSO years in figures 3, 4 and 5 the very strong El Niño events of 1983, 1998 and 2015 and the strong El Niño events 1957, 1965, 1972, 1987 and 1991 are marked by red circles and the strong La Niña events 1974, 1976, 1989, 1999, 2000, 2007 and 2010 are marked by blue squares in these years. A shoaling thermocline, such as occurs in the eastern Pacific during La Niña or cool (negative) PDO state, enhances nutrient supply and organic matter export in the eastern Pacific while simultaneously increasing the fraction of that organic matter that is respired in the low-oxygen water of the uplifted thermocline. The opposite occurs during El Niño or a warm (positive) PDO state; a deeper thermocline reduces both export and respiration in low-oxygen water in the eastern Pacific, allowing the hypoxic water volume to shrink (Deutsch et al., 2011; Fig. S7). ENSO also has some influence on the tropical Atlantic and Indian Oceans. The equatorial Atlantic oscillation is influenced by the Pacific ENSO with the equatorial Atlantic sea surface temperature lagging by about six months (Latif and Grötzner, 2000). In the Indian Ocean a recent weakening of the coupling between the ENSO and the IOD mode after the 2000s and 2010s compared to the previous two decades (1980s and 1990s) (Ham et al., 2017).

3 Trends in temperature, salinity, oxygen and nutrients

3.1 Trends in the 300 to 700 m depth layer

Nutrient data are sparse in the deeper part of the ocean and are less important than the near surface layer for the marine ecosystems and therefore are not presented here for the 300 to 700 m depth layer. Oxygen trends for the period 1960 to 2008 for the 300 to 700 m layer of the six areas investigated (Stramma et al., 2008) for the tropical oceans were all negative in the range -0.09 to -0.34 $\mu\text{mol kg}^{-1} \text{ year}^{-1}$ (Table 1). For the extended time period between 1950 and 2018 the oxygen trends were in the same order of magnitude for the areas A to F in the range -0.11 to -0.27 $\mu\text{mol kg}^{-1} \text{ year}^{-1}$ (Table 1). The 1950 to 2018 temperature trends were positive in the three Atlantic areas and the eastern tropical Pacific, but negative in the central Pacific and Indian Ocean areas (Table 1). In the eastern tropical Pacific (area D) and the eastern Indian Ocean (area F) there was even a reversed trend in temperature compared to the shorter time period between 1960 and 2008, although all temperature



trends are not within the 95% confidence interval difference from 0. The salinity of the 300 to 700 m layer increased for the Atlantic and Indian Ocean areas and decreased in the two Pacific areas (Table 1).

Table 1. Linear trends (300 to 700 m) of temperature in °C yr⁻¹, oxygen in μmol kg⁻¹ yr⁻¹ and salinity yr⁻¹ with 95% confidence intervals (p-values) where data are available for the entire period listed. Trends whose 95% confidence interval includes zero are shown in *italics*. Trends computed in Stramma et al. (2008) are shown for comparison.

Parameter	trend	time period	depth layer	(Stramma et al., 2008)
Area A	10°N-14°N, 20°W-30°W			
Temperature	+0.009 ± 0.005	1952-2018	300-700 m	+0.009 ± 0.008 1960-2006 300-700 m
Oxygen	-0.27 ± 0.12	1952-2018	300-700 m	-0.34 ± 0.13 1960-2006 300-700 m
Salinity	+0.0012 ± 0.0009	1952-2018	300-700 m	
Area B	3°S-3°N, 18°W-28°W			
Temperature	+0.005 ± 0.004	1952-2018	300-700 m	+0.005 ± 0.008 1960-2006 300-700 m
Oxygen	-0.25 ± 0.65	1952-2018	300-700 m	-0.19 ± 0.12 1960-2006 300-700 m
Salinity	+0.0001 ± 0.0005	1952-2018	300-700 m	
Area C	14°S-8°S, 4°E-12°E			
Temperature	+0.006 ± 0.004	1950-2018	300-700 m	+0.002 ± 0.011 1961-2008 300-700 m
Oxygen	-0.11 ± 0.100	1950-2018	300-700 m	-0.17 ± 0.11 1961-2008 300-700 m
Salinity	+0.0005 ± 0.0009	1950-2018	300-700 m	
Area D	5°S-5°N, 105°W-115°W			
Temperature	+0.003 ± 0.004	1955-2016	300-700 m	-0.001 ± 0.009 1962-2006 300-700 m
Oxygen	-0.24 ± 0.15	1957-2016	300-700 m	-0.13 ± 0.32 1962-2006 300-700 m
Salinity	-0.0001 ± 0.009	1950-2016	300-700 m	
Area E	5°S-5°N, 165°W-175°W			
Temperature	-0.001 ± 0.011	1950-2016	300-700 m	-0.010 ± 0.008 1961-2006 300-700 m
Oxygen	-0.18 ± 0.25	1950-2016	300-700 m	-0.19 ± 0.20 1961-2006 300-700 m
Salinity	-0.0003 ± 0.0009	1950-2016	300-700 m	
Area F	5°S-0°N, 90°E-98°E			



203	Temperature	-0.004 ± 0.010 1960-2016 300-700 m	$+0.005 \pm 0.007$ 1960-2007 300-700 m
204	Oxygen	-0.13 ± 0.17 1960-2016 300-700 m	-0.09 ± 0.21 1960-2007 300-700 m
205	Salinity	$+0.0001 \pm 0.0010$ 1960-2016 300-700 m	

206

207

208

209

210 For the area A (10° - 14° N, 20° - 30° W) the oxygen trend for 300 to 700 m for the period 1952 to 2018 (Figure 2a) was weaker
 211 ($-0.27 \pm 0.12 \mu\text{mol kg}^{-1} \text{yr}^{-1}$) than for the period 1960 to 2006 ($-0.34 \pm 0.13 \mu\text{mol kg}^{-1} \text{yr}^{-1}$). In the western subtropical and
 212 tropical Atlantic oxygen measurements from time series stations as well as shipboard measurements showed a significant
 213 relationship with the wintertime AMO index (Montes et al., 2016). During negative wintertime AMO years trade winds are
 214 typically stronger and these conditions stimulate the formation and ventilation of Subtropical Underwater (Montes et al.,
 215 2016) with higher oxygen content. Even in the 300 to 700 m layer of Area A (Figure 2a) as well as the 50 to 300 m layer
 216 (Figure 3a) the oxygen content is higher during the negative AMO period and lower during the positive AMO phase. For a
 217 section along 23° W between 6° - 14° N from 2006 to 2015 crossing area A an oxygen decrease in the 200 to 400 m layer and
 218 an increase in the 400 to 1000 m layer was described (Hahn et al., 2017) which can't be confirmed in area A due to the
 219 different geographical and temporal boundaries and the variable annual mean oxygen values after 2006 in area A.

220 The 1952 to 2018 oxygen trend in the equatorial Atlantic (area B) shows a large 95% confidence interval, different to the
 221 shorter time period 1960 to 2006 (Table 1). The larger confidence interval is caused by a low oxygen concentration in 1952
 222 and large variability after 2006 (Figure 2b). The equatorial Atlantic in the depth range 500 to 2000 m is influenced by
 223 Equatorial Deep Jets with periodically reversing flow direction influencing the transport of oxygen (Bastin et al., 2020)
 224 which might be one reason of the large oxygen variability. During the negative AMO the oxygen trend was slightly positive
 225 ($-0.034 \pm 1.39 \mu\text{mol kg}^{-1} \text{yr}^{-1}$) but negative after 1995 (Figure 2b).

226

227

228




Figure 2: Annual mean oxygen concentration for years available and trends for the layer 300 to 700 m in $\mu\text{mol kg}^{-1}$ plotted for the available years in the time period 1950 to 2018 (dashed red line) and for the positive and negative periods of the AMO in the Atlantic (a-c), the PDO in the Pacific (d,e) and the IOD in the Indian Ocean (f) as solid red lines. The AMO, PDO and IOD are shown as grey lines. The change of AMO status in 1963 and 1995, the change of the PDO phase in 1977, 1999 and 2013 and the IOD in 1990 are marked by dotted vertical lines.

The area C in the eastern tropical South Atlantic shows similar positive trends in temperature and salinity (Table 1) as in the two other Atlantic areas investigated. Area C is located in the region with the lowest oxygen content in the Atlantic Ocean



(Figure 1). Due to the already low oxygen concentration in this region the decrease in oxygen is weaker than in the two other Atlantic Ocean areas in the period 1950 to 2018, similar to the weaker decrease in area C for the shorter time period 1961 to 2008 (Table 1). Higher oxygen concentrations were also seen in the few oxygen profiles in area C during the negative AMO and lower oxygen concentrations were measured after the year 2000 (Figure 2c).

In the equatorial Pacific the two areas show a clear long-term oxygen decrease in the 300 to 700 m layer, but no clear changes related to the PDO phases before and after 1977 (Figure 2d,e). However, the PDO-index after 1977 was mainly positive until 1999 and mainly negative between 1999 and 2013. In case these time periods are looked at separately the oxygen concentration was higher during the period 1977 to 1990 and lower during 1999 to 2010 as expected for the PDO influence (e.g. Deutsch et al. 2011).

In the eastern Indian Ocean, the 300 to 700 m oxygen concentration was lower for the slightly positive IOD phase after 1990 leading to a long-term oxygen concentration decrease in area F although the trends for the shorter periods prior to 1990 and after 1990 showed a positive oxygen trend (Figure 2f), which are caused by high oxygen concentrations near the end of both measurement periods. The temperature in this area decreased and salinity showed barely any change (Table 1), hence the oxygen decrease is not coupled to temperature or hydrographic water mass changes.

3.2 Trends in the 50 to 300 m layer

The trend computations for the layer 50 to 300 m for temperature, salinity, oxygen and nutrients (Table 2) show different trends for the selected areas in the three tropical oceans. In the near surface layer 50 to 300 m the long-term oxygen trends were negative as in the deeper layer 300 to 700 m, except for area C in the eastern tropical South Atlantic (Figure 3c). However, this oxygen trend in area C is not stable due to the large variability in the time period 1960 to 1990. The upper layer of the area C is influenced by the Angola Dome centered at 10°S, 9°E (Mazeika, 1967) which might influence the larger variability near the surface. The area C shows the largest mean nitrate, silicate and phosphate concentrations in the Atlantic in the 50 to 300 m layer as well as the 300 to 700 m layer (Table 3) and shows the large nutrient availability in the eastern tropical South Atlantic. At 250 m and 500 m depth the region of area C was shown with the highest nitrate and phosphate concentrations of the tropical and subtropical Atlantic Ocean (Levitus et al. 1993). It was observed that in the Pacific Ocean nutrient are related to oxygen changes and climate variability (Stramma et al., 2020). The ENSO signal was apparent in most cases as in the tropical Atlantic and Indian Ocean (Nicholson, 1997) hence the oxygen distribution for the layer 50 to 300 m (Figure 3) is marked for El Niño and La Niña events to check for the possible influence of ENSO in the shallow depth layer. Most of the nutrient trends are due to sparse data coverage not statistically significant, nevertheless it is insightful to compare the nutrient trends with the oxygen trends as well as the climate signals.



Table 2. Linear trends (50-300 m) of temperature in °C yr⁻¹, salinity yr⁻¹ and solutes in μmol kg⁻¹ yr⁻¹ with 95% confidence intervals (p-values) where data are available for the entire period 1950 to 2018 (left rows) and for the earlier time period (center rows) and later time period (right rows) separated in 1995 in the Atlantic Ocean (areas A, B, C), in 1977 in the Pacific Ocean (areas D, E) and 1990 in the Indian Ocean (area F).. Trends whose 95% confidence interval includes zero are shown in *italics*.

Parameter	trend	time period	trend	time period	trend	time period
Area A 10°N-14°N, 20°W-30°W, 50-300 m						
Temperature	+0.007 ± 0.008	1952-2018	+0.004 ± 0.021	1952-1993	-0.001 ± 0.050	2001-2018
Salinity	+0.0009 ± 0.0012	1952-2018	+0.027 ± 0.0033	1952-1993	+0.006 ± 0.0083	2001-2018
Oxygen	-0.329 ± 0.231	1952-2018	-0.387 ± 0.639	1952-1993	+0.131 ± 1.120	2001-2018
Nitrate	+0.038 ± 0.077	1952-2018	+0.112 ± 0.116	1952-1993	-0.022 ± 0.581	2001-2018
Silicate	-0.066 ± 0.086	1952-2018	+0.002 ± 0.310	1952-1989	+0.029 ± 0.151	2001-2018
Phosphate	+0.001 ± 0.004	1952-2018	-0.002 ± 0.010	1952-1993	-0.024 ± 0.029	2001-2018
Area B 3°S-3°N, 18°W-28°W, 50-300 m						
Temperature	-0.007 ± 0.012	1952-2018	-0.013 ± 0.028	1952-1995	-0.017 ± 0.042	1997-2018
Salinity	+0.0003 ± 0.0011	1952-2018	+0.0001 ± 0.0030	1952-1994	+0.0010 ± 0.0040	1997-2018
Oxygen	-0.172 ± 0.421	1952-2018	-0.174 ± 0.874	1952-1994	-1.050 ± 2.010	1999-2018
Nitrate	+0.022 ± 0.075	1961-2018	+0.095 ± 0.111	1961-1994	+0.055 ± 0.369	1997-2018
Silicate	-0.061 ± 0.041	1961-2018	-0.079 ± 0.107	1961-1994	-0.056 ± 0.144	1999-2018
Phosphate	+0.001 ± 0.004	1952-2018	+0.007 ± 0.005	1952-1994	+0.003 ± 0.021	1997-2018
Area C 14°S-8°S, 4°E-12°E, 50-300 m						
Temperature	+0.006 ± 0.024	1950-2018	+0.018 ± 0.020	1950-1994	+0.04 ± 0.108	1995-2018
Salinity	+0.0008 ± 0.0020	1950-2018	-0.0019 ± 0.0025	1950-1994	+0.0039 ± 0.0070	1995-2018
Oxygen	+0.028 ± 0.474	1950-2018	-0.183 ± 1.190	1950-1994	-0.675 ± 0.819	1995-2018
Nitrate	+0.051 ± 0.088	1966-2018	+0.257 ± 0.220	1966-1988	-0.011 ± 0.530	1995-2018
Silicate	-0.052 ± 0.077	1968-2018	+0.020 ± 0.139	1968-1994	-0.161 ± 0.444	1995-2018
Phosphate	+0.002 ± 0.005	1957-2018	+0.011 ± 0.008	1957-1988	-0.001 ± 0.009	1995-2018
Area D 5°S-5°N, 105°W-115°W, 50-300 m						
Temperature	+0.003 ± 0.019	1955-2016	+0.076 ± 0.209	1955-1975	-0.004 ± 0.094	1979-2016
Salinity	-0.0000 ± 0.0018	1955-2016	-0.0017 ± 0.0068	1955-1975	+0.0001 ± 0.0022	1979-2016



306	Oxygen	-0.643 ± 0.367 1957-2016	-2.390 ± 3.100 1957-1971	-0.825 ± 0.825 1979-2016
307	Nitrate	$+0.033 \pm 0.166$ 1964-2016	$+0.329 \pm 14.90$ 1964-1968	$+0.223 \pm 0.272$ 1983-2016
308	Silicate	-0.001 ± 0.147 1967-2016	$+1.410 \pm 0.921$ 1967-1970	$+0.053 \pm 0.546$ 1983-2016
309	Phosphate	-0.002 ± 0.013 1957-1994	$+0.005 \pm 0.046$ 1957-1971	$+0.035 \pm 0.021$ 1983-1994
310				
311	Area E	5°S-5°N, 165°W-175°W, 50-300 m		
312	Temperature	-0.006 ± 0.020 1950-2016	$+0.026 \pm 0.060$ 1950-1976	-0.010 ± 0.051 1977-2016
313	Salinity	$+0.0005 \pm 0.0026$ 1950-2016	$+0.0005 \pm 0.0100$ 1950-1979	$+0.0000 \pm 0.0058$ 1977-2016
314	Oxygen	-0.361 ± 0.224 1950-2016	-0.192 ± 0.781 1950-1975	-0.570 ± 0.574 1977-2016
315	Nitrate	$+0.054 \pm 0.062$ 1961-2016	$+0.159 \pm 0.366$ 1961-1975	$+0.105 \pm 0.154$ 1977-2016
316	Silicate	-0.046 ± 0.148 1956-2016	$+0.172 \pm \text{NaN}$ 1956-1975	$+0.085 \pm 0.174$ 1977-2016
317	Phosphate	-0.003 ± 0.003 1950-2009	-0.002 ± 0.007 1950-1979	$+0.005 \pm 0.022$ 1990-2009
318				
319	Area F	5°S-0°N, 90°E-98°E, 50-300 m		
320	Temperature	-0.002 ± 0.028 1960-2016	$+0.004 \pm 0.056$ 1960-1990	$+0.033 \pm 0.163$ 1995-2016
321	Salinity	$+0.0020 \pm 0.0025$ 1960-2016	$+0.0049 \pm 0.0038$ 1960-1996	$+0.0043 \pm 0.0071$ 1995-2016
322	Oxygen	-0.221 ± 0.263 1960-2016	-0.098 ± 0.765 1960-1990	$+0.123 \pm 1.220$ 1995-2016
323	Nitrate	$+0.036 \pm 0.174$ 1962-2007	-0.130 ± 0.581 1962-1984	$-0.207 \pm \text{NaN}$ 1995-2007
324	Silicate	$+0.033 \pm 0.410$ 1960-2007	$+0.173 \pm 0.619$ 1960-1990	$-0.368 \pm \text{NaN}$ 1995-2007
325	Phosphate	$+0.003 \pm 0.009$ 1960-2007	$+0.003 \pm 0.014$ 1960-1989	$-0.015 \pm \text{NaN}$ 1995-2007

326

327

328

329

330

331




Figure 3: Annual mean oxygen concentration for years available and trends for the layer 50 to 300 m in $\mu\text{mol kg}^{-1}$ plotted for the available years in the time period 1950 to 2018 (dashed red line) and for the positive and negative periods of the AMO in the Atlantic (a-c), the PDO in the Pacific (d,e) and the IOD in the Indian Ocean (f) as solid red lines. The AMO, PDO and IOD are shown as grey lines. The change of AMO status in 1963 and 1995, the change of the PDO phase in 1977, 1999 and 2013 and the IOD in 1990 are marked by dotted vertical lines. El Niño years defined as strong are marked by an additional magenta circle, strong La Niña years by an additional blue square.




Figure 4: Annual mean nitrate concentration for years available and trends for the layer 50 to 300 m in $\mu\text{mol kg}^{-1}$ plotted for the available years in the time period 1950 to 2018 (dashed red line) and for the positive and negative periods of the AMO in the Atlantic (a-c), the PDO in the Pacific (d,e) and the IOD in the Indian Ocean (f) as solid red lines. For area A the nitrate measurements in 1974 were removed as the 50-300 m mean was much too low $2.93 \mu\text{mol kg}^{-1}$ and for area D the nitrate measurements were removed in 1970 which were too high ($30.28 \mu\text{mol kg}^{-1}$). The AMO, PDO and IOD are shown as grey lines. The change of AMO status in 1963 and 1995, the change of the PDO phase in 1977, 1999 and 2013 and the IOD in 1990 are marked by dotted vertical lines. El Niño years defined as strong are marked by an additional magenta circle, strong La Niña years by an additional blue square.



While oxygen decreased in all areas except for area C in the eastern tropical South Atlantic for the entire time period in the 50 to 300 m layer, nitrate increased in all areas (Figure 4). Phosphate also increased in the Atlantic and Indian Ocean areas, while it decreased in the 2 areas of the equatorial Pacific Ocean (Table 2). Silicate decreased in the Atlantic and Pacific areas but increased in the eastern Indian Ocean (area F). The temperature decreased in the central equatorial Pacific and the eastern Indian Ocean (areas E and F) as is the case for these areas also in the 300 to 700 m layer. Surprisingly at the equatorial area in the Atlantic (area B) the temperature in the 50 to 300 m layer decreased while it increased in the 300 to 700 m layer. The 50 to 300 m layer at the equator is governed by the eastward flowing Equatorial Undercurrent (EUC) while in the 300 to 700 m layer the westward flowing Intermediate Undercurrent (IUC) is located which might have an influence on the temperature change over time. The salinity in the 50 to 300 m layer increased in all areas except for a stagnant salinity concentration in the eastern tropical Pacific Ocean (area D; Table 2).

The largest amount of years with available nutrient data exists in area A in the Atlantic Ocean. The long-term trends in area A for temperature and oxygen for the 50 to 300 m layer (Table 2, Figure 5a,c) are similar as for the deeper layer 300 to 700 m (Table 1), however with increased variability near the surface most likely influenced by the seasonal cycle. For the 3 Atlantic areas A, B and C the long-term 50 to 300 m trend decreased for oxygen and silicate, and increased for salinity, nitrate, phosphate and temperature, the latter except for temperature in area B with a weak not significant temperature decrease. In the Atlantic, the equatorial station B shows higher mean 50 to 300 m layer temperature, salinity and oxygen and lower mean nitrate, silicate and phosphate values compared to the off-equatorial stations A and C (Table 3) and shows the eastward transport of oxygen-rich water with the EUC to the low oxygen regions in the eastern tropical Atlantic.




Figure 5: Annual mean oxygen concentration for years available and trends for the layer 50 to 300 m in $\mu\text{mol kg}^{-1}$ plotted for the available years in the time period 1950 to 2018 (dashed red line) and for the positive and negative periods of the AMO in the Atlantic at area A for temperature (a) in $^{\circ}\text{C}$, salinity (b), oxygen (c) in $\mu\text{mol kg}^{-1}$, nitrate (d) in $\mu\text{mol kg}^{-1}$, silicate (e) in $\mu\text{mol kg}^{-1}$ and phosphate (f) in $\mu\text{mol kg}^{-1}$. The AMO is shown as a grey line. The change of AMO status in 1963 and 1995 is marked by dotted vertical lines. El Niño years defined as strong are marked by an additional magenta circle, strong La Niña years by an additional blue square.



Table 3. Mean values for the layers 50-300 m and 300 to 700 m of temperature in °C, salinity and solutes in $\mu\text{mol kg}^{-1}$ in the Atlantic Ocean (areas A, B, C), in the Pacific Ocean (areas D, E) and in the Indian Ocean (area F).

Parameter	area A	area B	area C	area D	area E	area F
50-300 m						
Temperature	13.239	14.957	13.378	13.906	19.292	16.999
Salinity	35.350	35.432	35.333	34.888	35.116	35.010
Oxygen	91.045	109.888	55.178	70.059	141.116	96.157
Nitrate	22.518	17.754	27.438	23.629	13.771	18.755
Silicate	9.718	7.523	10.742	20.586	10.330	19.923
Phosphate	1.414	1.148	1.729	1.825	1.153	1.404
300 – 700 m						
Temperature	9.156	7.417	7.799	8.404	8.354	9.805
Salinity	35.020	34.673	34.726	34.656	34.646	34.966
Oxygen	62.194	104.494	47.251	32.991	72.027	67.922
Nitrate	34.457	31.280	39.284	35.662	32.967	31.437
Silicate	17.505	19.312	22.054	43.906	38.101	38.401
Phosphate	2.097	2.038	2.473	2.659	2.458	2.146

In the 50 to 300 m layer of area A despite the expected generally lower oxygen during positive AMO phase oxygen increased in the positive AMO phase after 1995 (Figure 3a) different to the decrease in the 300 to 700 m layer (Figure 2a). During the positive AMO phase after 1995 in the 50 to 300 m layer of area A trends in temperature, oxygen, nitrate, silicate and phosphate (Figure 5) changed sign compared to the long-term trend while salinity showed for this period the same continuous trend as the positive long-term trend. In contrast none of these parameters changed during positive AMO compared to the long-term trend at the 50 to 300 m layer in the equatorial Atlantic in area B. In the tropical North Atlantic (area A) and the equatorial Atlantic (area B) the La Niña events showed lower than normal oxygen concentrations especially for the years 1973/74, 1975/76 and 2010/11 (Figure 3a,b). These years were not covered in the eastern tropical South Atlantic (area C). In the equatorial area B, the El Niño years 1965/66, 1972/73, 1987/88 and 1991/92 showed slightly higher than normal oxygen concentrations (Figure 3b). Although not true for all ENSO events, there seems to be some influence of the La Niña and El Niño events in the eastern tropical and equatorial Atlantic, which might be due to the various types with different hydrographic impact of ENSO events described in literature.



In eastern Pacific regions near the Galapagos Islands (2-5°S, 84-87°W) and near the American continent in the CalCofi region (34-35°N, 121-122°W) and the Peru region (7-12°S, 78-83°W) oxygen increased and nutrient decreased in the 50 to 300 m layer during the negative PDO phase before 1977 with opposing trends during the positive PDO phase after 1977 (Stramma et al. 2020). Different to the eastern Pacific the eastern and central and equatorial areas D and E (Table 2) don't show the reversed trends in oxygen and nutrients, however temperature and salinity indicate a reversal with the PDO phase as the PDO index encapsulates the major mode of sea surface temperature variability in the Pacific. On a global scale the long-term SST trend 1901-2012 was positive everywhere except for a region in the North Atlantic (IPCC 2013, Fig. 2.21). For 1981 to 2012, while the western Pacific showed a warming trend, a large region with decreasing SST's was seen in the eastern and equatorial Pacific Ocean (IPCC 2013, Fig. 2.22). This agrees to the temperature reversal seen in areas D and E. However, if the time period after 1977 is looked at separately for the positive PDO phase 1977 to 1999 and the negative PDO phase 1999 to 2013 as in the layer 300 to 700 m also the layer 50 to 300 m shows the expected high oxygen concentrations in the period 1977 to 1990 and lower oxygen concentrations during 1999 to 2010 (Figure 3d,e).

Although ENSO is a signal originating in the Pacific the equatorial Pacific areas D and E show no obvious oxygen concentration changes related to ENSO events (Figure 3d,e). The central equatorial Pacific area E shows the highest mean 50 to 300 m temperature and oxygen concentrations and the lowest nitrate concentrations of all six areas investigated (Table 3). The low nitrate and phosphate and lower silicate compared to the eastern equatorial area D shows the nutrient concentration decreasing westward in the equatorial Pacific in the 50 to 300 m layer (Stramma et al., 2020; their Figure 2). The principal source of nutrients to surface water is vertical flux by diffusion and advection and by regeneration (Levitus et al., 1993). At the sea surface airborne nutrient supply from land is contributed as well as terrestrial runoff of fertilizer-derived nutrients and organic waste add nutrients to the ocean (Levin, 2018). The tongue of high nutrient concentrations at the equatorial Pacific compared to the subtropical Pacific results from upwelling near the American shelf (Levitus et al., 1993) and equatorial upwelling.

In the eastern Indian Ocean as in the 300 to 700 m layer the temperature in the 50 to 300 m layer (Table 2) decreases and indicates other processes related to the oxygen decrease instead of warming. In the Indian Ocean the IOD shows large variability on shorter time scales. Observations indicate that positive IOD events prevent anoxia off the west coast of India (Vallivattathillan et al. 2017). The IOD is very variable with a slightly higher index after 1990. The few oxygen measurements in the 50 to 300 m layer indicate in area F until 1990 high mean oxygen concentrations with a decrease in oxygen and after 1990 low oxygen concentrations with an increase in oxygen (Figure 3f). The higher oxygen concentrations before 1990 and lower oxygen concentrations afterwards are also visible in the 300 to 700 m layer (Figure 2f). The ENSO events don't indicate a visible influence on the oxygen concentration in area F. The four La Niña events between 1988 and 2008 were either below or above the mean trend-line, the same is true for the two El Niño events in 1973/74 and 1987/88 (Figure 3f).



4 Discussion and Summary

The time-series expansion of the six areas in the tropical oceans to the period 1950 to 2018 years showed a similar decrease in oxygen in the 300 to 700 m layer as described for the 1960 to 2008 period. Therefore, despite the overlying variability the long-term deoxygenation in the tropical oceans is continuous for the 68-year period. This confirms the indicated importance on the 48-year period (Stramma et al. 2008) of the oxygen trend for future oceanic scenarios. The salinity trends are weak and not statistically significant, except for a salinity increase of 0.0012 yr^{-1} in the 300 to 700 m layer of area A in the tropical Northeast Atlantic. A consistent pattern in vertical sections in the Pacific Ocean is that nitrate and phosphate increase with depth to about 500 m, with a slight maximum at intermediate depths of 500–1500 m, while silicate continues to increase with depth (Fiedler and Talley, 2006) which is well visible in the higher mean concentrations in the 300 to 700 m layer in comparison to the 50 to 300 m layer (Table 3).

The temperature trends were positive in the three Atlantic areas, but positive or negative in relation to the time period included in the Pacific and Indian Ocean areas, hence we can conclude that the decreasing oxygen is not fully coupled to the local temperature change. As the decline of oxygen in the tropical Pacific was not accompanied by a temperature increase, Ito et al. (2016) concluded that the cause of the oxygen decline must include changes in biological oxygen consumption and/or ocean circulation. Modelling the depth range 260 to 710 m depth range for 1990s-1970s the region of our areas D and E were mainly influenced by circulation variability (Ito et al., 2016).

Enhanced temperature differences between land and sea could intensify upwelling winds in eastern upwelling areas (Bakun, 1990). Observed and modelled changes in wind in the Atlantic and Pacific over the past 60 years appears to support the idea of increased upwelling winds (Sydeman et al., 2014). Coastal and equatorial upwelling enhance nutrients in the upper ocean, therefore the increase of nutrients in the eastern and equatorial oceans might be caused by winds intensifying upwelling. More nutrients in the surface layer enhances production and subsequently export and thus at greater depth its decay with increased respiration reduces the oxygen content. The sinking flux of organic matter, which over time depletes oxygen, while adding carbon and nutrients to subsurface waters, is known as the biological pump (Keeling et al., 2010) and could cause the often observed opposite trends in oxygen and nutrient trends in the 50 to 300 m layer investigated here. In the 50 to 300 m layer oxygen, temperature, salinity and nutrients showed long-term trends, which were different in the three ocean basins. Nitrate increased in all areas. Phosphate also increased in the Atlantic and Indian Ocean areas, while it decreased in the two areas of the equatorial Pacific Ocean. The phosphate increase in the Atlantic Ocean might be related to a continuous phosphate supply with the Saharan dust distributed over the Atlantic Ocean with the wind (Gross et al. 2015). Silicate decreased in the Atlantic and Pacific areas but increased in the eastern Indian Ocean. Often the expected inversely trend of oxygen and nutrients caused by remineralization of marine detritus (Whitney et al. 2013) was observed, however variations based on other drivers influence the nutrient trends.

An influence of ENSO years on the oxygen distribution with lower mean oxygen concentrations in the 50 to 300 m layer in La Niña years and larger oxygen concentrations in El Niño years was visible in the tropical North Atlantic and equatorial Atlantic. No clear impact of ENSO was observed in the tropical South Atlantic and the Pacific and Indian Ocean areas (C to



F). To construct time series in areas with low data availability measurements from larger areas had to be taken into account. As a result, there is a small possible bias due to the distribution of the measurements within the area and due to gaps in the time line. In addition, there might be variations due to the measurement techniques for oxygen and nutrients and the use of different reference material used for nutrient measurements or applied bias for nutrient measurements. Utilization of historical nutrient data to assess decadal trends has been hindered by their inaccuracy, manifested as offsets in deep water concentrations measured by different laboratories (Zhang et al., 2000). Although the trends are often not 95% significant the results indicate existing trends and climate related changes, which might be verified with additional data in the future. Although the data base is small especially for nutrients there is an indication that variability overlain on the long-term trends is connected to climate modes as was found in the eastern Pacific with reversing trends related to the PDO (Stramma et al., 2020). The six areas of the tropical ocean basins indicate some connection to the climate modes of the 3 ocean basins. In the tropical eastern North Atlantic (area A) there is some dependence with the AMO. In the equatorial Pacific areas D and E a connection to the PDO is visible when the positive PDO phase 1977 to 1999 and the negative PDO phase 1999 to 2013 are looked at separately. In the eastern tropical Indian Ocean there seems to be some dependence to the state of the IOD, despite the fact that the IOD varies more on shorter time scales and the IOD change in 1990 is weak. Future measurements of temperature, salinity, oxygen and nutrients could lead to more stable results determining trends and their variability to better understand the influence of climate change on the ocean ecosystem and prepare future predictions of ocean oxygen from Earth System Models (Frölicher et al., 2016). Making existing nutrient data public which are so far not in public data bases and modelling efforts on oxygen and nutrient changes would further improve the understanding of oxygen and nutrient variability and its biological influence e.g. on fishery. First ecosystem changes like habitat compression can be observed and negative impacts are expected on biological regulation, nutrient cycling and fertility, and sea food availability with an increasing risk of fundamental and irreversible ecological transformations (Hoegh-Guldberg and Bruno, 2010).

Data availability. The AMO time series was taken from <https://www.esrl.noaa.gov/psd/data/timeseries/AMO/> (ESRL, Climate time series, status 17.02.2020). The Indian Ocean Dipole Mode was taken from https://www.esrl.noaa.gov/psd/gcos_wgsp/Timeseries/Data/dmi.long.data on 3 March 2020. The yearly PDO data were taken from <http://ds.data.jma.go.jp/tcc/tcc/products/elnino/decadal/annpdo.txt> on 9 July 2020 from the Japan Meteorological Society covering the period 1901 to 2019.

The bottle data from cruises in 2016 at 170°W (096U2016426_hyd1.csv) and at 110°W (33RO20161119_hyd1.csv) were downloaded from the CCHDO at the University of California San Diego (<https://cchdo.ucsd.edu>, CCHDO, 2020) on 8 November 2018.



534 The added ship cruises are contained for CTD data in the data sets of <https://doi.org/10.1594/PANGAEA> for RV Meteor
 535 cruises M120, <https://doi.pangaea.de/10.1594/PANGAEA.868654> (Kopte and Dengler 2016), M130
 536 <https://doi.pangaea.de/10.1594/PANGAEA.903913> (Burmeister et al. 2019), M131
 537 <https://doi.pangaea.de/10.1594/PANGAEA.910994> (Brandt et al. 2020), M145
 538 <https://doi.pangaea.de/10.1594/PANGAEA.904382> (Brandt and Krahmann, 2019), and, for RV Meteor cruise M148
 539 <https://doi.pangaea.de/10.1594/PANGAEA.????>, and RV Merian 07 <https://doi.pangaea.de/10.1594/PANGAEA.????> and
 540 for nutrient data in the data sets of Merian MSM10/1 <https://doi.pangaea.de/10.1594/PANGAEA.775074> (Tanhua et al.
 541 2012), RV Poseidon 250 <https://doi.pangaea.de/10.1594/PANGAEA.????>, M68/2
 542 <https://www.ncei.noaa.gov/data/oceans/nci/ocads/data/0108078/>, M83/1
 543 <https://doi.pangaea.de/10.1594/PANGAEA.821729> (Tanhua 2013). M97
 544 <https://doi.pangaea.de/10.1594/PANGAEA.863119> (Tanhua 2016), Meteor M106
 545 <https://doi.pangaea.de/10.1594/PANGAEA.????>, Meteor M119 <https://doi.pangaea.de/10.1594/PANGAEA.????>, M120
 546 <https://doi.pangaea.de/10.1594/PANGAEA.????>, Meteor M130 <https://doi.pangaea.de/10.1594/PANGAEA.913986> (Tanhua
 547 2020), Meteor M131 <https://doi.pangaea.de/10.1594/PANGAEA.????>, Meteor M145
 548 <https://doi.pangaea.de/10.1594/PANGAEA.????>, and Meteor M148 <https://doi.pangaea.de/10.1594/PANGAEA.????>. Open
 549 references (shown as ????) will be made available before publication.

550

551 *Author contributions.* L. Stramma conceived the study and wrote the manuscript. S. Schmidtke compiled the data for the
 552 time series, collected further references and discussed and modified the manuscript.

553

554 *Competing interests.* The authors declare that they have no conflict of interest.

555

556 *Acknowledgements.* Financial support was received through GEOMAR and the Deutsche Forschungsgemeinschaft (DFG) as
 557 part of the “Sonderforschungsbereich 754: Climate-Biogeochemistry Interactions in the Tropical Ocean”.

558 References

- 559 Allison, E.H., and Bassett, H.R.: Climate change in the oceans: Human impacts and responses, *Science*,
 560 350, 778–782. <https://doi.org/10.1126/science.aac8721>, 2015.
- 561 Ayers, J. M., Strutton, P. G., Coles, V. J., Hood, R. R., and Matear, R. J.: Indonesian throughflow
 562 nutrient fluxes and their potential impact on Indian Ocean productivity, *Geophysical Research*
 563 *Letters*, 41, 5060–5067, doi:10.1002/2014GL060593, 2014.
- 564 Bakun A.: Global climate change and the intensification of coastal upwelling, *Science* 247,198–201,
 565 1990.



- 566 Bastin, S., Claus, M., Brandt, P., and Greatbatch, R. J.: Equatorial deep jets and their influence on the
 567 mean equatorial circulation in an idealized ocean model forced by intraseasonal momentum flux
 568 convergence, *Geophysical Research Letters*, 47, e2020GL087808.
 569 <https://doi.org/10.1029/2020GL087808>, 2020.
- 570 Bograd, S. J., Pozo Buil, M., Di Lorenzo, E., Castro, C. G., Schroeder, I. D., Goericke, R., Anderson, C.
 571 R., Benitez-Nelson, C., and Whitney, F. A.: Changes in source waters to the Southern California
 572 Bight, *Deep-Sea Res. II*, 112, 42-52, <https://doi.org/10.1016/j.dsr2.2014.04.009>, 2015.
- 573 Brandt, P., and Krahmann, G.: Physical Oceanography (CTD) during METEOR cruise M145,
 574 PANGAEA, <https://doi.org/10.1594/PANGAEA.904382>, 2019.
- 575 Brandt, P., Kopte, R., and Krahmann, G.: Physical oceanography (CTD) during METEOR cruise M131.
 576 PANGAEA, <https://doi.org/10.1594/PANGAEA.910994>, 2020.
- 577 Bretherton, F. P., Davis, R. E., and Fandry, C. B.: A technique for objective analysis and design of
 578 oceanographic experiments applied to MODE-73, *Deep-Sea Research*, 23, 559-582,
 579 [https://doi.org/10.1016/0011-7471\(76\)90001-2](https://doi.org/10.1016/0011-7471(76)90001-2), 1976.
- 580 Burmeister, K., Lübbecke, J., Brandt, P., Claus, M., and Hahn, J.: Ventilation of the eastern tropical
 581 North Atlantic by intraseasonal flow events of the North Equatorial Undercurrent. PANGAEA,
 582 <https://doi.org/10.1594/PANGAEA.903913>, 2019.
- 583 CCHDO: Welcome to the CCHDO, available at: <https://cchdo.ucsd.edu>, last access: 13 February 2020.
- 584 Cheung, W. W. L., Sarmiento, J. L., Dunne, J., Frölicher, T. L., Lam, V. W. Y., Palomares, M. L. D.,
 585 Watson, R., and Pauly, D.: Shrinking of fishes exacerbates impacts of global ocean changes of
 586 marine ecosystems, *Nature Climate Change*, 3, 254-258, <https://doi.org/10.1038/nclimate1691>,
 587 2013.
- 588 Deser, C., Alexander, M. A., Xie, S.-P., and Phillips, A. S.: Sea surface temperature variability: Patterns
 589 and mechanisms, *Annual Review of Marine Science*, 2, <https://doi.org/10.1146/annurev-marine-120408-151453>, 2010.
- 591 Deutsch, C., Brix, H., Ito, T., Frenzel, H., and Thompson, L.: Climate-forced variability of ocean
 592 hypoxia, *Science*, 333, 336-339, <https://doi.org/10.1126/science.1202422>, 2011.
- 593 Fiedler, P. C., and Talley, L. D.: Hydrography of the eastern tropical Pacific: A review, *Progress in*
 594 *Oceanography*, 69, 143-180, 2006.
- 595 Frölicher, T. L., Rogers, K. B., Stock, C. A., and Cheung, W. L. W.: Sources of uncertainties in 21th
 596 century projections of potential ecosystem stressors, *Global Biological Cycles*, 30, 1224-1243,
 597 <https://doi.org/10.1002/2015GB005338>, 2016.
- 598 Gilly, W. F., Beman, J. M., Litvin, S. Y., and Robinson, B. H.: Oceanographic and biological effects of
 599 shoaling of the oxygen minimum zone, *Annual Review of Marine Science*, 5,
 600 <https://doi.org/10.1146/annurev-marine-120710-100849>, 2013.



- Gross, A., Goren, T., Pio, C., Cardoso, J., Tirosh, O., Todd, M.C., Rosenfeld, D., Weiner, T., Custódio, D., and Angert, A.: Variability in Sources and Concentrations of Saharan Dust Phosphorus over the Atlantic Ocean, *Environmental Science & Technology Letters*, 2, 2, 31–37, 2015.
- Hahn, J., Brandt, P., Schmidtke, S., and Krahmann, G.: Decadal oxygen change in the eastern tropical North Atlantic, *Ocean Sci.*, 13, 551–576, <https://doi.org/10.5194/os-13-551-2017>, 2017.
- Ham, Y., Choi, J., and Kug, J.: The weakening of the ENSO–Indian Ocean Dipole (IOD) coupling strength in recent decades, *Climate Dynamics*, 49, 249–261, <https://doi.org/10.1007/s00382-016-3339-5>, 2017.
- Hoegh-Guldberg, O., and Bruno, J. F.: The impact of climate change on the world’s marine ecosystems, *Science*, 328, 1523–1528.
- Hurrell, J. W., and Deser, C.: North Atlantic climate variability: The role of the North Atlantic Oscillation, *Journal of Marine Systems*, 79, 231–244, <https://doi.org/10.1016/j.jmarsys.2009.11.002>, 2010.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M. (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp, 2013.
- Ito, T., Nenes, A., Johnson, M. S., Meskhidze, N., and Deutsch, C.: Acceleration of oxygen decline in the tropical Pacific over the past decades by aerosol pollutants, *Nature Geoscience*, 9, 443–448, <https://doi.org/10.1038/NGEO2717>, 2016.
- Ito, T., Minobe, S., Long, M. C., and Deutsch, C.: Upper ocean O₂ trends: 1958–2015, *Geophysical Research Letters*, 44, 4214–4223, doi:10.1002/2017GL073613, 2017.
- Keeling, R. F., Körtzinger, A., and Gruber, N.: Ocean deoxygenation in a warming world, *Annual Review of Marine Science*, 2, 199–229, 2010.
- Kopte, R., and Dengler, M.: Physical oceanography during METEOR cruise M120. PANGAEA, <https://doi.org/10.1594/PANGAEA.868654>, 2016.
- Latif, M., and Grötzner, A.: The equatorial Atlantic oscillation and its response to ENSO, *Climate Dynamics*, 16, 213–218, <https://doi.org/10.1007/s003820050014>, 2000.
- Levin, L. A.: Manifestation, drivers, and emergence of open ocean deoxygenation, *Annual Review of Marine Science*, 10, 229–260, <https://doi.org/10.1146/annurev-marine-12916-063359>, 2018.
- Levitus, S., Conkright, M. E., Reid, J. L., Najjar, R. G., and Mantyla, A.: Distribution of nitrate, phosphate and silicate in the world oceans, *Progress in Oceanography*, 31, 254–273, 1993.
- Li, G., Cheng, L., Zhu, J., Trenberth, K.E., Mann, M. E., and Abraham, J. P.: Increasing ocean stratification over the past half-century, *Nature Climate Change*, Online: DOI:10.1038/s41558-020-00918-2, 2020.



- 637 Louanchi, F., and Najjar, R.G.: A global monthly climatology of phosphate, nitrate, and silicate in the
638 upper ocean: Spring-summer export production and shallow remineralization, *Global*
639 *Biogeochemical Cycles*, 14, 957-977, 2000.
- 640 Mazeika, P. A.: Thermal domes in the eastern tropical Atlantic Ocean, *Limnology and Oceanography*,
641 12, 537–539, 1967.
- 642 McPhaden, M. J., Zebiak, S. E., and Glantz, M. H.: ENSO as an Integrating Concept in Earth Science,
643 *Science*, 314, 1740-1745, <https://doi.org/10.1126/science.1132588>, 2006.
- 644 Montes, E., Muller-Karger, F. E., Cianca, A., Lomas, M. W., and Habtes, S.: Decadal variability in the
645 oxygen inventory of North Atlantic subtropical underwater captured by sustained, long-term
646 oceanographic time series, *Global Biogeochemical Cycles*, 30, 460-476,
647 doi:10.1002/2015GB005183, 2016.
- 648 Nicholson, S. E.: An analysis of the ENSO signal in the tropical Atlantic and western Indian Oceans,
649 *International Journal of Climatology*, 17, 345-375, [https://doi.org/10.1002/\(SICI\)1097-](https://doi.org/10.1002/(SICI)1097-0088(19970330)17:4%3C345::AID-JOC127%3E3.0.CO;2-3)
650 [0088\(19970330\)17:4%3C345::AID-JOC127%3E3.0.CO;2-3](https://doi.org/10.1002/(SICI)1097-0088(19970330)17:4%3C345::AID-JOC127%3E3.0.CO;2-3), 1997.
- 651 Ono, T., Shiimoto, A., and Saino, T.: Recent decrease of summer nutrients concentrations and future
652 possible shrinkage of the subarctic North Pacific high-nutrient low-chlorophyll region, *Global*
653 *Biogeochemical Cycles*, 22, GB3027, <https://doi.org/10.1029/2007GB003092>, 2008.
- 654 Oschlies, A.: NAO-induced long-term changes in nutrient supply to the surface waters of the North
655 Atlantic, *Geophysical Research Letters*, 28, <https://doi.org/10.1029/2000GL012328>, 2001.
- 656 Palter, J.B., Lozier, M. S., and Barber, R. T.: The effect of advection on the nutrient reservoir in the
657 North Atlantic subtropical gyre, *Nature*, 437, 687–692, 2005.
- 658 Saji, N. N., Goswami, B. N., Vinayachandran, P. N., and Yamagata, T.: A dipole mode in the tropical
659 Indian Ocean, *Nature*, 401, 360-363, <https://doi.org/10.1038/43854>, 1999.
- 660 Schmidtko, S., Johnson, G. C., and Lyman, J. M.: MIMOC: A global monthly isopycnal upper-ocean
661 climatology with mixed layers, *J. Geophys. Res. Oceans*, 118, 1658-1672,
662 <https://doi.org/10.1002/jgrc.20122>, 2013.
- 663 Schmidtko, S., Stramma, L., and Visbeck, M.: Decline in global oceanic oxygen content during the past
664 five decades, *Nature*, 542, 335-339, doi:10.1038/nature21399, 2017.
- 665 Shepherd, J.G., Brewer, P.G., Oschlies, A., and Watson, A.J.: Ocean ventilation and deoxygenation in a
666 warming world: introduction and overview, *Philosophical Transactions of the Royal Society A:*
667 *Mathematical, Physical and Engineering Sciences*, 375, 20170240.
668 <https://doi.org/10.1098/rsta.2017.0240>, 2017.
- 669 Sigman, D. M., and Hain, M. P.: The biological productivity of the ocean, *Nature Education*, 3 (6), 1-
670 16, 2012.
- 671 Stramma, L., Johnson, G. C., Sprintall, J., and Mohrholz, V.: Expanding Oxygen-Minimum Zones in
672 the Tropical Oceans, *Science*, 320, 655-658, doi:10.1126/science.1153847, 2008.



- Stramma, L., Prince, E. D., Schmidtko, S., Luo, J., Hoolihan, J. P., Visbeck, M., Wallace, D. W. R., Brandt, P., and Körtzinger, A.: Expansion of oxygen minimum zones may reduce available habitat for tropical pelagic fishes, *Nature Climate Change*, 2, 33-37, <https://doi.org/10.1038/nclimate1304>, 2012.
- Stramma, L., Schmidtko, S., Bograd, S. J., Ono, T., Ross, T., Sasano, D., and Whitney, F.: Trends and decadal oscillations of oxygen and nutrients at 50 to 300 m depth in the equatorial and North Pacific, *Biogeosciences*, 17, 813-831, <https://doi.org/10.5194/bg-17-813-2020>, 2020.
- Sydeman, W. J., García-Reyes, M., Schoeman, D. S., Rykaczewski, R., R., Thompson, S., A., Black, B. A., & Bograd, S. J.: Climate change and wind intensification in coastal upwelling ecosystems, *Science*, 345, 77–80, 2014.
- Tanhua, T.: Hydrochemistry of water samples during METEOR cruise M83/1. PANGAEA, <https://doi.org/10.1594/PANGAEA.821729>, 2013.
- Tanhua, T.: Hydrochemistry of water samples during METEOR cruise M97. PANGAEA, <https://doi.org/10.1594/PANGAEA.863119>, 2016.
- Tanhua, T.: Hydrochemistry of METEOR cruise M130. PANGAEA, <https://doi.pangaea.de/10.1594/PANGAEA.913986> (dataset in review), 2020.
- Tanhua, T., van Heuven, S., Key, R. M., Velo, A., Olsen, A., and Schirnick, C.: Quality control procedures and methods of the CARINA database, *Earth Syst. Sci. Data*, 2, 35-49, 2010.
- Tanhua, T., Stramma, L., Desai, F., and Löscher, C. R.: Hydrochemistry and molecular biology during Maria S. Merian cruise MSM10/1. IFM-GEOMAR Leibniz-Institute of Marine Sciences, Kiel University, PANGAEA, <https://doi.org/10.1594/PANGAEA.775074>, 2012.
- Vallivattathillam, P., Iyyappan, S., Lengaigne, M., Ethé, C., Vialard, J., Levy, M., Suresh, N., Aumont, O., Resplandy, L., Naik, H., & Naqvi, W.: Positive Indian Ocean Dipole events prevent anoxia off the west coast of India, *Biogeosciences*, 14, 1541-1559, <https://doi.org/10.5194/bg-14-1541-2017>, 2017.
- Whitney, F. A., Bograd, S. J., and Ono, T.: Nutrient enrichment of the subarctic Pacific Ocean pycnocline, *Geophys. Res. Lett.*, 40, 2200-2205, <https://doi.org/10.1002/grl.50439>, 2013.
- Williams, R. G., and Follows, M. J.: Physical transport of nutrients and the maintenance of biological production. In: Fasham, M. J. R. (eds) *Ocean Biogeochemistry. Global Change – The IGBP Series* (closed), Springer, Berlin, Heidelberg, 2003.
- Wishner, K. F., Outram, D. M., Seibel, B. A., Daly, K. L., and Williams, R. L.: Zooplankton in the eastern tropical North Pacific: Boundary effects of oxygen minimum zone expansion, *Deep-Sea Research Part I: Oceanographic Research Papers*, 79, 122-140, <https://doi.org/10.1016/j.dsr.2013.05.012>, 2013.
- Zhang, J.-Z., Mordy, C. W., Gordon, L. I., Ross, A. and Garcia, H. E.: Temporal trends in deep ocean Redfield ratios, *Science*, 289, 1839, <https://doi.org/10.1126/science.289.5486.1839a>, 2000.