

The depth scales of the AMOC on a decadal timescale

Tim Rohrschneider^{1,2}, Johanna Baehr³, Veit Lüscho¹, Dian Putrasahan¹, and Jochem Marotzke^{1,4}

¹Max Planck Institute for Meteorology, Hamburg, Germany

²International Max Planck Research School on Earth System Modelling, Hamburg, Germany

³Institute of Oceanography, Center for Earth System Research and Sustainability, Universität Hamburg, Hamburg, Germany

⁴Center for Earth System Research and Sustainability, Universität Hamburg, Hamburg, Germany

Correspondence: Tim Rohrschneider (tim.rohrschnneider@mpimet.mpg.de)

Abstract. We use wind sensitivity experiments to understand the wind forcing dependencies of the level of no motion as the depth of maximum overturning and the e-folding pycnocline scale as well as their relationship to northward transport of the mid-depth Atlantic meridional overturning circulation (AMOC). In contrast to previous studies, we investigate the interplay of nonlocal and local wind effects on a decadal timescale. We use 30-year simulations with a high-resolution ocean general circulation model (OGCM) which is an eddy-resolving version of the Max Planck Institute Ocean Model (MPIOM). Our findings deviate from the common perspective that the AMOC is a nonlocal phenomenon only, because northward transport and its depth scales depend on both nonlocal Southern Ocean wind effects and local wind effects in the northern hemisphere downwelling region where Ekman pumping takes place. Southern Ocean wind forcing predominantly determines the magnitude of the pycnocline scale throughout the basin, whereas northern hemisphere winds additionally influence the level of no motion locally. In that respect, the level of no motion is a better proxy for northward transport and mid-depth velocity profiles than the pycnocline scale, since the wind forcing dependencies of the level of no motion and maximum overturning are equal. The changes in maximum overturning with wind forcing are explained by the changes in the level of no motion only. This is because wind-driven Ekman compensation is baroclinic and occurs above the level of no motion, and the internal vertical velocity shear that is not influenced by the external Ekman cells stays approximately constant. The analysis of the wind experiments suggests a hemisphere-dependent scaling of the strength of AMOC. We put forward the idea that the ability of numerical models to capture the spatial and temporal variations of the level of no motion is crucial to reproduce the mid-depth cell in an appropriate way both quantitatively and dynamically.

1 Introduction

To date and despite a wide range of theoretical and experimental studies, we do not fully understand inter-hemispheric over-
20 turning of the mid-depth cell in the Atlantic and the role of southern and northern hemisphere processes like the response to
changes in the surface winds. Current understanding of the Atlantic meridional overturning circulation (AMOC) suggests an
interplay between adiabatic pole-to-pole overturning (e.g. Toggweiler and Samuels, 1995; Wolfe and Cessi, 2011) and low-
latitude diabatic forcing that establishes a balance between downwelling of heat and upwelling of deep waters (e.g. Munk and
Wunsch, 1998; Marotzke, 1997). Especially in connection to Southern Ocean processes and the Antarctic circumpolar current
25 (ACC), the effect of winds on the AMOC and basin-wide density stratification has gained considerable attention during the
last two decades (e.g. Marshall and Speer, 2012; Johnson et al., 2019). Poorly understood, however, is the influence of winds
on inter-hemispheric overturning in the Atlantic away from the surface Ekman layer. This paper presents an analysis of wind
sensitivity experiments in order to provide insight into the wind forcing dependence of the inter-hemispheric circulation by
understanding the behavior of the depth scale(s) of the AMOC. We focus on the response of the AMOC to wind forcing on a
30 decadal timescale after the realization of major adjustments.

The research of the present study is inherently about depth scaling that reflects the wind forcing dependence of the AMOC.
Oceanographers use theoretical scaling relationships to provide conceptual understanding and to estimate the strength of the
AMOC in response to different forcings. Nowadays, the most common analytical model to describe meridional overturning
35 of the upper branch of the AMOC is the pycnocline model (Gnanadesikan, 1999). According to this model, the vertically in-
tegrated northward transport in the northern hemisphere is proportional to the basin-averaged e-folding pycnocline scale. The
pycnocline scale is a measure for density stratification and describes how density unfolds vertically. The deeper the pycnocline
scale, the stronger the transport, with the assumption that zonal or meridional density gradients are fixed. The depth scale
itself is determined by Southern Ocean winds and eddies, diapycnal upwelling in the tropics, and North Atlantic deep water
40 formation. In the same manner, a wide range of theoretical studies or scaling arguments rely on the assumption that the pyc-
nocline scale translates a zonal or meridional density gradient into horizontal force that drives northward flow (e.g. Robinson
and Stommel, 1959; Bryan, 1987; Marotzke, 1997; Marotzke and Klingler, 2000). The majority of the different scalings makes
use of the thermal wind relation $\partial v / \partial z = -\Delta \rho_x g / (\rho_0 f L_x)$ which is based on the geostrophic and hydrostatic approximations
of the momentum equations. The left-hand side represents the vertical shear of the zonal-mean meridional velocity, $\Delta \rho_x$ is
45 the west-east density difference, g is the gravitational force, ρ_0 is a reference density, f is the Coriolis parameter, and L_x
is the zonal extent. The thermal wind relation is also key for reconstructing the AMOC strength using observations which are
boundary densities (Hirschi et al., 2003; Baehr et al., 2004; Hirschi and Marotzke, 2007; Baehr et al., 2009). In general, the
theoretical and experimental studies have in common that they integrate vertical shear or zonal or meridional density gradients
twice in the vertical, $\psi \propto \Delta \rho \eta^2$, where ψ is the strength of northward overturning, $\Delta \rho$ is the density gradient, and η is the
50 depth scale. Different assumptions like advective-diffusive balance modify the simple scaling relationship considered here.

The nonequivalence of different depth scales of the AMOC has first been noted in Scott (2000), and recent studies put into question whether the pycnocline scale is the appropriate depth scale to estimate the AMOC strength. For instance, Lev-
ermann and Fuerst (2010) find that the pycnocline scale and meridional density gradients are mutually independent. Using a
55 coarse-resolution model, Griesel and Maqueda (2006) show that the pycnocline scale does not scale northward transport in
experiments in which density gradients are artificially modified. In this connection, DeBoer et al. (2010) find that the depth of
maximum overturning (level of no motion) is a more appropriate parameter to scale maximum overturning. Shakespeare and
Hogg (2012) use the depth scales of the overturning extrema to build an analytical model for both the mid-depth cell and the
abyssal ocean. Finally, Marshall and Johnson (2017) combine the depth of maximum overturning (level of no motion) and the
60 e-folding pycnocline scale to express the relative strength of the ACC and the AMOC. The present study addresses the wind
forcing dependencies of the level of no motion, pycnocline scale and northward transport. Understanding the wind forcing
dependence of the AMOC by understanding its depth scales makes the underlying research question twofold, in the sense that
we discuss the wind forcing dependence of the AMOC using the depth scales and we discuss whether the depth scales are
proxies for northward transport to understand the wind forcing dependence. We hypothesize that the level of no motion is a
65 proxy for northward transport in the inter-hemispheric cell because the background velocity shear of the meridional velocity
may stay constant under changing wind forcing. In this connection, the study is based on different ways or definitions which
describe meridional flow in order to analyze how the changes in wind forcing are translated into the changes in the AMOC.

We focus on the inter-hemispheric region 30S-30N and analyze the interplay of nonlocal wind forcing over the Southern
70 Ocean and local wind forcing in the northern hemisphere downwelling region where Ekman pumping takes place. The absence
of continental barriers in the Southern Ocean and the strong input of momentum at the surface establishes a deep reaching
Ekman overturning cell. The steepening of isopycnals, which is compensated by baroclinic instability that induces an eddy
field, is thought to influence deep stratification and northward transport throughout the basin (e.g. Vallis, 2000; Klingler and
Cruz, 2009; Allison et al., 2011). The zonal periodicity of the ACC boosts the strength of the AMOC in response to an increase
75 in Southern Ocean wind forcing (Klingler et al., 2003, 2004). By contrast, the local influence of northern hemisphere winds on
the AMOC is less understood. Tsujino and Sugimoto (1998) propose an enhancement of the thermohaline circulation due to a
wind-driven buoyancy gain in the upwelling region of the northern hemisphere north of the inter-hemispheric region. Recently,
Cessi (2018) finds that the inter-hemispheric cell weakens in response to increased westerlies at the northern high latitudes.
Considering the northern hemisphere downwelling region at low and mid-latitudes, the local influence of northern hemisphere
80 wind forcing on the AMOC is commonly ignored in the scientific literature on the AMOC. A study by Cabanes et al. (2008)
already indicates that wind stress curl variations may play a crucial role in setting the AMOC shear component which is altered
by the surface forcing according to this study. The wind stress curl is computed by the rotation of the surface wind vector and
establishes downwelling or upwelling regions. In this connection, vertical Ekman pumping emerges from the forcing imposed
by the wind stress curl at the surface. We address the question how changes in both nonlocal and local wind forcing influence
85 the AMOC. We hypothesize that the influence of northern hemisphere winds on the AMOC is substantial.

To answer the specific questions that are outlined above, we use 30-year wind experiments with a high-resolution ocean general circulation model (OGCM). We expect a robust response of the AMOC on a decadal timescale because major adjustments by basin-wide wave propagation are realized on this timescale. The basin-wide transmission of density signals by wave propagation occurs on a pentadal and decadal timescale. Limiting the analysis to the adjustment on a decadal timescale makes it possible to use presumably more realistic eddy-resolving wind experiments rather than fully equilibrated low-resolution model runs. In this way we avoid uncertainties introduced by eddy parameterization. High-resolution model output is needed because we analyze the relationships between depth scales as reference depths and meridional transport and its vertical velocity shear. These relationships depend sensitively on the vertical and horizontal model resolution as wave propagation of density signals does.

To the best of our knowledge and despite a wide range of theoretical and modeling studies on the Atlantic circulation, the research questions of this study have never been answered in an explicit way. That is to say, from an overarching perspective we analyze whether the depth scales are proxies for northward transport in the inter-hemispheric cell, and we ask whether we should adopt a more nonlocal or local perspective with respect to inter-hemispheric overturning and hemispheric differences.

In the following section we briefly describe the different experiments and the experimental strategy. In section 3 we describe the differences in density stratification with the changes in wind forcing as well as the wind forcing dependencies of the level of no motion, the pycnocline scale and northward transport. In section 4 we analyze the relationship between velocity profiles or shear and the depth scales.

2 Experiments and methods

2.1 Numerical model and experiments

We use wind experiments conducted with a vertically and horizontally high-resolution, eddy configuration of the Max Planck Institute for Meteorology ocean model (MPIOM). The version is called TP6ML80 and has been developed within the German Consortium project STORM (von Storch et al., 2012). It is based on a tripolar grid with a horizontal resolution of 0.1 degrees and a vertical resolution of 80 unevenly spaced levels. Compared to low-resolution MPIOM versions, we assume better model physics in terms of the resolution of mesoscale eddies as well as wave propagation. Gutjahr et al. (2019) already indicate that the high-resolution MPIOM configuration reveals the most realistic simulation compared to low-resolution MPIOM versions in terms of the mean state of the ocean. For instance, a cold bias in sea surface temperature over the Southern Ocean is strongly reduced, because the resolved eddies better influence the flattening and cropping of isopycnals compared to the Gent-McWilliams thickness diffusivity parameterization (Gent et al., 1995). We do not expect that the wind forcing dependence of the AMOC depth scales differs between high- and low-resolution model runs. However, the horizontal transmission of density signals is sensitive to the horizontal model grid and the accumulation of vertical shear is sensitive to the vertical model grid.




Figure 1. The zonal-mean zonal wind stress in the 1X experiment (black), 2XSH experiment (blue), and 2X experiment (red) (Lüschow et al., 2021).

120 We have available three different wind experiments with realistic geometry (Lüschow et al., 2021); Table 1. In the 1X experiment, the standard surface momentum and buoyancy fluxes from NCEP-NCAR reanalysis-1 (Kalnay et al., 2018) are applied. We are not interested in high-frequency variability and therefore change the monthly-mean climatology of the surface wind stress only. In the 2XSH experiment, the monthly climatology of the zonal and meridional surface wind stress is doubled only over the Southern Ocean. By applying a sine function which declines to zero towards the equator in the southern hemisphere, we obtain a smooth transition of the surface wind stress towards the original surface momentum fluxes. In the 2X experiment, we double the monthly climatology of the zonal and meridional surface wind stress in both the northern and southern hemispheres throughout the basin. Fig. 1 shows the climatology of the zonal surface wind stress of the different experiments. The wind experiments are initialized with the state of the 1X experiment, and the wind forcing is switched on at year 1980. Because the computational costs for an eddy-resolving simulation are still large, the simulation period is only until year 2010.

130 Using monthly-mean output, we focus on the time-window 1991 to year 2010 after the realization of major adjustments to the changes in the surface wind stress in order to analyze the AMOC response to forcing. That is, we allow for a decadal adjustment of the AMOC and density field towards stationarity before using the model time series. Analyzing the decadal response of the AMOC to Southern Ocean wind forcing, Klinger and Cruz (2009) show that the AMOC in the source region adjusts on an inter-annual timescale. Lüschow et al. (2021) discuss the adequacy of the wind experiments in terms of integration length.

135

Table 1. Overview of the experimental strategy

1X (1991-2010)	The reference experiment which mimics observed conditions
2XSH (1991-2010)	We double the monthly climatology of the zonal and meridional surface wind stress over the Southern Ocean only in order to analyze the nonlocal effects of Southern Ocean winds
2X (1991-2010)	We double the monthly climatology of the zonal and meridional surface wind stress throughout the basin in order to analyze the interplay between nonlocal and local wind effects and extract local wind forcing dependencies

2.2 Experimental strategy and quantities

The AMOC is described by the overturning streamfunction in the latitude-depth section, $\psi(t, z, y) = \int_z^D \int_{x_e}^{x_w} v(t, z, y, x) dx dz$, where v is the meridional velocity, z is the depth with D the depth at the bottom of the ocean column, and x_e and x_w are the eastern and western boundaries of the Atlantic basin.

140

We focus on the upper, northward flowing branch of the mid-depth cell which rotates clockwise, viewed from the east. Fig. 2a,b,c show the time-mean (1991-2010) overturning cells in the wind experiments. Analyzing the AMOC in the inter-hemispheric region 30S-30N south and north of the equator, we explore the nonlocal response to changes in Southern Ocean winds and local wind effects in the downwelling region of the northern hemisphere. In general, the mid-depth cell strengthens with higher wind forcing over the Southern Ocean. However, at this point, from Fig. 2 we cannot capture the details of the meridional velocities in terms of spatial variations that may influence the relationship between overturning and its depth. The northward flow seems to be continuous and contiguous throughout the basin, but regional dynamics may give rise to local wind forcing dependencies of the AMOC which can be hardly identified away from the surface Ekman layer. The surface Ekman flux, for instance, is compensated by an interior return flow which changes the meridional transport at different depths. Furthermore, the wind stress curl over the basin imposes a forcing that may change stratification and the meridional transport and its depth locally.

150

We limit our analysis of the different experiments to a set of quantities (Table 2). The level of no motion of the mid-depth cell is the depth of maximum overturning where zonally averaged velocities reverse in sign, $\eta_\psi = z_{\psi(\max)}$. We use the monthly-mean outcome of the related quantities and subsequently take the time-mean (1991-2010). In doing so, we describe in more detail the behavior of the level of no motion and its relationship to northward transport. The reader should notice that the level

155




Figure 2. The time-mean AMOC streamfunctions (1991-2010) in the (a) 1X experiment, (b) 2XSH experiment, and (c) 2X experiment.

Table 2. Experimental quantities

ψ	AMOC streamfunction
ρ	Potential density
η_ψ	Level of no motion
η_ρ	Pycnocline scale
η_w	Advective depth scale
ψ_t	Total maximum overturning streamfunction
ψ_g	Geostrophic approximation of the maximum overturning streamfunction
$\frac{\partial \psi}{\partial z}$	Vertical derivative of the overturning streamfunction (velocities)
$\frac{\partial^2 \psi}{\partial z^2}$	Second vertical derivative of the overturning streamfunction (shear)
ψ^*	Maximum overturning streamfunction associated with the level of no motion

of no motion incorporates nonlinearities because it is an integrated quantity and related to the zonal-mean meridional velocity. The level of no motion is not a zonal-mean quantity that is averaged over the zonal extent. Further, we avoid that the algorithm selects a model level within the depth range of the surface Ekman layer or perturbations that emerge from equatorial upwelling.

160

The pycnocline scale is the single mode e-folding scale for vertical density stratification, and the profile of the latter is assumed to be exponential and self-similar. We use the following fitting algorithm at each grid cell and subsequently take the zonal and temporal mean, $\eta_\rho = 2 \frac{\int_{z_r}^0 (\rho - \rho_r) z dz}{\int_{z_r}^0 (\rho - \rho_r) dz}$ (Gnanadesikan, 1999; Gnanadesikan et al., 2007; DeBoer et al., 2010). The reference depth is $z_r = -2500$ meters, and ρ_r is the potential density at the reference depth. Using this algorithm we take the

165 e-folding scale twice. In this way, 80 to 90 percent of the vertical density change is scaled, which scales the upper branch of the mid-depth cell. In some regions, especially at the lateral margins of the basin, the profile for vertical density stratification may become more linear and deviate from an exponential profile. However, such deviations from a perfectly exponential profile do not restrict the ability of the pycnocline scale to interpret density stratification in different regions in the sense that it is an independent measure as long as density increases with depth.

170

We make use of a third depth scale and compute the advective depth η_w in order to provide the linkages between the differences in the wind stress curl, the differences in the density field ρ , and the depth scales. The advective depth scale gives the vertical depth range for local Ekman pumping and likewise emerges from the forcing imposed by the wind stress curl at the surface. It thereby scales the maximum change in density stratification associated with the rotation of the surface wind vector.

175 Theory of wind driven changes in stratification can be traced back to the development of the planetary geostrophic equations (e.g. Welander, 1959; Bryan, 1987) and the related theory on the ventilated thermocline in the subtropical region (Luyten et al., 1983). Classical scaling of the thermocline equations suggests that the advective depth scale is directly proportional to the square-root of the local Ekman pumping velocity W_E (e.g. Vallis, 2000), $(\frac{W_E f L^2}{g'})^{0.5}$, where f is the Coriolis parameter, L is the basin width, and g' is the reduced gravity. We compute the local Ekman pumping velocity W_E from the wind stress data of

180 the wind experiments.

During the course of our study we analyze both the total maximum overturning streamfunction ψ_t and the geostrophic approximation of the maximum overturning streamfunction ψ_g ; that is, the total streamfunction minus the surface Ekman flux. We compute both ψ_t and ψ_g because their wind forcing dependencies should be fundamentally different due to the interior, geostrophic return flow of the surface Ekman flux. In this way, we analyze the relationship between overturning and its depth using vertically integrated transport that may depend on the ageostrophic surface component by vertical integration.

185

We use the zonal surface wind stress τ_x to compute the strength of the surface meridional Ekman transport, $\int_{x_e}^{x_w} -\frac{\tau_x(x,y,t)}{\rho_0 f} dx$, where ρ_0 is the reference density and f the Coriolis parameter. We do not subtract the interior return flow of the surface Ekman flux. For the sake of simplicity a range of studies assumes that the interior return flow of the surface Ekman flux is barotropic even on longer than a monthly timescale (e.g. Hirschi and Marotzke, 2007; Moreno-Chamarro et al., 2016). Based on an idealized experiment with an OGCM, a recent study demonstrates that the return flow is baroclinic and has strong contributions at the upper levels of the ocean on the timescale considered here (Williams and Roussenov, 2014). On a monthly timescale, anomalies of the return flow are barotropic because the density field does not adjust (e.g. Jayne and Marotzke, 2001), but these anomalies are negligible in the set-up presented in this study and by their nature do not change the time-mean outcome. Later on, we support the perspective that the interior return flow of the surface Ekman flux is baroclinic. This finding has been demonstrated theoretically by McCreary and Lu (1994).

190

195

Finally, we analyze whether there are causal linkages between the changes in the depth scales and northward transport at different depths. Therefore, we compute the vertical derivative of the AMOC streamfunction $\frac{\partial \psi}{\partial z}$, which represents the zonal-mean meridional velocity scaled by the basin-width. We then compute the second derivative of the AMOC streamfunction $\frac{\partial^2 \psi}{\partial z^2}$, which represents the vertical velocity shear scaled by the basin-width. In this way, we analyze to which degree the changes in the northward transport of the mid-depth cell are directly related to the displacement of the level of no motion. The computation of $\frac{\partial \psi}{\partial z}$ and $\frac{\partial^2 \psi}{\partial z^2}$ below the surface Ekman layer is independent of the way how we approximate the maximum overturning streamfunctions ψ_t and ψ_g . To finish our analysis, we compute the maximum overturning streamfunction ψ^* that

200

205

is associated with the level of no motion only. We use the time-mean vertical velocity shear $\frac{\partial^2 \psi}{\partial z^2}$ of the 1X reference experiment and integrate vertically from the level of no motion in the 2XSH and 2X experiments to the top. Before averaging over the 20 years, we use the highest temporal resolution and compute the integral with the reference depth for each month in order to ensure the robustness of the results.

210 3 Wind forcing dependencies

3.1 Wind-driven changes in stratification

As a starting point, we highlight the differences in density stratification in the Atlantic basin between the wind experiments. Afterwards, we relate these differences to the level of no motion η_ψ and the pycnocline scale η_ρ . We normalize the zonal-mean potential density to highlight the differences in density stratification between the wind experiments, $\frac{\rho_0 - \rho}{\rho_0}$, with the reference
 215 density $\rho_0 = 1025 \text{ kg m}^{-3}$. Fig. 3a,b show the time-mean difference in density stratification between the 2XSH and 1X experiment and the 2X and 1X experiment. Fig. 3c illustrates the advective depth scale η_w in the Atlantic. Within the range of the advective depth scale η_w , isopycnals shoal towards the equator due to equatorial divergence and deepen in the subtropical region towards higher latitudes due to the local forcing that is imposed by the wind stress curl. Deep stratification below η_w but within the depth range of the upper, northward flowing branch of the mid-depth cell reveals the same behavior in the southern
 220 hemisphere, but isopycnals rise constantly towards the region of North Atlantic deep water formation.

The time-mean differences in density stratification between the wind experiments mirror the experimental set-up of the present study. In general, Fig. 3a,b suggests that the difference in density stratification between the 2XSH experiment and the 1X experiment is driven by the wind stress curl over the Southern Ocean, which has a strong influence in the southern
 225 hemisphere, but deep isopycnals change even in the northern hemisphere. The change of the 2X experiment relative to the 1X experiment is driven by the change of the climatology of the wind stress curl in both hemispheres. Local Ekman pumping displaces isopycnals downward also north of the equator with a maximum change in the subtropical region, and this displacement is scaled by η_w (Fig. 3c). The influence of the local wind forcing prevails but deep isopycnals are also influenced nonlocally below η_w . Both nonlocal and local wind effects change the density field, and their relative influence on stratification depends
 230 on location and depth. In the following, we analyze how these changes in density stratification translate into changes in both depth scales η_ψ and η_ρ and maximum overturning ψ . Afterwards, in section 4, we discuss how the changes in wind forcing translate into changes in the level of no motion η_ψ from a generic point of view on the vertical velocity shear of the meridional velocity.

3.2 Level of no motion and pycnocline scale

235 Fig. 4 shows the meridional dependence of the level of no motion η_ψ and the zonal-mean pycnocline scale η_ρ in the wind experiments in order to explore the wind forcing dependencies of the depth scales. Considering the 1X experiment, the spatial




Figure 3. Differences in the zonal-mean density stratification averaged over the years 1991-2010: (a) the difference between the 2XSH and 1X experiments and (b) the difference between the 2X and 1X experiment. The black contour lines represent the zonal-mean density stratification (10^{-1}) in the 1X reference experiment. Density stratification is computed by $\frac{\rho_0 - \rho}{\rho_0}$, with the reference density $\rho_0 = 1025 \text{ kg m}^{-3}$. In (c) we show the advective depth scale η_w in meters depth in the 1X experiment (black) and the 2X experiment (red), with the reduced gravity set to $g' = 0.013$ (opaque) and $g' = 0.02 \text{ m s}^{-2}$ (transparent). The smaller the value for the reduced gravity, the deeper the advective depth scale η_w .




Figure 4. The time-mean (1991-2010) wind forcing dependencies of the level of no motion η_ψ (solid) and the pycnocline scale η_ρ (dashed) with respect to the 1X experiment (black), the 2XSH experiment (blue), and the 2X experiment (red).

variations of η_ψ and η_ρ coincide in the sense that both shoal towards the equator in the southern hemisphere and stay more or less constant or change slightly in the northern hemisphere. Although the general behavior of the level of no motion and the pycnocline scale coincides, the pycnocline scale η_ρ measures how density stratification unfolds over the ocean column. In contrast, the level of no motion is a single layer at a certain depth. By their nature, the level of no motion η_ψ and pycnocline scale η_ρ do not match exactly. Even though we know how density stratification unfolds, it is unclear whether the pycnocline scale represents the differences in stratification as found in the previous section.

The wind forcing dependencies of the level of no motion and the pycnocline scale differ. The pycnocline scale η_ρ deepens in the 2XSH and 2X experiments relative to the 1X reference experiments throughout the basin. The pycnocline scales η_ρ in the 2XSH and 2X experiments are congruent, and η_ρ is mainly determined by the wind stress over the Southern Ocean. Local changes in the wind forcing at lower latitudes and in the northern hemisphere do not change η_ρ significantly. In the northern hemisphere, density stratification approximately unfolds at the same scale despite the differences in density stratification at the depth of η_w . In the southern hemisphere, the wind forcing dependence of the level of no motion η_ψ is in line with the wind forcing dependence of pycnocline scale η_ρ . However, we find considerable differences between the wind experiments in the northern hemisphere due to an additional dependence on the local wind forcing. Wind forcing over the Southern Ocean nonlocally deepens the level of no motion η_ψ in the northern hemisphere. Yet, in addition to this nonlocal effect, a local effect acts




Figure 5. The time-mean (1991-2010) wind forcing dependencies of the total maximum overturning streamfunction ψ_t (solid) and the geostrophic maximum overturning streamfunction ψ_g (dashed) with respect to the 1X experiment (black), the 2XSH experiment (blue), and the 2X experiment (red). Considering ψ_g , we exclude the equatorial band where the approximation of the surface Ekman flux breaks down.

on the level of no motion η_ψ because we observe differences in the northern hemisphere subtropical region between the 2XSH and 2X experiments. The findings on the wind forcing dependencies of η_ψ correspond to those findings on the differences in density stratification between the wind experiments. The level of no motion η_ψ is connected to the advective depth scale η_w and Ekman pumping locally.

3.3 Maximum overturning and its depth

We now analyze the wind forcing dependence of the northward flowing branch of the mid-depth cell. We compute the total maximum overturning streamfunction ψ_t and the geostrophic maximum overturning streamfunction ψ_g . Conceptually, the differences between ψ_t and ψ_g provide insight on the degree to which the depth scale(s) are proxies for the strength of the AMOC. Computing the geostrophic maximum overturning streamfunction ψ_g , the level of no motion is unchanged, but the clockwise (upper) and counterclockwise (lower) rotating overturning cells are substantially altered. The maximum streamfunction ψ_t includes the surface Ekman flux and the maximum streamfunction ψ_g excludes the surface Ekman flux. However, the surface Ekman fluxes have to be compensated by an interior return flow that changes in relationship between overturning and its depth.

The wind forcing dependencies of the level of no motion and the total maximum overturning streamfunction are equal. By contrast, the wind forcing dependencies of the level of no motion and the geostrophic maximum overturning streamfunction are unequal. As a first step towards understanding, this behavior helps understand whether and why the level of no motion is a proxy for northward flow. Fig. 5 shows the meridional dependence of both the total and geostrophic maximum overturning streamfunction (ψ_t, ψ_g). Considering ψ_t , the southern latitudes of the southern hemisphere are strongly influenced by the surface Ekman flux which scales with the zonal wind stress. Northward of the Southern Ocean (30S-10S), the AMOC becomes increasingly geostrophic. In the southern hemisphere, the strength of the mid-depth cell increases with higher wind forcing, and we find a stronger transport in the 2XSH and 2X experiment than in the 1X experiment. In this region, the total maximum streamfunctions ψ_t of the 2XSH and 2X experiments are approximately equal. However, ψ_t intensifies north of the equator. In the subtropical region (10N-30N), ψ_t increases with higher wind forcing, and the integrated transport of the 2X experiment is stronger than the integrated transport of the 2XSH experiment.

In contrast to ψ_t , the geostrophic maximum overturning streamfunction ψ_g is relatively weak near the Southern Ocean but constantly increases towards the equator. North of the Southern Ocean (30S-10S), ψ_g is increasingly dependent on the geostrophic return flow that compensates the southward surface Ekman flux locally. As a result of the interior return flow, ψ_g becomes stronger with higher wind forcing, with markedly higher values in the 2X experiment compared the 2XSH experiment. At the low and mid-latitudes of the northern hemisphere (10N-30N), the interior return flow of the surface Ekman flux is directed southward and strongest at roughly 15°N. At these latitudes, ψ_g of the 2X experiment is weaker than both the geostrophic transport of the 1X reference experiment and the 2XSH experiment. Comparing ψ_t and ψ_g , the Ekman cells play a crucial role in setting the relationship between overturning and its depth. We discuss the role of the Ekman cells and the possible explanations for the differences in maximum overturning besides the Ekman return flow later on, in section 4. First, we highlight the relationships between the maximum overturning streamfunctions and the depth scales in order to bring together their spatial dependencies.

Combining our findings from Fig. 4 and 5, we describe the relationship between northward overturning and its depth from a more nonlocal perspective on hemispheric differences. To highlight hemispheric differences in the inter-hemispheric region, we show the meridional averages (30S-10S) and (10N-30N) (Fig. 6). In the southern hemisphere, the level of no motion η_ψ of the 2XSH and 2X experiment fall into the same model layer, and the associated maximum streamfunctions ψ_t are not notably different. In this region, ψ_g increases with higher wind forcing. South of the equator, the wind forcing dependencies of the level of no motion η_ψ as well as the pycnocline scale η_ρ (not shown) are in line with the wind forcing dependencies of ψ_t , in the sense that a broader depth scale is related to stronger northward overturning. Insights on the northern hemisphere are not as simple as in the southern hemisphere. The level of no motion η_ψ deepens with higher wind forcing due to the downward displacement of isopycnals at mid-depth, whereas η_ρ (not shown) reveals the same wind forcing dependencies as in its southern counterpart. The differences in the depth of the level of no motion η_ψ between the wind experiments are important although they are set by one model layer only, because even a small change in the model layers implies a change in the accumulation




Figure 6. The time-mean (1991-2010) relationship between maximum overturning and its depth, the level of no motion, in the southern hemisphere and the northern hemisphere. The markers represent the meridional averages (30S-10S) (stars) and (10N-30N) (triangles). We show both the total maximum overturning streamfunction (ψ_t , opaque) and the geostrophic maximum overturning streamfunction (ψ_g , transparent). We do not show the pycnocline scale η_ρ because the qualitative behavior is same as the behavior of the level of no motion η_ψ except that the pycnocline scales of the 2XSH and 2X experiment are congruent throughout the basin. The thin grey lines indicate the vertical model grid.

of vertical velocity shear, and the pycnocline scale η_ρ does not account for these details. As in the southern counterpart, the differences in the level of no motion η_ψ do not represent the changes in ψ_g , but the changes in ψ_t are consistent with the common assumption that the northward transport of the mid-depth cell becomes stronger with a deeper level of no motion η_ψ .

305

In general, the differences between ψ_t and ψ_g in the inter-hemispheric region suggest that the interior return flow of the surface Ekman flux is baroclinic and mostly compensated above the level of no motion η_ψ . If so, the level of no motion is a proxy for ψ_t rather than a proxy for ψ_g in the case that the vertical velocity shear stays approximately constant with wind forcing. This raises the questions how well the total and geostrophic maximum overturning streamfunctions represent interior

310 flow despite the surface Ekman flux or its geostrophic return, and how well the level of no motion represents transport at different depths. In order to answer these questions, we analyze meridional velocity profiles ($\frac{\partial\psi}{\partial z}$) and shear ($\frac{\partial^2\psi}{\partial z^2}$).

4 The relationship between the depth scales and velocity profiles

The depth scales can be related to vertical velocity profiles ($\frac{\partial\psi}{\partial z}$) and shear ($\frac{\partial^2\psi}{\partial z^2}$). To discuss changes in the velocity profiles with respect to the level of no motion and the pycnocline scale, we first compute the vertical derivative of the AMOC streamfunction
315 $\frac{\partial\psi}{\partial z}$ (the zonal-mean meridional velocity scaled by the basin width). Neglecting the surface Ekman layer and integrating from the level of no motion η_ψ to the top gives the geostrophic approximation of the maximum overturning streamfunction that approximately represents the geostrophic conditions in the interior. Analyzing $\frac{\partial\psi}{\partial z}$ is a perspective on force balance, and we disentangle different contributions.

320 Fig. 7 shows the zonal-mean meridional velocities ($\frac{\partial\psi}{\partial z}$) in the wind experiments and the difference in $\frac{\partial\psi}{\partial z}$ between these wind experiments. The differences in $\frac{\partial\psi}{\partial z}$ between the wind experiments are strongest near the equator at the upper levels where the vertical velocity shear changes drastically. Taking the difference between the 2X and 1X experiments, we find an increase in $\frac{\partial\psi}{\partial z}$ south of the equator and a decrease north of the equator. To a substantial extent, these changes can be attributed to the strengthening of the local Ekman cells. The differences in $\frac{\partial\psi}{\partial z}$ at the upper levels between the different experiments demonstrate
325 that the Ekman return flow is baroclinic and occurs mostly above η_ψ . The strong influence of the Ekman cells near the surface suggests that, at these levels, the external wind-driven flow associated with the Ekman cells superposes the internal flow that is associated with the level of no motion. These considerations support the perspective that η_ψ is a proxy for ψ_t rather than a proxy for ψ_g . Small differences in ψ_t emerge in case of weak compensation of the surface Ekman flux below η_ψ .

330 In Fig. 7 we further find differences in $\frac{\partial\psi}{\partial z}$ at mid-depth that are associated with the nonlocal wind forcing over the Southern Ocean. These changes can be inferred from the difference between the 2XSH and 1X experiments. The difference between these experiments is related to the enhanced inflow from the southern hemisphere into the northern hemisphere, which in turn can be related to the strengthening of the western boundary current with an increase in wind forcing (not shown). In the 2X experiment, however, the perturbation associated with the Ekman cells is stronger than the perturbation associated with the
335 strengthening of the western boundary current. The Ekman cells south and north of the equator further obscure the influence of the local Ekman pumping velocity W_E on stratification and transport at the advective depth η_w , which can be hardly identified in Fig. 7. The changes in the wind stress curl with changing wind forcing must be translated into the changes of the level of no motion η_ψ , and next we analyze explicitly how different depths of the level of no motion η_ψ result in different velocity profiles.

340 In this connection, we analyze whether there is a direct relationship between the vertical velocity profiles described by $\frac{\partial\psi}{\partial z}$ and the changes in the level of no motion η_ψ , by computing the meridional averages 30S-10S and 10N-30N (Fig. 8a,b). We analyze how much of the velocity profiles can be predicted by the level of no motion only. For this purpose, we assume vertical velocity shear that is constant under changing wind forcing. As indicated by the vertical bars, deep transport has a substantial contribution in both hemispheres because shear accumulates in the vertical over a wide depth range. In the southern hemi-
345 sphere, the changes in the level of no motion η_ψ and the pycnocline scale η_ρ correspond to the changes in the actual velocity




Figure 7. The time-mean (1991-2010) vertical derivative of the AMOC streamfunction $\frac{\partial\psi}{\partial z}$ in the (a) 1X experiment, (b) 2XSH experiment, and (c) 2X experiment. We further show the difference in $\frac{\partial\psi}{\partial z}$ between the (d) 2XSH and 1X experiments, (e) the 2X and 1X experiments, and (f) the 2X and 2XSH experiments. The black lines represent the level of no motion η_ψ . We exclude the surface Ekman layer.




Figure 8. The meridional averages in the southern hemisphere (a, 30S-10S) and in the northern hemisphere (b, 10N-30N) of the time-mean (1991-2010) vertical derivative of the AMOC streamfunction $\frac{\partial\psi}{\partial z}$ in the 1X experiment (black), 2XSH experiment (blue), and 2X experiment (red). We exclude the surface Ekman layer. The circles show the meridional averages of the level of no motion η_ψ and the stars show the meridional averages of the pycnocline scale η_ρ . The dashed lines represent the velocity profiles that arise from the displacements of the level of no motion η_ψ while the velocity shear $\frac{\partial^2\psi}{\partial z^2}$ is held constant (1X). Note that in the Southern hemisphere the blue dashed line and the red dashed line fall on top of one another. We indicate the transport (Sv) at the deeper levels by the annotation at the vertical bars. The thin grey lines indicate the vertical model grid.

profiles $\frac{\partial\psi}{\partial z}$ (solid lines) at deeper levels. In the northern hemisphere downwelling region, the pycnocline scale does not change with local changes in wind forcing, whereas the level of no motion deepens locally. To assess to which degree the level of no motion η_ψ represents changes in the velocity profiles, we vertically integrate the time-mean vertical velocity shear $\frac{\partial^2\psi}{\partial z^2}$ of the 1X reference experiment from the level of no motion η_ψ in the 2XSH and 2X experiments with zero reference velocity (dotted lines). This is not possible in the case of the pycnocline scale η_ρ , because it is a scale height and even a small layer difference results in differences in $\frac{\partial\psi}{\partial z}$. We find that the changes in deep velocity profiles are related to the changes in the level of no motion η_ψ in both hemispheres. To a considerable extent, even the velocity profile of the 2XSH experiment at the upper levels in both hemispheres is connected to the displacement of the level of no motion η_ψ . In the 2X experiment, however, the signal that arises from the interior return flow of the surface Ekman flux overcomes the signal that arises from the displacement of η_ψ . In the southern hemisphere at the upper levels, the velocities associated with the changes of the level




Figure 9. The total maximum overturning streamfunction ψ_t and the maximum overturning streamfunction ψ^* in the case that we hold the vertical velocity shear $\frac{\partial^2 \psi}{\partial z^2}$ constant (1X).

of no motion η_ψ are lower than the actual velocities in this experiment. In the northern hemisphere at the upper levels, the velocities associated with the changes of the level of no motion η_ψ are much higher than the actual velocities. Integrating the velocity profiles vertically, however, the Ekman cells should cancel out such that the level of no motion is a proxy for northward transport.

360

We therefore show the changes in maximum overturning that are associated with the level of no motion only by analyzing the vertically integrated transport. Fig. 9 shows the total maximum overturning streamfunction ψ_t and the maximum overturning streamfunction ψ^* in the case that we hold the vertical velocity shear constant. We find that the changes in total maximum overturning ψ_t are explained by the changes in the level of no motion to a very large degree. The maximum overturning streamfunctions ψ_t and ψ^* are approximately congruent, and the vertical velocity shear $\frac{\partial^2 \psi}{\partial z^2}$ stays approximately constant with wind forcing on the timescale considered here. There is deviation at the equator due to systematic errors that arise from perturbations in equatorial upwelling. Away from the equator, however, the differences between the 2XSH and 2X experiments arise solely from the differences in the level of no motion. The mechanism how the changes in wind forcing translate into changes in the level of no motion η_ψ is thus easy to comprehend from a generic point of view. Small changes in the zonal pressure gradients at the depth range scaled by advective depth scale η_w are related to substantial differences in the level of no motion η_ψ , because the internal velocity shear that is not influenced by the external Ekman cells hardly changes between the wind experiments.

370

5 Discussion

In line with the current understanding of the Atlantic circulation, Southern Ocean winds boost the strength of the AMOC and change density stratification throughout the basin (e.g. Vallis, 2000; Klinger et al., 2003, 2004; Klinger and Cruz, 2009). Northern hemisphere winds over the downwelling region additionally influence the meridional flow and density stratification locally, which is commonly ignored in the scientific literature on the AMOC. The present study is based on simulations with an eddy-resolving OGCM on a decadal timescale rather than a fully equilibrated experiment. We find a robust adjustment of the AMOC and density field, which demonstrates the realization of major adjustments due to wave propagation, and the 30-year simulations are long enough to analyze the wind forcing dependencies of the depth scales and northward transport. The wind forcing-dependence of the AMOC is reflected by the wind experiments.

The findings of the present study support the pycnocline model described in Gnanadesikan (1999) in the sense that Southern Ocean wind forcing deepens the pycnocline scale and the level of no motion and strengthens the AMOC. However, local wind forcing over the northern hemisphere downwelling region additionally influences the level of no motion and northward transport locally. In that respect, the level of no motion is more appropriate to scale northward transport than the pycnocline scale. By artificial modification of density gradients in OGCM experiments, Griesel and Maqueda (2006) and DeBoer et al. (2010) indicate that the pycnocline scale does not scale northward transport at all. By contrast, we provide insight on the scaling behavior of the depth scales from a conceptual point of view, and the pycnocline scale fails to scale northward transport in the northern hemisphere.

Wind stress curl variations at the surface translate into changes in the AMOC. The changes of the AMOC with changing wind forcing in the inter-hemispheric region are explained by the changes in the level of no motion. The internal velocity shear that is not influenced by the external Ekman cells remains constant on the timescale considered here. In contrast to what is stated in Cabanes et al. (2008) who analyze interannual variability, the forcing imposed by the wind stress curl at the surface does not substantially change the vertical shear but the reference depth of the AMOC shear component. Our findings also deviate from Levermann and Fuerst (2010) who evaluate the pycnocline model using a model of intermediate complexity. They analyze equilibrated experiments which reproduce the response to Southern Ocean wind forcing and focus on meridional density gradients instead of zonal density gradients to represent vertical velocity shear. Using meridional density gradients instead of zonal density gradients is based on the assumption that these gradients are proportional and have the same order of magnitude, and zonal and meridional velocities compare well with one another. According to their findings, both the pycnocline scale and meridional density gradients vary, while according to our study the internal velocity shear remains fixed. We speculate that our high-resolution simulation better simulates velocity shear.

The displacement of the level of no motion in the MPIOM wind experiments approximates the conditions in the interior with the Ekman cells mainly cancelled out. Comparing the wind experiments, the ocean response at the upper levels is much more

complex than the response at the deeper levels, which is mostly related to the baroclinicity of the interior return flow of the surface Ekman flux. However, integrating vertically, the changes that are associated with the level of no motion give approximately the changes in the total maximum overturning streamfunction with changing wind forcing. As a general contribution and supporting the theoretical considerations made in McCreary and Lu (1994), our findings give baroclinic Ekman compensation which has been demonstrated in an idealized way by Williams and Roussenov (2014). Baroclinic Ekman compensation may depend sensitively on the resolution of an OGCM.

The wind forcing dependence of the AMOC suggests that the temporal adjustment of the AMOC to global warming is not independent of location. Both nonlocal Southern Ocean wind forcing and local wind forcing in the northern hemisphere downwelling region are likely to influence the adjustment of the level of no motion and northward transport in the inter-hemispheric region.

6 Summary and Conclusions

We use wind sensitivity experiments to analyze the wind forcing dependence of depth scales of the inter-hemispheric cell in the Atlantic and their relationship to northward transport. We focus on the inter-hemispheric region in order to analyze the interplay of nonlocal and local wind effects, and our perspective deviates from the common view that the AMOC is a nonlocal phenomenon only. The dynamics of the inter-hemispheric cell can only be understood by analyzing both Southern Ocean wind effects and local wind effects in the northern hemisphere downwelling region which arises from the forcing imposed by the wind stress curl at the surface.

We find different wind forcing dependencies of the pycnocline scale and the level of no motion. Southern Ocean processes determine the magnitude of the pycnocline scale, whereas northern hemisphere wind stress additionally influences the level of no motion. The pycnocline scale is insensitive to the local wind stress over the northern hemisphere and cannot capture the details of deep velocity profiles and mid-depth stratification. Local wind forcing changes density stratification and displaces isopycnals downward at an advective depth. In that respect, the level of no motion is a better proxy for velocity profiles than the pycnocline scale, because the level of no motion accounts for the changes in the surface winds over the northern hemisphere.

To a large extent, the changes in transport at deeper levels below the surface layers between the wind experiments can be related to the changes of the level of no motion in the case that we hold the vertical velocity shear of the meridional velocity constant. The changes in the level of no motion between the wind experiments explain a large fraction of the changes in the meridional velocities. Near the surface, however, the signal that arises from the interior return flow of the surface Ekman flux overcomes the signal that arises from the displacement of the level of no motion. In this regard, the changes in the meridional flow which are associated with the displacement of the level of no motion and the actual transport can differ considerably at these levels. However, the changes in maximum overturning with changing wind forcing are explained by the changes in

440 the level of no motion only, because the internal velocity shear that is not influenced by the external Ekman cells stays constant.

There is no unique way to describe and quantify interior geostrophic flow that is not directly influenced by the local Ekman cells, and in this sense both the total maximum overturning streamfunction and the geostrophic maximum overturning streamfunction are approximations for the conditions in the interior. Our findings suggest that the differences in the total maximum
445 overturning streamfunction are related to the differences in the level of no motion, since the surface Ekman fluxes are compensated mostly above the level of no motion. Compared to the total maximum overturning streamfunction, the geostrophic approximation makes a scaling more complex, but it is the result of the force balance below the surface Ekman layer. The hemispheric differences in the level of no motion and the associated meridional transport suggest a hemispheric scaling rather than a single depth scale approximation for the entire basin.

450

The present manuscript relies on experiments that are conducted with a single but horizontally high-resolution model. Low-resolution models may differ significantly from high-resolution models. We put forward the idea that the ability of numerical models to capture the spatial and temporal variations of the level of no motion is crucial to reproduce the mid-depth cell in an appropriate way. Changes in the relationships between the level of no motion and vertical velocity shear change the AMOC
455 both quantitatively and dynamically.

Data availability. Data will be published via MPG.PuRe. We publish the data associated with the present study during the external review process. The codes associated with the present study are available upon reasonable request.

Author contributions. TR conducted the research and developed the coding scripts associated with the present study. TR developed the text of the present manuscript. JB, VL, DP, and JM reviewed the manuscript and checked the consistency of the research results.

460 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. We thank Oliver Gutjahr for the internal review on the present manuscript. This work was funded by the Max Planck Society (MPG) and the International Max Planck Research School on Earth System Modelling (IMPRS ESM). JB received funding under Germany's Excellence Strategy, EXC 2037 'Climate, Climatic Change and Society' CLICCS, Project Number: 390683824, as contribution to the Center for Earth System Research and Sustainability (CEN) of Universität Hamburg. DP received funding under PRIMAVERA, a
465 Horizon 2020 project funded by the European Commission, with grant number 641727. We also thank the German consortium project STORM for supporting the realization of high-resolution model simulations. We further thank Deutsches Klima Rechenzentrum (DKRZ) for providing the computational resources. Finally, we thank the anonymous reviewer.

References

- Allison, L. C., Johnson, H. L., and Marshall, D. P.: Spin-up and adjustment of the Antarctic circumpolar current and global pycnocline, *Journal of Marine Research*, 69, 167–189, <https://doi.org/10.1357/002224011798765330>, 2011.
- Baehr, J., Hirschi, J., Beismann, J. O., and Marotzke, J.: Monitoring the meridional overturning circulation in the North Atlantic: a model-based array design study, *Journal of Marine Research*, 62, 283–312, <https://doi.org/10.1357/0022240041446191>, 2004.
- Baehr, J., Stroup, A., and Marotzke, J.: Testing concepts for continuous monitoring of the meridional overturning circulation in the South Atlantic, *Ocean Modelling*, 29, 147–153, <https://doi.org/10.1016/j.ocemod.2009.03.005>, 2009.
- Bryan, F.: Parameter sensitivity of primitive-equation ocean general circulation models, *Journal of Physical Oceanography*, 17, 970–985, 1987.
- Cabanes, C., Lee, T., and Fu, L.-L.: Mechanisms of Interannual Variations of the Meridional Overturning Circulation of the North Atlantic Ocean, *Journal of Physical Oceanography*, 38, 467–480, <https://doi.org/10.1175/2007JPO3726.1>, 2008.
- Cessi, P.: The Effect of Northern Hemisphere Winds on the Meridional Overturning Circulation and Stratification, *Journal of Physical Oceanography*, 48, 2495–2506, <https://doi.org/10.1175/JPO-D-18-0085.1>, 2018.
- DeBoer, A. M., Gnanadesikan, A., Edwards, N. L., and Watson, A. J.: Meridional Density Gradients Do Not Control the Atlantic Overturning Circulation, *Journal of Oceanography*, 40, 368–380, <https://doi.org/10.1175/2009JPO4200.1>, 2010.
- Gent, P. R., Willebrand, J., McDougall, T. J., and McWilliams, J. C.: Parameterizing Eddy-Induced Tracer Transports in Ocean Circulation Models, *Journal of Physical Oceanography*, 25, 463–474, [https://doi.org/10.1175/1520-0485\(1995\)025<0463:PEITTI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1995)025<0463:PEITTI>2.0.CO;2), 1995.
- Gnanadesikan, A.: A Simple Predictive Model for the Structure of the Oceanic Pycnocline, *Science*, 283, 2077–2079, <https://doi.org/10.1126/science.283.5410.2077>, 1999.
- Gnanadesikan, A., DeBoer, A. M., and Mignone, B. K.: A Simple Theory of the Pycnocline and Overturning Revisited, *Ocean Circulation: Mechanisms and Impacts (Geophysical Monograph Series)*, 173, <https://doi.org/10.1029/173GM04>, 2007.
- Griesel, A. and Maqueda, M. A. M.: The relation of meridional pressure gradients to North Atlantic deep water volume transport in an ocean general circulation model, *Climate Dynamics*, 26, 781–799, <https://doi.org/10.1007/s00382-006-0122-z>, 2006.
- Hirschi, J. and Marotzke, J.: Reconstructing the meridional overturning circulation from boundary densities and the zonal wind stress, *Journal of Physical Oceanography*, 47, 743–763, <https://doi.org/10.1175/JPO3019.1>, 2007.
- Hirschi, J., Baehr, J., Marotzke, J., Stark, J., Cunningham, S., and Beismann, J. O.: A monitoring design for the Atlantic meridional overturning circulation, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2002GL016776>, 2003.
- Jayne, S. R. and Marotzke, J.: The dynamics of ocean heat transport variability, *Reviews of Geophysics*, 39, 385–411, <https://doi.org/10.1029/2000RG000084>, 2001.
- Johnson, H. L., Cessi, P., Marshall, D. P., Schloesser, F., and Spall, M. A.: Recent Contributions of Theory to Our Understanding of the Atlantic Meridional Overturning Circulation, *Journal of Geophysical Research*, 124, 5376–539, <https://doi.org/10.1029/2019JC015330>, 2019.
- Kalnay, E. et al.: The NCEP/NCAR 40-year reanalysis project, *Bulletin of the American Meteorological Society*, 77, 347–471, [https://doi.org/10.1175/1520-0477\(1996\)077<0437:TNYRP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2), 2018.
- Klinger, B. A. and Cruz, C.: Decadal Response of Global Circulation to Southern Ocean Zonal Wind Stress Perturbation, *Journal of Physical Oceanography*, 39, 1888–1904, <https://doi.org/10.1175/2009JPO4070.1>, 2009.

- 505 Klinger, B. A., Drijfhout, S., Marotzke, J., and Scott, J. R.: Sensitivity of Basinwide Meridional Overturning to Diapycnal Diffusion and Remote Wind Forcing in an Idealized Atlantic–Southern Ocean Geometry, *Journal of Physical Oceanography*, 33, 249–266, [https://doi.org/10.1175/1520-0485\(2003\)033<0249:SOBMOT>2.0.CO;2](https://doi.org/10.1175/1520-0485(2003)033<0249:SOBMOT>2.0.CO;2), 2003.
- Klinger, B. A., Drijfhout, S., Marotzke, J., and Scott, J. R.: Remote Wind-Driven Overturning in the Absence of the Drake Passage Effect, *Journal of Physical Oceanography*, 34, 1036–1049, [https://doi.org/10.1175/1520-0485\(2004\)034<1036:RWOITA>2.0.CO;2](https://doi.org/10.1175/1520-0485(2004)034<1036:RWOITA>2.0.CO;2), 2004.
- 510 Levermann, A. and Fuerst, J. J.: Atlantic pycnocline theory scrutinized using a coupled climate model, *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL044180>, 2010.
- Luyten, J. R., Pedlosky, J., and Stommel, H.: The ventilated Thermocline, *Journal of Physical Oceanography*, 13, 292–309, [https://doi.org/10.1175/1520-0485\(1983\)013<0292:TVT>2.0.CO;2](https://doi.org/10.1175/1520-0485(1983)013<0292:TVT>2.0.CO;2), 1983.
- Lüschow, V., von Storch, J.-S., and Marotzke, J.: Overturning response to a doubling of the surface wind stress in an eddying and a non-eddying ocean, *Journal of Physical Oceanography*, 51, 1007–1020, <https://doi.org/10.1175/JPO-D-20-0176.1>, 2021.
- 515 Marotzke, J.: Boundary Mixing and the Dynamics of Three-Dimensional Thermohaline Circulations, *Journal of Physical Oceanography*, 27, 1713–1728, [https://doi.org/10.1175/1520-0485\(1997\)027<1713:BMATDO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<1713:BMATDO>2.0.CO;2), 1997.
- Marotzke, J. and Klinger, B. A.: The Dynamics of Equatorially Asymmetric Thermohaline Circulations, *Journal of Physical Oceanography*, 30, 955–968, [https://doi.org/10.1175/1520-0485\(2000\)030<0955:TDOEAT>2.0.CO;2](https://doi.org/10.1175/1520-0485(2000)030<0955:TDOEAT>2.0.CO;2), 2000.
- Marshall, D. P. and Johnson, H. L.: Relative strength of the Antarctic Circumpolar Current and Atlantic Meridional Overturning Circulation, *Tellus A: Dynamic Meteorology and Oceanography*, 69, <https://doi.org/10.1080/16000870.2017.1338884>, 2017.
- 520 Marshall, J. and Speer, K.: Closure of the meridional overturning circulation through Southern Ocean upwelling, *Nature Geoscience*, 5, 171–180, <https://doi.org/10.1038/ngeo1391>, 2012.
- McCreary, J. P. and Lu, P.: Interaction between the Subtropical and Equatorial Ocean Circulations: The Subtropical Cell, *Journal of Physical Oceanography*, 24, 466–497, [https://doi.org/10.1175/1520-0485\(1994\)024<0466:IBTSAE>2.0.CO;2](https://doi.org/10.1175/1520-0485(1994)024<0466:IBTSAE>2.0.CO;2), 1994.
- 525 Moreno-Chamorro, E., Ortega, P., Gonzalez-Rouco, F., and Montoya, M.: Assessing reconstruction techniques of the Atlantic Ocean circulation variability during the last millennium, *Climate dynamics*, <https://doi.org/10.1007/s00382-016-3111-x>, 2016.
- Munk, W. and Wunsch, C.: Abyssal recipes II: energetics of tidal and wind mixing, *Deep Sea Research Part I: Oceanographic Research Papers*, 45, 1977–2010, [https://doi.org/10.1016/S0967-0637\(98\)00070-3](https://doi.org/10.1016/S0967-0637(98)00070-3), 1998.
- Robinson, A. and Stommel, H.: The Oceanic Thermocline and the Associated Thermohaline Circulation, *Tellus*, <https://doi.org/10.1111/j.2153-3490.1959.tb00035.x>, 1959.
- 530 Scott, J. R.: The Roles of Mixing, Geothermal Heating, and Surface Buoyancy Forcing in Ocean Meridional Overturning Dynamics, PhD thesis, Massachusetts Institute of Technology, 2000.
- Shakespeare, C. J. and Hogg, A. M.: An Analytical Model of the Response of the Meridional Overturning Circulation to Changes in Wind and Buoyancy Forcing, *Journal of Physical Oceanography*, 42, 1270–1287, <https://doi.org/10.1175/JPO-D-11-0198.1>, 2012.
- 535 Toggweiler, J. R. and Samuels, B.: Effect of Drake Passage on the global thermohaline circulation, *Deep Sea Research Part I: Oceanographic Research Papers*, 42, 477–500, [https://doi.org/10.1016/0967-0637\(95\)00012-U](https://doi.org/10.1016/0967-0637(95)00012-U), 1995.
- Tsujino, H. and Suginohara, N.: Thermohaline Circulation Enhanced by Wind Forcing, *Journal of Physical Oceanography*, 29, 1506–1516, [https://doi.org/10.1175/1520-0485\(2003\)033<0249:SOBMOT>2.0.CO;2](https://doi.org/10.1175/1520-0485(2003)033<0249:SOBMOT>2.0.CO;2), 1998.
- 540 Vallis, G. K.: Large-Scale Circulation and Production of Stratification: Effects of Wind, Geometry, and Diffusion, *Journal of Physical Oceanography*, 30, 933–953, [https://doi.org/10.1175/1520-0485\(1997\)027<1713:BMATDO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<1713:BMATDO>2.0.CO;2), 2000.

- von Storch, J.-S., Eden, C., Fast, I., Haak, H., Deckers, D., Maier-Reimer, E., Marotzke, J., and Stammer, D.: An Estimate of the Lorenz Energy Cycle for the World Ocean Based on the 1/10 STORM/NCEP Simulation, *Journal of Physical Oceanography*, 42, 2185–2205, <https://doi.org/10.1175/JPO-D-12-079.1>, 2012.
- Welander, P.: An advective model of the ocean thermocline, *Tellus*, <https://doi.org/10.1111/j.2153-3490.1959.tb00036.x>, 1959.
- 545 Williams, R. G. and Roussenov, V.: Decadal Evolution of Ocean Thermal Anomalies in the North Atlantic: The Effects of Ekman, Overturning, and Horizontal Transport, *Journal of Climate*, 27, 698–719, <https://doi.org/10.1175/JCLI-D-12-00234.1>, 2014.
- Wolfe, C. L. and Cessi, P.: The Adiabatic Pole-to-Pole Overturning Circulation, *Journal of Physical Oceanography*, 41, 1705–1810, <https://doi.org/10.1175/2011JPO4570.1>, 2011.