

1 A protocol for quantifying mono- and polysaccharides in 2 seawater and related saline matrices by electro-dialysis (ED) – 3 combined with HPAEC-PAD

4 Sebastian Zeppenfeld¹, Manuela van Pinxteren¹, Anja Engel², Hartmut Herrmann^{1,*}

5 1 Atmospheric Chemistry Department (ACD), Leibniz-Institute for Tropospheric Research (TROPOS),
6 Leipzig, Germany

7 2 GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

8 *Correspondence to: Hartmut Herrmann (herrmann@tropos.de)

9

10 Submitted to Ocean Science, ACP/OS special issue: 'Marine organic matter: from biological production in
11 the ocean to organic aerosol particles and marine clouds'

12 Abstract

13 An optimized method is presented to determine dissolved free (DFCHO) and dissolved combined
14 carbohydrates (DCCHO) in saline matrices, such as oceanic seawater, Arctic ice core samples or brine using
15 a combination of a desalination with electro-dialysis (ED) and high performance anion exchange
16 chromatography coupled to pulsed amperometric detection (HPAEC-PAD). Free neutral sugars, such as
17 glucose and galactose, were found with 95-98% recovery rates. Free amino sugars and free uronic acids
18 were strongly depleted during ED at pH=8, but an adjustment of the pH could result in higher recoveries
19 (58-59% for amino sugars at pH=11; 45-49% for uronic acids at pH=1.5). The applicability of this method
20 for the analysis of DCCHO was evaluated with standard solutions and seawater samples compared with
21 another established desalination method using membrane dialysis. DFCHO in field samples from different
22 regions on earth ranged between 11-118 nM and DCCHO between 260-1410 nM. This novel method has
23 the potential to contribute to a better understanding of biogeochemical processes in the oceans and sea-
24 air transfer processes of organic matter into the atmosphere in future studies.

25

26 Introduction

27 The majority of organic matter (OM) in oceanic seawater can be assigned to the chemical classes of
28 proteinogenic amino acids, carbohydrates and lipids (Benner and Kaiser, 2003; Kaiser and Benner, 2009;
29 Kuznetsova and Lee, 2002; Marty et al., 1979; Skoog et al., 1999; Wakeham et al., 1997). Previous studies
30 uncovered that combined amino acids are the most abundant organic substances in fresh particles sinking

31 within the water column (Wakeham et al., 1997), although more recent studies indicate that
32 carbohydrates may be equally abundant (Cisternas-Novoa et al., 2019), while hydrolysable carbohydrates
33 dominate the chemical composition of dissolved organic matter (DOM) (Kaiser and Benner, 2009;
34 McCarthy et al., 1996). Marine carbohydrates also appear in high concentrations in other related saline
35 matrices, such as ice cores, brine and melt ponds in the Arctic (Ewert and Deming, 2013; Underwood et
36 al., 2013; Zeppenfeld et al., 2019). Hence, a reliable analysis of carbohydrates is essential for understanding
37 biogeochemical processes in the ocean and their impact on Earth's atmosphere.

38 Most marine carbohydrates exist as polysaccharides or combined carbohydrates (CCHO), which are linear
39 or branched chains of monosaccharides, including hexoses, pentoses, deoxy sugars, amino sugars and
40 uronic acids. In aquatic environments, CCHO either appear in a particulate (PCCHO) or dissolved form
41 (DCCHO). In living marine microorganisms including prokaryotes, polysaccharides assume their functions
42 as structural compounds or as energy storage (Skoog and Benner, 1997). Storage carbohydrates mainly
43 consist of glucose, such as laminarans and other glucans, while structural heteropolysaccharides (e.g.
44 galactans) such as occurring in algal cell walls can contain a lot of galactose, mannose and rhamnose
45 (McCarthy et al., 1996). Furthermore, an elevated release of polysaccharides by phytoplankton, mostly of
46 gelatinous nature, has been associated with stress situations, such as a deficiency of nutrients, freezing or
47 fluctuating water potential (Berman-Frank et al., 2007; Bianchi and Canuel, 2011; Borchard and Engel,
48 2012, 2015; Ittekkot et al., 1981; Krembs et al., 2002; Krembs and Deming, 2008). These exuded
49 polysaccharides are relatively depleted in glucose and galactose and mainly contain acidic sugars, fucose,
50 rhamnose and arabinose in their chemical structure (Borchard and Engel, 2012; Passow, 2002). Even
51 though polysaccharides are ubiquitous in nature, a recent study revealed that the individual sugar pattern
52 is different between algae and terrestrial plants (Hepp et al., 2016) and may allow a source apportionment
53 of carbohydrates in seawater.

54 Dissolved free carbohydrates (DFCHO) have been found to form another fraction of marine carbohydrates
55 (Engel and Händel, 2011; Ittekkot et al., 1981; Kirchman et al., 2001). DFCHO are considered to be either
56 directly released by phytoplankton cells or be the product of enzymatic degradation of CCHO (Pakulski and
57 Benner, 1994). DFCHO are mostly found in lower concentrations than CCHO, since marine microbes utilize
58 them with high turnover rates (Engbrodt, 2001; Engel and Händel, 2011; Ittekkot et al., 1981; Thornton et
59 al., 2016) as it has been reported for amino acids analogously as well (Kuznetsova and Lee, 2002). From
60 the concentrations of DFCHO, or rather the ratio between CCHO and DFCHO, information about in situ
61 activities of local phytoplankton and bacteria in the seawater can be obtained (Pakulski and Benner, 1994;
62 Sakugawa and Handa, 1985). Recently, correlations between the concentrations of free glucose in Arctic

63 surface water samples and their ice nucleating activity (INA) suggested a potential link between the
64 formation of INA and marine carbohydrates (Zeppenfeld et al., 2019).

65 At the ocean surface, wind and wave interactions lead to bubble bursting. The emitted sea spray aerosol
66 contains marine carbohydrates, including hydrogels, which contribute to the chemical and physical
67 properties of these particles (Bigg and Leck, 2008; Frossard et al., 2014; Hawkins and Russell, 2010;
68 Rosenørn et al., 2006). They have been detected in particles at different marine environments on earth,
69 including the North Atlantic, the Arctic and Antarctica (Barbaro et al., 2015; Frossard et al., 2014; Gao et
70 al., 2011, 2012; Leck et al., 2013; Russell et al., 2010). However, understanding the quantitative fluxes of
71 marine carbohydrates from the ocean to the atmosphere is still challenging, since chemical analysis of
72 sugars in seawater strongly suffers from matrix effects, especially caused by sea salt.

73 The concentrations of individual monosaccharides in seawater, related saline matrices and aerosol
74 particles can be determined with different kinds of chromatographic methods, such as high performance
75 liquid chromatography and gas chromatography. These methods require a prior derivatization in order to
76 enable the chromatographic separation and detectability of these carbohydrates (Panagiotopoulos and
77 Sempéré, 2005). In the last decades, high performance anion exchange chromatography coupled to pulsed
78 amperometric detection (HPAEC-PAD) has been established as a reliable alternative, since it facilitates a
79 sensitive quantification of sugar compounds both in seawater and in airborne particles without a prior
80 derivatization (Iinuma et al., 2009; Panagiotopoulos and Sempéré, 2005; van Pinxteren et al., 2012; Skoog
81 and Benner, 1997). However, the high ionic content in seawater samples strongly affects the
82 chromatographic performance of the HPAEC-PAD and needs to be removed before analysis.

83 Several procedures are available for the desalination of seawater. The desalination using anion exchange
84 resins AG2-X8 and the cation exchange resin AG50W-X8 exhibits strong drawbacks such as the complete
85 loss of charged sugars (amino sugars, uronic acids) and quite low recovery rate of neutral sugars between
86 20-80% depending on the individual monosaccharide (Borch and Kirchman, 1997; Mopper et al., 1992;
87 Rich et al., 1996). The use of silver cartridges (Dionex OnGuard II Ag/H Cartridges) is faster and easier, but
88 requires very expensive consumables and the capacity of removable sea salt per cartridge is strongly
89 limited (Mopper et al., 1992; Panagiotopoulos and Sempéré, 2005). The desalination applying dialysis
90 membranes achieves reproducible and very high recovery rates of hydrolysable polysaccharides (> 90%).
91 However, this method does not allow the analysis of DFCHO, since these small molecules pass the
92 membrane during dialysis (Engel and Händel, 2011).

93

94 Electro-dialysis (ED) is a fast way to remove ions by applying an electrical field. The use of two different
95 chemo-selective ion exchange membranes allows the exclusive removal of small anions, or cations
96 respectively. Hence, uncharged small substances (neutral DFCHO) and macromolecules (CCHO) can, in
97 principle, be recovered in high quantities. Amongst others, ED is being used for the desalination of salty
98 water to generate potable water and the denitrification of wastewater and soil remediation (Gain et al.,
99 2002; Ottosen et al., 2000; Sadrzadeh and Mohammadi, 2008; Tsiakis and Papageorgiou, 2005; Wisniewski
100 et al., 2001). For analytical sample preparation, ED has been reported as a powerful desalination, e.g. for
101 the analysis of DOM and marine neutral DFCHO (Josefsson, 1970; Koprivnjak et al., 2009; Mopper et al.,
102 1980; Vetter et al., 2007; Wirth et al., 2019). However, following biases, which have hitherto not been
103 discussed in this analytical context, can occur during the application of ED and might falsify the determined
104 concentration of the analytes in the sample. In contact with ion exchange membranes, the passive
105 transport of water (*osmosis*) and solutes with a low molecular weight (*diffusion*), such as DFCHO, can occur
106 triggered by a concentration gradient between the sample and concentration channels (Galama et al.,
107 2014; Galier et al., 2012). Additionally, the active transport of charged molecules (*migration*) and water
108 bound to ions in their hydration sphere (*electro-osmosis*) takes place by operating an electrical field
109 (Galama et al., 2014; Han et al., 2015, 2017). While osmosis and electro-osmosis induce an unavoidable
110 loss of water and hence of the total volume of the sample, diffusion and migration of the analytes result
111 in a loss of analyzable molecules. Furthermore, water splitting and associated pH fluctuations have been
112 reported, when a limiting current is exceeded during an ED desalination (Cowan, 1962; Martí-Calatayud et
113 al., 2018; Ottosen et al., 2000; Vetter et al., 2007).

114
115 Within the present study, a novel protocol for the desalination of seawater samples and other saline
116 samples (e.g. ice cores and brine from Arctic sea ice), applying ED and HPAEC-PAD is presented, accounting
117 for the biases described above. With a low need of consumables, this method allows the analysis of
118 monosaccharides with (DCCHO) and without hydrolysis (DFCHO), including the possible determination of
119 free amino sugars and free uronic acids. This developed technique was applied to analyze a diverse set of
120 carbohydrates in different kinds of ambient seawater samples.

121

122

123

124 **2. Experimental**

125 2.1 Chemicals and materials

126 Prior to the analysis of carbohydrates in seawater, all used laboratory glassware had been washed with
127 ultrapure water (resistivity >18.2 MΩ·cm) thoroughly and pre-heated in a muffle furnace at 550 °C for 4 h.
128 All plastic equipment was first rinsed with 10% HCl solution for three times and then washed with ultrapure
129 water another three times. For calibrating the HPAEC-PAD and determining the recovery of individual
130 monosaccharides, a mixed stock solution was prepared from fucose (Roth, 95%), galactosamine (Sigma,
131 99%), rhamnose (Sigma, 99%), arabinose (Sigma, 99%), glucosamine (Fluka), galactose (Fluka, 99%),
132 glucose (99%), xylose (Fluka, 99%), mannose (Fluka,99%), fructose (Aldrich, 99%), ribose (Aldrich, 98%),
133 muramic acid (Sigma, 95%), galacturonic acid (Sigma-Aldrich, 97%), glucuronic acid (Sigma, 97%),
134 mannuronic acid (Sigma, 90%). Synthetic seawater samples were made from commercially available sea
135 salts (Sigma) achieving four solutions with salinities *S* of 10, 20, 30 and 40 on the practical salinity scale.
136 The salinity and the pH of water aliquots was measured by using a conductivity meter (pH/Cond 3320,
137 WTW).

138
139 **Table 1** Sampling details of discussed saline samples including seawater samples (SWS), ice cores (IC) and
140 brine (B). SML stands for surface microlayer (Engel et al., 2017). *(Wendisch et al., 2019)

| Saline sample (SS) | Location | Sampling date | Campaign | Latitude | Longitude | Depth (m) |
|-----------------------|-------------------|------------------|--------------|----------|-----------|--------------|
| SWS 1 | Tropical Atlantic | 13.11.2011 | Cape Verde | 16.935°N | 024.915°W | 0 (SML) |
| SWS 2 | Tropical Atlantic | 13.11.2011 | Cape Verde | 16.935°N | 024.915°W | 2 |
| SWS 3 | Raunefjorden | 16.05.2011 | Raunefjorden | 60.274°N | 005.181°E | 2 |
| SWS 4 | North Atlantic | 07.05.2012 | ANT-XXVIII-5 | 33.3°N | 013.5°W | 0 (SML) |
| SWS 5 | Arctic Ocean | 13.07.2017 | PS 106* | 81.229°N | 018.744°E | 1 |
| SWS 6 | Arctic Ocean | 14.07.2017 | PS 106* | 81.015°N | 026.883°E | 1 |
| SWS 7 | North Sea | 25.05.2017 | PS 106* | 57.288°N | 005.213°E | 1 |
| IC 1 | Arctic Ocean | 12.06.2017 | PS 106* | 81.824°N | 011.571°E | 0-0.8 |
| B 1 | Arctic Ocean | 12.06.2017 | PS 106* | 81.824°N | 011.571°E | 0-0.8 |
| B 2 | Arctic Ocean | 12.06.2017 | PS 106* | 81.824°N | 011.571°E | 0-1.5 |

141
142
143

144 2.2 Field samples

145 Seven different seawater samples, one ice core from Arctic sea ice and two brines collected within Arctic
146 ice cores (**Table 1**) were measured and used for evaluating recovery rates and comparability of the method
147 presented here. These saline samples were collected during different sampling campaigns and stored at -
148 20 °C. All sampling bottles had been rinsed with dilute hydrochloric acid (10% v/v) prior to the campaign.
149 Field blanks (ultra-pure water filled up in sampling bottles during the campaign) were collected during
150 each campaign and treated in the same way as the samples.

151

152 2.3 The ED system

153 The centerpiece of the PCell Micro Bench ED system for small sample volumes consisted of three
154 separated compartments (**Figure 1**): The sample compartment was an open chamber that was filled up
155 with 9 mL of the standard solution or seawater sample. The functionalized anion exchange membrane
156 (quaternary ammonium aliphatic polyether) and cation exchange membrane (sulfonated aromatic
157 polyether) bordered this compartment on both sides. Depending on their chemical properties, the
158 membranes allowed exclusively the migration of either positively or negatively charged ions. For that
159 matter, the anion exchange membrane bordering the sample chamber was oriented to the anode and the
160 cation exchange membrane to the cathode. The contact surface with the sample was 7.8 cm² for each
161 membrane. For maintaining the conductivity within the system and receiving the sea salt from the sample,
162 the next compartment contained the concentration circuit, a 16 g·L⁻¹ NaCl solution (Merck). This
163 concentration was chosen in order to minimize the osmotic water transfer as discussed below. This
164 solution was circulated at a rate of 60 mL·min⁻¹. Two end membranes on each side divided the
165 concentration circuit from the third compartment containing the electrodes. The end membranes were
166 cation exchange membranes with an increased chemical durability and an additional reinforcement in
167 order to withstand the strong differential pressure within the ED system. The mixed metal oxide (MMO)
168 anode was made of a titanium base body coated with RuO₂, IrO₂ and TiO₂. The MMO cathode was made
169 of stainless steel. The electrodes were permanently surrounded by a circulating 0.25 M Na₂SO₄ (Fluka)
170 electrolyte circuit for avoiding unwanted redox reactions (e.g. the generation of corrosive elemental
171 chlorine from chloride). Spacers were inserted between each membrane for keeping the electrolyte and
172 concentration circuits well mixed. The sample solution was homogenized during each desalination by
173 drawing some liquid into a Pasteur pipette and draining it immediately back to the sample compartment.
174 The electrolyte and the concentration solutions were renewed after every tenth desalination. The maximal
175 electrical current I_{max} within the ED cell was adjusted by an automatic online adaption of the voltage, which

176 never exceeded 25 V. The maximal current was set on 0.6 A in order to perform a fast desalination, but
177 also to avoid a scaling of the membranes due to water splitting and is discussed more in detail below. The
178 desalination was stopped when the electric current dropped to a value of 0.20 A.

179 The used ion-exchange membranes have a quite long lifetime as long as they are not damaged
180 mechanically. However, very high attention needs to be given to remove residues of previous desalinated
181 samples in order to avoid carry-over phenomena and obtain good reproducibility. Hence, every time
182 before a new sample was desalinated, the sample chamber was always first filled with ultrapure water for
183 ten minutes and then flushed once with an aliquot of the new sample, which was disposed after.




Figure 1 Schematic setup of the used ED cell. Red circles represent cations, green circles anions, and blue hexagons are carbohydrates. CEM and AEM stand for cation exchange membrane and anion exchange membrane respectively.

184

185 2.4 HPAEC-PAD system

186 HPAEC-PAD was applied for the analysis of marine carbohydrates in seawater samples. Here, we used an
187 Dionex ICS-3000 ion chromatography system coupled to an autosampler AS-1 as it has been already
188 described for the analysis of saccharidic biomass burning markers in atmospheric particles (Iinuma et al.,
189 2009). Several neutral monosaccharides, amino sugars and uronic acids were separated on a Dionex
190 CarboPac PA20 analytical column (3x150mm) combined with a Dionex CarboPac PA20 guard column
191 (3x30mm). The column oven temperature was maintained at 30 °C. The separation of these saccharides

192 was conducted by applying the gradient profile shown in **Table 2**, which was an adaption of Meyer et al.
 193 (2008). Neutral and amino sugars were eluted in 4 mM NaOH within the first 19 min. By adding sodium
 194 acetate, sugar acids eluted. At the same time, organic and inorganic contaminants were flushed from the
 195 column. After the removal of the remaining acetate by 250 mM NaOH, the system was equilibrated with
 196 4 mM NaOH for the next sample injection. The flow rate of the eluent was 0.5 mL·min⁻¹. The retention
 197 times, peak widths and resolution factors of the measured monosaccharides are shown in **Table 3**. For the
 198 injection of a sample aliquot, a 25 µL loop was used. Each sample was measured in duplicate and each
 199 standard in triplicate. Limits of detection (LOD) of individual monosaccharides ranged from 2-12 nM, which
 200 is in good agreement with reported data (Engel and Händel, 2011; Panagiotopoulos and Sempéré, 2005).

201
 202 For the preparation of eluents A-D, filtered ultra-pure water (resistivity >18.2 MΩ·cm) was degassed with
 203 helium for 20 min. Eluents A and B were made by adding a defined volume of low-carbonate NaOH solution
 204 (Fisher Chemical, 50% w/w) to the degassed water. Eluent C was prepared by dissolving sodium acetate
 205 (Thermo scientific, anhydrous) in ultra-pure water, filtering it through a nylon membrane (0.2 µm, Thermo
 206 Scientific), degassing the solution with helium for 20 min and adding the corresponding volume of NaOH
 207 solution.

208

209 **Table 2** Gradient profile applied on the CarboPac PA20 column during HPAEC-PAD analysis

| Time (min) | %A (250 mM NaOH) | %B (20 mM NaOH) | %C (1 M Na-acetate/ 250 mM NaOH) | %D (H ₂ O) |
|------------|---------------------|--------------------|--|--------------------------|
| 0 | 0 | 20 | 0 | 80 |
| 19 | 0 | 20 | 0 | 80 |
| 20 | 5 | 0 | 15 | 80 |
| 35 | 5 | 0 | 15 | 80 |
| 38 | 20 | 0 | 40 | 40 |
| 39 | 100 | 0 | 0 | 0 |
| 44 | 100 | 0 | 0 | 0 |
| 45 | 0 | 20 | 0 | 80 |
| 78 | 0 | 20 | 0 | 80 |

210

211 **Table 3** Averaged peak characteristics of monosaccharide standards separated by the presented elution averaged over 24 runs.
 212 Resolution factors are calculated between peak and following peak in chromatogram.

| | Retention time (min) | Peak width (min) | Resolution factor |
|-------------------|----------------------|------------------|-------------------|
| Fucose | 3.79±0.01 | 0.27 | 10.5 |
| Galactosamine | 7.28±0.01 | 0.40 | 1.0 |
| Rhamnose | 7.70±0.02 | 0.42 | 1.1 |
| Arabinose | 8.15±0.02 | 0.42 | 2.2 |
| Glucosamine | 9.23±0.02 | 0.56 | 2.1 |
| Galactose | 10.33±0.02 | 0.51 | 2.8 |
| Glucose | 11.81±0.03 | 0.56 | 3.4 |
| Xylose | 13.89±0.03 | 0.65 | 0.8 |
| Mannose | 14.43±0.04 | 0.77 | 2.8 |
| Fructose | 16.52±0.05 | 0.75 | 2.0 |
| Ribose | 18.20±0.05 | 0.93 | 13.2 |
| Muramic acid | 25.84±0.01 | 0.23 | 9.0 |
| Galacturonic acid | 29.27±0.02 | 0.53 | 4.6 |
| Glucuronic acid | 31.49±0.02 | 0.45 | 2.8 |
| Mannuronic acid | 32.83±0.02 | 0.49 | |

213

214 **2.5 Protocol for the analysis of DFCHO and DCCHO in seawater and other saline samples**

215 Stored frozen samples were defrosted in a fridge at 4 °C. 9 mL of the filtered sample (0.2 µm, Millex, PTFE)

216 was desalinated with ED as described above. At the end of each desalination, electro-(osmotic) water loss

217 was replenished with ultra-pure water and mixed thoroughly. In order to analyze free amino sugars or

218 uronic acids, pH could be adapted with concentrated HCl or 1 M NaOH.

219 A concentration step using a vacuum concentrator (MiVac) at 55 °C allowed the detection of expected

220 DFCHO concentrations in seawater. For this purpose, a round-bottom glass vial was filled with an aliquot

221 of 6 mL desalted sample, which was weighed empty and filled. After reaching a volume of approximately

222 600 µL, the glass vial was weighed again (in order to calculate the concentration factor) and the

223 concentrated aliquot was pipetted in the autosampler vial for HPAEC-PAD analysis. This step allowed a

224 decrease of LOD by a factor of 10. Each sample was prepared and measured in duplicate.

225 In order to measure DCCHO, marine polysaccharides need to be cleaved into their monomeric compounds
226 by acid hydrolysis. We applied the optimized conditions described by Engel and Händel (2011) with slight
227 modifications. An aliquot of 1 mL desalted sample was hydrolyzed with hydrochloric acid (HCl
228 concentration in sample= 0.8 M) in pre-heated (550 °C, 4 h) glass ampules for 20 h at 100 °C. Neutralization
229 was performed by evaporating all liquid under vacuum at 55 °C until dryness. The dry residue was dissolved
230 in 700 µL ultra-pure water, treated with a vortex homogenizer (IKA MS 3 basic) and filled in the
231 autosampler vial for HPAEC-PAD analysis. Each sample was prepared and measured in duplicate.

232

233 2.6 Parameter optimization and assessment of method

234 Impact of osmosis and electro-osmosis during ED desalination

235 For quantifying the loss of water in the sample due to osmosis and electro-osmosis, a synthetic sea salt
236 solution was pipetted into the desalination chamber, which was desalinated for 0, 5, 10, 15, 20 and 25 min
237 with a maximal voltage of 25 V and a maximal current of 0.6 A. After the lapse of time, the total remaining
238 volume was pipetted quantitatively into a glass vial and weighed (Mettler Toledo, XS105 DualRange).
239 These measurements were repeated in triplicate for four different sea salt solutions (with S of 10, 20, 30
240 and 40). The recovery of the sample mass was calculated as the ratio between the mass after the
241 corresponding desalination time and the averaged mass at 0 min.

242 Recovery of DFCHO within the ED membrane system

243 Standard addition experiments with seawater samples were performed, for quantifying the recovery of
244 monosaccharides due to diffusion and migration under consideration of all matrix effects. For that reason,
245 sample 7 was filtered (0.2 µm, Millex, PTFE) and spiked with a sugar standard mix (neutral sugars, amino
246 sugars, uronic acids) resulting in a concentration increase of 10 µg L⁻¹ and 100 µg L⁻¹. These samples were
247 desalinated using ED ($I_{max}=0.6$ A, stop at 0.2 A). At the end of each run, (electro-) osmotic water loss was
248 either replenished or not, and the sample directly measured with the HPAEC-PAD. These measurements
249 were repeated in triplicate for each concentration. In order to account for possible wasting phenomena,
250 repetitions were performed with new membranes and membranes which already had been used for some
251 time before. Given recovery rates for neutral monosaccharides are the average of the results for 10 µg
252 L⁻¹ and 100 µg L⁻¹. For sugar acids and amino sugars, only the averaged recovery rates for 100 µg L⁻¹ are
253 given for avoiding determinations close to the LOD.

254 In order to investigate the influence of pH on the migration of charged monosaccharides, this experiment
255 was repeated for three different pH values: At pH=8 (natural pH of seawater), pH of 1.5 (acidified with
256 concentrated HCl) and pH of 11 (addition of 1 M NaOH). Since high pH in seawater leads to precipitation
257 of hydroxides of alkaline earth metals, an additional filtering (0.2 μm) was performed for these runs.

258 Recovery of CCHO within the ED membrane system

259 Recovery experiments were performed with solutions and a suspension of the polysaccharide standards
260 sodium alginate (Aldrich), laminarin from Laminaria digitate (Sigma) and cellulose powder from spruce
261 (Fluka) at natural pH. Stock solutions were added to filtered sample SWS 7 resulting in concentrations of
262 10 mg L⁻¹. Aliquots of 1 mL with and without desalinations were hydrolyzed (HCl 0.8 M, 100 °C, 20 h) and
263 neutralized by evaporation of the volatile liquid with the vacuum concentrator (55 °C) until dryness. The
264 residue was reconstituted in 700 μL , treated with a vortex homogenizer (IKA MS 3 basic) and filled in the
265 autosampler vial for HPAEC-PAD analysis. Recovery rates were calculated as a ratio between the
266 determined monosaccharide concentrations after hydrolysis of the standard solutions with and without
267 desalination.

268 In order to compare our method on the recovery of DCCHO with another established method, aliquots of
269 four filtered (0.2 μm , Millex, PTFE) seawater samples were treated following the ED protocol presented
270 here and the protocol by Engel and Händel (2011) using membrane desalination, an acid hydrolysis with
271 HCl (0.8 M, 100 °C, 20h), neutralization by evaporation (nitrogen, 50 °C) and an elution on a Dionex
272 CarboPac PA10 column.

273

274 **3. RESULTS AND DISCUSSION**

275 A reproducible quantification of carbohydrates in seawater samples using HPAEC-PAD requires prior
276 removal of sea salt. Here, we present ED as a reliable desalination method, its parameter optimization and
277 the discussion of arising phenomena resulting in a protocol for the analysis of marine carbohydrates.

278

279 3.1 Kinetics and efficiency of desalination

280 During the desalination of seawater by ED, anions and cations migrate through an electrical field and pass
281 chemo-selective membranes. Depending on their electrical charge, they move either to the positively

282 charged anode or to the negatively charged cathode. In this process, the salt flux through the membranes
 283 j_s ($\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), which determines the desalination time, is proportional to the applied electrical current I
 284 (Han et al., 2017; Vanoppen et al., 2015). **Figure 2** shows the current within the used ED system and the
 285 salinity of the seawater sample during a typical desalination of an artificial seawater sample ($S=40$) for two
 286 different applied maximal currents I_{max} within the system. For almost the entire desalination run, the
 287 current I was maintained at I_{max} due to automatic adjustment of the voltage. During this time, the salt flux
 288 was approximately constant. Towards the end of the desalination, when almost all salt ions were removed,
 289 the current dropped down and the salt flux became lower. Since a direct salinity measurement was not
 290 possible in the sample chamber without contaminating the sample, the end of each desalination was
 291 defined, when the current I reached a value of 0.2 A. At this point, the salinity of the sample typically
 292 ranged between 0.2 and 0.4, which was found to be sufficiently low for the carbohydrate analysis at the
 293 HPAEC-PAD. This reduction in salinity represents an overall desalination of more than 99% of the initial
 294 salt concentration. A desalination reaching a salinity below 0.1 was possible, but was not necessary for this
 295 application and would have resulted in longer desalination times. Consequently, for minimizing the
 296 required desalination time, a high I_{max} is favorable.

297 However, it was observed that the application of an I_{max} of more than 0.8 A during the desalination,
 298 resulted in a large pH increase and a white precipitation in the (synthetic) seawater solution, apparently
 299 due to the formation of hydroxides of alkaline earth metals. This uncontrolled precipitation strongly
 300 disturbed the efficiency of the desalination and the reproducibility of the carbohydrate measurements and




Figure 2 Measured current within the ED system and salinity of a synthetic seawater sample ($S=40$) versus desalination time with two different maximal applied currents I_{max} (0.6 A solid line, 0.3 A dashed line). The red line represents the current and the corresponding salinity, when desalination was stopped.

301 caused a scaling of the membranes. Previous studies explained these unfavorable changes of pH by a
302 strong concentration polarization at the membranes surface leading to water splitting to H^+ and OH^- ions,
303 when a certain limiting current is exceeded. This phenomenon has been preferably observed at anion
304 exchange membranes with quaternary amino groups in the presence of divalent cations, such as Mg^{2+} and
305 Ca^{2+} (Cowan, 1962; Martí-Calatayud et al., 2018; Ottosen et al., 2000). The precipitation exclusively
306 occurred when (synthetic) seawater was desalinated and not during the desalination of NaCl standard
307 solutions. This finding shows the importance of performing parameter optimization tests with synthetic
308 seawater standards that include all important seawater constituents such as divalent cations. In summary,
309 the optimum maximal current I_{max} of 0.6 A was found for the used ED system for avoiding scaling effects
310 and performing desalination as fast as possible.

311

312 3.2 Possible biases during the application of ED

313 Osmotic and electro-osmotic transport of water

314 Osmosis describes the passive transport of free water molecules through a partially permeable membrane
315 caused by large differences of the osmotic pressures between the concentrate circuit and the sample
316 solution (Sata, 2007). The direction and the quantity of the water transport depends on the residence time
317 t_R of the sample solution within the membrane system, the difference between the concentrations of
318 solutes in the sample solution and the concentration circuit ($c_s - c_c$) and membrane specific parameters,
319 such as the osmotic water transfer coefficient the membrane area and the membrane thickness (Galama
320 et al., 2014). The quantitative effect of osmosis can be reduced by minimizing t_R and $(c_s - c_c)$. Hence, c_c was
321 set at 16 g NaCl L^{-1} , which is approximately in the middle between the concentrations of a typical seawater
322 sample before (salinity=30-39) and after the desalination (salinity=0.2-0.4) for balancing the positive and
323 negative contribution of osmosis on the total sample volume during a typical desalination.

324 In aqueous solutions, water molecules form a hydration shell around ions (Ohtaki and Radnai, 1993).
325 Whenever ions pass through membranes during ED, a cotransport of these hydrating water molecules
326 occurs, known as electro-osmosis (Galama et al., 2014). The electroosmotic water transfer j_W ($m^3 \cdot m^{-2} \cdot s^{-1}$)
327 is proportional to the salt flux j_s in the system and can be expressed by formula (1) with the molar volume

328 of water V_M ($1.8 \cdot 10^{-5} \cdot \text{m}^3 \cdot \text{mol}^{-1}$) and the salt hydration number n_H (mol water \cdot mol $^{-1}$ salt) (Galier and
 329 Balmann, 2015; Han et al., 2015).

$$330 \quad j_W = n_H \cdot V_M \cdot j_S \quad (1)$$

331 The salt hydration number of NaCl, as the major compound of sea salt, has been reported with values
 332 between 11 and 14 (Han et al., 2015; Rutgers and Hendriks, 1962; Singlande et al., 2006; Walker et al.,
 333 2014). Assuming a NaCl concentration of 30 g \cdot L $^{-1}$ and n_H to be 14, a maximal reduction of the sample
 334 volume by 13 % due to electro-osmosis was expected, additionally to osmosis.

335 The recovery of the sample volume due to electro-osmosis and osmosis during the desalination was
 336 characterized for four different salinities for the used ED system (**Figure 3**). During the active removal of
 337 sea salt, electro-osmosis is the dominating force causing the water loss in the sample. The electroosmotic
 338 water loss is continual as long as the salt flux stays constant. However, in the final stages of each
 339 desalination, the salt flux decreases and consequently the electroosmotic water transfer decreases, too.
 340 For a synthetic seawater sample with a salinity of 30, 84% of sample mass was recovered. This is in good




Figure 3 Combined effect of electro-osmosis and osmosis (solid lines) and osmosis (dashed lines) on the recovery of sample mass as a function of the desalination time within the described membrane system ($I_{max}=0.6$ A, $c_c=16$ g NaCl L $^{-1}$) for artificial sea salt solutions with four different initial salinities.

341 agreement with the estimation mentioned above considering a small additional contemporaneous
342 contribution of osmosis. Once the sea salt is removed, osmotic water transfer remained at a constant rate
343 of approximately $0.1\% \cdot \text{min}^{-1}$.

344 The overall water loss resulting from osmosis and electro-osmosis needs to be taken into account since it
345 falsifies the determined concentrations of marine carbohydrates. For its compensation, the chamber was
346 replenished with ultra-pure water at the end of each desalination until the initial sample volume was
347 reached. This procedure was performed with a maximal overall error of 0.5%.

348

349 Analysis and recovery of DFCHO in seawater samples

350 The recovery of neutral monosaccharides during ED is impacted by diffusion and convection processes
351 (Galier and Balmann, 2015). Additionally, free amino sugars and uronic acids migrate through an electrical
352 field due to their charge and pass the ion exchange membranes. Recovery tests were performed with
353 standard solutions spiked to a seawater samples with concentrations, which have been typically reported
354 for seawater samples before (Kirchman et al., 2001; Mopper et al., 1980; Skoog et al., 1999; Zeppenfeld et
355 al., 2019). Recovery rates of neutral sugars (Glc, Man, Xyl, Gal, Ara, Fuc, Rha, Fru) ranged between 95-98%
356 at the natural pH of seawater (approx. pH=8) (**Table 4**). Hence, the overall impact of diffusion and
357 convection on the recovery of monosaccharides seems to be quite low for the short contact time with
358 these membranes. However, a higher loss of neutral monosaccharides due to diffusion was observed,
359 when the sample solution remained within the membrane system for a longer period of time, which calls
360 for a fast desalination. This is in agreement with Vanoppen et al. (2015) who concluded that diffusion and
361 affinity for the membrane are the main drivers for losses of uncharged, low-molecular organics during a
362 ED desalination. An overestimation of the determined concentrations was avoided by performing a
363 correction of the water loss in the end of each desalination. Charged monosaccharides were found with
364 much lower recoveries of 25-31% for uronic acids and 16-19% for amino sugars at pH=8. This is due to
365 their weak acidic/basic properties (pK_a (amino sugars) = 7.6-8.5 (Bichsel and von Gunten, 2000; Sinnott,
366 2007), pK_a (uronic acids) = 3.3-3.5 (Kohn and Kovác, 1978)) and hence their partially ionic state, which
367 makes them migrate through the electrical field. However, a low pH can protonate the carboxylic group of
368 uronic acids and a high pH deprotonates the amino group of amino sugars for reducing this effect. Here,
369 we found that an initial pH of 1.5 before desalination could increase the recovery of free uronic acids up
370 to 45-49%, while a high pH of 11 resulted into a higher recovery of free amino sugars up to 58-59%. The
371 recovery of neutral sugars seemed to be quite unaffected within the range of the tested pH, with the

372 exception of fructose, which was 89% recovered at pH 1.5, certainly due to its instability within acid
 373 conditions.

374 **Table 4** Recovery of individual free monosaccharides (neutral, amino sugars, uronic acids) after desalination with ED including
 375 correction of (electro-) osmotic water loss. n.d.= not determined

| Monosaccharide | Recovery rate (%) | | |
|-----------------|---------------------------|------------------------------------|--------------------------|
| | pH _{Start} = 1.5 | pH _{Start} = 8 (seawater) | pH _{Start} = 11 |
| Galactose | 97.6±1.0 | 96.0±0.9 | 97.2±1.4 |
| Fucose | 97.2±2.4 | 97.9±1.6 | 97.0±1.0 |
| Glucose | 96.9±1.5 | 97.2±1.4 | 97.1±2.1 |
| Mannose | 95.8±1.8 | 95.8±1.3 | 95.8±2.1 |
| Xylose | 95.6±2.3 | 95.9±2.2 | 95.2±2.0 |
| Rhamnose | 93.5±1.6 | 94.2±1.6 | 95.1±1.4 |
| Arabinose | 93.4±2.3 | 95.1±1.1 | 94.5±1.9 |
| Fructose | 89.3±3.4 | 94.6±2.7 | 94.1 ±2.2 |
| Glucuronic acid | 48.7±2.7 | 31±1.9 | n.d. |
| Mannuronic acid | 44.9±0.6 | 25±1.9 | n.d. |
| Muramic acid | 24.7±1.7 | n.d. | n.d. |
| Glucosamine | 1.5±0.4 | 18.9±1.2 | 59±3.2 |
| Galactosamine | 1.2±0.3 | 15.9±1.5 | 58±3.4 |

376
 377 Analysis and recovery of CCHO in seawater samples with standard polysaccharides
 378 Recovery experiments with standard solutions of common polysaccharides were performed with and
 379 without desalination by ED. The neutral, water-soluble polysaccharide laminarin was recovered with
 380 91.0±5.4%. The acidic polysaccharide alginic acid was recovered with 93.2±5.3%. Even though alginic acid
 381 might move within the electrical field due to its acidic molecular structure, its molecular weight does not
 382 allow passing the membrane and it does not leave the sample solution. Standard desalination experiments
 383 with a suspension of the water insoluble cellulose as a representative of PCCCHO resulted in much lower
 384 recoveries of 48±19%. The reason for this high, less reproducible loss of polysaccharides was likely caused
 385 by sedimentation within the sample chamber. Engel and Händel (2011) described adsorption processes
 386 during the desalination with dialysis membranes and tackled this problem with sonification of the
 387 membranes. However, sonification could not be applied in our apparatus. In this study, flushing the
 388 chamber several times with a defined volume of ultra-pure water after desalination and reuniting the

389 washing water with the desalinated sample could increase the yield of cellulose up to $85.2\pm 6.9\%$ under
 390 consideration of dilution factors. This procedure was not found to be feasible, since a dilution of a natural
 391 sample reduces the sensitivity of low concentrated sugars in seawater in the analysis by HPAEC-PAD.
 392 Rather, we recommend ED only for the application to filtered samples (DCCHO), while PCCHO might be
 393 better analyzed from filters after filtration.

394

395 Comparison of ED and dialysis method for the determination of DCCHO in seawater samples

396 In order to evaluate the presented procedure for the analysis of DCCHO, comparison studies were
 397 performed measuring four ambient seawater samples (SWS 1-4) with the established membrane dialysis
 398 protocol after Engel and Händel (2011) and with the method presented here. **Figure 4** shows a comparison
 399 of the results of the individual monosaccharides after hydrolysis with HCl. The slope of the regression line
 400 using all sugar data together with the determination coefficient ($R^2=0.89$) indicate a good overall
 401 agreement between both methods. Major concentrated sugars, such as glucose, galactose and
 402 xylose/mannose, were determined at similar concentrations. Furthermore, a good agreement was
 403 observed for minor concentrated sugars, such as fucose and galactosamine. Discrepancies were found for
 404 rhamnose and arabinose, which was overestimated by the ED method, and glucosamine, which was
 405 determined at lower concentrations. These variations might be explained by statistical uncertainties or co-




Figure 4 Determined monosaccharide concentrations in four seawater samples (SWS 1-4) after hydrolysis (DCCHO) comparing the desalination by dialysis (Engel&Händel, 2011) and ED (presented in this study).The full line represents the line of equality. The dotted line represents the regression line between the data of both methods.

406 elution of unknown substances. In summary, the here presented method using ED has shown to be in good
407 agreement with the established membrane dialysis method regarding the analysis of DCCHO. In addition,
408 the ED offers the major advantage of analysing the full spectrum of DFCHO as well - which comprise a
409 group of hardly investigated but potentially important marine compounds.

410

411 3.3 Chromatographic performance with HPAEC PAD after desalination.

412 Several kinds of saline samples were desalinated with ED and analyzed on the CarboPac PA20 column.
413 **Figure 5** shows some examples for DFCHO and DCCHO chromatograms in a brine and a seawater sample
414 after desalination with ED. The insufficient chromatographic separation of mannose and xylose using
415 previous kinds of analytical columns has been frequently described in literature (Borch and Kirchman,
416 1997; Engbrodt, 2001; Engel and Händel, 2011; Kirchman et al., 2001). Therefore, xylose and mannose
417 have been reported as sum concentrations frequently. The elution of the sugars on a CarboPac20 column,
418 applied in the present study, strongly improved the separation between the both sugars mannose and
419 xylose (resolution factor=0.8), and allowed the individual determination of these two sugars. However,
420 most of the analyzed samples showed high concentrations of xylose, which strongly overlapped the
421 smaller peak of mannose. For these cases, we kept reporting a sum value for Xyl/Man.

422

423 3.4 DFCHO and DCCHO in saline field samples from different regions.

424 Several samples were analyzed on DFCHO and DCCHO (**Table 5** and **6**). In both sugar fractions, glucose was
425 the most abundant monosaccharide, as it has been reported before (Panagiotopoulos and Sempéré,
426 2005). In some of the samples, free fructose could be determined reaching concentrations comparable to
427 glucose. However, fructose cannot be determined in DCCHO, since hydrolysis leads to complete
428 destruction of this sugar. High DFCHO was found in the samples from the Arctic including brine and ice
429 core samples reaching up to 118 nM in comparison to seawater samples from the Atlantic SWS 1-4 (11-
430 15 nM). However, a clear regional trend could not be identified for DCCHO with concentrations, ranging
431 between 260-1410 nM. Traces of free amino sugars and uronic acids were found after neutral
432 desalinations. However, a stronger enrichment is required in order to determine them quantitatively and
433 will be the focus of further studies.

434

435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457




Figure 5 a) Full chromatogram of standard solution $100 \mu\text{g}\cdot\text{L}^{-1}$; b-e) chromatograms of neutral sugars and amino sugars of b) standard solution $100 \mu\text{g}\cdot\text{L}^{-1}$; c) DFCHO in brine (B2) desalinated at natural pH; d) DCCHO in a seawater sample (SWS 3); e) DCCHO in Arctic brine (B2). 1 fructose, 2 galactosamine, 3 rhamnose, 4 arabinose, 5 glucosamine, 6 galactose, 7 glucose, 8 xylose, 9 mannose, 10 fructose, 11 ribose, 12 muramic acid, 13 galacturonic acid, 14 glucuronic acid, 15 mannuronic acid.

458
 459
 460

461 **Table 5** Mol percentages of individual neutral monosaccharides within DFCHO in seawater, Arctic brine and ice core samples.
 462 <LOD stands for below detection limit.

| | Glc mol% | Gal mol% | Xyl/Man mol% | Rha mol% | Fuc mol% | Ara mol% | Fru mol% | Total DFCHO nM |
|-------|-------------|-------------|-----------------|-------------|-------------|-------------|-------------|-------------------|
| SWS 1 | 57 | 8 | <LOD | <LOD | 7 | 8 | 20 | 15 |
| SWS 2 | 52 | <LOD | <LOD | <LOD | <LOD | <LOD | 48 | 14 |
| SWS 3 | 80 | <LOD | 15 | <LOD | <LOD | 5 | <LOD | 11 |
| SWS 4 | 87 | <LOD | 5 | <LOD | <LOD | 8 | <LOD | 14 |
| SWS 5 | 89 | 3 | 3 | <LOD | <LOD | 4 | <LOD | 35 |
| SWS 6 | 16 | 8 | 22 | <LOD | 34 | 20 | <LOD | 27 |
| IC 1 | 50 | 6 | 2 | 3 | 15 | <LOD | 25 | 118 |
| B 1 | 23 | 27 | 13 | 8 | 29 | <LOD | <LOD | 15 |
| B 2 | 85 | 2 | 3 | 2 | 6 | 1 | <LOD | 53 |

463

464

465 **Table 6** Mol percentages of individual monosaccharides within DCCHO in seawater, Arctic brine and ice core samples. <LOD
 466 stands for below detection limit.

| | Glc mol% | Gal mol% | Xyl/Man mol% | Rha mol% | Fuc mol% | Ara mol% | GalN mol% | GluN mol% | Gal-ac mol% | Gluc-ac mol% | Total DCCHO nM |
|-------|-------------|-------------|-----------------|-------------|-------------|-------------|--------------|--------------|----------------|-----------------|-------------------|
| SWS 1 | 20 | 15 | 22 | 7 | 7 | 8 | 3 | 4 | 12 | 3 | 680 |
| SWS 2 | 25 | 12 | 24 | 8 | 7 | 7 | 2 | 3 | 11 | 1 | 290 |
| SWS 3 | 20 | 14 | 21 | 9 | 10 | 9 | 3 | 6 | 7 | 2 | 580 |
| SWS 4 | 31 | 12 | 26 | 5 | 6 | 7 | 2 | 3 | 6 | 2 | 410 |
| SWS 5 | 84 | 3 | 2 | 2 | 2 | 2 | 1 | 1 | <LOD | 3 | 1410 |
| SWS 6 | 48 | 7 | 10 | 4 | 2 | <LOD | 2 | 4 | 15 | 7 | 260 |
| IC 1 | 54 | 9 | 11 | 1 | <LOD | 1 | 2 | 1 | 10 | 10 | 330 |
| B 1 | 47 | 11 | 13 | 4 | 4 | 3 | 3 | 5 | <LOD | 10 | 420 |
| B 2 | 65 | 8 | 10 | 3 | 4 | 1 | 2 | 3 | <LOD | 3 | 640 |

467

468 5. Summary and conclusion

469 In this study, a novel protocol was presented for the analysis of both DFCHO and DCCHO in saline aqueous
 470 samples by applying HPLC-PAD with prior desalination by ED. Recovery rates for neutral
 471 monosaccharides ranged between 95-98%. By adjusting pH, charged monosaccharides such as free amino
 472 sugars and uronic acids could be recovered with 58-59% at pH = 11 and 45-49% at pH = 1.5, respectively.
 473 Dissolved polysaccharide standards, such as laminarin and alginic acid showed good recovery rates of 91-
 474 93%, while a suspension of insoluble cellulose was quite difficult to recover reproducibly. Hence, ED for
 475 carbohydrate analysis is recommended to be used for filtered samples or for samples with low amount of
 476 particulate matter. In this study, the osmotic and electro-osmotic loss of water was considered in order to

477 avoid an overestimation of the determined concentrations. In seawater from different locations, Arctic
478 brine and sea ice core samples, DCCHO was found in concentrations between 260 and 1410 nM. DFCHO
479 ranged in much lower concentrations of 11-118 nM. Within both, DFCHO and DCCHO, the most dominant
480 monosaccharide was glucose, followed by other neutral sugars.

481 In this study, the successful application of ED in combination with HPAEC-PAD for the analysis of marine
482 carbohydrates (both free and combined) in marine matrices, such as seawater, ice cores and brine could
483 be demonstrated. The application of ED for other more salt sensitive analyses should be the focus of
484 further research, e.g. the reported interference of suspended sea spray aerosol in Arctic snow samples
485 during the quantification of insoluble light absorbing impurities such as black carbon and dust performed
486 via nebulization. Hence, this developed method has the potential to contribute strongly in further research
487 studies understanding biogeochemical processes in the oceans and related saline matrices and sea-air
488 exchange processes, especially for studying hot spot regions of climate change, such as the Arctic.

489

490 **Acknowledgements**

491
492 We gratefully acknowledge the funding by the Deutsche Forschungsgemeinschaft (DFG, German Research
493 Foundation) – Project-ID 268020496 – TRR 172, within the Transregional Collaborative Research Center
494 “ArctiC Amplification: Climate Relevant Atmospheric and SurfaCe Processes, and Feedback Mechanisms
495 (AC)³” in sub-projects B04. Additional support through the Leibniz Association SAW funding of the project
496 “Marine biological production, organic aerosol particles and marine clouds: a Process Chain
497 (MarParCloud)”, (SAW-2016-TROPOS-2) is also gratefully acknowledged. We thank for the opportunities
498 to use aqueous samples from various sampling campaigns in order to develop the method presented here.
499 We thank Swantje Pöge for supporting work in the laboratory. We thank Jon Roa for DCCHO analysis
500 following the Engel & Händel (2011) protocol. We thank Susanne Fuchs, Kristin Recklies and Christian
501 Weller of Thermo Fisher in his in-house collaborations for fruitful discussions. We thank Patrick Altmeier
502 and Philip Kalkhoff from PCell GmbH (Heusweiler, Germany) for guidance and technical support for
503 helping to put ED to operation in our laboratory.

504

505 *Data availability.* All data are available on the public repository PANGAEA;
506 <https://doi.pangaea.de/10.1594/PANGAEA.910575>.

507 *Author contributions.* SZ wrote the manuscript with contributions from MvP, HH and AE. SZ and MvP
508 collected seawater samples during different field campaigns. SZ optimized the presented method and
509 performed the chemical measurements. AE performed DCCHO analysis with supplied samples following
510 her published protocol for evaluation purposes. All co-authors proofread and commented the manuscript.

511

512 *Competing interest.* The authors declare that they have no conflict of interest.

513

514 **References**

515 Barbaro, E., Kirchgeorg, T., Zangrando, R., Vecchiato, M., Piazza, R., Barbante, C. and Gambaro, A.: Sugars
516 in Antarctic aerosol, *Atmospheric Environment*, 118, 135–144, doi:10.1016/j.atmosenv.2015.07.047,
517 2015.

518 Benner, R. and Kaiser, K.: Abundance of amino sugars and peptidoglycan in marine particulate and
519 dissolved organic matter, *Limnology and Oceanography*, 48(1), 118–128, doi:10.4319/lo.2003.48.1.0118,
520 2003.

521 Berman-Frank, I., Rosenberg, G., Levitan, O., Haramaty, L. and Mari, X.: Coupling between autocatalytic
522 cell death and transparent exopolymeric particle production in the marine cyanobacterium
523 *Trichodesmium*, *Environmental Microbiology*, 9(6), 1415–1422, doi:10.1111/j.1462-2920.2007.01257.x,
524 2007.

525 Bianchi, T. S. and Canuel, E. A.: *Chemical biomarkers in aquatic ecosystems*, Princeton University Press.,
526 2011.

527 Bichsel, Y. and von Gunten, U.: Formation of Iodo-Trihalomethanes during Disinfection and Oxidation of
528 Iodide-Containing Waters, *Environ. Sci. Technol.*, 34(13), 2784–2791, doi:10.1021/es9914590, 2000.

529 Bigg, E. K. and Leck, C.: The composition of fragments of bubbles bursting at the ocean surface, *Journal of*
530 *Geophysical Research: Atmospheres*, 113(D11), doi:10.1029/2007JD009078, 2008.

531 Borch, N. H. and Kirchman, D. L.: Concentration and composition of dissolved combined neutral sugars
532 (polysaccharides) in seawater determined by HPLC-PAD, *Marine Chemistry*, 57(1), 85–95,
533 doi:10.1016/S0304-4203(97)00002-9, 1997.

534 Borchard, C. and Engel, A.: Organic matter exudation by *Emiliana huxleyi* under simulated future ocean
535 conditions, *Biogeosciences*, 9(8), 3405–3423, doi:10.5194/bg-9-3405-2012, 2012.

536 Borchard, C. and Engel, A.: Size-fractionated dissolved primary production and carbohydrate composition
537 of the coccolithophore *Emiliana huxleyi*, *Biogeosciences*, 12(4), 1271–1284, doi:10.5194/bg-12-1271-
538 2015, 2015.

539 Cisternas-Novoa, C., Le Moigne, F. A. C. and Engel, A.: Composition and vertical flux of particulate organic
540 matter to the oxygen minimum zone of the central Baltic Sea: impact of a sporadic North Sea inflow,
541 *Biogeosciences*, 16(4), 927–947, doi:10.5194/bg-16-927-2019, 2019.

- 542 Cowan, D. A.: Research in the problem of scaling of electro dialysis demineralizers, Texas Water Comm.
543 Bull., 6206, 29, 1962.
- 544 Engbrodt, R.: Biogeochemistry of dissolved carbohydrates in the Arctic, Berichte zur Polar-und
545 Meeresforschung (Reports on Polar and Marine Research), 396, 106pp, 2001.
- 546 Engel, A. and Händel, N.: A novel protocol for determining the concentration and composition of sugars
547 in particulate and in high molecular weight dissolved organic matter (HMW-DOM) in seawater, Marine
548 Chemistry, 127(1), 180–191, doi:10.1016/j.marchem.2011.09.004, 2011.
- 549 Engel, A., Bange, H. W., Cunliffe, M., Burrows, S. M., Friedrichs, G., Galgani, L., Herrmann, H., Hertkorn,
550 N., Johnson, M., Liss, P. S., Quinn, P. K., Schartau, M., Soloviev, A., Stolle, C., Upstill-Goddard, R. C., van
551 Pinxteren, M. and Zäncker, B.: The Ocean's Vital Skin: Toward an Integrated Understanding of the Sea
552 Surface Microlayer, Front. Mar. Sci., 4, 165, doi:10.3389/fmars.2017.00165, 2017.
- 553 Ewert, M. and Deming, J. W.: Sea Ice Microorganisms: Environmental Constraints and Extracellular
554 Responses, Biology, 2(2), 603–628, doi:10.3390/biology2020603, 2013.
- 555 Frossard, A. A., Russell, L. M., Burrows, S. M., Elliott, S. M., Bates, T. S. and Quinn, P. K.: Sources and
556 composition of submicron organic mass in marine aerosol particles, Journal of Geophysical Research:
557 Atmospheres, 119(22), 12,977–13,003, doi:10.1002/2014JD021913, 2014.
- 558 Gain, E., Laborie, S., Viers, P., Rakib, M., Durand, G. and Hartmann, D.: Ammonium nitrate wastewater
559 treatment by coupled membrane electrolysis and electro dialysis, Journal of Applied Electrochemistry,
560 32(9), 969–975, 2002.
- 561 Galama, A. H., Saakes, M., Bruning, H., Rijnaarts, H. H. M. and Post, J. W.: Seawater predesalination with
562 electro dialysis, Desalination, 342, 61–69, doi:10.1016/j.desal.2013.07.012, 2014.
- 563 Galier, S. and Balmann, H. R.: Demineralization of glucose solutions by electro dialysis: Influence of the
564 ionic composition on the mass transfer and process performances, The Canadian Journal of Chemical
565 Engineering, 93(2), 378–385, doi:10.1002/cjce.22076, 2015.
- 566 Galier, S., Courtin, M. and Balmann, H. R.: Influence of the Ionic Composition on the Demineralisation of
567 Saccharide Solutions by Electro dialysis, Procedia Engineering, 44, 826–829,
568 doi:10.1016/j.proeng.2012.08.587, 2012.
- 569 Gao, Q., Nilsson, U., Ilag, L. L. and Leck, C.: Monosaccharide compositional analysis of marine
570 polysaccharides by hydrophilic interaction liquid chromatography-tandem mass spectrometry, Anal
571 Bioanal Chem, 399(7), 2517–2529, doi:10.1007/s00216-010-4638-z, 2011.
- 572 Gao, Q., Leck, C., Rauschenberg, C. and Matrai, P. A.: On the chemical dynamics of extracellular
573 polysaccharides in the high Arctic surface microlayer, Ocean Science, 8(4), 401–418, 2012.
- 574 Han, L., Galier, S. and Roux-De Balmann, H.: Ion hydration number and electro osmosis during
575 electro dialysis of mixed salt solution, Desalination, 373, 38–46, doi:10.1016/j.desal.2015.06.023, 2015.
- 576 Han, L., Galier, S. and Roux-De Balmann, H.: A phenomenological model to evaluate the performances of
577 electro dialysis for the desalination of saline water containing organic solutes, Desalination, 422, 17–24,
578 doi:10.1016/j.desal.2017.08.008, 2017.

579 Hawkins, L. N. and Russell, L. M.: Polysaccharides, Proteins, and Phytoplankton Fragments: Four
580 Chemically Distinct Types of Marine Primary Organic Aerosol Classified by Single Particle
581 Spectromicroscopy, *Advances in Meteorology*, 2010, 1-14, doi:10.1155/2010/612132, 2010.

582 Hepp, J., Rabus, M., Anhäuser, T., Bromm, T., Laforsch, C., Sirocko, F., Glaser, B. and Zech, M.: A sugar
583 biomarker proxy for assessing terrestrial versus aquatic sedimentary input, *Organic Geochemistry*, 98,
584 98–104, doi:10.1016/j.orggeochem.2016.05.012, 2016.

585 Iinuma, Y., Engling, G., Puxbaum, H. and Herrmann, H.: A highly resolved anion-exchange
586 chromatographic method for determination of saccharidic tracers for biomass combustion and primary
587 bio-particles in atmospheric aerosol, *Atmospheric Environment*, 43(6), 1367–1371,
588 doi:10.1016/j.atmosenv.2008.11.020, 2009.

589 Ittekkot, V., Brockmann, U., Michaelis, W. and Degens, E. T.: Dissolved free and combined carbohydrates
590 during a phytoplankton bloom in the northern North Sea, *Marine Ecology Progress Series*, 4, 299–305,
591 1981.

592 Josefsson, B. O.: Determination of soluble carbohydrates in sea water by partition chromatography after
593 desalting by ion-exchange membrane electro dialysis, *Analytica Chimica Acta*, 52(1), 65–73,
594 doi:10.1016/S0003-2670(01)80042-8, 1970.

595 Kaiser, K. and Benner, R.: Biochemical composition and size distribution of organic matter at the Pacific
596 and Atlantic time-series stations, *Marine Chemistry*, 113(1–2), 63–77,
597 doi:10.1016/j.marchem.2008.12.004, 2009.

598 Kirchman, D. L., Meon, B., Ducklow, H. W., Carlson, C. A., Hansell, D. A. and Steward, G. F.: Glucose fluxes
599 and concentrations of dissolved combined neutral sugars (polysaccharides) in the Ross Sea and Polar
600 Front Zone, Antarctica, *Deep Sea Research Part II: Topical Studies in Oceanography*, 48(19–20), 4179–
601 4197, doi:10.1016/S0967-0645(01)00085-6, 2001.

602 Kohn, R. and Kováč, P.: Dissociation constants of D-galacturonic and D-glucuronic acid and their O-methyl
603 derivatives, *Chem. zvesti*, 32(4), 478–485, 1978.

604 Koprivnjak, J.-F., Pfromm, P. H., Ingall, E., Vetter, T. A., Schmitt-Kopplin, P., Hertkorn, N., Frommberger,
605 M., Knicker, H. and Perdue, E. M.: Chemical and spectroscopic characterization of marine dissolved
606 organic matter isolated using coupled reverse osmosis–electrodialysis, *Geochimica et Cosmochimica*
607 *Acta*, 73(14), 4215–4231, doi:10.1016/j.gca.2009.04.010, 2009.

608 Krembs, C. and Deming, J. W.: The role of exopolymers in microbial adaptation to sea ice, in
609 *Psychrophiles: from biodiversity to biotechnology*, pp. 247–264, Springer., 2008.

610 Krembs, C., Eicken, H., Junge, K. and Deming, J. W.: High concentrations of exopolymeric substances in
611 Arctic winter sea ice: implications for the polar ocean carbon cycle and cryoprotection of diatoms, *Deep*
612 *Sea Research Part I: Oceanographic Research Papers*, 49(12), 2163–2181, doi:10.1016/S0967-
613 0637(02)00122-X, 2002.

614 Kuznetsova, M. and Lee, C.: Dissolved free and combined amino acids in nearshore seawater, sea surface
615 microlayers and foams: Influence of extracellular hydrolysis, *Aquatic sciences*, 64(3), 252–268, 2002.

616 Leck, C., Gao, Q., Mashayekhy Rad, F. and Nilsson, U.: Size-resolved atmospheric particulate
617 polysaccharides in the high summer Arctic, *Atmospheric Chemistry and Physics*, 13(24), 12573–12588,
618 doi:<https://doi.org/10.5194/acp-13-12573-2013>, 2013.

619 Martí-Calatayud, M. C., García-Gabaldón, M. and Pérez-Herranz, V.: Mass Transfer Phenomena during
620 Electrodialysis of Multivalent Ions: Chemical Equilibria and Overlimiting Currents, *Applied Sciences*, 8(9),
621 1566, doi:[10.3390/app8091566](https://doi.org/10.3390/app8091566), 2018.

622 Marty, J. C., Saliot, A., Buat-Ménard, P., Chesselet, R. and Hunter, K. A.: Relationship between the lipid
623 compositions of marine aerosols, the sea surface microlayer, and subsurface water, *J. Geophys. Res.*,
624 84(C9), 5707, doi:[10.1029/JC084iC09p05707](https://doi.org/10.1029/JC084iC09p05707), 1979.

625 McCarthy, M., Hedges, J. and Benner, R.: Major biochemical composition of dissolved high molecular
626 weight organic matter in seawater, *Marine Chemistry*, 55(3), 281–297, doi:[10.1016/S0304-4203\(96\)00041-2](https://doi.org/10.1016/S0304-4203(96)00041-2), 1996.

628 Meyer, A., Fischer, H., Kuzyakov, Y. and Fischer, K.: Improved RP-HPLC and anion-exchange
629 chromatography methods for the determination of amino acids and carbohydrates in soil solutions, *J.*
630 *Plant Nutr. Soil Sci.*, 171(6), 917–926, doi:[10.1002/jpln.200700235](https://doi.org/10.1002/jpln.200700235), 2008.

631 Mopper, K., Dawson, R., Liebezeit, G. and Ittekkot, V.: The monosaccharide spectra of natural waters,
632 *Marine Chemistry*, 10(1), 55–66, doi:[10.1016/0304-4203\(80\)90058-4](https://doi.org/10.1016/0304-4203(80)90058-4), 1980.

633 Mopper, K., Schultz, C. A., Chevolut, L., Germain, C., Revuelta, R. and Dawson, R.: Determination of
634 sugars in unconcentrated seawater and other natural waters by liquid chromatography and pulsed
635 amperometric detection., *Environ. Sci. Technol.*, 26(1), 133–138, doi:[10.1021/es00025a014](https://doi.org/10.1021/es00025a014), 1992.

636 Ohtaki, H. and Radnai, T.: Structure and dynamics of hydrated ions, *Chem. Rev.*, 93(3), 1157–1204,
637 doi:[10.1021/cr00019a014](https://doi.org/10.1021/cr00019a014), 1993.

638 Ottosen, L. M., Hansen, H. K. and Hansen, C. B.: Water splitting at ion-exchange membranes and
639 potential differences in soil during electro-dialytic soil remediation, *Journal of Applied Electrochemistry*,
640 30(11), 1199–1207, doi:[10.1023/A:1026557830268](https://doi.org/10.1023/A:1026557830268), 2000.

641 Pakulski, J. D. and Benner, R.: Abundance and distribution of carbohydrates in the ocean, *Limnology and*
642 *Oceanography*, 39(4), 930–940, 1994.

643 Panagiotopoulos, C. and Sempéré, R.: Analytical methods for the determination of sugars in marine
644 samples: A historical perspective and future directions, *Limnology and Oceanography: Methods*, 3(10),
645 419–454, doi:[10.4319/lom.2005.3.419](https://doi.org/10.4319/lom.2005.3.419), 2005.

646 Passow, U.: Transparent exopolymer particles (TEP) in aquatic environments, *Progress in Oceanography*,
647 55(3), 287–333, doi:[10.1016/S0079-6611\(02\)00138-6](https://doi.org/10.1016/S0079-6611(02)00138-6), 2002.

648 van Pinxteren, M., Müller, C., Iinuma, Y., Stolle, C. and Herrmann, H.: Chemical Characterization of
649 Dissolved Organic Compounds from Coastal Sea Surface Microlayers (Baltic Sea, Germany), *Environ. Sci.*
650 *Technol.*, 46(19), 10455–10462, doi:[10.1021/es204492b](https://doi.org/10.1021/es204492b), 2012.

- 651 Rich, J. H., Ducklow, H. W. and Kirchman, D. L.: Concentrations and uptake of neutral monosaccharides
652 along 14°W in the equatorial Pacific: Contribution of glucose to heterotrophic bacterial activity and the
653 DOM flux, *Limnology and Oceanography*, 41(4), 595–604, doi:10.4319/lo.1996.41.4.0595, 1996.
- 654 Rosenørn, T., Kiss, G. and Bilde, M.: Cloud droplet activation of saccharides and levoglucosan particles,
655 *Atmospheric Environment*, 40(10), 1794–1802, doi:10.1016/j.atmosenv.2005.11.024, 2006.
- 656 Russell, L. M., Hawkins, L. N., Frossard, A. A., Quinn, P. K. and Bates, T. S.: Carbohydrate-like composition
657 of submicron atmospheric particles and their production from ocean bubble bursting, *Proc. Natl. Acad.
658 Sci. U.S.A.*, 107(15), 6652–6657, doi:10.1073/pnas.0908905107, 2010.
- 659 Rutgers, A. J. and Hendriks, Y.: IONIC HYDRATION, *Transactions of the Faraday Society (England)*
660 Superseded by *J. Chem. Soc., Faraday Trans., I and II*, Vol: 58, doi:10.1039/TF9625802184, 1962.
- 661 Sadrzadeh, M. and Mohammadi, T.: Sea water desalination using electrodialysis, *Desalination*, 221(1–3),
662 440–447, 2008.
- 663 Sakugawa, H. and Handa, N.: Chemical studies on dissolved carbohydrates in the water samples collected
664 from the North Pacific and Bering Sea, *Oceanologica acta*, 8(2), 185–196, 1985.
- 665 Sata, T.: *Ion Exchange Membranes: Preparation, Characterization, Modification and Application*, Royal
666 Society of Chemistry, 2007.
- 667 Singlande, E., Roux-de Balman, H., Lefevbre, X. and Sperandio, M.: Improvement of the treatment of
668 salted liquid waste by integrated electrodialysis upstream biological treatment, *Desalination*, 199(1), 64–
669 66, doi:10.1016/j.desal.2006.03.020, 2006.
- 670 Sinnott, M.: *Carbohydrate Chemistry and Biochemistry: Structure and Mechanism*, Royal Society of
671 Chemistry., 2007.
- 672 Skoog, A. and Benner, R.: Aldoses in various size fractions of marine organic matter: Implications for
673 carbon cycling, *Limnology and Oceanography*, 42(8), 1803–1813, doi:10.4319/lo.1997.42.8.1803, 1997.
- 674 Skoog, A., Biddanda, B. and Benner, R.: Bacterial utilization of dissolved glucose in the upper water
675 column of the Gulf of Mexico, *Limnology and Oceanography*, 44(7), 1625–1633,
676 doi:10.4319/lo.1999.44.7.1625, 1999.
- 677 Thornton, D. C. O., Brooks, S. D. and Chen, J.: Protein and Carbohydrate Exopolymer Particles in the Sea
678 Surface Microlayer (SML), *Front. Mar. Sci.*, 3, 135–143, doi:10.3389/fmars.2016.00135, 2016.
- 679 Tsiakis, P. and Papageorgiou, L. G.: Optimal design of an electrodialysis brackish water desalination plant,
680 *Desalination*, 173(2), 173–186, 2005.
- 681 Underwood, G. J. C., Aslam, S. N., Michel, C., Niemi, A., Norman, L., Meiners, K. M., Laybourn-Parry, J.,
682 Paterson, H. and Thomas, D. N.: Broad-scale predictability of carbohydrates and exopolymers in Antarctic
683 and Arctic sea ice, *PNAS*, 110(39), 15734–15739, doi:10.1073/pnas.1302870110, 2013.
- 684 Vanoppen, M., Bakelants, A. F. A. M., Gaublumme, D., Schoutteten, K. V. K. M., Bussche, J. V.,
685 Vanhaecke, L. and Verliefde, A. R. D.: Properties Governing the Transport of Trace Organic Contaminants
686 through Ion-Exchange Membranes, *Environ. Sci. Technol.*, 49(1), 489–497, doi:10.1021/es504389q, 2015.

- 687 Vetter, T. A., Perdue, E. M., Ingall, E., Koprivnjak, J.-F. and Pfromm, P. H.: Combining reverse osmosis and
688 electro dialysis for more complete recovery of dissolved organic matter from seawater, *Separation and*
689 *Purification Technology*, 56(3), 383–387, doi:10.1016/j.seppur.2007.04.012, 2007.
- 690 Wakeham, S. G., Lee, C., Hedges, J. I., Hernes, P. J. and Peterson, M. J.: Molecular indicators of diagenetic
691 status in marine organic matter, *Geochimica et Cosmochimica Acta*, 61, 5363–5369, doi:10.1016/S0016-
692 7037(97)00312-8, 1997.
- 693 Walker, W. S., Kim, Y. and Lawler, D. F.: Treatment of model inland brackish groundwater reverse
694 osmosis concentrate with electro dialysis — Part II: Sensitivity to voltage application and membranes,
695 *Desalination*, 345, 128–135, doi:10.1016/j.desal.2014.04.026, 2014.
- 696 Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Dethloff, K., Velasco, C. B.,
697 Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer, U., Engelmann,
698 R., Engler, C., Eppers, O., Gehrman, M., Gong, X., Gottschalk, M., Gourbeyre, C., Griesche, H., Hartmann,
699 J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel,
700 E., Järvinen, E., Jourdan, O., Kästner, U., Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L.,
701 Leroy, D., Maturilli, M., Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T.,
702 Notholt, J., Palm, M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder,
703 K., Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P., Shupe, M. D., Siebert, H., Spreen, G., Stapf,
704 J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M. and Zeppenfeld, S.: The Arctic
705 Cloud Puzzle: Using ALOUD/PASCAL Multiplatform Observations to Unravel the Role of Clouds and
706 Aerosol Particles in Arctic Amplification, *Bull. Amer. Meteor. Soc.*, 100(5), 841–871, doi:10.1175/BAMS-
707 D-18-0072.1, 2019.
- 708 Wirth, M. A., Sievers, M., Habedank, F., Kragl, U., Schulz-Bull, D. E. and Kanwischer, M.: Electro dialysis as
709 a sample processing tool for bulk organic matter and target pollutant analysis of seawater, *Marine*
710 *Chemistry*, 217, 103719, doi:10.1016/j.marchem.2019.103719, 2019.
- 711 Wisniewski, C., Persin, F., Cherif, T., Sandeaux, R., Grasmick, A. and Gavach, C.: Denitrification of drinking
712 water by the association of an electro dialysis process and a membrane bioreactor: feasibility and
713 application, *Desalination*, 139(1–3), 199–205, 2001.
- 714 Zeppenfeld, S., van Pinxteren, M., Hartmann, M., Bracher, A., Stratmann, F. and Herrmann, H.: Glucose
715 as a Potential Chemical Marker for Ice Nucleating Activity in Arctic Seawater and Melt Pond Samples,
716 *Environ. Sci. Technol.*, 53(15), 8747–8756, doi:10.1021/acs.est.9b01469, 2019.

717

718