

Reviewer 1

This paper presents a comparison of eddy fields in two models: a ROMS configuration and a FESOM configuration. The models seem to agree in general but differ some in the details. The analysis focuses more on eddy statistics and there is little dynamical analysis. The main conclusion is that the two model configurations simulate roughly similar eddy statistics.

I recommend major revision because I think the paper would be much better if it demonstrated a connection between the dynamical analysis (regions of baroclinic or barotropic instability) and the eddy statistics (e.g. generation regions of long-lived eddies). The individual pieces for analyzing this connection mostly exist so this seems reasonable to expect. Otherwise, the paper is well written and easy to understand. Some of the diagnostics used (e.g. T, S anomalies) were not described clearly —these need some major improvement (see below). The figures were nice to look at and illustrate the main points well though some extra panels would be useful (see below). Most of my other comments are minor.

We would like to thank you for the constructive comments, which significantly improved the manuscript. A major comment is about the connection between eddy statistics and dynamics. However, there is no 1:1 correspondence between (a) sites of eddy generation through local instability, (b) the energy conversion as a result of the presence and interaction of an eddy field with the background mean state and (c) the statistics of the diagnosed eddy trajectories (only considering coherent vortices, but not all the filaments and eddy genesis in-between). In the revised version, we discuss the connection between these points (Section 6.1). A deeper analysis, e.g. computing unstable growth rates as done by Isachsen (2015, <https://doi.org/10.1002/2014JC010448>), would be out of the scope of this paper.

As suggested, we improved the description of the diagnostics. Below are detailed answers to your comments.

1. deformation radius \approx 4x grid spacing. Is this "eddy resolving" or "eddy permitting"? Some discussion with references to literature would be useful background here.

We added this sentence in the introduction:

"Hallberg (2013) showed that in ocean models, a resolution of two grid points per Rossby radius of deformation can be considered as threshold between "non-eddying" and "eddy-permitting" regimes, and thus higher resolution is needed for a model to be considered as "eddy-resolving"."

2. Line 90: Does Figure 1 show the domain for the ROMS simulation? Is this the region with refined resolution for the FESOM run?

As you also suggested in 19., we added a second panel in Figure 1 showing a map of the Arctic Ocean. The red box indicates the area (76°N-82°N, 10°W-20°E) used for eddy detection. This region comprises the domain of the ROMS simulation (except for very small parts in the north-west and south-west corners, which can be seen in Fig. 2a). It is also the region with refined resolution in FESOM.

3. Line 100: Please list the 4 constraints.

We now listed the 4 constraints in section 2.3.

4. Line 100: Do a and b have units? What is the minimum possible detected eddy size in km? How does this compare to grid spacing?

We added a better description of a and b in section 2.3:

“Parameter a defines over how many grid points the increases in magnitude of v along the EW axes and u along the NS axes are checked, and its unit is grid points. It also defines the minimum size of detectable eddies, which is a-1 grid points. Parameter b defines the size (also in grid points) of the area used to find the local minimum of velocity.”

5. Line 167: What was the longest gap you had to interpolate over?

For the WSC time series, the longest gap was 14 days. For the EGC time series, we used the time period Sep 8 2006 - Dec 31 2009 to avoid a long gap in the beginning of 2006. We added this description in the text.

6. Line 162: Did you average the spectra from the 3 moorings?

Yes, we averaged the velocity components u and v over the moorings, and then computed the spectrum from this time series. We added this clarification.

7. Since the eddies in FESOM are weaker (see spectra), should you adjust your criteria to be more appropriate for these weaker eddies? In other words, are you undercounting eddies in FESOM because they are weaker than the thresholds you are using?

We added a new panel in Figure 5 showing a spatial map of eddy kinetic energy (EKE) in both models. As also visible from the seasonal cycle of EKE, FESOM shows a higher

energy level in the WSC than ROMS, and in contrast, the energy level in the EGC is higher in ROMS than in FESOM. This is also reflected in the spectra.

Even if eddies were weaker in FESOM, they are defined by a circular velocity structure which is detected by the algorithm. Thus, the Nencioli method does not depend on the strength of the velocity field, but on the geometry of the flow field (closed contours). Both models have the same resolution (interpolated to the same regular grid), so there is no risk of undercounting.

8. Figure 6: How does this figure compare to a map of EKE averaged over the 3 years?

1. Is the sign or caption wrong in Figure 6c, d? It seems to say there are more anticyclones on the shelves contrary to line 189. 2. Also are the topographic contours the same as in previous figures? They are hard to see and it is hard to identify the 1000m isobath referred to in the text.

We added maps of EKE averaged over 2006-2009 to Figure 5. The comparison of the EKE maps and eddy occurrences is discussed now in section 6.1. In fact, Figure 6c,d shows that there are more cyclones on the shelves than anti-cyclones (red color, where bottom topography is shallower than 200 m). We changed the topographic contours and increased the linewidth.

9. Section 4.3: This "eddy intensity" is really a Rossby number (relative vorticity / f). Can you call it that?

Yes, you are right. We take the Rossby number as an index for the eddy intensity. We added the description in section 4.3. This was also done by Kang et al 2014 (<https://doi.org/10.1002/jgrc.20318>) in their study of Gulf Stream eddies.

10. Section 4.5: These eddy pathways should be interpreted with the help of mean velocity vectors. Jet instabilities generate cyclones on the anticyclonic side of the jet and anticyclones on the cyclonic side of the jet. For the Svalbard Branch for example, one would see anticyclones on the deeper side but cyclones in shallower side which would partially explain the asymmetry in trajectories, i.e. cyclones and anticyclones are being generated in different places. I agree that the long term tendency for anticyclones to cluster in Boreas basin is likely a topographic effect.

Thank you for this explanation. We included an extended version of explanation that resorts to the fact that there is a front between different water masses: "Along the Svalbard coast, the Svalbard Coastal Current transports cold and fresh waters northward, close to the salty and warm AW which is carried northward by the WSC a little offshore. The meandering between the two water masses, light water on the eastern side and

denser water on the western side (roughly indicated by the 200 m isobath) leads to the generation of cyclones on the eastern side and anti-cyclones on the western (offshore) side.”

The situation is comparable to the Gulf Stream, where dense (cold) water is located on the northern side of the main path, and light (warm) water is located south of the main path. This leads to the generation of cyclones on the southern side, and anticyclones on the northern side of the main path. We added this description in section 4.5.

11. Line 255: How do you define the vertical extent?

For every eddy center location (of the eddy detected in 100 m depth), we compute relative vorticity, temperature and salinity in the whole water column at that location, i.e. in every model layer. We added this description in the text: “...by calculating relative vorticity/ f at the location of the eddy centres...”

12. Line 255: I also didn't understand how you define T, S anomalies. Are these T, S at all detected eddy center locations minus a monthly mean at the eddy center? Is this a climatological monthly mean or a monthly mean for a particular year the eddy was tracked? This definition should be clearly written.

Right, we compute T and S at the detected eddy center locations, and then subtract the monthly mean value at this location of the particular month. Description added in section 4.6.

13. Figure 8,9: Why is there a large difference in number of cyclone tracks between ROMS and FESOM in "FS Central South"

Indeed, more cyclones are generated in FESOM in "FS Central South" than in ROMS (269 and 107 generated cyclones in FESOM and ROMS, respectively, during the years 2006-2009). Not only cyclones, but also anticyclones are more abundant in FESOM than in ROMS in this region. This is consistent with the fact that baroclinic energy conversion in this region is stronger in FESOM (Fig. 12).

14. Lines 285-290: There are other regions with large conversions (e.g. just north and south of 77N) that are stable in Figure 12b.

· Would this disappear if you calculated the barotropic PV gradient properly accounting for u also?

We computed the barotropic PV gradient by also taking into account u . In particular, we defined segments of length 0.25° along the 1000 m isobath and approximated the along-stream velocity as the velocity tangential to the segments. The across-stream derivative

was computed in the normal direction of the segments. However, the difference to the term d^2v/dx^2 is only minor, and does not reveal visible difference.

Nonetheless, there are small regions where the barotropic PV gradient changes sign near 77°N , which are maybe hard to see in Figure 12b. We modified the colormap slightly, and hope that the figure is more clear now.

- Did you use depth-averaged and monthly mean v ? or monthly mean v at 100m?

We used depth-averaged monthly mean v velocity. We added this description in section 5.1.

- Why did you only use FESOM for Figure 12b?

Since the energy transfer terms in Figure 11a,b are very similar, we do not expect significant differences. For simplicity and also practically to reduce the workload of coauthors on the ROMS side, we decided to show it only for FESOM.

15. "EPE" — this should really be APE (available potential energy) since eddies are deriving their KE from the APE of the /mean/ state (Vallis 2006 textbook).

Corrected.

16. Line 315: This is a nice demonstration that the flow is baroclinically unstable but is it unstable in the regions necessary to explain the eddy tracks in Figures 8,9? Some connection between the instability analysis and the eddy tracks is needed.

Eddy tracks are determined by both eddy generation and eddy movement. The latter as depicted in Figures 8,9 makes it hard to link the tracks to eddy generation associated with instability. Instead, we can compare the number of detected eddies (Figure 6a,b) with the maps of EKE (Figure 5) and the energy conversion rates (Figure 12). We added a subsection (Section 6.1) where we discuss this connection:

"Although baroclinic instability is the main driver of mesoscale eddy variability, the connection between eddy occurrences and EKE as well as the APE to EKE conversion rate ($w'b'$) is very non-local. For one thing, the eddies form as the result of nonlinear evolution of baroclinic instability waves and jet meanders. For another, mean circulation transports and modifies all eddy-like features, moving them away from the sites where their 'seeds' originally appeared. As a result, the observed pattern of eddy occurrences (Figure 6a,b) differ from the EKE and $w'b'$ distributions (Figures 5 and 12)...."

17. Lines 345, 346: I'm not sure there was any discussion of "EKE→MKE conversion associated with steepening isopycnals". Please describe this in more detail. The "energy backscatter" parameterizations being currently developed would be useful literature to reference (e.g.Jansen et al 2015)

We reformulated the last paragraph in the discussion, and hope to explain it better now. Thanks for pointing us to the study by Jansen et al. We also cite the papers by Juricke et al, which describe the implementation of the method by Jansen et al. into FESOM and its application to realistic test cases.

18. Discuss sensitivity of eddy maps to choice of 100m depth

We added a justification of the choice of the depth of 100 m:

“We decided to choose the depth of 100 m because both main water masses of the Fram Strait, AW and PW, are present at this depth (e.g. Wekerle et al., 2017, their Figure 9). In addition, the maximum relative vorticity is also located close to this depth for different regions of the Fram Strait (Figure 10).”

19. Figure 1: Showing a second panel with a larger region would be nice for readers not familiar with this place.

Thanks for this suggestion. We added a second panel in Figure 1 showing the Arctic Ocean and our study region.

20. Figure 4: It is more conventional to show logarithm axis scales instead of showing $\log_{10}(\text{quantity})$ on a linear scale. Why did you choose to not do that? Please mark 1/30 days since that's the frequency you chose to separate "mean" and "eddies"

As you suggested, we changed Figure 4 and now show logarithm axis scales. We also marked the frequency at 1/30 days.

Reviewer 2

The present paper reports results from a sensitivity study on the ability of two different numerical circulation models (ROMS and FESOM) in reproducing comparable eddy dynamic properties in terms of generation sites and propagation pathway in Fram Strait, a region characterized by a complex bathymetric configuration. Main difference between these models lies on their numerics and formulations, which includes numerical grid discretization, horizontal and vertical mesh resolution, parameterizations, and coverage (i.e. global vs. regional models).

The study is generally well written, and the analysis performed are appropriate. The figures are neat and clean. The discussion of the results is superficial. The authors should inspect more thoroughly the results presented. Nevertheless, the study stands as a good

contribution, as it adds knowledge to the region. However, unfortunately it doesn't add new knowledge to the subject field whether in ocean modelling or in geophysical fluid dynamics. If it does, then perhaps it has not been clearly presented.

Find below few/minor points which the authors should address before its acceptance for publication.

We would like to thank you for your helpful comments. As you suggested, we improved the Discussion section. In particular, we added subsections discussing the connection between eddy occurrences and EKE (Section 6.1), the differences between models and observations (Section 6.2), and implications for contributing to future model development (Section 6.3). Below, we addressed your comments point by point.

The reference for ROMS in the introduction section may not be the most appropriate here!

In the introduction section, we added references to the papers by Shchepetkin and McWilliams (2005) and Budgell (2005).

Line 105 the sentence seems incomplete. In part it reads as "we also the Okubo-Weiss". Do you mean "we also used the Okubo-Weiss"?

Corrected.

Are there any criteria on the choice of the number of days taken as threshold to discard eddies with lifetime lesser than 3 days? Please note that the caption in Figure 10 and in other parts of the manuscript indicate a lifetime of 30 days. Is there any inconsistency??

We added a justification of the choice of the 3 days in section 2.3:

"We decided to use a threshold of 3 days mainly because the temporal resolution of the model output data is daily, and the eddy should form a track. This also helps to make sure that the eddies detected are real and not an over-detection due to uncertainties in the detection method. Eddies with a lifetime of at least three days are also required when computing the translation velocity needed to compute the eddy nonlinearity parameter, for which centred differences are used."

We analyzed all detected eddies (with lifetime > 2 days) and discussed them in section 4.1, 4.2, 4.3 and 4.4. However, in sections 4.5 and 4.6 we decided to focus on long-lived eddies only. Analysis of pathways only makes sense for long-lived eddies. Otherwise the displacements are too short to be informative. In both sections, we made it clear that we only analyse "eddies with lifetime of more than 30 days".

Paragraph 130. What do you mean by eddy detected by experts?? Please make it clear.

Chaigneau et al. (2008) compared two methods of eddy detection, the 'winding-angle' method and Okubo-Weiss method. To validate their results, they asked oceanographic experts to detect eddies from sea level anomaly maps. Compared to the eddies detected by experts, the Okubo-Weiss method over-detected eddies. To avoid confusion, we just removed this sentence in the new version.

Did you use any filtering on the field of Okubo-Weiss parameter?

No, we do not use any filtering to compute the Okubo-Weiss parameter.

Perhaps the differences between the model maps shown in Figures 2 and 3 should be quantified to highlight geographical/spatial sites where the models converge and where they diverge (e.g. $r = \text{ROMS} - \text{FESOM}$ for the parameters presented in Fig2 and Fig3).

Since the thermo-haline properties are not the main focus of this paper, we decided not to show the difference between ROMS and FESOM in T and S. Instead, we added contours of the 1°C and 2°C isotherms and the 34 and 35 isohalines, so that simulated T and S in both models can be more easily compared. Note that Figure 2 shows snapshots in time, so it would not make sense to plot the difference.

Under the Model assessment section, paragraph 155, the authors indicate the model simulates similar spatial distribution of the water masses. However, no T/S diagrams have been shown. Perhaps replacing the term "water-masses" with "thermo-haline properties" would be more appropriate. The same is true wherever this term appears in this manuscript.

As suggested, we added T/S diagrams for ROMS and FESOM in Figure 3. The differences in the simulated thermohaline properties are better illustrated with it. Model differences are described now in more detail in section 3. We also exchanged the term "water-masses" with the term "thermo-haline properties" where it is appropriate.

Lines 170 – 173, is the difference only related to the effects of tides simulated in ROMS? Could not be also somehow related to differences in the surface forcing fields between the models?

This is correct. The surface atmospheric forcing as well plays a role. We added this in Section 3.

Lines 180 – 184: Are the eddy statistics computed in a Lagrangian or Eulerian frame of reference? Please clarify this important aspect.

We added this clarification in Section 4.2:

“Eddy properties such as their radius are determined at the locations where they are detected. In this sense, the eddy statistics are computed in a Lagrangian framework.”

An important eddy property which could be included in this study is the eddy non-linearity parameter. This parameter would give a good insight on the eddy’s coherence and ability to trap and transport material along their pathway of propagation.

Thanks for this suggestion. We added a new section (4.7) to analyse the eddy nonlinearity. Chelton et al. (2011) describe three different parameters to study eddy non-linearity, the advective nonlinearity parameter, the quasi-geostrophic nonlinearity parameter and the upper-layer thickness nonlinearity parameter. Here we decided to focus on the advective nonlinearity parameter, defined as the ratio of maximum rotational speed and translation speed.

Lines 248 – 249. Are the tides the only difference between the models? What about the vertical discretization of the water column??

We changed the sentence to:

“One of the many differences between the models...”

There are much more differences, which are discussed in Section 7. In this paragraph, we just speculate that tides are the main reason for the difference in eddy occurrences on the Yermak plateau in the models.

In section vertical extension and hydrographic properties: Are the values of vorticity in the eddy centers, a single point values or are averaged values within the eddies? Please make this aspect clear.

The relative vorticity in the water column is computed in the eddy center, and thus it is a single point value for every depth layer. We added in section 4.6: “...by calculating relative vorticity/ f at the location of the eddy centres...”

Line 279: is the superscript number after (Figure 11a, b)¹ meant to be there?

The superscript refers to a footnote.

Comments by the Editor

Thank you for this interesting manuscript. As you see, both reviewers have some constructive comments. I wanted to drop this editor comment so that you can consider the points below when preparing your final response and the revised version. My major criticism is that the discussion section is not well developed and must be improved.

Thank you for your detailed comments. We improved the discussion section, and hope that it is better now.

Here is a list of minor issues which must be clarified or written differently:

Li 25: “strongly turbulent”: presence of mesoscale eddies does not make the oceanic conditions strongly turbulent. If this is supported by microstructure measurements, please use and cite; if not please choose a different wording (energetic?).

Corrected. “Energetic” sounds much better.

Li 33: do eddies lead to vertical eddy fluxes? I thought they would lead to lateral fluxes. Otherwise, please clarify the pathway from lateral to vertical.

Eddies lead to both lateral and vertical fluxes, and the reason is that fluid particles move along isopycnals which are inclined. When eddies flatten isopycnals converting APE to EKE, lighter water moves upward, and denser downward along mean isopycnals. This is the main part of the GM parameterization. Therefore, eddies contribute to vertical heat flux. We reformulated point (2) in the introduction section:

“As shown by Hattermann et al. (2016), this region is characterised by negative values of vertical eddy temperature flux. Thus, eddy processes likely play an important role for the subduction of AW.”

Li 38. How is the MIZ “shaped” by eddies?

We reformulated this sentence: “Eddies play an important role for sea ice-ocean interaction. The marginal ice zone is influenced by eddies...”

Li 47-48: I do not think Teigen et al or Johannessen et al are the key references for the theory or dynamics related to barotropic instability or topographic steering/trapping of eddies, respectively.

We added the reference to Cushman-Roisin's book, and modified the sentence about barotropic instability: "... and eddies can be formed by receiving kinetic energy from the mean flow as shown for the Fram Strait by Teigen et al. 2011." We also added the reference to the study by Smith et al. (1984) which describes the topographic generation of an eddy in the EGC.

Li 92: Sundfjord et al (2017) is about a Svalbard fjord and I am not sure how it is relevant to the ability of the model to reproduce the slope boundary current.

We removed this reference.

Li 100: What are "a and b". Please clarify.

We added a description of parameters a and b.

Li 105: we also use

Corrected.

Li 117 and 118: please justify the choices of 3 day and 100 m depth

We added a justification of the choice of the 3 days and depth of 100 m in section 2.3: "We decided to use a threshold of 3 days mainly because the temporal resolution of the model output data is daily, and the eddy should form a track. This also helps to make sure that the eddies detected are real and not an overdetection due to uncertainties in the detection method. Eddies with a lifetime of at least three days are also required when computing the translation velocity needed to compute the eddy nonlinearity parameter, for which centred differences are used."

"We decided to choose the depth of 100 m since both main water masses of the Fram Strait, AW and PW, are present at this depth (e.g. Wekerle et al., 2017, their Figure 9)."

Li 130: by experts? (please clarify)

We removed this sentence to avoid confusion.

Heading 2.4, remove one "and"

Corrected.

Li 145: these terms do not indicate instability but rather conversion from MKE to EKE and EPE to EKE (which can be related to the instabilities you mention).

We corrected this sentence: "... can be related to.."

Li 169-171: the syntax of frequencies and slopes are difficult to follow for a reader

We restructured the sentence, and hope that it is clearer now.

Li 193: did you introduce a stream function?

This was described in the methods section 2.3: "Eddy boundaries around each detected centre are determined by the outermost closed contour of the stream function field."

Li 200: Isn't this the Rossby number?

Right, this is the Rossby number. We used the Rossby number as an index for eddy intensity. We added a better description of the eddy intensity in section 4.3.

Li 229: Molloy

Corrected.

Li 293: The Ghaffari analysis is from a 2-layer model? I am not sure how this is directly comparable (at least you might want to point this out). About the instability of the slope current along the Lofoten escarpment, please see some recent conversion rate calculations similar to yours, using high resolution ROMS fields (Section 9, in a otherwise mooring observation paper): Ocean Sci., <https://doi.org/10.5194/os-16-685-2020>. Not that I authored this paper, so feel free to ignore this suggestion. However, the conversion rate fields (Fig 11 in both papers) are directly comparable.

Thanks a lot for hinting us to this recent study, it fits much better here and we thus cited it. It is also very valuable to be able to compare the magnitude of conversion rates to our estimates. We removed the reference to the paper by Ghaffari et al.

Li 309: Need a dot product before the buoyancy gradient?

Corrected.

Discussion section is not appropriate and must be improved. Also I note that the last paragraph (on providing information to develop GM type parameterization) is not really supported by your results or built upon them in a convincing way. Please improve this part or remove.

We improved the Discussion section. In particular, we added a new paragraph which discusses the differences between eddy occurrences and EKE. Moreover, we added a paragraph about “Implications for contributing to future model development”. Here we also point to the energy backscatter scheme mentioned by Reviewer 1. We also improved the paragraph about the GM parameterization.

Opening paragraph of the Conclusions is not conclusions (or findings from your study), and could be integrated to discussion or removed.

We removed this paragraph.

Fram Strait: (If I’m not wrong) in the English usage, you should drop “the” in front of Fram Strait (except referring to a specific feature associated with Fram Strait, say, the Fram Strait circulation etc.) This must be corrected throughout, including the title.

Corrected.

Fig 4: I think it is not very meaningful to show the spectra of daily averaged speed (this is what you mean by absolute velocity?). It would be better to show the sum of spectra from u and v components (this corresponds to distribution of double the horizontal kinetic energy, or divide by two and call it HKE spectrum). In any case, you need units on y axis ($(\text{m/s})^2/(1/\text{day})$?). It would be helpful with 95% confidence intervals on the spectra.

As suggested, we computed spectra for u and v separately and show now in Figure 4 the sum of the two spectra divided by 2. Now units are shown on the y-axis, and the 95% confidence intervals are indicated too.

Properties and dynamics of mesoscale-eddies in Fram Strait from a comparison between two high-resolution ocean-sea ice models

Claudia Wekerle¹, Tore Hattermann^{2,3}, Qiang Wang¹, Laura Crews⁴, Wilken-Jon von Appen¹, and Sergey Danilov¹

¹Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany

²Norwegian Polar Institute, Tromsø, Norway

³Energy and Climate Group, Department of Physics and Technology, The Arctic University of Tromsø, Norway

⁴School of Oceanography and Applied Physics Laboratory, University of Washington, Seattle, USA

Correspondence: Claudia Wekerle (Claudia.Wekerle@awi.de)

Abstract. Fram Strait, the deepest gateway to the Arctic Ocean, is strongly influenced by eddy dynamics. Here we analyse the output from two eddy-resolving models (ROMS and FESOM) with around 1 km mesh resolution in Fram Strait, with focus on their representation of eddy properties and dynamics. A comparison with mooring observations shows that both models reasonably simulate hydrography and eddy kinetic energy. Despite differences in model formulation, they show relatively similar eddy properties. The eddies have a mean radius of 4.9 km and 5.6 km in ROMS and FESOM, respectively, with slightly more cyclones (ROMS: 54%, FESOM: 55%) than anticyclones. The mean lifetime of detected eddies is relatively short in both simulations (ROMS: 10 days, FESOM: 11 days), and the mean travel distance is 35 km in both models. More anticyclones are trapped in deep depressions or move toward deep locations. The two models show comparable spatial patterns of baroclinic and barotropic instability. ROMS has relatively stronger eddy intensity and baroclinic instability, possibly due to its smaller grid size, while FESOM has stronger eddy kinetic energy in the West Spitsbergen Current. Overall, the relatively good agreement between the two models strengthens our confidence in their ability to realistically represent the Fram Strait ocean dynamics, and also highlights the need for very high mesh resolution.

1 Introduction

Fram Strait, located between Svalbard and Greenland (Figure 1), is the deepest gateway that connects the Arctic Ocean and the North Atlantic via the Nordic Seas. Many important processes of climate relevance take place in this region. On the one hand, Atlantic Water (AW) carried northward by the West Spitsbergen Current (WSC, e.g. von Appen et al., 2016) enters the Arctic Ocean as its largest oceanic heat source. In the last decades, an increase in AW temperature has been observed in Fram Strait, with implications for the Arctic Ocean's sea ice decline (Beszczynska-Möller et al., 2012; Polyakov et al., 2012). On the other hand, a part of the AW recirculates in Fram Strait and continues southward in the East Greenland Current (EGC, e.g. de Steur et al., 2009). This water mass, which was densified on its way north to Fram Strait, contributes to the Denmark Strait overflow, which forms the dense part of the North Atlantic Deep Water, a key component of the Atlantic meridional overturning circulation (Eldevik et al., 2009). Furthermore, cold and fresh Polar Water (PW) carried southward by the EGC is injected into

the cyclonic Greenland Sea Gyre, impacting convection there (Rudels, 1995), and thus also the overflow across the Greenland Scotland Ridge.

25 The oceanic conditions in Fram Strait are strongly [energetic](#). Already in the 1980's it was revealed by measurement campaigns such as the Marginal Ice Zone Experiments that eddies are abundant there (Johannessen et al., 1987b; Smith et al., 1984). They play an important role in shaping the ocean circulation and hydrography, sea ice and ecosystem:

(1) Some eddies are shed from the WSC and travel westward, driving the recirculation of warm and salty AW. This was shown by mooring measurements (Schauer et al., 2004; von Appen et al., 2016) and model simulations (Hattermann et al.,
30 2016; Wekerle et al., 2017), which revealed high levels of eddy kinetic energy (EKE) in the WSC and along the recirculation pathway. [It is found that EKE in the WSC is much stronger than in the Arctic interior \(Wang et al., 2020\).](#)

(2) As AW recirculates, it subducts underneath cold and fresh Polar Water (PW) carried by the East Greenland Current (EGC). [As shown by Hattermann et al. \(2016\), this region is characterised by negative values of vertical eddy temperature flux. Thus, eddy processes likely play an important role for the subduction of AW.](#)

35 (3) Once the Return Atlantic Water (RAW) crosses (likely eddy mediated) the Northeast Greenland continental shelf break, part of it travels through a trough system towards the Northeast Greenland glaciers (Schaffer et al., 2017). An increase in its temperature might lead to the glaciers' destabilisation (Wilson et al., 2017), and it has been shown that eddy overturning is important for lifting AW onto the continental shelf in Fram Strait (Tverberg and Nøst, 2009; Cherian and Brink, 2018).

(4) [Eddies play an important role for sea ice-ocean interaction. The marginal ice zone is influenced by eddies](#) (Johannessen
40 et al., 1987b) and submesoscale features (von Appen et al., 2018). By means of idealised model experiments, Manucharyan and Thompson (2017) showed that cyclonic eddies can trap sea ice and carry it to warm waters, leading to enhanced melting rates.

(5) Eddy and filamentary structures are important features for the marine ecosystem. Among other effects, they play an important role in transporting nutrients into the euphotic zone for phytoplankton production, and can cause stratification within
45 days, thereby increasing light exposure for phytoplankton trapped close to the surface (Mahadevan, 2016).

Eddies can be generated through both baroclinic and barotropic instabilities (e.g. Cushman-Roisin, 1994). In the presence of horizontal density gradients and baroclinic instability, mesoscale eddies develop through the conversion of the available potential energy (APE) to EKE. Barotropic instability in contrast is associated with horizontal shear in jet-like currents, and eddies can be formed by receiving kinetic energy from the mean flow [as shown for the Fram Strait by Teigen et al. \(2011\).](#)
50 [Eddies can also be steered or trapped by topography, as observed for an eddy generated in the EGC \(Smith et al., 1984\).](#) This steering modulates the conversion between eddy and mean kinetic energy, which can be directed in both ways. Fram Strait, featured with its complex topography, strong lateral gradients in temperature and salinity (warm and saline AW in the eastern part, cold and fresh PW in the western part) and thus steep isopycnal slopes across the strait, strong convective events in the winter months and strong boundary currents (WSC and EGC), is thus a highly active and interesting region for studying eddy
55 dynamics.

The Rossby radius of deformation, which characterises the spatial scale of eddies, is small in Fram Strait, around 4–6 km in summer and 3–4 km in winter (von Appen et al., 2016). [Hallberg \(2013\) showed that in ocean models, a resolution of two](#)

grid points per Rossby radius of deformation can be considered as threshold between "non-eddy" and "eddy-permitting" regimes, and thus higher resolution is needed for a model to be considered as "eddy-resolving". This poses problems for ocean models which typically operate on coarser grids. Recently, high resolution ocean models focused on the Fram Strait region have emerged, which perform well in reproducing the observed eddy activity (Kawasaki and Hasumi, 2016; Hattermann et al., 2016; Wekerle et al., 2017).

Given the possible sensitivity of simulations to model numerics, to the complex bottom topography and ocean currents in Fram Strait, it is not known whether the above cited models have a broad agreement on the representation of eddy dynamics in terms of eddy generation and propagation. Answering this question will not only add credence to our understanding of eddy dynamics, but also create a reference for developing parameterisations required by coarse resolution ocean models. The aim of this study is two-fold. First, we compare the output of two high-resolution, eddy-resolving ocean-sea ice models to answer the above question. We will show that there is good agreement in energy conversion that maintains eddy dynamics and in simulated eddy statistics as well, despite the fact that these models, namely ROMS (Shchepetkin and McWilliams, 2005; Budgell, 2005; Hattermann et al., 2016) and FESOM (Wang et al., 2014; Wekerle et al., 2017), differ in many aspects such as numerical discretisation, horizontal and vertical mesh resolution, parameterisations, global vs. regional configurations. Second, we explore and describe the properties of eddies in Fram Strait. We use an eddy-following approach to generate regional statistics focusing on the following questions: How are eddies spatially distributed? Are anticyclones or cyclones dominating? What is their typical size, lifetime and what are their main travel pathways?

2 Methods

2.1 Model description FESOM

Model output from the Finite-Element Sea-ice Ocean Model (FESOM) version 1.4 (Wang et al., 2014; Danilov et al., 2015) is used for eddy detection and tracking in this study. FESOM is an ocean-sea ice model which solves the hydrostatic primitive equations in the Boussinesq approximation and is discretised with the finite element method (Wang et al., 2008). In the vertical, z-levels are used. We use a global FESOM configuration that was optimised for Fram Strait with regional resolution (grid size) refined to 1 km in this area, and a coarser resolution elsewhere (1° resolution throughout most of the world's oceans, 24 km resolution north of 40°N and 4.5 km resolution in the Nordic Seas and Arctic Ocean; Wekerle et al. (2017)). By comparing with the local Rossby radius of deformation (around 3–6 km in Fram Strait, see above), this configuration can be considered as "eddy-resolving". It is forced with atmospheric reanalysis data from COREv.2 (Large and Yeager, 2008), and river runoff is taken from the interannual monthly data set provided by Dai et al. (2009). Tides are not taken into account in the FESOM configuration used here. The simulation covers the time period 2000–2009, and has daily output. In this study, we analyse model output for the years 2006–2009.

2.2 Model description ROMS

The second high-resolution model simulation used in this study is based on the Regional Ocean Modeling System (ROMS) (Budgell, 2005; Haidvogel et al., 2008; Shchepetkin and McWilliams, 2005, 2009) with a configuration optimised for Fram Strait and the waters around Svalbard (called S800). With 800 m x 800 m horizontal resolution, S800 is eddy resolving in Fram Strait. S800 was initialised with and forced at the ocean boundaries with daily ocean and sea ice data from a 4 km resolution pan-Arctic model called A4, together with tidal elevations from global TPXO tidal model (Egbert and Erofeeva, 2002). A4's initial state and boundary conditions were taken from monthly-averaged global reanalyses (Storkey et al., 2010). Atmospheric forcing in A4 and S800 used 6-hourly ERA-Interim reanalysis (Dee et al., 2011). A4 was initialised in 1993, and following A4 spin-up S800 was initialised in January 2005. Analyses in this paper are done for the period of 2006–2009. Model characteristics of ROMS, and also of FESOM, are summarised in Table 1. Additional information about S800, including discussions of its ability to reproduce boundary current observations in Fram Strait and along the continental slope north of Svalbard, is given in Hattermann et al. (2016), Crews et al. (2018) and Crews et al. (2019).

2.3 Eddy detection and tracking

Eddy detection and tracking algorithms are important tools to understand eddy properties such as their size, strength, lifetime and travel pathways. For datasets as large as the output of ocean models, automated methods need to be used. Eddy detection methods can be assigned to two categories, based either on (1) geometrical or on (2) physical characteristics of the flow field, or on a combination of both. In this study, we apply a method developed by Nencioli et al. (2010) to detect and track eddies simulated with ROMS and FESOM, which is based on the geometry of velocity vectors and thus belongs to the first category of methods. The eddy detection is based on four constraints derived from the general characteristics of velocity fields in the presence of eddies (Nencioli et al., 2010):

1. Along an east-west (EW) section, v has to reverse in sign across the eddy centre and the magnitude of v has to increase away from it.
2. Along a north-south (NS) section, u has to reverse in sign across the eddy centre and the magnitude of u has to increase away from it. The sense of rotation has to be the same as for v .
3. The velocity magnitude has a local minimum at the eddy centre.
4. The sign of vorticity cannot change around the eddy centre.

Two parameters, a and b , which determine the minimum size of detectable vortices, have to be set in the algorithm. Parameter a defines over how many grid points the increases in magnitude of v along the EW axes and u along the NS axes are checked, and its unit is "grid points". It also defines the size of detectable eddies, which is $a-1$ grid points. Parameter b defines the size (also in grid points) of the area used to find the local velocity minimum. After some sensitivity tests, we set $a = 4$ and $b = 3$, which equals the values used in the test case of Nencioli et al. (2010). Note that our mesh resolutions (800 m and 1 km in

ROMS and FESOM, respectively) are similar to theirs (1 km). Eddy boundaries around each detected centre are determined
 120 by the outermost closed contour of the stream function field, across which velocity magnitudes are still increasing in the
 radial direction. This definition is different than the one used by Bashmachnikov et al. (2020) where the eddy boundary is
 approximated by the zero relative vorticity contour with a circle or an ellipse. Note that the method used in this study results in
 smaller eddy radii than the one used by Bashmachnikov et al. (2020).

To cross-validate our results, we also used the Okubo-Weiss criterion, which belongs to the second category of methods
 125 (Okubo, 1970; Weiss, 1991). Eddies are identified as areas where vorticity dominates over strain. More precisely, the area
 where the Okubo-Weiss parameter

$$OW = \underbrace{(\partial_x u - \partial_y v)^2 + (\partial_x v + \partial_y u)^2}_{\text{normal and shear component of strain}} - \underbrace{(\partial_x v - \partial_y u)^2}_{\text{relative vorticity}} \quad (1)$$

is below a threshold of $OW_0 = -0.2\sigma_{OW}$ with same sign of vorticity, where σ_{OW} is the spatial standard deviation of OW, is
 considered as an eddy (Isern-Fontanet et al., 2006). Here (u, v) is the horizontal velocity field, and f is the Coriolis parameter.

After eddies are detected, eddy tracks are computed by comparing eddy centres in successive time steps. More precisely,
 130 if two eddies at successive time steps lie within a search radius and have the same sense of rotation, they form a track. The
 eddy tracking scheme is thus sensitive to the prescribed search radius. A too small value might lead to a false splitting of the
 track, whereas a too large value would lead to more than one eddy within the searching area. As a first approximation, eddies
 are advected with the mean current. Considering a mean velocity of around 0.2 m/s (see e.g. Figure 5 in Wekerle et al. (2017))
 and a daily mean model velocity field, a possible choice would be a search radius of 17 km. After performing sensitivity
 135 tests with different radii, we chose a radius of 14 km. This value reduced the number of occasions when several eddies were
 detected in the searching area. Furthermore, eddies with a lifetime shorter than 3 days were discarded. We decided to use this
 threshold because the temporal resolution of the model output data is daily, and the eddy should form a track. This also helps
 to make sure that the eddies detected are real and not an over-detection due to uncertainties in the detection method. Eddies
 with a lifetime of at least three days are also required when computing the translation velocity needed to compute the eddy
 140 nonlinearity parameter (Section 4.7), for which centred differences are used.

For the eddy detection and tracking, we use daily model output for the time period 2006–2009 at 100 m depth. At this
 depth, the water mass lateral distribution is characterised by warm and salty AW in the eastern part of Fram Strait (in the
 WSC), and by cold and fresh PW in its western part (in the EGC). We decided to choose the depth of 100 m because both
 main water masses of the Fram Strait, AW and PW, are present at this depth (e.g. Wekerle et al., 2017, their Figure 9). In
 145 addition, we found that eddy vorticity has largest magnitudes at about 100 m depth (see section 4.6). Output from both models
 is interpolated to a regular grid (0.05° longitude x 0.01° latitude) which has approximately the same resolution as the original
 grids. Relative vorticity normalised by f at 100 m depth on Jan 1st 2006 is shown in Figure 2, as well as eddies detected by the
 Nencioli et al. (2010) method overlaid on the simulated Okubo-Weiss parameter. Note that the colour only shows the area with
 $OW < -0.2\sigma_{OW}$, i.e. the area considered as vortices. In both models, the relative vorticity field exhibits strong eddy activity,
 150 particularly along the pathway of the main currents, WSC and EGC, along the Yermak and Svalbard branches and in the AW
 recirculation area. Apart from well defined eddies, the relative vorticity fields show lots of elongated filamentary structures

reminiscent to what was found by von Appen et al. (2018). They seem to have a smaller scale in ROMS than in FESOM. Most of the eddies can be detected by both the Okubo-Weiss method and the Nencioli et al. (2010) method, while the Okubo-Weiss method detected a bit more eddies.

155 2.4 Reynolds decomposition of eddy fluxes and kinetic energy

To estimate the contributions of mesoscale eddy field to the flow variability, we decompose a variable x which can stand for velocity (u) or tracers (c) into a monthly mean (\bar{x}) and a daily-averaged fluctuating (x') component, $x = \bar{x} + x'$. We derive the time-mean eddy flux of the tracer c in the u velocity direction from the equality $\overline{c'u'} = \overline{c'u} - \bar{c}\bar{u}$. Similarly, time-averaged eddy kinetic energy (EKE) is computed as

$$EKE = \frac{1}{2}(\overline{u'^2} + \overline{v'^2}) = \frac{1}{2}(\overline{u^2} + \overline{v^2} - \bar{u}^2 - \bar{v}^2). \quad (2)$$

160 2.5 Energy budget

An energy budget can be obtained by expressing velocity as $\mathbf{u} = \bar{\mathbf{u}} + \mathbf{u}'$ as described in the previous section, inserting it in the momentum equation in the Boussinesq approximation, multiplying the equation with \mathbf{u}' , and time-averaging it. This leads to a conservation equation for EKE. The change of EKE in time is governed by the advection of eddies, energy transfer from mean kinetic energy (MKE) and available potential energy (APE) to EKE, and energy dissipation (vertical mixing and horizontal diffusion) (e.g. Olbers et al., 2012, chapter 12):

$$\frac{\partial \frac{1}{2}(\overline{u_1'^2} + \overline{u_2'^2})}{\partial t} + \underbrace{\frac{\partial}{\partial x_j} \left(\frac{1}{2} \overline{u_j u_i'^2} + \frac{1}{2} \overline{u_j' u_i'^2} + \frac{1}{\rho_0} \overline{u_j' p'} \right)}_{\text{Transport}} = \underbrace{-\overline{u_j' u_i'} \frac{\partial \bar{u}_i}{\partial x_j}}_{\text{MKE} \leftrightarrow \text{EKE}} + \underbrace{\overline{w' b'}}_{\text{APE} \leftrightarrow \text{EKE}} + \underbrace{\overline{V_i' u_i'} + \overline{D_i' u_i'}}_{\text{Dissipation}}, \quad (3)$$

where $b = -\frac{g\rho}{\rho_0}$ is the buoyancy and D_i and V_i are horizontal and vertical dissipation terms. Cartesian tensor notation with summation convention has been used, with $i = 1, 2$ and $j = 1, 2, 3$. u_i is thus the horizontal component of the velocity vector u_j , and $u_3 = w$ is the vertical velocity. In this study, we diagnose the first two terms on the right hand side of the equation. They are the main source terms of EKE, and can be related to barotropic and baroclinic instability.

170 3 Model assessment

For more than two decades, mooring measurements have been conducted across Fram Strait at around 79°N to monitor the exchange of water masses through this gateway (e.g. Beszczynska-Möller et al., 2012; von Appen et al., 2016; von Appen et al., 2019). To assess the overall model performance in reproducing the mean state and resolving the flow variability, we use the observed hydrography as well as the velocity field and compare the latter in terms of power density spectra (PDS) and EKE to the model results.

The two models simulated **relatively** similar spatial distributions of **thermo-haline properties**. The simulated mean temperature and salinity at 100 m depth reveal that the warm ($>5^{\circ}\text{C}$) and narrow WSC closely follows the 1000 m isobath along the Svalbard shelf break (Figure 3). Recirculation of AW mainly occurs north of the Boreas Basin (north of 78°N). The western part of Fram Strait is characterised by cold and fresh polar outflow. The two models differ more significantly north of 80°N ,
180 with much warmer and saltier waters on the Yermak Plateau in FESOM. **The front between cold PW and warm AW, indicated by 1°C and 2°C isotherms, is sharper in FESOM than in ROMS. This can also be seen in T/S diagrams (Figure 3e and f).** Compared to the mooring observations across Fram Strait, both models **relatively well** represent the **thermo-haline properties**. ROMS shows a slightly cold bias which is not present in FESOM (ROMS: root mean square (rms) error of 1.28°C , FESOM: rms error of 0.49°C), and has earlier been identified to be associated with a cold bias in the A4 model that provides the inflow
185 boundary conditions for S800 (Hattermann et al., 2016). The simulated **thermo-haline properties** in FESOM, particularly in central and eastern Fram Strait, are slightly too saline, whereas they are slightly too fresh in ROMS. The overall rms error in salinity is 0.26 and 0.31 in ROMS and FESOM, respectively.

For the comparison of velocity, current meter data from two moorings located in the WSC and three moorings located in the EGC (for locations see Figure 1) deployed during the time period 2006–2009 were used (von Appen et al., 2019). **Time series
190 of the u and v components of the velocity in the WSC and EGC were created by averaging over the two WSC and three EGC moorings, respectively.** Daily averages of measured velocity at 75 m depth were calculated. Note that there are slight variations in the depth between the individual deployment years. The observed mean speed averaged over WSC and EGC moorings at 75 m depth is 0.22 m/s and 0.13 m/s respectively, while the mean speed of ROMS/FESOM at the mooring locations is 0.24/0.20 m/s and 0.16/0.12 m/s respectively.

195 Power density spectra of the **horizontal kinetic energy** from the observations and from the models were estimated via the Thomson multitaper method (Figure 4). **For the WSC time series, we used linear interpolation to fill some mooring data gaps (maximum gap of 14 days). For the EGC time series, we used the time period Sep 8 2006 – Dec 31 2009 due to too long data gaps in early 2006. Spectra were computed for u and v components separately, summed and divided by 2. Slopes of the spectra between frequencies of $1/(14 \text{ days})$ and $1/(3 \text{ days})$ were computed by determining the median in $\log_{10}(0.05/\text{day})$ frequency
200 steps and then fitting the slopes to those binned values.** The slopes of the observations are both ~ -1.6 for WSC and EGC moorings, respectively, while ROMS/FESOM showed slopes of $\sim -1.7/\sim -2.0$ and $\sim -2.4/\sim -2.7$ respectively. The difference between the models is larger at high frequency, which might be related to the fact that tides were simulated in ROMS **and the models apply different atmospheric forcing. The differences between the models will be further discussed in Section 6.**

Maps of the simulated EKE reveal high energy levels along the pathways of the WSC, the recirculation area and the EGC
205 (Figure 5a and b). In both models, there is a lateral gradient from west to east, with a higher level of EKE in the eastern part of Fram Strait, the WSC region. This gradient is even more pronounced in FESOM than in ROMS. In the WSC region, FESOM shows a higher EKE level than ROMS. In the EGC, this is opposite, with a more energetic EGC in ROMS than in FESOM. This is also reflected in the power density spectra described above. A seasonal cycle of EKE in 75 m depth computed from current meter data of moorings deployed across Fram Strait is shown in Figure 5c. The highest level of EKE is reached in the
210 winter months (January–March), and lowest values are reached in early autumn (September–November). Both models well

reproduce the observed seasonal and spatial variations of EKE (Figure 5b and c), except that the observation shows a higher EKE level in the central Fram Strait than the models.

4 Eddy properties

4.1 Eddy spatial distribution and polarisation

215 During the time period 2006–2009, altogether 218,213 eddies were detected in the area 8°W – 20°E / 76°N – 82°N in ROMS (and thus 149 eddies per day), with slightly more cyclones (54%) than anticyclones. The result is very similar in FESOM, with 55% of the 244,811 detected eddies (168 per day) being cyclones. The tracking algorithm then revealed that these eddies belong to 30539 and 39040 tracks for ROMS and FESOM, respectively. In both simulations, the eddy density is highest in the eastern and central part of Fram Strait (Figure 6a,b). In contrast, the eddy density is low in the western part of Fram Strait and on the
220 East Greenland continental shelf, that is, in areas covered by sea ice year-round. Comparing FESOM and ROMS, there are fewer eddies detected in that region in FESOM, which is also reflected in lower EKE values in the western part of Fram Strait than in ROMS (Figure 5). Both models show a consistent pattern in the distribution of cyclones vs. anticyclones, which has strong regional differences (Figure 6c and d). Over the Svalbard shelf and along the East Greenland continental shelf break, cyclones are predominant. Anticyclones dominate along the main pathway of the WSC (along the 1000 m isobath), over the
225 Yermak Plateau, and along the Svalbard branch.

4.2 Eddy size

In this study we compute the eddy radius as average distance from the eddy centre to the eddy boundary, which is defined by the outermost closed contour of the stream function field. Eddy properties such as their radius are determined at the locations where they are detected. In this sense, the eddy statistics are computed in a Lagrangian framework. Eddies detected in both
230 models are relatively small, with 95%/92% of cyclones and 92%/87% of anticyclones in ROMS/FESOM having a radius below 10 km (Figure 7a). Averaged over the whole Fram Strait region, the mean/median radius for ROMS and FESOM is 4.9/4.1 and 5.6/4.7 km, respectively (Table 2). Eddies simulated in FESOM are thus slightly larger than in ROMS. The eddy radius compares well with the Rossby radius of deformation (~ 4 – 6 km in summer and smaller values in winter (von Appen et al., 2016)). This suggests that baroclinic instability is likely the main mechanism of eddy generation, which will be further
235 investigated in Section 5. In both simulations, cyclones are slightly smaller than anticyclones (Table 2).

4.3 Eddy intensity

Here we take the Rossby number, the absolute value of relative vorticity divided by the Coriolis parameter f , as an index for the eddy intensity. A Rossby number of ~ 1 indicates that the eddy is in cyclogeostrophic balance. The maximum value of daily mean relative vorticity within the eddy boundary is computed and averaged over all detected eddies. The mean/median
240 intensity of eddies simulated by ROMS and FESOM is 0.4/0.36 and 0.28/0.24, respectively. Eddies simulated by FESOM are

thus weaker than eddies simulated by ROMS (see also Figure 7b and Table 2). The proportion of eddies with absolute values below 0.3 is larger for FESOM (63%) than for ROMS (38%). Cyclones are slightly more intensive and have a larger standard deviation than anticyclones in both models (Table 2).

4.4 Eddy lifetime and travel distance

245 The duration over which eddies are continuously detected by the employed method is on average 10 and 11 days in ROMS and FESOM, respectively (Figure 7e). 85%/82% of eddies detected in ROMS/FESOM have lifetimes below 15 days, whereas only 4%/6% of eddies detected in ROMS/FESOM have lifetimes above 30 days. Pathways of these long-living eddies will be analysed in the next section. Note that the eddy lifetime may be longer if one considers that eddies likely can exist for some time before and after being detected as an eddy by the tracking method. Also, a false splitting of the track could occur if the eddy
250 moved relatively fast in combination with a too small searching area. In both simulations, there is no significant difference in lifetime regarding polarisation. They are very similar regarding travel distance. On average, eddies travel around 34 and 35 km in ROMS and FESOM, respectively (Table 2). Again, there is no significant difference in travel distance regarding polarisation (Figure 7f). Compared to eddies generated e.g. in the Gulf Stream region, the lifetime of Fram Strait eddies is rather short (Kang and Curchitser, 2013).

255 4.5 Eddy pathways

Eddy pathways are investigated by focusing only on long-living eddies, e.g. eddies with lifetime of more than 30 days, and by classifying them by generation areas (Figures 8 and 9). In both simulations, eddies generated on the Svalbard shelf have very distinct travel pathways for cyclones and anticyclones, which is consistent with their distribution (Figure 6e and f). Cyclones tend to stay on the shelf, and populate the narrow Svalbard fjords. Anticyclones in contrast leave the shallow shelf area and
260 tend to travel westward into the deep basin. As shown in Figure 7c, more cyclones (31% and 25% in ROMS and FESOM, respectively) are detected in shallow areas with water depths less than 500 m than anticyclones (21% and 19% in ROMS and FESOM, respectively). Note that as the number of detected eddies on the East Greenland shelf is relatively small in both simulations, most eddies detected in shallow areas are located on the Svalbard shelf.

Anticyclones generated in the WSC core region, here defined approximately as the area between the 500 m and 2000 m
265 isobaths, show longer travel pathways than cyclones. In both simulations, most of them travel westward along the recirculation pathway north of the Molloy Deep (Hattermann et al., 2016), and some even continue southward along the East Greenland continental shelf break. Some eddies travel northward along the western rim of the Yermak plateau or recirculate around the Molloy Deep, while only few trajectories deviate westward south of 79°N in both models.

The asymmetric pathways of eddies generated on the Svalbard shelf and in the WSC core region can have dynamical reasons.
270 As described by Cushman-Roisin (1994, Chapter 17), fluid parcels surrounding a rotating eddy are stretched when they move to deeper waters and thus acquire relative vorticity. In contrast, when moving to shallower waters, on the flank of the eddy the surrounding fluid is squeezed and thus relative vorticity is decreased. This results in a secondary drift of the vortices, with cyclones moving towards shallower regions and anticyclones moving to deeper regions. Morrow et al. (2004), based on satellite

altimetry, showed that this dynamical reasoning can explain the diverging pathways of cyclones and anticyclones in different
275 ocean basins. The asymmetry can also be explained by the different water masses present along the Svalbard continental shelf.
Along the Svalbard coast, the Svalbard Coastal Current transports cold and fresh waters northward, close to the salty and warm
AW which is carried northward by the WSC a little offshore. The meandering between the two water masses, light water on the
eastern side and denser water on the western side (roughly indicated by the 200 m isobath in Figure 6c,d) leads to the generation
of cyclones on the eastern side and anti-cyclones on the western (offshore) side, which is comparable to eddy shedding along
280 the Gulf Stream (e.g. Olson, 1991).

Tracks of long-living eddies generated in southern central Fram Strait, in particular those simulated in ROMS, show a high
density of anticyclones in the Boreas Basin, the region between $0^{\circ}\text{EW}-5^{\circ}\text{E}$, $76^{\circ}\text{N}-77^{\circ}\text{N}$. More anticyclones appear to be
trapped in this depression, a similar situation as occurring in the Lofoten Basin (Raj et al., 2016; Volkov et al., 2015). As in the
case of eddies generated along the Svalbard shelf break, the clustering of anticyclones can be explained by the dynamical reason
285 described above (anticyclones move towards the deeper basin, thus the centre of a depression). Eddies generated in northern
central Fram Strait tend to travel westward, then follow the East Greenland continental shelf break. Particularly, anticyclones
travel westward between the northern rim of the Boreas Basin and the Molloy Deep, contributing to the AW recirculation.

Regarding eddies present in northern Fram Strait, both ROMS and FESOM show a high density along the western flank
of the Yermak Plateau. Additionally, ROMS shows more long-living (>30 days lifetime) eddies west of the Plateau (Figure
290 6a-d) than FESOM (Figures 9 and 8). Eddies in this region have earlier been identified to occur with a different seasonality
than would be expected from changes in baroclinic instability of the boundary current that explains the seasonality in eddy
occurrence along other parts of the shelf break (Crews et al., 2019). One of the many differences between the two models is
the inclusion of tidal forcing in ROMS. The circulation and water mass transformations above the Yermak plateau are known
to be strongly influenced by barotropic to baroclinic tidal conversion and mixing at the semi-diurnal critical latitude (Fer et al.,
295 2015), that may also explain the enhanced eddy generation in this region in ROMS. As revealed by FESOM, more cyclones
tend to follow the Svalbard Branch, whereas more anticyclones tend to follow the Yermak Branch.

4.6 Vertical extent and hydrographic properties

We determined the vertical extent of eddies detected in 100 m depth with lifetime above 30 days by calculating relative
vorticity/ f at the location of the eddy centres in the water column (Figure 10). In addition, temperature and salinity anomalies
300 were calculated in the same way to study the hydrographic properties of eddies, with anomalies computed relative to the
mean value for the month. This was done for eddies generated in the five different regions shown in Figure 1. Profiles of
relative vorticity are relatively similar in ROMS and FESOM, with most negative/positive (i.e. strongest vortices) values for
anticyclones/cyclones generated in the WSC region and central Fram Strait.

The hydrographic conditions in regions WSC, central southern Fram Strait and Yermak/Svalbard Branch are characterised by
305 warm and salty AW (Figure 3). These regions are temperature-stratified and unstable in salinity. Anticyclones generated there
carry anomalously warm and salty and thus lighter waters and have depressed isopycnals, whereas cyclones carry anomalously
cold and fresh and thus denser waters and have raised isopycnals (Figure 10). Western Fram Strait is characterised by cold and

fresh PW, and is salinity stratified. The transition from a temperature stratified to a salinity stratified regime in the different regions may partly explain the difference in properties between ROMS and FESOM.

310 4.7 Eddy nonlinearity

We assessed the nonlinearity of eddies by computing the advective nonlinearity parameter U/c , where U is the maximum rotational speed estimated as the maximum speed inside the eddy defined by the outer boundaries and c is the translation speed of the eddy estimated at each point along the eddy trajectory from centered differences (Chelton et al., 2011). Eddies with a value of $U/c > 1$ can trap fluid in their interior and transport water properties, and are considered as nonlinear. In ROMS, 315 86% of the simulated eddies have a value of $U/c > 1$, and the percentage is quite similar in FESOM with 83% (see also the histogram of U/c shown in Figure 7d). When considering long-living eddies only (lifetime > 30 days), the percentage of nonlinear eddies is higher (94% and 92% in ROMS and FESOM, respectively). This is different in comparison with the global study of Chelton et al. (2011) who find all of the observed mesoscale eddies outside the tropics are nonlinear. However, they only consider long-living eddies with lifetime above 16 weeks. The most highly nonlinear eddies are found on the offshore side 320 of the strongly meandering WSC and in the AW recirculation area (Figure 11). This indicates that ocean heat is transported from the main current into the deeper basin.

5 Energetics in eastern Fram Strait

We now analyse the source of EKE as simulated in ROMS and FESOM. We focus here on the eastern side of Fram Strait, which is the most energetic region (Figure 5). As described in section 2.5, the change of EKE in time is governed by the advection 325 of eddies, energy transfer from mean kinetic energy (MKE) and eddy available potential energy (APE) to EKE, and energy dissipation. In this study we analyse only the first two terms on the right hand side of the EKE conservation equation (Eq. 3), which are the main source terms for EKE and are related to barotropic and baroclinic instability.

5.1 Barotropic instability

The transfer of MKE to EKE is related to barotropic instability. It can be expressed as the sum of two terms, the product of 330 horizontal eddy Reynolds stress and horizontal mean shear, and the product of vertical eddy Reynolds stress and vertical mean shear. Strong velocity shear thus support barotropic instability. Here we consider only terms that contain horizontal derivatives, and assume that the terms with vertical derivatives play a minor role (as shown for the Gulf Stream region by Gula et al. (2015)). In the two models, the energy conversion between MKE and EKE is directed in both ways: it shows an alternating pattern, with positive values indicating conversion from MKE to EKE and negative values indicating conversion from EKE to MKE 335 (Figure 12a,b)¹. The alternating pattern is very similar between the both models, with consistent locations and magnitude of positive and negative energy transfer. The energy transfer occurs mainly along the pathway of the WSC core, which is located

¹Note that there has been an error in the computation of the MKE to EKE conversion term shown in Figure 14a of Wekerle et al. (2017). Figure 12b shows the correct pattern.

approximately along the 500–1000 m isobaths (see also Figure 3). This is comparable to the Norwegian continental slope off the Lofoten islands, where barotropic instability is particularly important in the presence of steep bottom slopes as shown in a recent study by Fer et al. (2020). The magnitude of the depth averaged barotropic energy transfer of around (0–1) 10^{-4} Wm^{-3} obtained from a high-resolution ROMS simulation in Fer et al. (2020) compares well to the values in the WSC estimated from ROMS and FESOM shown in this study.

The relatively similar pattern in both models suggests that there is a strong influence of bathymetry, which determines positive and negative spots of energy conversion. A necessary condition for barotropic instability is that $\beta - \partial_{yy}\bar{u}$ vanishes within the domain, where $u = \bar{u}(y)$ is a zonal current with arbitrary meridional profile (e.g. Cushman-Roisin, 1994). The planetary potential vorticity is weak and can be ignored in polar regions, so we only consider the topographic β , with $\beta = -\frac{f}{H}\nabla H$ quantifying the change in potential vorticity across the bathymetry and H and ∇H being the water depth and its horizontal gradient. A map of the topographic β west of Svalbard reveals large values along the Svalbard shelf break (Figure 13a). We take the depth-averaged monthly mean meridional velocity v from FESOM as an approximation of the along-stream velocity, and compute $\beta - \partial_{xx}v$ (Figure 13b). In many places along the Svalbard shelf break, β is much larger than $\partial_{xx}v$. However, in some places, e.g. at the entrance of Kongsfjorden (79°N) and Isfjorden ($78^\circ10'\text{N}$) and along the 250 m isobath at around 80°N , $\beta - \partial_{xx}v$ changes sign. These regions are characterised by positive values of energy conversion in both models, indicating active barotropic instability there.

5.2 Baroclinic instability

For baroclinic instability to be active, a horizontal density gradient must be present to provide available potential energy which can be converted to EKE. This transfer from APE to EKE can be expressed as the mean vertical eddy buoyancy flux (Eq. 3). In contrast to barotropic instability, the energy conversion between APE and EKE in FESOM and ROMS is directed mostly one way, with mainly positive values revealing conversion from APE to EKE (Figure 12c,d). As eastern Fram Strait is temperature-stratified, it is mainly the vertical eddy temperature flux that contributes to vertical eddy buoyancy flux (Hattermann et al., 2016, their Figure 3d). In both models, baroclinic instability is strongest between the 1000 m and 2000 m isobaths in eastern Fram Strait. The values are slightly weaker in FESOM than in ROMS. The weaker baroclinic instability in FESOM is also reflected by the fact that detected eddies are characterised by lower values of absolute relative vorticity (Figure 7b). Between the 500 m and 1000 m isobaths, both models show patches of negative vertical eddy buoyancy fluxes. Usually, those patches indicate regions where eddy fluxes interact with the sloping topography to lift dense water onto the continental shelf (Tverberg and Nøst, 2009), with upward sloping isopycnals near the seafloor that locally enhance the APE of the mean field.

A necessary condition for baroclinic instability is that the cross-stream gradient of Ertel potential vorticity (PV) changes sign with depth (e.g. Spall and Pedlosky, 2008). Ertel PV Π is defined as

$$\begin{aligned} \Pi &= (f\mathbf{k} + \nabla \times \mathbf{u}) \cdot \nabla b \\ &= \underbrace{f \partial_z b}_{\text{vertical stretching}} + \underbrace{(\partial_y w - \partial_z v) \partial_x b + (\partial_x w - \partial_z u) \partial_y b}_{\text{tilting vorticity}} + \underbrace{(\partial_x v - \partial_y u) \partial_z b}_{\text{relative vorticity}}. \end{aligned}$$

Here we compute Π from simulated long-term mean velocity $\mathbf{u} = (u, v, w)$ and buoyancy b , and neglect the small terms
370 containing derivatives of vertical velocity w . Figure 14 shows the Ertel PV and its gradient in zonal direction for two sections
across the Svalbard shelf break (78°N and $78^\circ50'\text{N}$) for the FESOM simulation. The dominant term is the vertical stretching
term, with a smaller contribution from the relative vorticity terms. The tilting terms are one order of magnitude smaller (Figure
not shown). At both sections, the cross-stream gradient reveals a change in sign with depth, indicating that the mean current is
baroclinically unstable. This is in agreement with studies by Teigen et al. (2011) and von Appen et al. (2016), and our simulated
375 energy conversions (Figure 12c,d).

6 Discussion

6.1 Connection between eddy occurrences and EKE

Although baroclinic instability is the main driver of mesoscale eddy variability, the connection between eddy occurrences
and the APE to EKE conversion rate ($w'b'$) is very non-local. For one thing, eddies form as a result of nonlinear evolution
380 of baroclinic instability waves and jet meanders. For another, mean circulation transports and modifies all eddy-like features,
moving them away from the sites where their 'seeds' originally appeared. Therefore, the observed pattern of eddy occurrences
(Figure 6a,b) differ from the $w'b'$ distributions (Figures 5 and 12).

According to Martínez-Moreno et al. (2019), EKE can be divided into a part containing energy related to eddies and a
part related to other effects such as meandering of the current. The non-negligible potential contribution of meandering to the
385 calculated EKE fields can be seen in the maps of eddy occurrences. Along the main pathway of the WSC, which is roughly
along the 1000 m isobath, the eddy occurrences are rather low in both ROMS and FESOM, whereas the EKE in both models
shows a maximum along this isobath. In general, the spatial correspondence between high EKE and high eddy occurrence is
not very strong. This mismatch could also be due to the fact that different individual eddies can have different levels of EKE.
We do not expect to use the level of EKE to predict the number of eddies. As a result, the pattern of eddy occurrences fills the
390 basin. The regions of high EKE and $w'b'$ are at the periphery, but they supply the perturbations that evolve into eddies.

6.2 Differences and similarities between observations, ROMS and FESOM

Despite their very fine resolution, ROMS and FESOM simulate a weaker variability in velocity than the observed in terms of
the power density spectrum (Figure 4). This might indicate that the model resolution used is still insufficient to well resolve all
the mesoscale eddies in the presence of numerical dissipation. Part of the variability revealed by the power density spectrum
395 can also be attributed to the atmospheric forcing. Although the forcing datasets are different in the two cases, both of them
are derived from relatively coarse reanalysis products (in particular, COREv.2 used in the FESOM simulation has a zonal
resolution of approximately 1.875°) and may miss part of small-scale variability. A topic for further research is to clarify the
importance of these factors.

A recent study by Bashmachnikov et al. (2020) compared properties of eddies detected from FESOM sea surface height fields, AVISO altimetry and spaceborne synthetic aperture radar (SAR), and revealed the difficulty of comparing model results to satellite data. The study showed that AVISO and SAR form two complementary data sets of large mesoscale eddies and of small mesoscale/submesoscale eddies, respectively. The mean FESOM eddy radius lies in-between of AVISO and SAR results. The resolution of 1 km in FESOM is thus too coarse to well resolve the submesoscale eddies detected from SAR data.

Snapshots of simulated relative vorticity (Figure 2) and the histogram of eddy intensity (Figure 7b) suggest that ROMS simulated finer and more intensive eddies and filaments. This indicates that the model effective resolution (Soufflet et al., 2016) in FESOM might be slightly lower than in ROMS. First, the grid size is slightly larger for FESOM (1 km vs 800 m for ROMS). This small difference in the grid size (20%) might matter as both numerical dissipation and explicit viscosity decrease with the grid size. In both models, biharmonic viscosity which scales with grid size cubed is applied. Second, FESOM1.4 is based on a collocated discretisation (an analog of Arakawa A-grid), whereas a staggered Arakawa C-grid is employed by ROMS. Because of pressure gradient averaging required by collocated discretisations, the effective resolution could be reduced. The collocated discretisation of FESOM also requires to use the no-slip boundary condition, which implies more dissipation along the boundary as well. Third, FESOM relies on implicit time stepping for external mode whereas ROMS uses a specially selected split-explicit method (see, e.g. Soufflet et al. (2016)) which is less dissipative. However, maps of simulated EKE and its seasonal cycle (Figure 5) reveal that FESOM has a higher energy level in the WSC than ROMS, and in contrast, the energy level in the EGC is higher in ROMS than in FESOM. This is also reflected in the horizontal kinetic energy spectra (Figure 4). Therefore, there could be certain energy dissipation in ROMS, the source of which is not identified. This can be the case for the WSC region considering that the baroclinic energy conversion to EKE is even stronger in ROMS (Figures 12c,d). There might be other reasons for the difference in the simulated EKE in certain regions between the two models. In particular, a higher EKE level in western Fram Strait in ROMS might be related to the difference in the simulated sea ice. Sea ice could damp eddies through the ocean-ice stress.

Apart from the differences, both models do show a high similarity in eddy properties such as eddy lifetime, size, pathways and travel distance (Figures 7, 8 and 9). In addition, both models exhibit a very similar pattern in barotropic energy conversion in eastern Fram Strait (Figure 12a,b). The degree of similarity is quite surprising, given that FESOM uses z-levels in the vertical whereas ROMS relies on terrain-following coordinates, which might lead to differences in topographic steering. Topography is bi-linearly interpolated to grid points and only smoothed over the 2d-stencil of nearest vertices in FESOM. In contrast, ROMS requires a smoother bathymetry.

6.3 Implications for contributing to future model development

Our instability analysis indicates that the GM parameterisation (Gent and McWilliams, 1990; Griffies, 1998) traditionally used in coarse resolution climate models does not fully account for the effect of eddies. This parameterisation accounts for eddy-induced flattening of isopycnals, it thus parameterises the effect of baroclinic instability. However, our analysis shows that barotropic instability plays an important role in some regions, too. In particular, there are areas with conversion from EKE to MKE (blue patches in Figure 12a,b) indicating a strengthening of the mean flow, which is not taken into account by the GM

parameterisation. Furthermore, over sloping bottom topography, the interaction of mesoscale eddies with the mean flow will be governed by a balance between the dissipation of APE and the homogenization of potential vorticity (Adcock and Marshall, 2000). Hence, it has been shown that interactions with sloping topography may locally increase the APE, e.g. by lifting dense water upward along the continental slope in Fram Strait as shown by Tverberg and Nøst (2009). Our analysis shows consistent patches of such reversed APE to EKE conversion along the Svalbard continental shelf break in both FESOM and ROMS, corroborating these theoretical considerations, which indicate that the GM parameterization that traditionally is used in coarse resolution climate models does not fully account for the effect of eddies. A similar result was shown recently in the study by Lüschow et al. (2019) investigating the vertical structure of the Atlantic deep western boundary current (DWBC). They find that below the core of the DWBC, eddy fluxes steepen isopycnals and thus feed potential energy to the mean flow, which is not represented in the GM framework.

7 Conclusions

Based on the results of two eddy-resolving ocean-sea ice models, ROMS and FESOM, we examined the properties and generation mechanisms of mesoscale eddies in Fram Strait. We found that the models agree with each other with respect to the modelled circulation, hydrography and eddy characteristics. They simulate rather short-living eddies (lifetime is on average 10–11 days), with a very slight dominance of cyclones (ROMS: 54%, FESOM: 55%). Cyclones and anticyclones show very distinct travel pathways, e.g., cyclones generated on the shallow Svalbard shelf tend to stay there, whereas anticyclones tend to travel offshore into the deep basin. More anticyclones tend to be trapped in deep depressions. Mean eddy radius is 5.0–6.0 km, which compares well with the first baroclinic Rossby radius of deformation in this region. On average, eddies travel around 35 km in both models. Eddy cores are located at about 100 m depth on average. Cyclones are predominantly cold eddies, while anticyclones are predominantly warm eddies.

The models also agree on mechanisms driving eddy generation, with consistent patterns of conversions to EKE from the mean kinetic and eddy available potential energies. The small size of eddies explains why a very high (1 km or finer) resolution is needed to simulate them. The good agreement on eddy generation and properties despite the very different numerics of FESOM (unstructured horizontal grid with vertical z-levels) and ROMS (regular horizontal grid with a terrain following vertical coordinate) gives us confidence in their ability to realistically simulate eddy processes. [The similarities of the simulated eddy fields also provide confidence in the eddy properties presented in this paper.](#) Some differences between the two models are also identified in this work, including the intensity of eddies and the rates of energy conversion, which require more dedicated research to better understand the reasons.

Author contributions. CW, QW and SD contributed the FESOM data and analysis, TH and LC contributed the ROMS data, and WJvA contributed the observational data. All authors discussed the content of the manuscript and contributed to the interpretation of data and writing of the manuscript.

Competing interests. We declare that no competing interests are present.

465 *Acknowledgements.* This work was supported by the AWI FRAM (FRontiers in Arctic marine Monitoring) program (CW and WJvA). TH acknowledges financial support from the Norwegian Research Council project 280727. QW is supported by the German Helmholtz Climate Initiative REKLIM (Regional Climate Change). LC acknowledges support from the Fram Centre "Arctic Ocean" flagship project "Mesoscale modeling of Ice, Ocean, and Ecology of the Arctic Ocean (ModOIE)" and Office of Naval Research grant number N00014-18-1-2694. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under
470 Grant No. DGE-1762114. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation. The FESOM simulation was performed at the North-German Supercomputing Alliance (HLRN). The ROMS simulation was performed under the NOTUR hpc project nn9238k. We are grateful to the two anonymous reviewers and the editor for their constructive comments.

References

- 475 Adcock, S. T. and Marshall, D. P.: Interactions between Geostrophic Eddies and the Mean Circulation over Large-Scale Bottom Topography, *Journal of Physical Oceanography*, 30, 3223–3238, [https://doi.org/10.1175/1520-0485\(2000\)030<3223:IBGEAT>2.0.CO;2](https://doi.org/10.1175/1520-0485(2000)030<3223:IBGEAT>2.0.CO;2), 2000.
- Bashmachnikov, I. L., Kozlov, I. E., Petrenko, L. A., Glok, N. I., and Wekerle, C.: Eddies in the North Greenland Sea and Fram Strait From Satellite Altimetry, SAR and High-Resolution Model Data, *Journal of Geophysical Research: Oceans*, 125, <https://doi.org/10.1029/2019JC015832>, 2020.
- 480 Beszczynska-Möller, A., Fahrback, E., Schauer, U., and Hansen, E.: Variability in Atlantic water temperature and transport at the entrance to the Arctic Ocean, 1997–2010, *ICES J. Mar. Sci.*, 69, 852–863, <https://doi.org/10.1093/icesjms/fss056>, 2012.
- Budgell, W. P.: Numerical simulation of ice-ocean variability in the Barents Sea region, *Ocean Dynamics*, 55, 370–387, <https://doi.org/10.1007/s10236-005-0008-3>, 2005.
- Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Progress in Oceanography*, 91, 167
485 – 216, <https://doi.org/10.1016/j.pocean.2011.01.002>, 2011.
- Cherian, D. A. and Brink, K. H.: Shelf Flows Forced by Deep-Ocean Anticyclonic Eddies at the Shelf Break, *Journal of Physical Oceanography*, 48, 1117–1138, <https://doi.org/10.1175/JPO-D-17-0237.1>, 2018.
- Crews, L., Sundfjord, A., Albreetsen, J., and Hattermann, T.: Mesoscale Eddy Activity and Transport in the Atlantic Water Inflow Region North of Svalbard, *Journal of Geophysical Research: Oceans*, 123, 201–215, <https://doi.org/10.1002/2017JC013198>, 2018.
- 490 Crews, L., Sundfjord, A., and Hattermann, T.: How the Yermak Pass Branch Regulates Atlantic Water Inflow to the Arctic Ocean, *Journal of Geophysical Research: Oceans*, 124, 267–280, <https://doi.org/10.1029/2018JC014476>, 2019.
- Cushman-Roisin, B.: *Introduction to Geophysical Fluid Dynamics*, Prentice Hall, 1994.
- Dai, A., Qian, T., Trenberth, K. E., and Milliman, J. D.: Changes in Continental Freshwater Discharge from 1948 to 2004, *J. Climate*, 22, 2773–2792, <https://doi.org/10.1175/2008JCLI2592.1>, 2009.
- 495 Danilov, S., Wang, Q., Timmermann, R., Iakovlev, N., Sidorenko, D., Kimmritz, M., Jung, T., and Schröter, J.: Finite-Element Sea Ice Model (FESIM), version 2, *Geosci. Model Dev.*, 8, 1747–1761, <https://doi.org/10.5194/gmd-8-1747-2015>, 2015.
- de Steur, L., Hansen, E., Gerdes, R., Karcher, M., Fahrback, E., and Holfort, J.: Freshwater fluxes in the East Greenland Current: A decade of observations, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2009GL041278>, 2009.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer,
500 P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- 505 Egbert, G. D. and Erofeeva, S. Y.: Efficient Inverse Modeling of Barotropic Ocean Tides, *Journal of Atmospheric and Oceanic Technology*, 19, 183–204, [https://doi.org/10.1175/1520-0426\(2002\)019<0183:EIMOBO>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2), 2002.
- Eldevik, T., Nilsen, J. E. Ø., Iovino, D., Olsson, K. A., Sandø, A. B., and Drange, H.: Observed sources and variability of Nordic Seas overflow, *Nature Geoscience*, 2, 406–410, <https://doi.org/10.1038/NGEO518>, 2009.
- Fer, I., Müller, M., and Peterson, A. K.: Tidal forcing, energetics, and mixing near the Yermak Plateau, *Ocean Science*, 11, 287–304,
510 <https://doi.org/10.5194/os-11-287-2015>, 2015.

- Fer, I., Bosse, A., and Dugstad, J.: Norwegian Atlantic Slope Current along the Lofoten Escarpment, *Ocean Science*, 16, 685–701, <https://doi.org/10.5194/os-16-685-2020>, 2020.
- Gent, P. and McWilliams, J.: Isopycnal Mixing in Ocean Circulation Models, *J. Phys. Oceanogr.*, 20, 150–155, 1990.
- Griffies, S.: The Gent-McWilliams Skew Flux, *J. Phys. Oceanogr.*, 28, 831–841, [https://doi.org/10.1175/1520-0485\(1998\)028<0831:TGMSF>2.0.CO;2](https://doi.org/10.1175/1520-0485(1998)028<0831:TGMSF>2.0.CO;2), 1998.
- 515 Gula, J., Molemaker, M. J., and McWilliams, J. C.: Gulf Stream Dynamics along the Southeastern U.S. Seaboard, *J. Phys. Oceanogr.*, 45, 690–715, <https://doi.org/10.1175/JPO-D-14-0154.1>, 2015.
- Haidvogel, D., Arango, H., Budgell, W., Cornuelle, B., Curchitser, E., Lorenzo, E. D., Fennel, K., Geyer, W., Hermann, A., Lanerolle, L., Levin, J., McWilliams, J., Miller, A., Moore, A., Powell, T., Shchepetkin, A., Sherwood, C., Signell, R., Warner, J., and Wilkin, J.:
520 Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the Regional Ocean Modeling System, *Journal of Computational Physics*, 227, 3595 – 3624, <https://doi.org/10.1016/j.jcp.2007.06.016>, 2008.
- Hallberg, R.: Using a resolution function to regulate parameterizations of oceanic mesoscale eddy effects, *Ocean Modelling*, 72, 92 – 103, <https://doi.org/10.1016/j.ocemod.2013.08.007>, 2013.
- Hattermann, T., Isachsen, P. E., von Appen, W.-J., Albrechtsen, J., and Sundfjord, A.: Eddy-driven recirculation of Atlantic Water in Fram
525 Strait, *Geophys. Res. Lett.*, 43, <https://doi.org/10.1002/2016GL068323>, 2016.
- Isern-Fontanet, J., Garcia-Ladona, E., and Font, J.: Vortices of the Mediterranean Sea: An Altimetric Perspective, *Journal of Physical Oceanography*, 36, 87–103, <https://doi.org/10.1175/JPO2826.1>, 2006.
- Johannessen, J. A., Johannessen, O. M., Svendsen, E., Shuchman, R., Manley, T., Campbell, W. J., Josberger, E. G., Sandven, S., Gascard, J. C., Olausen, T., Davidson, K., and Van Leer, J.: Mesoscale eddies in the Fram Strait marginal ice zone during the 1983 and 1984
530 Marginal Ice Zone Experiments, *J. Geophys. Res. Oceans*, 92, 6754–6772, <https://doi.org/10.1029/JC092iC07p06754>, 1987a.
- Johannessen, O. M., Johannessen, J. A., Svendsen, E., Shuchman, R. A., Campbell, W. J., and Josberger, E.: Ice-Edge Eddies in the Fram Strait Marginal Ice Zone, *Science*, 236, 427–429, <https://doi.org/10.1126/science.236.4800.427>, 1987b.
- Kang, D. and Curchitser, E. N.: Gulf Stream eddy characteristics in a high-resolution ocean model, *Journal of Geophysical Research: Oceans*, 118, 4474–4487, <https://doi.org/10.1002/jgrc.20318>, 2013.
- 535 Kawasaki, T. and Hasumi, H.: The inflow of Atlantic water at the Fram Strait and its interannual variability, *J. Geophys. Res. Oceans*, 121, 502–519, <https://doi.org/10.1002/2015JC011375>, 2016.
- Large, W. and Yeager, S.: The global climatology of an interannually varying air-sea flux data set, *Clim. Dynam.*, 33, 341–364, <https://doi.org/10.1007/s00382-008-0441-3>, 2008.
- Lüschow, V., Storch, J.-S. v., and Marotzke, J.: Diagnosing the Influence of Mesoscale Eddy Fluxes on the Deep Western Boundary Current in
540 the 1/10° STORM/NCEP Simulation, *Journal of Physical Oceanography*, 49, 751–764, <https://doi.org/10.1175/JPO-D-18-0103.1>, 2019.
- Mahadevan, A.: The Impact of Submesoscale Physics on Primary Productivity of Plankton, *Annual Review of Marine Science*, 8, 161–184, <https://doi.org/10.1146/annurev-marine-010814-015912>, 2016.
- Manucharyan, G. E. and Thompson, A. F.: Submesoscale Sea Ice-Ocean Interactions in Marginal Ice Zones, *Journal of Geophysical Research: Oceans*, 122, 9455–9475, <https://doi.org/10.1002/2017JC012895>, 2017.
- 545 Martínez-Moreno, J., Hogg, A. M., Kiss, A. E., Constantinou, N. C., and Morrison, A. K.: Kinetic Energy of Eddy-Like Features From Sea Surface Altimetry, *Journal of Advances in Modeling Earth Systems*, 11, 3090–3105, <https://doi.org/10.1029/2019MS001769>, 2019.
- Morrow, R., Birol, F., Griffin, D., and Sudre, J.: Divergent pathways of cyclonic and anti-cyclonic ocean eddies, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL020974>, 2004.

- Nencioli, F., Dong, C., Dickey, T., Washburn, L., and McWilliams, J. C.: A Vector Geometry-Based Eddy Detection Algorithm and Its
550 Application to a High-Resolution Numerical Model Product and High-Frequency Radar Surface Velocities in the Southern California
Bight, *Journal of Atmospheric and Oceanic Technology*, 27, 564–579, <https://doi.org/10.1175/2009JTECHO725.1>, 2010.
- Okubo, A.: Horizontal dispersion of floatable particles in the vicinity of velocity singularities such as convergences, *Deep Sea Research and
Oceanographic Abstracts*, 17, 445 – 454, [https://doi.org/10.1016/0011-7471\(70\)90059-8](https://doi.org/10.1016/0011-7471(70)90059-8), 1970.
- Olbers, D., Willebrand, J., and Eden, C.: *Ocean Dynamics*, Springer-Verlag Berlin Heidelberg, <https://doi.org/10.1007/978-3-642-23450-7>,
555 2012.
- Olson, D. B.: Rings in the ocean, *Annual Review of Earth and Planetary Sciences*, 19, 283–311, 1991.
- Polyakov, I. V., Pnyushkov, A. V., and Timokhov, L.: Warming of the Intermediate Atlantic Water of the Arctic Ocean in the 2000s, *J.
Climate*, 25, 8362–8370, <https://doi.org/10.1175/JCLI-D-12-00266.1>, 2012.
- Raj, R. P., Johannessen, J. A., Eldevik, T., Nilsen, J. E. A., and Halo, I.: Quantifying mesoscale eddies in the Lofoten Basin, *Journal of
560 Geophysical Research: Oceans*, 121, 4503–4521, <https://doi.org/10.1002/2016JC011637>, 2016.
- Rudels, B.: The thermohaline circulation of the Arctic Ocean and the Greenland Sea, *Philosophical Transactions of the Royal Society of
London A: Mathematical, Physical and Engineering Sciences*, 352, 287–299, <https://doi.org/10.1098/rsta.1995.0071>, 1995.
- Schaffer, J., von Appen, W.-J., Dodd, P. A., Hofstede, C., Mayer, C., de Steur, L., and Kanzow, T.: Warm water path-
ways toward Nioghalvfjerdingsfjorden Glacier, Northeast Greenland, *Journal of Geophysical Research: Oceans*, 122, 4004–4020,
565 <https://doi.org/10.1002/2016JC012462>, 2017.
- Schauer, U., Fahrbach, E., Østerhus, S., and Rohardt, G.: Arctic warming through the Fram Strait: Oceanic heat transport from 3 years of
measurements, *J. Geophys. Res.*, 109, 2004.
- Shchepetkin, A. F. and McWilliams, J. C.: The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-
following-coordinate oceanic model, *Ocean Modelling*, 9, 347 – 404, <https://doi.org/10.1016/j.ocemod.2004.08.002>, 2005.
- 570 Shchepetkin, A. F. and McWilliams, J. C.: Correction and commentary for "Ocean forecasting in terrain-following coordinates: Formu-
lation and skill assessment of the regional ocean modeling system" by Haidvogel et al., *J. Comp. Phys.* 227, pp. 3595–3624, *Journal of
Computational Physics*, 228, 8985 – 9000, <https://doi.org/10.1016/j.jcp.2009.09.002>, 2009.
- Smith, D. C., Morison, J. H., Johannessen, J. A., and Untersteiner, N.: Topographic generation of an eddy at the edge of the East Greenland
Current, *J. Geophys. Res. Oceans*, 89, 8205–8208, <https://doi.org/10.1029/JC089iC05p08205>, 1984.
- 575 Soufflet, Y., Marchesiello, P., Lemarié, F., Jouanno, J., Capet, X., Debreu, L., and Benshila, R.: On effective resolution in ocean models,
Ocean Modelling, 98, 36 – 50, <https://doi.org/10.1016/j.ocemod.2015.12.004>, 2016.
- Spall, M. A. and Pedlosky, J.: Lateral Coupling in Baroclinically Unstable Flows, *Journal of Physical Oceanography*, 38, 1267–1277,
<https://doi.org/10.1175/2007JPO3906.1>, 2008.
- Storkey, D., Blockley, E. W., Furner, R., Guivarc’h, C., Lea, D., Martin, M. J., Barciela, R. M., Hines, A., Hyder, P., and Sid-
580 dorn, J. R.: Forecasting the ocean state using NEMO: The new FOAM system, *Journal of Operational Oceanography*, 3, 3–15,
<https://doi.org/10.1080/1755876X.2010.11020109>, 2010.
- Teigen, S. H., Nilsen, F., Skogseth, R., Gjevik, B., and Beszczynska-Möller, A.: Baroclinic instability in the West Spitsbergen Current, *J.
Geophys. Res. Oceans*, 116, <https://doi.org/10.1029/2011JC006974>, 2011.
- Tverberg, V. and Nøst, O. A.: Eddy overturning across a shelf edge front: Kongsfjorden, west Spitsbergen, *Journal of Geophysical Research:*
585 *Oceans*, 114, <https://doi.org/10.1029/2008JC005106>, 2009.

- Volkov, D. L., Kubryakov, A. A., and Lumpkin, R.: Formation and variability of the Lofoten basin vortex in a high-resolution ocean model, *Deep Sea Research Part I*, 105, 142–157, <https://doi.org/10.1016/j.dsr.2015.09.001>, 2015.
- von Appen, W.-J., Schauer, U., Hattermann, T., and Beszczynska-Möller, A.: Seasonal Cycle of Mesoscale Instability of the West Spitsbergen Current, *J. Phys. Oceanogr.*, 46, 1231–1254, <https://doi.org/10.1175/JPO-D-15-0184.1>, 2016.
- 590 von Appen, W.-J., Wekerle, C., Hehemann, L., Schourup-Kristensen, V., Konrad, C., and Iversen, M. H.: Observations of a Submesoscale Cyclonic Filament in the Marginal Ice Zone, *Geophysical Research Letters*, 45, 6141–6149, <https://doi.org/10.1029/2018GL077897>, 2018.
- von Appen, W.-J., Beszczynska-Möller, A., Schauer, U., and Fahrbach, E.: Physical oceanography and current meter data from moorings F1-F14 and F15/F16 in the Fram Strait, 1997-2016, <https://doi.org/10.1594/PANGAEA.900883>, 2019.
- Wang, Q., Danilov, S., and Schröter, J.: Finite element ocean circulation model based on triangular prismatic elements, with application in
595 studying the effect of topography representation, *J. Geophys. Res.*, 113, 2008.
- Wang, Q., Danilov, S., Sidorenko, D., Timmermann, R., Wekerle, C., Wang, X., Jung, T., and Schröter, J.: The Finite Element Sea Ice-Ocean Model (FESOM) v.1.4: formulation of an ocean general circulation model, *Geosci. Model Dev.*, 7, 663–693, <https://doi.org/10.5194/gmd-7-663-2014>, 2014.
- Wang, Q., Koldunov, N. V., Danilov, S., Sidorenko, D., Wekerle, C., Scholz, P., Bashmachnikov, I. L., and Jung, T.: Eddy Kinetic Energy in
600 the Arctic Ocean from a Global Simulation with a 1-km Arctic, *Geophysical Research Letters*, <https://doi.org/10.1029/2020GL088550>, 2020.
- Weiss, J.: The dynamics of enstrophy transfer in two-dimensional hydrodynamics, *Physica D: Nonlinear Phenomena*, 48, 273 – 294, [https://doi.org/10.1016/0167-2789\(91\)90088-Q](https://doi.org/10.1016/0167-2789(91)90088-Q), 1991.
- Wekerle, C., Wang, Q., von Appen, W.-J., Danilov, S., Schourup-Kristensen, V., and Jung, T.: Eddy-Resolving Simulation of the Atlantic
605 Water Circulation in the Fram Strait With Focus on the Seasonal Cycle, *J. Geophys. Res. Oceans*, <https://doi.org/10.1002/2017JC012974>, 2017.
- Wilson, N., Straneo, F., and Heimbach, P.: Satellite-derived submarine melt rates and mass balance (2011–2015) for Greenland’s largest remaining ice tongues, *The Cryosphere*, 11, 2773–2782, <https://doi.org/10.5194/tc-11-2773-2017>, 2017.

Table 1. Characteristics of the FESOM and ROMS configurations used in this study.

	FESOM	ROMS
Numerical method	finite elements	finite differences
Horizontal grid	A-grid (P1-P1 scheme)	C-grid
Vertical coordinate	z-levels	terrain following levels
Domain	global	regional
Horizontal mixing scheme	biharmonic Smagorinsky	biharmonic
Vertical mixing scheme	KPP	KPP
Tides	no	yes

Table 2. Mean properties and their standard deviation (in brackets) for all eddies generated in the area 8°W – 20°E / 76°N – 82°N in the years 2006–2009 in ROMS and FESOM.

Eddy type	Radius (km)	Abs. rel. vorticity (normalised by f)	Lifetime (days)	Travel distance (km)
<i>ROMS</i>				
All eddies	4.9 (2.8)	0.40 (0.22)	10 (14)	34 (44)
Cyclones	4.6 (2.6)	0.41 (0.25)	10 (12)	33 (38)
Anticyclones	5.2 (3.0)	0.39 (0.18)	10 (16)	35 (51)
<i>FESOM</i>				
All eddies	5.6 (3.3)	0.28 (0.18)	11 (16)	35 (44)
Cyclones	5.3 (3.1)	0.29 (0.19)	11 (15)	35 (43)
Anticyclones	6.0 (3.4)	0.27 (0.16)	11 (16)	35 (45)




Figure 1. Bathymetry of the Arctic Ocean (left, red box indicates our study region) and of the Fram Strait (right). Coloured polygons in right panel indicate regions used for analysis: Svalbard shelf (green), West Spitsbergen Current (cyan), central southern Fram Strait (yellow), central northern Fram Strait (blue), and Yermak and Svalbard Branch (magenta). Coloured dots indicate moorings deployed across Fram Strait at $78^{\circ}50'N$; red and magenta dots show moorings used to compute velocity time series representative for the EGC and WSC, respectively. Black arrows show major currents in the Fram Strait (WSC: West Spitsbergen Current, EGC: East Greenland Current, YB: Yermak Branch, SB: Svalbard Branch). MD and YP indicate the locations of the Molloy Deep and the Yermak Plateau, respectively.




Figure 2. Simulated relative vorticity at 100 m depth on Jan 1st, 2006 for ROMS (a) and FESOM (b). Simulated Okubo-Weiss parameter (s^{-2}) at 100 m depth on Jan 1st, 2006 in a region west of Svalbard (grey box in the top panel) for ROMS (c) and FESOM (d). Shown are only values with $OW < -0.2\sigma_{OW}$, where σ_{OW} is the spatial standard deviation of OW at that day. Red arrows show the velocity, with only every 8th vector plotted. Cyan and magenta contours show anticyclonic and cyclonic eddies respectively identified by the Nencioli algorithm (Nencioli et al., 2010). Grey contour lines indicate bathymetry at 1000 m intervals.




Figure 3. Temperature (a, b) and salinity (c, d) at 100 m depth averaged over the time period 2006–2009 simulated by ROMS (left) and FESOM (right). Black contour lines show the 1°C and 2°C isotherms and the 34 and 35 isohalines. Dots show mooring measurements in 75 m depth for the same time period (von Appen et al., 2019). Grey contour lines indicate bathymetry at 1000 m intervals. T/S diagram of simulated temperature and salinity in 100 m depth in the region 10°W–20°E / 76°N–82°N in ROMS (e) and FESOM (f). The colour shading indicates the longitude of data points.




Figure 4. Power density spectra of [horizontal kinetic energy](#) from daily averaged velocity in 75 m depth in the (a) West Spitsbergen Current and (b) East Greenland Current from mooring measurements (blue), and models FESOM (red) and ROMS (yellow), [computed as the sum of spectra of \$u\$ and \$v\$ components divided by 2](#). Thick lines indicate slopes of the spectra, and the shaded area indicates the 95% confidence intervals.




Figure 5. Eddy kinetic energy (EKE). Top panel: Maps of EKE at 100 m depth from (a) FESOM and (b) ROMS for the years 2006–2009. Gray contour lines indicate bathymetry at 1000 m intervals. Bottom panel: Seasonal cycle of EKE at 75 m depth across Fram Strait at 78°50'N from (a) mooring measurements (von Appen et al., 2019), (b) FESOM, and (c) ROMS for the years 2006–2009.




Figure 6. (a,b) Total number of eddy occurrences in the years 2006–2009 for (left) ROMS and (right) FESOM, binned in a $1/24^\circ$ grid and smoothed with a 3 point Hanning window kernel. (c,d) Difference between numbers of cyclonic and anticyclonic eddies (cyclones minus anticyclones). Black contour lines indicate bathymetry at 1000 m intervals and at 200 m.




Figure 7. Histogram of (a) radius, (b) maximum relative vorticity normalised by f , (c) water depth, (d) eddy nonlinearity parameter U/c , (e) eddy lifetime and (f) travel distance for anticyclonic (blue) and cyclonic eddies (red) normalised by the number of eddies/tracks tracked in the area $8^{\circ}\text{W}-20^{\circ}\text{E}/76^{\circ}\text{N}-82^{\circ}\text{N}$ in the years 2006–2009 in ROMS (dark colours) and FESOM (light colours).




Figure 8. Eddy tracks of cyclones (red lines, a) and anticyclones (blue lines, b) with lifetimes of more than 30 days that are generated in five different regions indicated by coloured polygons, see Fig. 1, detected in simulation ROMS from 2006–2009. Light and dark colours of the lines indicate the beginning and end of the track, respectively.




Figure 9. The same as Figure 8, but for FESOM.




Figure 10. Vertical extent of eddies tracked in 100 m depth during the years 2006–2009 in ROMS (dark colours) and FESOM (light colours) with lifetimes >30 days for cyclones (red) and anticyclones (blue) generated in regions a) Svalbard shelf, b) West Spitsbergen Current, c) southern central Fram Strait, d) northern central Fram Strait and e) Yermak and Svalbard Branch. Left, middle and right panels show relative vorticity, temperature anomaly and salinity anomaly, respectively. Anomalies are calculated by taking the value in the eddy centre relative to the mean value of the month.




Figure 11. Maps of averaged nonlinearity parameter U/c , where U and c are maximum rotational and translation speeds, respectively, for eddies detected between 2006–2009 in a) ROMS and b) FESOM. Values of U/c were averaged on a 1° longitude x 0.2° latitude grid. Gray contours show bathymetry contours at 1000 m intervals.




Figure 12. Simulated depth integrated energy transfer from (a,b) mean kinetic to eddy kinetic energy (HRS, product of horizontal Reynolds stress and mean shear, $\int_H \left(-\overline{\mathbf{u}'\mathbf{u}'} \cdot \frac{\partial \overline{\mathbf{u}}}{\partial x} - \overline{\mathbf{u}'\mathbf{v}'} \cdot \frac{\partial \overline{\mathbf{u}}}{\partial y} \right) dz$) and (c,d) eddy available potential to eddy kinetic energy (vertical eddy buoyancy flux, $\int_H \overline{w'b'} dz$) averaged for 2006–2009 in (a,c) ROMS and (b,d) FESOM. Black contours show bathymetry contours at 1000 m intervals and at 200 m and 500 m depth.




Figure 13. a) Topographic $\beta = -\frac{f}{H}\nabla H$ and b) $\beta - \partial_{xx}v$, where v is the simulated meridional velocity from FESOM. The second derivative of v is computed from monthly means, and then averaged over the years 2006–2009. A change of sign of $\beta - \partial_{xx}v$ is a necessary condition for barotropic instability. Black contours show bathymetry contours at 1000 m intervals and at 250 m and 500 m depth. Note that values in b) are only shown in the vicinity of the WSC main pathway (within a distance of 50 km to the 250 m isobath).




Figure 14. Ertel potential vorticity (left panel) and its gradient in zonal direction (right panel) across Fram Strait at (a) 78°N and (b) 78°50'N computed from long-term mean FESOM data (2006–2009). Black lines show simulated meridional velocity contours (0.1 and 0.2 m/s), and white lines show the simulated 27.9, 28, 28.1, 28.2, and 28.22 kg/m³ isopycnals.