

For both reviewers, the response is organised as follows:

- **Reviewer's comment in bold font;**
- Answer in plain text;
- *When relevant, new or modified text in italics (with line number for new text)*

Reviewer 1

The author sincerely thanks Reviewer 1 for their really helpful comments: not only did they point out issues, they also constructively suggested solutions. The role of the reviewers has been duly acknowledged:

L.554-555: "The author thanks the two anonymous reviewers whose comments greatly improved the quality of this manuscript"

1. While the 6x6 postage stamp figure format is quite useful, especially for the modelers, it may be a bit much for some readers. Please consider putting all the postage stamp figures into the appendix and switching the text figures to 3-panel figures (water property reference, mean or median difference of water property from reference, std deviation or interquartile range of water property difference from reference). This would keep 4 figures in the main text:

Two 3x3 figures (3 panels each of reference, mean, and standard deviation for potential temperature, practical salinity, and sigma-4) in the S. Ocean and N. Atlantic.

Two 1x3 figures of reference, mean differences, and standard deviations of AABW thickness and NADW thickness

This change would mean eight postage stamp "difference" figures in the appendix (NADW thickness, AABW thickness, and temper, salinity, and sigma-4 for both the Southern Ocean and North Atlantic.

This suggested change would also require some additional text to describe the means and standard deviations of the differences of the models from observations, but it might be worth it. The statistical figures would probably be much more widely used than the postage stamp figures. They would be great in a general presentation on climate model evaluation.

This comment from the reviewer echoes the first comment made by reviewer 2, that the target or even the aim of this paper was somewhat unclear. This paper was written with model users in mind, with the aim to be the go-to reference when they need to justify their choice of models for their study. As I had to do at that time for another study with the CMIP6 models. Consequently, I am reluctant to send the postage stamp figures to the appendix, as at least to me they are the most useful.

Nevertheless, I also really liked the reviewer's suggestion of what could be called "summary figures". So I created them, and in fact expanded on the reviewer's idea to make it two 4 x 4 figures with reference, multi model mean, mean difference in property from reference, and standard deviation, for density, temperature, salinity and mixed layer depth, in the Southern Ocean and North Atlantic, complete with the median value quoted on each panel. As the aim of this change is to make the paper

useful to as wide a range of readers as possible, I decided to include both the multi model mean and the mean difference in properties, even though they show the same thing.

These two figures now are in the text as (new) Figs. 2 and 4, and their description has been added in the text where relevant.

I chose to not add such figures for the AABW and NADW thickness, as the results of these postage stamps figures are already summarised in Tables B1 and B2.

Major comments:

2. L1. The first sentence is arguably inaccurate, as the global ocean circulation is mostly wind driven. Maybe “Deep and bottom water formation are an important part of the global ocean circulation” would be better.

Sentence changed to:

“Deep and bottom water formation are crucial components of the global ocean circulation”

3. Throughout. Please use the commonly accepted terminology “bottom water” to distinguish Antarctic Bottom Water from “deep water” (North Atlantic Deep Water). The manuscript may be slightly longer as a result (as sometimes the use of “deep and bottom water” will be required), but will be more in line with the field, and easier to read.

Terminology changed throughout the manuscript to:

“deep and bottom water” when referring to both AABW and NADW in the same sentence;

“bottom water formation” when referring to AABW only.

4. L19. The opening sentence of the introduction doesn’t seem accurate, and it is difficult to parse the meaning. How about “Bottom water formation around Antarctica and deep water formation in the North Atlantic ventilate the global abyssal and deep ocean.”?

Sentence modified as suggested.

5. L23. This sentence does not seem quite right, as AABW influence spreads quite far into the N. Atlantic, albeit in a form highly diluted by mixing with NADW above. How about “In a substantial portion of the Atlantic, Antarctic Bottom Water spreading north is overlain by North Atlantic Deep Water spreading south.”

Sentence modified as suggested.

6. L63-65. This sentence conflates two different issues, and perhaps should be broken in two at the “and”.

Sentence modified to clarify how the two issues are related:

“Furthermore, as some models are not fully independent as they share similar codes (Table 1), using different ensemble sizes would have accentuated the bias towards one model family”

7. L140-142. This portion is overly concise. How about something along the lines of: “Presently, Antarctic Bottom water is primarily formed in several locations (including the Weddell Sea, the Ross Sea, and the Adelie Lands) as water is cooled, made saltier, and denser on the continental

shelves, then cascades down the continental slopes, entraining deep waters on its way to the sea floor.”

The sentence suggested by the reviewer has been added.

8. L160-161. The recent Polynya in the Weddell Sea that is mentioned later in the paper should also be mentioned here.

Sadly, the recent Maud Rise polynya was not associated with any deep convection (e.g. Campbell et al., 2019), so it cannot be mentioned here.

9. Tables 2 and 3. Consider that instead of years out of 30, a percentage would be more quickly and easily interpreted by most readers.

I agree with the reviewer that it would be quicker and easier, but it would also be misleading. The CMIP6 runs contain more than 100 years; presenting these values as a percentage would imply for many readers that I worked with 100 years. Hence I prefer keeping the values in the table as “years out of the 30 studied”.

10. L205. How is it possible that “No model has regional biases”? Perhaps just delete the part of this sentence on L206, and start the sentence with “CMIP6 models tend to be either biased light or biased dense”.

Here I tried to mean that unlike some models in CMIP5, there is no difference in the sign of the bias between e.g. the Weddell and Ross seas. I cannot find a better reformulation than what the reviewer suggests, so the sentence was modified as suggested.

11. L265. Consider changing “The picture is less grim regarding bottom property biases” to “CMIP6 water property biases at the bottom of the North Atlantic are smaller than those at the bottom of the Southern Ocean”.

Sentence modified as suggested.

12. L298. Here “convenient” seems like the wrong word. This reviewer can’t guess at what to suggest for a replacement.

The word has been removed:

“a strong interhemispheric correlation”

13. L329-330. Consider changing “are rather accurately represented” to “often agree within observational uncertainties”.

The agreement with observational uncertainties is mentioned several times in the previous sentences. The last sentence “in summary, [they] are rather accurately represented” stays.

14. L469-470. This is a problematic sentence, because NADW does not occupy the global ocean, and in the real world the signatures of upper (subpolar gyre) and lower (GIN sea overflow) NADW are both traceable for substantial distances from their formation regions. How about “NADW formed in the subpolar gyre of the models clearly spreads southward, but the signature of the portions formed in the Nordic seas is less evident.”?

Sentence modified as suggested

Minor comments (typos, debatable word choices, and grammatical errors):

All addressed

Reviewer 2

The author thanks Reviewer 2 for their scientific comments. The role of the reviewers has been duly acknowledged:

L.554-555: "The author thanks the two anonymous reviewers whose comments greatly improved the quality of this manuscript"

The response is organised as follows: first, the comment from the reviewer; then, my answer; finally, when relevant, new or modified text.

The paper presents a comparison of time mean fields between new, CMIP6 models and observations. The focus is the formation and distribution of deep water formed in the North Atlantic and Southern Ocean. The aim (I assume) is to document these biases as a basis for further study. In general, I did not feel that the paper presented much that was convincing in terms of new scientific interpretation. [...] Nonetheless, the potential lack of new understanding is not necessarily a strong negative as documenting the models can be a useful exercise in and of itself.

Both reviewers raised this important point: the target audience and aim of the paper were not as clearly defined as I thought. This paper is indeed designed as a reference for model users to justify their choice of models for further studies. Although attempts are made at explaining these biases, the emphasis is on quantifying these biases.

This sentence was added to the introduction to clarify this objective, lines 51-52:

"The primary objective of this paper is to quantify and discuss biases of each model, so that model users can make informed model selections."

In particular many of the reported correlations seem small given that the many of the models are not very independent, which I don't think has been accounted for or even acknowledged.

The lack of independence of the models is acknowledged as early as line 67 (slightly modified from the previous version in response to a comment by reviewer 1):

"Furthermore, as some models are not fully independent as they share similar codes (Table 1)..."

In response to the reviewer's comment, this methodological clarification was also added line 68:

"To account for this lack of independence, the correlations quoted throughout the text have been verified with different model numbers"

Scientific issues:

L77-78. Why is a different threshold used to the observations? How can you then fairly compare with the observations? Please explain this.

I chose neither threshold. The threshold of the models is the “official” threshold of the CMIP6 procedure; that is, models that wanted to participate in CMIP6 had to use that threshold. Likewise, the threshold used in observations is the one that was chosen by the creators of this observational product. The literature on the impact of one threshold rather than another is plentiful (I even wrote a PhD dissertation chapter on this), and the conclusion is that for detecting spurious modelled deep convection, this difference is not critical. In fact, choosing a larger threshold for the models than for the observations means that we would underestimate deep convection in models. As the objective of this publication is primarily to compare models with each other, the most important is that all models use the same threshold.

The following was added to summarise this discussion, lines 80-88:

“As is requested for CMIP6, the MLD is then detected as the depth where σ_θ differs from that at 10 m depth by more than 0.125 kg m^{-3} . [new text associated with the new supplementary figure] Furthermore, a different threshold of 0.03 kg m^{-3} is used in the observational reference (de Boyer Montégut et al., 2004), which could lead to an underestimation of mixed layer depths in the models (as we show in section 3, it does not).”

L224-226. So what can we actually determine or learn from this? Is there a relationship, particularly after controlling for the fact that several of the models are nearly identical (or assimilate observations, which I am surprised is included as it seems fundamentally different to the other models)

Sentence expanded to make the point clearer:

“The models that convect the least or not at all tend to be the most accurate. For the CESM2 family, accurate bottom properties and lack of deep convection may both be the result of their overflow parameterisation (Briegleb et al., 2010; Snow et al., 2015). For another model, NorCPM1, the accuracy in all properties may come from its observation assimilation rather than accurate model physics (Counillon et al., 2016).”

L250-251. This link surely only makes sense if the climate sensitivity is driving the DMV. But previously you suggest the logic is the other way around (i.e. larger DMV can sequester more heat and thus reduce climate sensitivity). If the DMV drives the climate sensitivity, why should DMV in the Weddell Sea and SPG themselves be linked? Isn't it more likely that the DMV in these two regions correlate due to some global model bias?

Does the DMV impact the climate sensitivity, or does the sensitivity impact the DMV? Less mixing means less heat absorbed by the ocean, so more in the atmosphere and a larger sensitivity. The opposite is true: if the sensitivity is somehow controlled by another “global model bias”, a high sensitivity will lead to more ocean surface warming and stratification, and hence less mixing. As is obvious from just these two sentences, what we really have is a feedback loop, and investigating which comes first, or what that other global bias can be, are beyond the scope of this paper.

I added a sentence to reflect the point raised by the reviewer lines 277-279:

“As already mentioned, no causation can be inferred: deciphering whether global biases in DMV are responsible for the models' sensitivities, or in contrast sensitivities are set by other processes and impact the DMV, is beyond the scope of this analysis”

L364, L367. Are these correlations really robust given the real number of degrees of freedom is likely far fewer than the total number of models

All the correlations have been verified using different model numbers, in particular using only one member per family, and the results remained. This precision has been added to the Methods section, lines 68-69.

Minor issues:

To summarise, all the issues highlighted by the reviewer that impeded the understanding have been corrected. Only the ones for which a response longer than “corrected” was necessary are presented here. I leave it to the copy editor to decide whether grammar rules that we learnt at school can be bent in order to make the manuscript more dynamic and pleasant to read.

L1. “Deep water formation is the driver of the global ocean circulation” - on what timescale?

This sentence has already been modified in response to a comment by Reviewer 1.

L5. Large majority - can you be more quantitative?

Sentence modified to *“28 models in the Southern Ocean and all 35 models in the North Atlantic”*

L31. Is an accurate representation really needed for climate predictions? Over what time scales are you referring? Of what variable? Predictions in CMIP usually refers to initialised decadal simulations, whereas projections usually refer to century time scale uninitialised simulations.

I am not sure which point the reviewer is trying to make here. I changed “prediction” to “projection”, as I meant long term, IPCC-report type results.

Table 1. The horizontal resolution here doesn’t take into account any local grid refinement, which could also be noted

They could indeed, but too many cases would need to be considered to fit in the table. Instead, table caption has been modified to “nominal” grid resolution.

L58. Is this the actual variable name or is it mlotst?

The actual variable name is mlotst. There was a typo.

L58. For the models where you have MLD directly, can you show as a supplementary figure that your method and the online one are equivalent.

This is a very good idea. A new supplementary Figure (A1) has been added to show where and by how much they differ, along with a discussion in the Methods section.

L72. Is the deboyer montegut MLD data just to 2004 or is it updated?

It is updated, as is indicated on the data download website.

L104. “Hardly a third” is colloquial. Please be specific.

The number of models changed since the initial submission. Sentence now reads:

“it is provided by only 18 of the models (from 10 families)”

L173. “(Thin?)” - this made me a bit annoyed. If you want to hypothesize at the reason then please spell this out in a sentence rather than like this

I am confused by the reviewer’s reaction as the hypothesis is spelt out in that sentence. I removed the “(thin?)” that brought nothing to the sentence.

L183. Does the pipe physically suck the water in the model physics? Could the description of this parameterization be described more formally?

The following sentence was added lines 196-198:

“If the water on the shelf exceeds a critical density, a pipe artificially transports this dense water from the shelves to the deep basin. Without having to cascade, the dense shelf water keeps its properties.”

L243. Here (and elsewhere) the referencing is a bit unclear. You’ve already cited this paper, and here it seems as if you’re citing it as a reference showing a link between convection and climate sensitivity here.

I assume the reviewer meant the reference to Zelinka et al. (2020) of (previous version’s) line 248. Sentence modified to clarify that I refer to this paper as the source of the sensitivity values I used:

“There is a relationship with the climate sensitivities of Zelinka et al. (2020) though”

L257. Please define quantitatively what a “tolerable” bias is?

I do not understand what the reviewer means as the quantitative value is given in the same sentence, two words later.

L289. Is such a small correlation actually significant, especially when accounting for the limited degrees of freedom (i.e. many similar models)

This point has been addressed twice already in this response.

L297. “The question remains”. What question? Please specify.

Sentence now reads:

“the cause of NorCPM’s and other models’ bottom density bias in the GIN seas remains unknown”

L321. The AMOC can’t be said to be overestimated given the quoted uncertainty on both the model and observations.

Agreed. Sentence removed.

Antarctic Bottom Water and North Atlantic Deep Water in CMIP6 models

Céline Heuzé¹

¹Department of Earth Sciences, University of Gothenburg, Gothenburg, Sweden

Correspondence: Céline Heuzé (celine.heuze@gu.se)

Abstract. Deep ~~water formation is the driver~~ and bottom water formation are crucial components of the global ocean circulation, yet ~~it was~~ they were poorly represented in the previous generation of climate models. We here quantify biases in Antarctic Bottom Water (AABW) and North Atlantic Deep Water (NADW) formation, properties, transport, and global extent in 35 climate models that participated in the latest Climate Model Intercomparison Project (CMIP6). Several CMIP6 models are correctly forming AABW via shelf processes, but ~~in both hemispheres, the large majority of climate models form deep~~ 28 models in the Southern Ocean and all 35 models in the North Atlantic form deep and bottom water via open ocean deep convection, too deep, too often, over too large an area. Models that convect the least form the most accurate AABW, but the least accurate NADW. The four CESM2 models with their “~~pipe~~”-overflow parameterisation are among the most accurate models. In the Atlantic, the colder the AABW, the stronger the abyssal overturning at 30°S, and the further north the AABW layer extends. The saltier the NADW, the stronger the Atlantic Meridional Overturning Circulation (AMOC), and the further south the NADW layer extends. In the Indian and Pacific oceans in contrast, the fresher models are the ones ~~who~~ which extend the furthest regardless of the strength of their abyssal overturning, most likely because they also are the models with the weakest fronts in the Antarctic Circumpolar Currents. There are clear improvements since CMIP5: several CMIP6 models correctly represent or parameterise Antarctic shelf processes, fewer models exhibit Southern Ocean deep convection, more models con-
15 vect at the right location in the Labrador Sea, bottom density biases are reduced, and abyssal overturning is more realistic. But more improvements are required, e.g. by generalising the use of overflow parameterisations or by coupling to interactive ice sheet models, before deep and bottom water formation, and hence heat and carbon storage, are represented accurately.

Copyright statement. TEXT

1 Introduction

20 ~~At both poles,~~ Bottom water formation around Antarctica and deep water formation ~~sets the global ocean ventilation in motion in~~ the North Atlantic ventilate the global abyssal and deep ocean. Ocean-ice-atmosphere interactions by the Antarctic ice shelves (Orsi, 2010; Drucker et al., 2011; Ohshima et al., 2013) or, more rarely, in open ocean polynyas (Killworth, 1983; Campbell et al., 2019), create the coldest, densest ~~deep-water~~ water mass: the Antarctic Bottom Water (AABW). AABW does not stay

around Antarctica, but rather travels north on the sea floor as a 2000 m thick layer, filling all three basins (Johnson, 2008).

25 In a substantial portion of the Atlantic, ~~the progression of AABW is stopped by the other deep water (Johnson, 2008), the~~
~~AABW spreading north is overlain by~~ North Atlantic Deep Water (NADW) ~~spreading south (Johnson, 2008)~~. NADW forms in
the Labrador Sea and Nordic Seas ~~because of strong winds and haline convection respectively (Killworth, 1983)~~. It is the
saltiest of the two water masses, but is also warmer and lighter than AABW and hence ~~quits-leaves~~ the sea floor to continue
circulating above AABW ~~when-where~~ the two meet (Johnson, 2008). NADW production has long been linked to the strength
30 of the Atlantic Meridional Overturning Circulation (AMOC, e.g. Broecker, 1995), although observations from the recently
deployed OSNAP line (Lozier et al., 2019) ~~reveal that it suggest that this link~~ is more complex than “~~just~~ more deep water
formation = ~~stronger AMOC~~” ~~equals stronger AMOC~~. More crucially maybe, AABW and NADW formation provide a direct
path from the atmosphere to the bottom of the ocean, and as such, a conduit for heat and carbon storage (Chen et al., 2019;
Zanna et al., 2019). An accurate representation of deep water formation in climate models is thus a necessary precondition for
35 trustworthy future climate ~~predictions~~ ~~projections~~.

The Climate Model Intercomparison Project phase 6 (CMIP6, Eyring et al., 2016) is the latest release of CMIP, the organised
effort to make global climate models comparable, notably by running them with the same forcings. In the previous installment,
CMIP5 (Taylor et al., 2012), Southern Ocean mixed layers were poorly represented (Sallée et al., 2013) and models were
forming the majority of their AABW wrongly via open ocean convection, mostly in the ~~occurring-too-often-overly frequent~~
40 Weddell Polynya (Heuzé et al., 2013). They would only stop doing so once the ocean surface had freshened enough, by 2200
(De Lavergne et al., 2014) and, consequently, underestimated 21st century bottom property changes (Heuzé et al., 2015).
NADW formation was more accurately represented, although due to ~~too-overly~~ large sea ice extents in the North Atlantic, it
occurred more in the Irminger Sea than in the Labrador Sea (Menary et al., 2015; Heuzé, 2017). So far, results on CMIP6
models have shown that sea ice representation has improved in both hemispheres, but the intermodel spread remains large
45 (Roach et al., 2020; Shu et al., 2020). The Weddell Polynya is still opening too often, but only in half of the models (Mohrmann
et al., ~~subm.~~) ~~---,~~ those that now have the most accurate Antarctic Circumpolar Current (Meijers et al., 2012; Beadling et al.,
2020). CMIP6 models also have a higher climate sensitivity than CMIP5 models (Zelinka et al., 2020), more in line with
the observed sensitivity (Armour, 2017; Cox et al., 2018). Their AMOC however is too sensitive to the new aerosol forcings
(Menary et al., 2020). CMIP6 resolution is still coarse, with most models having a horizontal resolution of 1°, but recent results
50 showed that NADW formation is in fact less accurate with higher resolution (Koenigk et al., 2020). In summary, by improving
these crucial other processes but keeping a low resolution, CMIP6 models should have more realistic AABW and NADW than
CMIP5 models. We here investigate whether this is the case.

In this paper, we determine the characteristics of Antarctic Bottom Water (section 3.1) and North Atlantic Deep Water
(section 3.2) in CMIP6 models, focussing first on their respective formation processes, properties, and biases. ~~The primary~~
55 ~~objective of this paper is to quantify and discuss biases of each model, so that model users can make informed model selections.~~
~~Multi model means are also presented at the end of each subsection.~~ We then study the global transport of these two water
masses (section 3.3), in particular, how their properties determine their global extent. Finally (section 4), we conclude this
paper by a discussion on what -if anything- has improved since CMIP5.

2 Methods

2.1 CMIP6 models and observation-based reference data

We use the 35 CMIP6 models listed in Table 1. The only criterion for choosing them was the availability of at least their seawater salinity and temperature monthly output “so” and “thetao” for the entire historical run (January 1850 to December 2014) at the latest date of download (20 May 2020). When available, we also directly used their monthly mixed layer depth “~~mlostst~~mlostst”; if not, we computed it from the monthly salinity and temperature as detailed in section 2.2. We also made use of each model’s bathymetry “deptho” and grid cell area “areacello” files to accelerate our computations. Finally, for the transports calculations, we used the monthly meridional velocity “vo”. All output were obtained on the model’s native grid, except for NorESM2-LM and -MM that submitted their temperature and salinity on an isopycnic vertical grid; for these two models, we used the regularised z-level outputs.

We used only one ensemble member per model as even by the latest date of download, the majority of models had provided only one. ~~We acknowledge that~~ Furthermore, as some models are not fully independent as they share similar codes (Table 1), ~~and did not want to accentuate using different ensemble sizes would have accentuated~~ the bias towards one model family ~~by using different ensemble sizes~~. To account for this lack of independence, the correlations quoted throughout the text have been verified with different model numbers (not shown). For most models, the ensemble member we used is referred to as r1i1p1f1. It was not available for CNRM-CM6-1, CNRM-ESM2-1, MIROC-ES2L and UKESM1-0-LL, for which we used r1i1p1f2. Neither ~~was were~~ available for HadGEM3-GC31-LL, for which we used r1i1p1f3.

Although we used the full historical run for robustness verifications, we present only the results for the period January 1985 to December 2014, for consistency with the observational products. Note that we neither detrended the CMIP6 historical run nor substracted the pre-industrial control run, again for consistency with observations (which feature the climate change trend). These observations are the full-depth ocean temperature and salinity climatologies from the World Ocean Atlas 2018 (Locarnini et al., 2018; Zweng et al., 2018, respectively), the annual mixed layer depth climatology ~~of~~ first described by de Boyer Montégut et al. (2004), and the global bathymetry GEBCO (GEBCO Compilation Group, 2019).

2.2 Computations: deep and bottom water properties, transports and extents

To start with, when necessary, we computed the monthly mixed layer depth (MLD) of the CMIP6 models as per the CMIP6 procedures by first computing the monthly mean potential density σ_θ from their monthly practical salinity and potential temperature. ~~The~~ As is requested for CMIP6, the MLD is then ~~detected~~ diagnosed as the depth where σ_θ differs from that at 10 m depth by more than 0.125 kg m^{-3} . ~~Note that~~ Because of the non-linearity of the equation of state, this MLD computed from monthly temperature and salinity differs slightly from mlostst, which is the monthly average of the daily or higher resolution MLD outputted by the model. As shown for CanESM5 on supplementary Fig. A1, both mlostst and our recomputed MLD have the same spatial patterns but their largest values can differ by up to 300 m. As 1) the same regions are detected as having MLD exceeding the thresholds listed below and 2) the alternative is to not use the models that do not provide mlostst, we consider the difference acceptable. Furthermore, a different threshold of 0.03 kg m^{-3} is used ~~in~~ by the observational reference

Table 1. The 35 CMIP6 models used in this study; their ocean component; [nominal](#) horizontal resolution in ° latitude x ° longitude; vertical grid type (ρ means isopycnic, σ terrain-following, several symbols a hybrid grid) and number of vertical levels; and official reference. N/A indicates that no paper has been published yet for the CMIP6 configuration.

	Model name	Ocean component	Horizontal	Vertical	Reference
1	ACCESS-CM2	MOM5	1 x 1	z^* 50	N/A
2	ACCESS-ESM1-5	MOM5	1 x 1	z^* 50	Ziehn et al. (2017)
3	BCC-CSM2-MR	MOM4-L40	1 x 1	z 40	Wu et al. (2019)
4	BCC-ESM1	MOM4-L40	1 x 1	z 40	Wu et al. (2019)
5	CAMS-CSM1-0	MOM4	1 x 1	z 50	Rong et al. (2019)
6	CESM2	POP2	1 x 1	z 60	Danabasoglu et al. (2020)
7	CESM2-FV2	POP2	1 x 1	z 60	Danabasoglu et al. (2020)
8	CESM2-WACCM	POP2	1 x 1	z 60	Danabasoglu et al. (2020)
9	CESM2-WACCM-FV2	POP2	1 x 1	z 60	Danabasoglu et al. (2020)
10	CNRM-CM6-1	NEMO3.6	1 x 1	z^* 75	Volodire et al. (2019)
11	CNRM-ESM2-1	NEMO3.6	1 x 1	z^* 75	Séférian et al. (2019)
12	CanESM5	NEMO3.4.1	1 x 1	z 45	Swart et al. (2019)
13	EC-Earth3	NEMO3.6	1 x 1	z^* 75	N/A
14	EC-Earth3-Veg	NEMO3.6	1 x 1	z^* 75	N/A
15	GFDL-CM4	MOM6	0.25 x 0.25	$\rho - z^*$ 75	Held et al. (2019)
16	GFDL-ESM4	MOM6	0.5 x 0.5	$\rho - z^*$ 75	N/A
17	GISS-E2-1-G	GISS Ocean	1.25 x 1	z 40	N/A
18	GISS-E2-1-G-CC	GISS Ocean	1.25 x 1	z 40	N/A
19	GISS-E2-1-H	HYCOM	1 x 1	$z - \rho - \sigma$ 32	N/A
20	HadGEM3-GC31-LL	NEMO-HadGEM3-GO6.0	1 x 1	z^* 75	Kuhlbrodt et al. (2018)
21	INM-CM5-0	INM-OM5	0.5 x 0.25	σ 40	Volodin and Gritsun (2018)
22	IPSL-CM6A-LR	NEMO3.6	1 x 1	z^* 75	Lurton et al. (2020)
23	MCM-UA-1-0	MOM1	2 x 2	z 18	N/A
24	MIROC-ES2L	COCO4.9	1 x 1	$z - \sigma$ 62	Hajima et al. (2020)
25	MIROC6	COCO4.9	1 x 1	$z - \sigma$ 62	Tatebe et al. (2019)
26	MPI-ESM1-2-HAM	MPIOM1.6.3	1.5 x 1.5	z 40	Mauritsen et al. (2019)
27	MPI-ESM1-2-HR	MPIOM1.6.3	0.4 x 0.4	z 40	Müller et al. (2018)
28	MPI-ESM1-2-LR	MPIOM1.6.3	1.5 x 1.5	z 40	Mauritsen et al. (2019)
29	MRI-ESM2-0	MRI.COM4.4	1 x 0.5	z^* 60	Yukimoto et al. (2019)
30	NESM3	NEMO3.4	1 x 1	z 46	Cao et al. (2018)
31	NorCPM1	MICOM	1 x 1	$z - \rho$ 53	Counillon et al. (2016)
32	NorESM2-LM	MICOM	1 x 1	$z - \rho$ 53	Tjiputra et al. (2020)
33	NorESM2-MM	MICOM	1 x 1	$z - \rho$ 53	Tjiputra et al. (2020)
34	SAM0-UNICON	POP2	1 x 1	z 60	Park et al. (2019)
35	UKESM1-0-LL	NEMO-HadGEM3-GO6.0	1 x 1	z^* 75	Sellar et al. (2020)

(de Boyer Montégut et al., 2004), [which could lead to an apparent shallow bias for the models' mixed layer depths \(as we show in section 3, it does not\)](#). We could then quantify [deep-bottom](#) water formation in the three sectors of the Southern Ocean (borders at 65°W, 50°E and 130°E, south of 50°S, orange contours on Fig. 1), in the North Atlantic subpolar gyre (SPG, 70°W to 20°W, 50°N to 66°N) and in the Nordic seas (GIN, 30°W to 20°E, 65°N to 80°N, orange contours on Fig. 3) by computing the deep mixed volume (DMV) of each region as in Brodeau and Koenigk (2016). That is, for each month and each region, we keep only those grid cells where the MLD exceeds a critical value and sum the product MLD x cell area. We work with the maximum value of each year. As in Brodeau and Koenigk (2016) and Koenigk et al. (2020), we use a critical value of 700 m in the Nordic seas as it is the depth of the sill that connects them to the rest of the world ocean, and 1000 m in the Labrador Sea. As in e.g. Heuzé et al. (2013); De Lavergne et al. (2014), we use 2000 m in all three Southern Ocean sectors.

We quantify biases in the models by computing the root mean square error (model minus reference) in temperature, salinity and density σ_θ at the sea floor grid cell. To do so, all models had to be interpolated onto the reference's grid. [After interpolation we also computed the multi model mean properties, mean bias, and standard deviation of the bias for each grid cell.](#) Note

that we purposely keep σ_θ instead of σ_4 as σ_θ is the density used in the models' code to notably compute the MLD. For later
105 calculations, we also compute the temperature and salinity of the water masses AABW and NADW by taking their average
properties over a specific region. As we will show in section 3.1, the AABW formation region really differs from model to
model; as such, instead of using a limited region as in Johnson (2008), we detect AABW as having the temperature minimum
anywhere deeper than 2000 m and south of 50°S. For NADW, we produce two flavours: NADW_{SPG} as having the salinity
maximum anywhere deeper than 1000 m in the small area of SPG defined by Johnson (2008, 55°W to 54°W, 53°N to 63°N,
110 yellow box on Fig. 3); and NADW_{GIN}, the salinity maximum anywhere deeper than 1000 m in the GIN sector defined above.

We not only investigate the properties of AABW and NADW by their formation region, but also their transport into the rest
of the global ocean. For AABW, we hence compute each model's Southern Meridional Overturning ~~Circulation~~ Circulation or
SMOC, using the same method as Heuzé et al. (2015) for comparison purposes. That is, in each ocean basin, we first integrate
the meridional velocity v_o from the west coast to the east coast at 30°S, then we integrate this value from the sea floor to the
115 surface. The SMOC then is the northward deep maximum. We use a similar method for the AMOC, computing it at 35°N
instead for comparison with the CMIP6 results of Menary et al. (2020). After integration from coast to coast and from sea
floor to surface of the velocity v_o , the AMOC is defined as the southward subsurface maximum. We could not directly use the
meridional overturning circulation output "msftmz" as

- it is provided by ~~hardly a third~~ only 18 of the models (from 10 families);
- 120 – it is in kg s^{-1} instead of $\text{m}^3 \text{s}^{-1}$, requiring division by the density, which we only have the monthly mean of;
- for most models, the Indian and Pacific oceans are provided as one joint region, so we could not have obtained the SMOC
in each basin.

Having to interpolate the irregular model grids onto the sections instead of directly using the model output may have introduced
some errors. But as the AMOC results of this manuscript and that of Menary et al. (2020) for the models and experiment we
125 have in common are similar, we are confident in our MOC values. Note that two models, GFDL-ESM4 and NorCPM1, did not
provide v_o , limiting our transport analysis to 33 CMIP6 models.

Finally, to investigate in CMIP6 the link found in Heuzé et al. (2015) between the SMOCs and the northward extent of
AABW layer, we chose to re-create for CMIP6 models the Johnson (2008) maps of AABW and NADW volumes in the global
ocean. However, using the same approach as Johnson (2008) whereby we would have to determine the characteristics of every
130 water mass in each basin for each model is beyond the scope of this paper. Instead, as below the core of NADW the global
ocean (excluding the Arctic) is a mixture of NADW and AABW only, we determine at each depth the NADW and AABW
contents from a conservative property χ using the mixture equation of Jenkins (1999):

$$NADW_{content} = \frac{\chi_{AABW} - \chi}{\chi_{AABW} - \chi_{NADW}}, \quad (1)$$

and

$$135 \quad AABW_{content} = \frac{\chi_{NADW} - \chi}{\chi_{NADW} - \chi_{AABW}}. \quad (2)$$

Here, as in Johnson (2008), we consider the practical salinity and potential temperature as conservative enough to be used for these calculations. We then take the 50% content depth as the border between the NADW and AABW layers, i.e. anything with more than 50% AABW or less than 50% NADW is in the AABW layer, and the AABW thickness is the difference between the depth of that border and the sea floor. We finally take the median of all the combinations: temperature or salinity, NADW properties from SPG or GIN, and AABW or NADW contents. For the NADW layer, we detect the NADW core as the maximum NADW content with an extra criterion that the maximum must be larger than 80% NADW. Tests with values ranging from 60 to 100% yield similar values (not shown). Then the so-called NADW thickness is the thickness from the depth of the core to the NADW-AABW boundary (or to the sea floor if there is no AABW). By working with a mixture of two water masses only, we could not try and detect the top of the NADW layer. Note that traditional methods of using a fixed temperature and/or salinity for water mass determination cannot be applied to potentially biased climate models. The northward extent of AABW in each basin is defined as the northernmost latitude of the uninterrupted contour of thickness = 2000 m that starts in the Southern Ocean. We do the same for the southward extent of NADW in the Atlantic Ocean. For this part of the analysis, we show only the results for NADW that originated in SPG; NADW that originated in GIN seems to leave the Nordic Seas in no model (not shown).

150 3 Results

In this section, we first look at deep-bottom water formation and properties in the Southern Ocean, then deep water formation and properties in the North Atlantic. It is only in the last section that we analyse both water masses together, by determining their global transports and volumes. In this section, we talk only about CMIP6. The comparison with CMIP5 will come in section 4.

155 3.1 Southern Ocean bottom water characteristics in CMIP6 models

3.1.1 Shelf overflow and open ocean deep convection in the Southern Ocean

The Presently in the “real” ocean, Antarctic Bottom Water forms on the continental shelves of the Weddell and is primarily formed in several locations (including the Weddell Sea, the Ross Sea, and then flows into the deep basins by Adélie Land) as water is cooled, made saltier, and denser on the continental shelves, then cascades down the continental slopes, entraining deep waters on its way to the sea floor (visible as the densest areas on Fig. 1). The CMIP6 models’ bottom density bias on the shelves suggests that 19/35 models may form dense water on the shelf: ACCESS-CM2 (Weddell and Ross), ACCESS-ESM1-5 (Ross), CAMS-CSM1-0 (Ross), the four CESM2 (Ross), CanESM5 (Weddell and Ross), GFDL-ESM4 (Weddell and Ross), the three GISS (Ross mainly), HadGEM3-GC31-LL (Weddell and Ross), INM-CM5 (Weddell and Ross), IPSL-CM6A-LR (Weddell and Ross), the two NorESM2 (Ross mainly), SAM0-UNICON (Ross mainly), and UKESM1-0-LL (Weddell and Ross). The other 16 models are too light (strong negative bias on Fig. 1). Mean biases over 30 years are not enough to determine whether

the dense shelf water flows into the deep ocean; we instead created movies of the monthly bottom density over the entire historical run for these 19 models, of which two are available as video supplement.




Figure 1. Southern Ocean reference bottom density σ_θ (top left panel, top colorbar), and for each CMIP6 model, bottom density bias (model minus reference) averaged over 1985-2014. White number for each model is its RMSE over the entire Southern Ocean deeper than 1000 m. Thick black line indicates maximum mixed layer deeper than 2000 m. Thin grey line, the 2000 m isobath. Orange lines on the reference panel delimitate the Weddell (W), Amery (A) and Ross (R) sectors for the DMV calculation (see methods).

The movies let us split these 19 models into three groups:

- INM-CM5 and NorESM2-LM show overflowing from the Ross shelf to the deep basin; so does NorESM2-MM, as well as in the Amery sector (see video).
- GFDL-ESM4, HadGEM3-GC31-LL, IPSL-CM6A-LR, SAM0-UNICON and UKESM1-0-LL may overflow in the Ross sector, but we would need higher temporal resolution data to be certain;
- the other 11 models occasionally have a plume of dense water leaving the shelf, but it is nowhere near as dense as the shelf water it originates from (see video example of ACCESS-CM2).

In summary, in no model is there any (obvious) shelf export in the Weddell Sea. INM-CM5 (terrain following, high horizontal resolution model) and the two NorESM2 (hybrid isopycnic models) are the only ones forming AABW accurately via shelf processes, in the Ross sector only. The other high resolution models are not dense enough on the shelf or not clearly exporting their shelf water. There is no clear link between shelf processes and resolution (horizontal or vertical), vertical grid type, or ocean model component.

~~Southern Ocean reference bottom density σ_θ (top left panel, top colorbar), and for each CMIP6 model, bottom density bias (model minus reference) averaged over 1985-2014. White number for each model is its RMSE over the entire Southern Ocean deeper than 1000 m. Thick black line indicates maximum mixed layer deeper than 2000 m. Thin grey line, the 2000 m isobath. Orange lines on the reference panel delimitate the Weddell (W), Amery (A) and Ross (R) sectors for the DMV calculation (see methods).~~

If up to 8 models may export dense water from the shelf, how do the other 27 models of our study form their AABW? Deep convection, so far observed only once in the real ocean, in response to the 1974-1976 Weddell Polynya (Killworth, 1983), is the next obvious process to investigate. Mixed layer depths exceeding 2000 m are most prevalent in the Weddell sector (black contours on Fig. 1, DMVs in Table 2, MLD in supp. table A1). Of our 35 models, 24 exhibit deep convection in the Weddell Sea, of which 19 do so for most years of our study period. Most of these models also have a too large, ~~re-occurring-too-often~~ too frequent Weddell Polynya (Mohrmann et al., subm.), except for GFDL-CM4 and IPSL-CM6A-LR ~~who-which~~ may be convecting under sea ice cover, and the two MIROC who are ice-free (Mohrmann et al., subm.; Roach et al., 2020). In the Amery and Ross sectors, we need to distinguish between the models that have non zero DMV because of open ocean deep convection, and those with coastal polynyas. In the Amery sector, aside from MCM-UA-1-0 whose deep convection area is ~~but-just~~ a continuation of the Weddell one, we have 10/35 models with open ocean deep convection: the two CNRM, the two EC-Earth3, GISS-E2-1-H, MIROC6, MPI-ESM-1-2-HAM, MPI-ESM1-2-LR, and the two NorESM2. In the Ross sector, we have 15 models with open ocean deep convection, or to be more exact, 14 in the Ross Sea and NESM3 in the Amundsen Sea. In the Amery and Ross sectors, there is no link anymore between DMVs and the polynya activity by Mohrmann et al. (subm.), suggesting that ~~deep-bottom~~ water formation occurs under ~~(thin?)~~ ice cover. There is however a strong correlation of +0.47 between DMV in the Ross sector and DMV in the Weddell sector, i.e., models that convect a lot do it in both sectors. Behrens et al. (2016) suggests that a strong DMV is associated with a strong Antarctic Circumpolar Current (ACC), while Cabré et al.

(2017) find that strong DMV would weaken the westerly winds, i.e. may weaken the ACC. Here the only relationship between DMV and ACC, using the values from Beadling et al. (2020), is in agreement with Cabré et al. (2017): the more deep convection in the Amery sector, the weaker the ACC (correlation of -0.54, significant at 95% level). We find no relationship with DMV in the Weddell or Ross sectors.

205 Up to now, there are still 7 models that have no open ocean deep convection and no shelf overflow: the four CESM2, GISS-E2-1-G and -G-CC, and NorCPM1. GISS-E2-1-G and -G-CC have non zero DMV when considering the entire historical run (1850-2014, not shown), with GISS-E2-1-G convecting once in the Weddell sector and once in the Ross sector, and GISS-E2-1-G-CC thrice in the Weddell sector. The four CESM2 models do not, but they have an overflow parameterisation that artificially moves dense water from the shelf to the deep basins (Briegleb et al., 2010). ~~A pipe sucks the dense~~ If the water on the ~~shelves~~
 210 ~~and releases it in shelf exceeds a critical density, a pipe artificially transports this dense water from the shelves to~~ the deep basin ~~without~~. Without having to cascade. ~~This is why, the dense shelf water keeps its properties. Consequently, the absence of cascade means that~~ we cannot detect it on the overflow movies. For NorCPM1 though, ~~the mystery remains. Maybe it formed its AABW~~ we could not determine how its AABW is formed; maybe it formed before 1850. In ~~conclusions~~ conclusion, most models form their AABW by open ocean deep convection. Even the models that seem to represent shelf processes
 215 accurately exhibit open ocean deep convection. Somewhat surprisingly, the only relationship between the DMV and the climate sensitivities of Zelinka et al. (2020) is in the Weddell sector (correlation of -0.36): models that convect a lot there have a low sensitivity, which is to be expected as heat and CO₂ are sent to the deep ocean. What is surprising is that the relationship holds only in the Weddell sector. Hence, the sensitivity might be more linked to polynya activity, which is linked to deep convection only in the Weddell sector.

220 3.1.2 AABW properties

Does the way CMIP6 models form their AABW impact its characteristics, as it did in CMIP5 (Heuzé et al., 2013)? Only density biases are shown on Fig. 1, but salinity and temperature biases are provided as supplementary Figs A2 and A3 respectively. Ten models have a negligible bottom density bias (RMSE lower than 0.05 kg m⁻³, white numbers on Fig. 1): UKESM1-0-LL, CanESM5, IPSL-CM6A-LR, CESM2-WACCM-FV2, CESM2-FV2, CESM2, CESM2-WACCM, GISS-E2-1-H, CNRM-CM6
 225 and HadGEM3-GC31-LL. Twelve more models have an acceptable bottom density bias (RMSE lower than 0.1 kg m⁻³), including all the other models that are based on NEMO. That means that 22/35 models have acceptable biases. Let us investigate rather what may be common to the 13 models that are performing poorly.

INM-CM5 is the only model that is biased dense. Its bottom salinity is extremely high (RMSE of 0.42, supp. Fig. A2) while its bottom temperature is rather accurate (RMSE of 0.8°C, supp. Fig. A3). Its predecessor INM-CM4 had a similar
 230 issue, although whether this was caused by a too short spin-up or non-conservation of salt could not be determined (A. Gusev, personal communication, July 2014). All the other 12 models are biased light (Fig. 1). For BCC-CSM2-MR, BCC-ESM1, MCM-UA-1-0, MIROC-ES2L, MIROC6 and MRI-ESM2-0, it is because of a fresh bias (supp. Fig. A2). The other 6 models have relatively accurate bottom salinity, but are biased warm (supp. Fig. A3). ~~No model has regional biases, which means that~~ CMIP6 models are overall biased light or biased dense in the entire deep Southern Ocean (excluding the shelves).

Table 2. Median and maximum deep mixing volume (DMV, see methods) for the Southern Ocean sectors (orange contours on Fig. 1) for each CMIP6 model over 1985-2014. Values given in 10^{13} m^3 , which is approximately the DMV of a $1 \times 1^\circ$ grid cell with a 1000 m mixed layer. Number in brackets indicates how many years out of 30 is the DMV different from zero, i.e. the number of years with deep convection

model	Weddell			Amery			Ross		
	median	max	(nb years)	median	max	(nb years)	median	max	(nb years)
ACCESS-CM2	161	526	(30)	0	0	(0)	178	311	(30)
ACCESS-ESM1-5	408	588	(30)	0	0	(0)	66	266	(24)
BCC-CSM2-MR	432	764	(27)	1	11	(16)	0	0	(0)
BCC-ESM1	596	1108	(30)	0	0	(0)	0	11	(3)
CAMS-CSM1-0	128	415	(30)	1	7	(22)	0	13	(2)
CESM2	0	0	(0)	0	0	(0)	0	0	(0)
CESM2-FV2	0	0	(0)	0	0	(0)	0	0	(0)
CESM2-WACCM	0	0	(0)	0	0	(0)	0	0	(0)
CESM2-WACCM-FV2	0	0	(0)	0	0	(0)	0	0	(0)
CNRM-CM6-1	0	0	(0)	108	182	(28)	0	0	(0)
CNRM-ESM2-1	0	0	(0)	67	300	(30)	0	0	(0)
CanESM5	0	0.5	(2)	0	0	(0)	0	0	(0)
EC-Earth3	0	403	(12)	0	86	(11)	0	0	(0)
EC-Earth3-Veg	0	212	(12)	17	78	(20)	0	0	(0)
GFDL-CM4	1077	1334	(30)	0	6	(9)	16	53	(30)
GFDL-ESM4	0	0	(0)	0	0	(0)	0	46	(14)
GISS-E2-1-G	0	0	(0)	0	0	(0)	0	0	(0)
GISS-E2-1-G-CC	0	0	(0)	0	0	(0)	0	0	(0)
GISS-E2-1-H	114	205	(17)	0	61	(6)	0	8	(2)
HadGEM3-GC31-LL	21	80	(22)	0	0	(0)	0	0.5	(1)
INM-CM5-0	20	360	(27)	0	20	(6)	34	81	(30)
IPSL-CM6A-LR	56	168	(30)	0	0	(0)	0	2	(6)
MCM-UA-1-0	11	13	(30)	0	1	(26)	3	5	(30)
MIROC-ES2L	169	581	(27)	0	0	(0)	280	581	(30)
MIROC6	825	1117	(30)	29	80	(28)	930	1151	(30)
MPI-ESM-1-2-HAM	134	380	(23)	0	15	(7)	23	142	(20)
MPI-ESM1-2-HR	113	349	(30)	0	0	(0)	5	86	(30)
MPI-ESM1-2-LR	66	257	(30)	0	22	(10)	14	49	(29)
MRI-ESM2-0	0	2	(3)	1	5	(16)	0	5	(2)
NESM3	0	151	(9)	0	0	(0)	0	155	(10)
NorCPM1	0	0	(0)	0	0	(0)	0	0	(0)
NorESM2-LM	678	873	(30)	9	50	(19)	0	24	(12)
NorESM2-MM	650	882	(30)	79	146	(30)	1	130	(17)
SAM0-UNICON	0	0	(0)	0	0	(0)	0	29	(4)
UKESM1-0-LL	1	69	(17)	0	0.4	(2)	0	0	(0)

235 ~~The models-~~

240 Finally, the multi model mean bottom water is too light (mean bias of 0.06 kg m^{-3}), too warm (0.85°C) and too fresh (0.02 psu) throughout the entire Southern Ocean (Fig. 2). Technically, as the multi model mean bottom temperature is 0.57°C , we can even say that on average CMIP6 models do not have AABW (defined as having a temperature below 0°C) at the bottom of the Southern Ocean. The multi model mean biases in mixed layer depth exceed 2000 m in all sectors, especially in the Weddell Sea, but they do so over a small region so that the average bias over the entire deep Southern Ocean is low (151 m). As mentioned before, the biases in properties are moderate in the open ocean but exceed 0.1 kg m^{-3} or 0.1 psu on the shelves: the multi model mean shelves are too light and too fresh as many models do not form High Salinity Shelf Water. This is also the reason why the standard deviation (Fig. 2 last column) is largest on the shelves. In the open ocean, the standard deviation is larger in the Weddell Sea than in other sectors for the temperature, salinity and MLD, as it is where DMVs differ most (Table 2).

245




Figure 2. Multi model summary of the biases for the Southern Ocean. Columns: observational reference and multi model mean property (same colorbar for both), mean bias (model minus reference) and standard deviation of the difference; rows: bottom density, bottom potential temperature, bottom salinity and mixed layer depth. White numbers are the median value over the deep Southern Ocean, defined as per Fig. 1. Grey lines indicate the 1000 m and 2000 m isobaths.

The models with low biases in bottom density also tend to have zero to low DMVs in the Weddell Sea, but the relationship does not hold for maximum DMVs larger than $200 \cdot 10^{13} \text{ m}^3$ (Table 2). NorESM2-LM and -MM notably have low biases

but very high DMVs, but they also do shelf overflows. Open ocean deep convection leads to a warming and salinification of bottom waters (Zanowski et al., 2015); one hypothesis is then that models that hardly convect stay closer to the bottom density value they were initialised with. In the case of the CESM2 suite, the overflow parameterisation may help form accurate bottom water. Biases ~~as RMSE~~ are not the whole story though. As expected, we do find significant relationship (95% level) between the actual temperature and salinity of AABW and the DMV: in the Amery and Ross sectors, more deep convection leads to warmer AABW (correlation of +0.33 and +0.29 respectively) as in Wang et al. (2017). In the Weddell sector however, more deep convection leads to fresher AABW (correlation of -0.35), which in fact is consistent with the short-term response of the Southern Ocean to deep convection in Zanowski et al. (2015). The multimodel mean AABW salinity is 34.606 ± 0.154 ; the reference value from Johnson (2008), 34.641. The multimodel mean AABW temperature is -0.45 ± 0.73 °C; the reference value from Johnson (2008), -0.88 °C. That is, the multimodel mean AABW is warmer and fresher than the reference, and more DMV worsens these biases. Note that the values of the individual models are given in supp. Table. A2.

To summarise, in the Southern Ocean, most models form their AABW by open ocean deep convection. In the Weddell Sea, this convection seems tied to the Weddell Polynya activity, and impacts the AABW salinity most: more deep convection, fresher bottom salinity. In the Amery and Ross sectors, it is linked more to the bottom temperature: more deep convection, warmer bottom salinity. Models which seem to form dense water via shelf processes also exhibit deep convection, so we cannot determine whether overflows alone would make the Southern Ocean more accurate. ~~Models~~ The models that convect the least or not at all tend to be the most accurate; ~~four or these,~~ For the CESM2 ~~suite, may be aided by family, accurate~~ bottom properties and lack of deep convection may both be the result of their overflow parameterisation (Briegleb et al., 2010; Snow et al., 2015); ~~another one.~~ For another model, NorCPM1, ~~assimilates observations~~ the accuracy in all properties may come from its observation assimilation rather than accurate model physics (Counillon et al., 2016).

We will study the impact of these biases on the global transport of AABW in section 3.3. But as we cannot do so without investigating the AABW - NADW ~~tug-of-war~~ balance in the Atlantic basin, let us first evaluate the representation of NADW in CMIP6 models.

3.2 North Atlantic deep water in CMIP6 models

3.2.1 Deep water formation in the North Atlantic subpolar gyre and Nordic ~~Seas~~ seas

~~North Atlantic reference bottom density σ_θ (top left panel, top colorbar), and for each CMIP6 model, bottom density bias (model minus reference) averaged over 1985-2014. Orange lines on the reference panel delimitate the subpolar gyre (SPG) and Nordic Seas (GIN) areas for RMSE and DMV calculation; yellow dotted line, the SPG sector of Johnson (2008). White numbers for each model is its RMSE over the GIN (top) and SPG (bottom) areas, for depths over 1000 m. Thick black line indicates maximum mixed layer deeper than 1000 m; cyan dotted line in GIN, deeper than 700 m. Thin grey line, the 1000 m isobath.~~

In the North Atlantic, all 35 CMIP6 models of our study exhibit deep convection in the subpolar gyre (black contours on Fig. 3 and Table 3). As in CMIP5 (Heuzé, 2017), a large proportion of them ~~conveets~~convect not only in the Labrador Sea as the reference, but also intensely south of Iceland (Irminger Sea):

- 6/35 models convect only in the Labrador Sea (Fig. 3): CNRM-CM6-1, CNRM-ESM2-1, EC-Earth3-Veg, IPSL-CM6A-LR, NorCPM1 and NorESM2-LM;
- 9/35 models convect both in the Labrador and Irminger seas, but the two regions are not connected: ACCESS-CM2, BCC-ESM1, CESM2-WACCM-FV2, EC-Earth3, HadGEM3-GC31-LL, INM-CM5, MPI-ESM1-2-LR, NorESM2-MM and UKESM1-0-LL;
- 17/35 models convect both in the Labrador and Irminger seas, as one large SPG deep convection area: ACCESS-ESM1-5, BCC-CSM2-MR, CESM2, CESM2-FV2, CESM2-WACCM, GFDL-CM4, GFDL-ESM4, GISS-E2-1-G, GISS-E2-1-G-CC, GISS-E2-1-H, MCM-UA-1-0, MIROC6, MPI-ESM-1-2-HAM, MPI-ESM1-2-HR, MRI-ESM2-0, NESM3 and SAM0-UNICON;
- the last 3/35 models convect only in the Irminger Sea: CAMS-CSM1-0, CanESM5 and MIROC-ES2L.




Figure 3. North Atlantic reference bottom density σ_θ (top left panel, top colorbar), and for each CMIP6 model, bottom density bias (model minus reference) averaged over 1985-2014. Orange lines on the reference panel delimitate the subpolar gyre (SPG) and Nordic seas (GIN) areas for RMSE and DMV calculation; yellow dotted line, the SPG sector of Johnson (2008). White numbers for each model is its RMSE over the GIN (top) and SPG (bottom) areas, for depths over 1000 m. Thick black line indicates maximum mixed layer deeper than 1000 m; cyan dotted line in GIN, deeper than 700 m. Thin grey line, the 1000 m isobath.

Table 3. Median and maximum deep mixing volume (DMV, see methods) for the subpolar gyre (SPG) and Nordic Seas (GIN, see Fig. 3) for each CMIP6 model over 1985-2014. Values given in 10^{13} m^3 , which is approximately the DMV of a $1 \times 1^\circ$ grid cell with a 1000 m mixed layer. Number in brackets indicates how many years out of 30 is the DMV different from zero, i.e. the number of years with deep convection

model	SPG			GIN		
	median	max	(nb years)	median	max	(nb years)
ACCESS-CM2	77	181	(30)	57	102	(30)
ACCESS-ESM1-5	50	171	(30)	42	104	(30)
BCC-CSM2-MR	222	406	(30)	26	52	(30)
BCC-ESM1	190	302	(30)	23	60	(30)
CAMS-CSM1-0	107	225	(30)	12	35	(24)
CESM2	73	238	(30)	6	24	(30)
CESM2-FV2	118	206	(30)	6	15	(30)
CESM2-WACCM	76	226	(30)	8	23	(30)
CESM2-WACCM-FV2	107	263	(30)	6	22	(30)
CNRM-CM6-1	0	161	(14)	61	125	(30)
CNRM-ESM2-1	3	78	(25)	51	134	(30)
CanESM5	0	3	(7)	22	92	(30)
EC-Earth3	20	141	(26)	60	103	(30)
EC-Earth3-Veg	18	201	(20)	65	103	(30)
GFDL-CM4	417	548	(30)	36	70	(30)
GFDL-ESM4	152	383	(30)	40	128	(30)
GISS-E2-1-G	303	476	(30)	32	103	(30)
GISS-E2-1-G-CC	316	418	(30)	29	86	(29)
GISS-E2-1-H	402	509	(30)	102	199	(30)
HadGEM3-GC31-LL	17	156	(24)	29	124	(30)
INM-CM5-0	2	16	(18)	0	0	(0)
IPSL-CM6A-LR	0	43	(12)	47	109	(30)
MCM-UA-1-0	2	3	(30)	1	1	(30)
MIROC-ES2L	1	56	(19)	122	208	(30)
MIROC6	152	365	(30)	168	260	(30)
MPI-ESM-1-2-HAM	66	219	(30)	42	124	(30)
MPI-ESM1-2-HR	62	162	(30)	15	37	(30)
MPI-ESM1-2-LR	37	97	(28)	28	58	(30)
MRI-ESM2-0	132	269	(30)	37	96	(30)
NESM3	141	273	(30)	6	39	(26)
NorCPM1	0	0	(1)	0	0	(0)
NorESM2-LM	150	278	(30)	34	97	(30)
NorESM2-MM	112	236	(30)	93	131	(30)
SAM0-UNICON	333	456	(30)	151	225	(30)
UKESM1-0-LL	25	129	(20)	109	177	(30)

As in Koenigk et al. (2020), the higher resolution NorESM2-MM and MPI-ESM1-2-HR have larger deep convection areas than their corresponding lower resolution NorESM2-LM and MPI-ESM1-2-LR. Note that the difference between NorESM2-MM and -LM is in the atmospheric component resolution. There is however no robust relationship across models between the horizontal resolution and the DMV. There is a relationship with the climate sensitivity though (Zelinka et al., 2020) sensitivities of Zelinka et al. (2020) though: the larger the DMV in SPG, the least sensitive the model (correlation of -0.36) which, as already discussed in the Southern Ocean part, is not surprising. Consequently, there is also a strong correlation between the DMV in the Weddell Sea and in SPG (+0.57): models that convect a lot in the Weddell Sea convect a lot in SPG as well. As already mentioned, no causation can be inferred: deciphering whether global biases in DMV are responsible for the models' sensitivities, or in contrast sensitivities are set by other processes and impact the DMV, is beyond the scope of this analysis.

All models except INM-CM5-0 and NorCPM have deep convection in the GIN seas as well. Moreover, in GIN, models convect most years, with a minimum as high as 24/30 years (Table 3). There is more variability in the SPG, but likewise the

majority of models convect all years. Besides, they convect too deep. While in the Southern Ocean, deep convection to the sea floor can happen (Killworth, 1983), in the North Atlantic it should not go much beyond 1000 m (e.g. Våge et al., 2009). In the SPG, only the three models CanESM5, INM-CM5 and NorCPM1 have maximum mixed layer depths just exceeding 1000 m (supp table A1). An extra four models, ACCESS-CM2, CESM2, CESM2-WACCM and MIROC-ES2L, have tolerable depths up to 2500 m. But the vast majority convects too deep, often to the sea floor. It is the same in GIN, albeit with different models: this time, the four CESM2, MPI-ESM1-2-LR and NESM3 have MLDs up to 2000 m, and all the other models go to 3000 m or even the sea floor. There is no significant correlation between the DMV in SPG and that in GIN. In summary, CMIP6 models exhibit deep convection in the North Atlantic too often, too deep, and over too large an area. It is not possible to determine the one most accurate model in the North Atlantic or even in each subregion; model users must choose a compromise between correct representation of the variability, location, depth or extent.

3.2.2 North Atlantic bottom properties

~~The picture is less grim regarding bottom properties biases~~ CMIP6 water property biases at the bottom of the North Atlantic are of the same order of magnitude than those at the bottom of the Southern Ocean (shading on Fig. 3). Three models have bottom density biases resulting in an RMSE lower than 0.05 kg m^{-3} in both SPG and GIN: CESM2-WACCM, HadGEM3-GC31-LL and UKESM1-0-LL. And extra 9 models have an RMSE lower than 0.1 kg m^{-3} in both SPG and GIN: CESM2, CESM2-FV2, CNRM-CM6-1, EC-Earth3-Veg, GFDL-CM4, GFDL-ESM4, IPSL-CM6A-LR, MRI-ESM2-0 and SAM0-UNICON. As for the other 23 models, it depends on the region:

- INM-CM5, NorCPM1, NorESM2-M and NorESM2-MM are biased dense in both regions because they are biased salty (supp. Fig. A4). The magnitude of the bottom cell biases in INM-CM5 is very grid-cell dependent, maybe because of faulty regularisation of the sigma grid (even though it did not have this problem in the Southern Ocean).
- ACCESS-ESM1-5 is accurate in the SPG sector but biased dense (salty) in GIN.
- CESM2-WACCM-FV2, GISS-E2-1-G and GISS-E2-1-G-CC are accurate in the SPG sector but biased light in GIN. For CESM2-WACCM-FV2, it is because of a warm bias (supp Fig. A5); for GISS-E2-1-G and GISS-E2-1-G-CC, because of a fresh bias.
- ACCESS-CM2, BCC-CSM2-MR, CNRM-ESM2-1, EC-Earth3, MPI-ESM1-2-HAM and MPI-ESM1-2-LR are biased light in SPG but accurate in GIN. EC-Earth3 is the only one that is biased fresh. The other models are biased salty but warm.
- The last 9 models are biased light in both regions. For CanESM5, GISS-E2-1-H, MCM-UA-1-0, MIROC-ES2L and NESM3, this is caused by a salty bias; for BCC-ESM1, CAMS-CSM1-0, MIROC6 and MPI-ESM1-2-HR, a warm bias.

All models except CanESM5 and IPSL-CM6A-LR are in fact biased warm compared to the World Ocean Atlas 2018 bottom temperature. The evolution of the bottom properties throughout the entire historical run is complex, with significantly different

variabilities depending on the model (not shown), and their analysis is beyond the scope of this paper. All that we can say that
335 the warm bias is not a result of only the modelled climate change or any drift.

~~In~~

The multi model mean biases reflect the individual model biases that we just described (Fig. 4): in the deep ocean, in both GIN and SPG, the multi model mean is biased salty (0.07 in GIN, 0.15 in SPG) but warm (1.68°C and 1.91°C respectively), hence slightly fresh (0.04 and 0.07 kg m⁻³). The across model spread, given by the standard deviation (last column, Fig. 4), is
340 in the order of 0.1 kg m⁻³ or psu and 1°C in both regions, similar to what we found for the Southern Ocean. The multi model mean MLD is too deep over too large an area compared to the reference (bottom row, Fig. 4), corresponding to a bias of nearly 800 m on average in both regions. The standard deviation is relatively high though, also reaching 800 m on average, indicative of model disagreement over the location of the deep MLD. The largest bias in standard deviation, i.e. the location where the across-model agreement is lowest, is in the Greenland Sea, probably a consequence of the across model differences in sea ice representation (Shu et al., 2020).
345

In the SPG, there is a strong relationship between the bottom temperature RMSE and the climate sensitivity of Zelinka et al. (2020): the more sensitive the model, the least biased in temperature in SPG (correlation of -0.68). There is a somewhat significant (at the 90% level) relationship between the DMV and the bottom density bias in SPG only: the more the model convects, the ~~least-less~~ biased (correlation of -0.29). There is however no relationship with the location of deep convection
350 itself, e.g. MIROC-ES2L that convects only in the Irminger Sea has a similar bias (magnitude and sign) as MPI-ESM1-2-LR that convects only in the Labrador Sea; MPI-ESM1-LR in turns has a large bias and convects at the same location as UKESM1-0-LL, which has a low bias. The four CESM2 models and their overflow parameterisation by Denmark Strait are again among the most accurate, which was in fact the original motivation for that parameterisation (Briegleb et al., 2010). NorCPM1 is somewhat disappointing; it is built on NorESM2-LM and is supposed to have improved performances thanks
355 to data assimilation (Counillon et al., 2016). Its bottom density is indeed better than NorESM2-LM's in SPG, but is biased even denser (saltier) in GIN. There is no across-model relationship between the sensitivity or the DMV and the salinity or GIN though, so the ~~question for NorCPM~~cause of NorCPM's and other models~~remains'~~bottom density bias in the GIN seas remains unknown. It may be linked to their respective biases in the representation of the Atlantic Multidecadal Oscillation, as suggested by Lin et al. (2019). There is however a strong ~~,-convenient,-~~interhemispheric correlation: the more biased the
360 bottom density in the Southern Ocean, the more biased it is in SPG (+0.81)

Finally, looking at the NADW properties instead of the biases (individual values in supp. table A2), we find that the multi-model mean NADW_{SPG}, which Johnson (2008) refers to as upper NADW (UNADW) or Labrador Sea Water (LSW) is too warm and too salty: $4.86 \pm 0.81^\circ\text{C}$ and 35.163 ± 0.143 in CMIP6, instead of 3.32°C and 34.894 in Johnson (2008). The multimodel mean NADW_{GIN} in contrast is accurate: $0.77 \pm 0.99^\circ\text{C}$ and 35.001 ± 0.169 compared to 1.30°C and 34.878
365 for the water mass called lower NADW (LNADW) or Iceland-Scotland Overflow Water (ISOW) in Johnson (2008), despite 5 models having a temperature below ~~0~~0.0°C. Again in SPG, models with a high sensitivity have a lower salinity and temperature (correlations of -0.31 and -0.45 respectively), which is consistent with the links previously found with the DMV. But again, no relationship can be found in GIN. In GIN, we found no relationship between the biases or properties and the horizontal or




Figure 4. [Multi model summary of the biases for the North Atlantic. Columns: observational reference and multi model mean property \(same colorbar for both\), mean bias \(model minus reference\) and standard deviation of the difference; rows: bottom density, bottom potential temperature, bottom salinity and mixed layer depth. White numbers are the median value over the deep GIN \(top\) and SPG \(bottom\), defined as per Fig. 3. Grey lines indicate the 1000 m and 2000 m isobaths.](#)

vertical resolution, nor the grid type or ocean model component. In CMIP5, Heuzé and Årthun (2019) had found strong across
 370 model biases in the inflow to the Nordic Seas caused by the large scale oceanic and atmospheric circulations as well as

the bathymetry, while Lin et al. (2019) showed that GIN property biases can be linked to the representation of multidecadal variability. Investigating the exact cause of the biases in GIN is beyond the scope of this paper, not least because in the next section, we will show that NADW_{GIN} does not contribute to the global NADW in CMIP6 models. For now, we can conclude that the bottom property biases in GIN are not related to deep water formation in the region.

3.3 Global transport of NADW and AABW in CMIP6 models

In this last section, we shall determine the global fate of NADW and AABW once they leave their source regions. For NADW, this fate is tied to the strength of the Atlantic Meridional Overturning Circulation. The mean AMOC value lies at 18 Sv ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$), although observations both at the RAPID/MOCHA-array at 26.5°N (e.g. Ducheze et al., 2016) and in the more recently deployed OSNAP-lines in the subpolar North Atlantic (Lozier et al., 2019), reveal a strong interannual variability of up to 5 Sv. Aside from INM-CM5 and its AMOC of $63 \pm 19 \text{ Sv}$, all models fall in that range (individual values in supp. Table B1), resulting in a multimodel median of $19.5 \pm 9.5 \text{ Sv}$. ~~Which means that like Menary et al. (2020), we find that the AMOC is overestimated in CMIP6 models on average, but not dramatically.~~ Observations of the southern MOC at 30°S are rarer. From box inverse modelling, Lumpkin and Speer (2007) estimated the Atlantic SMOC at $5.6 \pm 3 \text{ Sv}$; apart from the three GISS models and MIROC-ES2L that are too weak, all models are in that range (supp table B2), giving a multimodel median of 2.8 ± 1.4 . Observational values in the Indian Ocean range between 3 and 27 Sv (Huussen et al., 2012), so unsurprisingly, from the weak MCM-UA-1-0 ($1.5 \pm 1.6 \text{ Sv}$) to the strong GFDL-CM4 ($11 \pm 18 \text{ Sv}$), all models are in that range and the multimodel median is $3.0 \pm 2.5 \text{ Sv}$. This is a remarkable improvement since CMIP5, where a majority of models had an Indian SMOC close to 0 (Heuzé et al., 2015). In the Pacific finally, Lumpkin and Speer (2007) estimated the MOC to be $11 \pm 5 \text{ Sv}$. MCM-UA-1-0 is again the weakest ($3.9 \pm 1.9 \text{ Sv}$), and the only model that falls out of the observational range, resulting in a multimodel median of $5.9 \pm 3.0 \text{ Sv}$. In summary, the AMOC and southern MOCs are rather accurately represented in CMIP6 models.

The across-model correlations between the transports are strong and significant (95% level): the stronger the SMOC in the Indian Ocean, the stronger as well in the Pacific Ocean (correlation of +0.37). In contrast, a strong SMOC in either of these basins corresponds to a weak SMOC in the Atlantic (Atlantic-Indian, correlation of -0.45; Atlantic-Pacific, -0.34). And a weak SMOC in the Atlantic corresponds to a strong AMOC (correlation of -0.30), as previously found by Patara and Böning (2014) in the NEMO model. We are ~~obviously~~ not implying causation from the correlations, but it is interesting to find relationships between the biases quantified in sections 3.1 and 3.2 and the transports. In agreement with Patara and Böning (2014), a stronger Atlantic SMOC is associated with lower temperature biases (correlation of 0.29), that is, colder AABW (-0.35), whereas a stronger Pacific SMOC is associated with stronger density biases (+0.36). A stronger AMOC is associated with larger biases in temperature and salinity in SPG (correlations of +0.33 and +0.37 respectively), and in particular a saltier NADW_{SPG} (+0.34, as in the paleoclimate simulations of Menviel et al., 2020). The Atlantic SMOC is the only transport that is linked to the strength of the Antarctic Circumpolar Current (ACC, values from Beadling et al., 2020): the stronger the ACC, the stronger the Atlantic SMOC (correlation of +0.37). There is no significant direct relationship between the transports and the DMV, which, at least for the AMOC, is in agreement with the recent observations of Lozier et al. (2019) and modelling work of Årthun

et al. (2019). Unlike e.g. Menary et al. (2015) or Koenigk et al. (2020), we find no link between the MOCs and the models' horizontal resolution.

In line with Heuzé et al. (2015), we expect the transports to impact the interbasin spread of NADW and AABW, that is, that the stronger the transport, the further from its source the water mass will travel. To investigate this, we recreated the AABW and NADW thickness maps of Johnson (2008) as Figs 5 and 6 respectively. For Fig. 6, we show only NADW_{SPG}; surprisingly, in no model could we find NADW_{GIN} beyond the Nordic Seas-seas (not shown). In agreement with observations and Johnson (2008), AABW occupies the majority of the water column in most of the Indian and Pacific oceans, but its northward extent is limited in the Atlantic Ocean. Said extent is highly model dependent in the Atlantic, whereas it extends as far north as the basin limits permit in most models in the Indian and Pacific oceans. Finally, in most models, AABW in the Indian Ocean seems to come from the Pacific. The NADW southward expansion in the Atlantic is also model-dependent, with some reaching to the Antarctic Circumpolar Current (e.g. BCC-ESM1) and others not even leaving the North Atlantic subpolar gyre (e.g. UKESM1-0-LL). As explained in the methods, the NADW layer in the Indian and Pacific oceans is most likely biased by our calculation method that takes into account only two water masses, and thus shall not be discussed further.

After extracting the southernmost extent of NADW and northernmost extents of AABW for each model (see supp. tables B1 and B2), we do find, as expected, that the stronger the AMOC, the further south NADW extends in the Atlantic (correlation of 0.32). And the stronger the Atlantic SMOC, the further north AABW extends in the Atlantic (correlation of 0.40). As we previously found an anticorrelation between the AMOC and the Atlantic SMOC across CMIP6 models, the Atlantic balance is complete: models with strong AMOC and weak SMOC have their Atlantic dominated by NADW (e.g. CESM2-WACCM), whereas those with a weak AMOC and strong SMOC are filled with AABW (e.g. IPSL-CM6A-LR). And although there was no significant correlation between the DMV and the transports, we do find that the larger the DMV, the further the extent of NADW (DMV SPG, correlation of +0.34) or AABW (DMV Weddell, +0.51). We found no significant correlation between the northward extent in the Indian or Pacific oceans and either the SMOCs or DMVs, or with the strength of the ACC. There are however relationships with their bottom properties: the northward extent of salty models is less than that of fresh models (correlations of -0.31 in the Indian and -0.44 in the Pacific). As we also find a strong positive relationship (correlation of +0.72) between the salinity of AABW and the salinity gradient across the ACC computed by Beadling et al. (2020), i.e., we find that the fresh models have a weak gradient to overcome, this result is not surprising. We can even speculate that in the absence of NADW, AABW would expand further north in the fresher models regardless of their SMOC.

~~Thickness of the Antarctic Bottom Water layer in observations (top-left panel) and in each CMIP6 models. See Methods.~~
~~Thickness of the North Atlantic Deep Water layer in observations (top-left panel) and in each CMIP6 models, from the NADW-core to its bottom. See Methods.~~

In conclusion, in CMIP6 models as in the real ocean, deep convection impacts bottom water characteristics and biases: in the Southern Ocean, deep convection seems associated with more biased deep-bottom waters; in the North Atlantic, the more the models convect, the least-less biased they are. Either way, these biases then impact the deep and bottom water transport: a saltier NADW is associated with a stronger AMOC; colder AABW, stronger Atlantic SMOC. These transports then impact the location of the “NADW - AABW border” in the Atlantic: stronger AMOC and weaker Atlantic SMOC (the two transports are

440 anticorrelated), further southward extent of NADW and less northward extent of AABW. In the Indian and Pacific oceans, the northward extent is larger in the fresher models, which are the ones with weak fronts in the ACC. To summarise, deep ~~water formation is~~ and bottom water formation are crucial for an accurate representation of the global deep ocean. We conclude this paper with a discussion of changes in deep and bottom water modelling since CMIP5, and what we can expect from the next generation(s) of simulations.




Figure 5. Thickness of the Antarctic Bottom Water layer in observations (top left panel) and in each CMIP6 models. See Methods.




Figure 6. Thickness of the North Atlantic Deep Water layer in observations (top left panel) and in each CMIP6 models, from the NADW core to its bottom. See Methods.

In CMIP5 models, no model assessed by Heuzé et al. (2013) could represent dense shelf overflows correctly. Consequently, models relied on open ocean deep convection for their ~~deep-bottom~~ water formation. The right amount of deep convection in the Weddell Sea was required for accurate bottom properties; models that convected too little or too much were the most biased. This relationship does not hold anymore for CMIP6, and it is the models that convect the least that tend to be the most accurate (Fig. 1 and ~~table-Table~~ Table 2). It may be because many models are now artificially prevented from opening polynyas and convecting in the Weddell Sea (Mohrmann et al., subm.). However, as the Weddell Polynya has now reopened in the real ocean (Campbell et al., 2019), future models may remove their “polynya-prevention” schemes again. Another reason for CMIP6 models seemingly not needing Southern Ocean deep convection to have accurate bottom properties may be that, as we showed in this paper, several CMIP6 models successfully represent shelf processes. This was an unexpected result considering that horizontal resolutions have not increased much since CMIP5, suggesting that models have improved their parameterisations instead (Danek et al., 2019). Regardless of the formation process, bottom density biases are smaller in CMIP6 than they were in CMIP5 (RMSEs on Fig. 2 of Heuzé et al., 2013). The new version of the models that performed well in CMIP5 also performs well in CMIP6 (e.g. the IPSL and NorESM families), and the others have improved (CanESM4 had a bias of 0.17 kg m^{-3} ; CanESM5, 0.03). The worst performing model of CMIP5 was INMCM4. The worst performing model of CMIP6 with respects to Southern Ocean bottom properties is its successor, INM-CM5-0, but even this model saw its bias halve. INM-CM5-0 has both shelf processes and open ocean deep convection whereas INMCM4 had neither, which probably contributed to ridding the model of its cold bottom bias (Zanowski et al., 2015).

In the North Atlantic, to the best of our knowledge, most CMIP5 studies focussed on the relationship between deep water formation and the AMOC or the warming hole (e.g. Menary and Wood, 2018) but did not investigate bottom property biases. The one exception is Ba et al. (2014) who found a recurrent cold bias; with the World Ocean Atlas 2018 as reference, we find in contrast that most CMIP6 models have a warm bias at the bottom of the North Atlantic. Deep water formation in the North Atlantic in the majority of CMIP5 models occurred too often, too deep, over too large an area (Heuzé, 2017). ~~This sentence is~~ These findings are still valid for CMIP6 (Fig. 3 and ~~table-Table~~ Table 3). One noticeable improvement (+) is that the models whose CMIP5 predecessor convected only in the Irminger Sea now convect in the entire subpolar gyre, including the Labrador Sea. Unfortunately, some of the models that performed well in CMIP5 when considering the location of deep convection in the SPG, i.e. had a relatively small area in the Labrador Sea, have also expanded to the entire SPG (e.g. the CNRM family). That is, the inaccurate models may be on the way to improvement, most likely because the Arctic sea ice is better represented in CMIP6 than in CMIP5 (Shu et al., 2020), but the ones that were relatively accurate have degraded. The same holds for the Nordic seas: CMIP6 models are convecting even more than CMIP5 models did, and they already were convecting too much. In an increasingly warmer and ice-free climate, Lique and Thomas (2018) predict that deep water formation would migrate from the North Atlantic subpolar gyre to its subtropical gyre, and from the Nordic seas to the Arctic. Liu et al. (2019) adds that this will depend on whether meltwaters will most strongly impact the stratification, shutting down deep convection, or the horizontal gradients and hence the winds, pushing meltwater away from convection areas. For now, we observe that from the

very icy CMIP5 to the more accurately de-iced CMIP6 models, deep water formation regions just expanded to occupy most of the space available in SPG and GIN. It is unclear whether increasing the resolution of future models would solve this issue: Danek et al. (2019) dramatically reduced mixed layer depths in SPG by using an adaptive mesh with 5-15 km resolution, while Koenigk et al. (2020) finds that DMVs in the SPG become even larger in the high resolution versions of the models that participated to HighResMIP. Without changing the horizontal resolution, a more systematic inclusion and better representation of the stratosphere may be enough to reduce deep convection in the North Atlantic (Haase et al., 2018).

Regarding the transports, as noted by Menary et al. (2020) the AMOC is stronger in CMIP6 than in CMIP5, which they ~~blame on~~ attribute to the aerosol forcing. Except for INM-CM5 that is now way too strong, or which uploaded incorrect velocity fields, this increase is not that strong and most models are in the observational range. In the case of the CNRM family, a stronger AMOC is in fact a much more accurate AMOC (from 12 Sv in CMIP5 to 19 Sv in CMIP6). The NorESM models have a weaker AMOC in CMIP6, which is more accurate than their CMIP5 version (from 32 Sv in CMIP5 to 21 Sv in CMIP6). The two highest resolution models have weakened so much that their AMOC is too low (GFDL-CM4 and MPI-ESM1-2-HR). This seems in contradiction to Koenigk et al. (2020) who found that increased resolution in HighResMIP leads to a stronger AMOC, but their result is mostly true when the models reach an eddy-resolving resolution. Which they do not, here, in CMIP6. It is harder to determine whether the Southern MOCs at 30°S have improved since the values from inverse modelling (Lumpkin and Speer, 2007) and observations (Huussen et al., 2012) have very large uncertainties. All that we can say is that the Atlantic SMOC is stronger in CMIP6, so that only the GISS family continues having an Atlantic SMOC around 0 Sv. In the Indian Ocean, no model has a transport of 0 anymore, which resulted in a doubling of the multimodel mean from 1.6 Sv in CMIP5 to 3 Sv in CMIP6, giving it the same importance as the Atlantic SMOC. The Pacific SMOC remains the strongest of the three and sees no significant difference between CMIP5 and CMIP6 except for the two models that used to be around 0, INMCM4 (INM-CM5-0 is now at 10 Sv) and GISS-E2-H (GISS-E2-1-H now at 7 Sv). As in CMIP6 the Southern Ocean representation from the ~~bottom-sea floor~~ (this manuscript) to the ~~top-surface~~ (Beadling et al., 2020) has improved, as well as the ACC (also Beadling et al., 2020), it is no surprise that more models are now capable of exporting AABW to the rest of the world ocean. To the best of our knowledge, the global extent of AABW and NADW, presented here for CMIP6 on Figs 5 and 6 respectively, was not assessed in CMIP5, so we cannot determine whether improved Southern Ocean characteristics lead to an improved global water mass distribution.

What can we expect from a hypothetical CMIP7? Higher resolution, most likely, although that was already expected from CMIP6 and did not happen. As explained above and by Koenigk et al. (2020) or Danek et al. (2019), a higher resolution would not necessarily improve deep water formation. Holt et al. (2017) goes as far as stating that shelf processes will not be correctly represented until the horizontal resolution remains lower than 1/72°, which they expect might be reachable by the most advanced computers within 10 years. Unfortunately, we do not all have access to these computers, so that even now, computing the global monthly mixed layer depth of the highest resolution model (GFDL-CM4, 1/4°) required over 600 core hours for the 165 years of the historical run. Higher resolution output will be impossible to manage, unless cloud-computing solutions such as PANGEO become the norm (Odaka et al., 2020). Instead of increasing the resolution, a seemingly easier solution would be to improve parameterisations (Holt et al., 2017), especially overflow parameterisations (Snow et al., 2015).

Briegleb et al. (2010) first showed that an overflow parameterisation to transport water from the Nordic ~~Seas-seas~~ to the rest of the North Atlantic resulted in an improved representation of the ocean there. In CMIP6, the CESM2 models with their “pipes” in the North Atlantic and Antarctic shelves were among the most accurate models, especially for AABW. It would be interesting to see whether such a parameterisation on a different model would yield the same results, or whether the CESM2 models are just very accurate. Efforts could also concentrate on improving other components of the climate model, for example the atmosphere, as an improved representation of the stratosphere would supposedly decrease unrealistic deep and bottom water formation (Haase et al., 2018). But where most progress can probably be made is in the cryosphere. As deep water formation is tied to the sea ice behaviour in both hemispheres, efforts such as sea ice MIP (SIMIP, Notz et al., 2016) dedicated to the modelling and coupling of sea ice may be the way forward. Likewise, the results of ice sheet MIP (ISMIP6, Nowicki et al., 2016) may shed a light on the debated impact of glacial meltwater on deep and bottom water formation (De Lavergne et al., 2014; Liu et al., 2019).

525 5 Conclusions

In this paper, we determined the characteristics of Antarctic Bottom Water and North Atlantic Deep Water in 35 models that participated in the latest installment of the Climate Model Intercomparison Project, CMIP6: their formation, properties, transports and extent in the global ocean. We focussed on the last thirty years of the historical run, January 1985 to December 2014. In the Southern Ocean (section 3.1), deep-bottom water formation is now more accurate, with several models representing shelf processes. Open ocean deep convection in the Weddell Polynya still happens in more than half of the models, but it is not a requirement for accurate bottom water properties. In fact, the most accurate models were the ones with little to no open ocean convection, especially the CESM2 family that has an overflow parameterisation. In the North Atlantic (section 3.2), models convect too often, too deep, over too large an area, but in the subpolar gyre that area has migrated from the Irminger Sea (in CMIP5 models) to the more accurate Labrador Sea. The models that convect the most in the North Atlantic subpolar gyre also have the least biased NADW. NADW ~~that forms formed~~ in the subpolar gyre ~~is the only one that occupies the world ocean;~~ NADW from the Nordic seas appears to stay of the models clearly spreads southward, but the signature of the portions formed in the Nordic seas is less evident. The saltier NADW, the stronger the AMOC, and the further south the extent of NADW (section 3.3). That extent is limited by the strength of the abyssal overturning in the southern Atlantic or SMOC, with stronger Atlantic SMOC (caused by colder AABW) resulting in a further northward extent of AABW. In the Indian and Pacific oceans, the extent is directly related to the AABW properties, not the SMOCs: models with a comparatively fresh AABW are also the ones with weak fronts across the Antarctic Circumpolar Current, and hence can travel the furthest north. In summary, for ~~both~~ deep-the deep and bottom water masses in CMIP6, their formation impacts their properties, which impact their transport and global extent, which in turns will have large impacts on global predictions of thermal expansion and sea level rise (Zickfeld et al., 2017), carbon storage (Tatebe et al., 2019), ecosystem changes (Sweetman et al., 2017) etc. Although CMIP6 models represent AABW and NADW more accurately than CMIP5 models did, a lot still need to be improved, especially deep and bottom water formation (section 4).

How to improve deep water formation in climate models then? A higher horizontal resolution may not be the answer as, depending on the model, it either reduces (Danek et al., 2019) or increases even further deep convection (Koenigk et al., 2020). In the ocean component, one solution could be a more systematic inclusion of overflow parameterisation (Snow et al., 2015);
550 in this study, it seems very effective for CESM2. The one data-assimilating model, NorCPM (Counillon et al., 2016), also proposes an interesting option. In the rest of the model, improving the representation of the stratosphere seems effective at reducing open ocean deep convection (Haase et al., 2018). Whatever the future holds, we hope it will feature a more systematic archiving of useful parameters. The situation has improved since CMIP5, but there are still CMIP6 models that do not provide their monthly mixed layer depth, and overturning streamfunctions (especially in density space) are a rarity. Making output
555 directly available on cloud computing based systems such as PANGEO (Odaka et al., 2020) should also be a priority, to let researchers work on heavy CMIP data as soon as they are released, regardless of their computing and storage capacities.

Code availability. Codes can be provided upon reasonable request

Data availability. CMIP6 data are freely available via any portal of the Earth System Grid Federation; for this manuscript, we mostly used <https://esgf-data.dkrz.de/projects/cmip6-dkrz/>. The World Ocean Atlas 2018 data can be accessed freely at <https://www.nodc.noaa.gov/OC5/woa18/woa18data.html>; the de Boyer Montégut et al. (2004) mixed layer depth reference data, at http://www.ifremer.fr/cerweb/deboyer/mld/Surface_Mixed_Layer_Depth.php; the GEBCO reference bathymetry, at <https://www.gebco.net/>.
560

Video supplement. Two videos of monthly bottom density around Antarctica over the entire historical run are available as supplement: in ACCESS-CM2 that has no overflow (<https://doi.org/10.5446/47545>), and in NorESM2-MM, which exhibits overflows very clearly (<https://doi.org/10.5446/47544>)

In this appendix you will find:

- [Fig A1, comparison between 'mlostst' and the mixed layer depth computed from 'thetao' and 'so' for one model;](#)
- Figs A2 and A3, Southern Ocean bottom salinity and temperature respectively, to complement the bias discussion of section 3.1;
- 570 – Figs A4 and A5, North Atlantic bottom salinity and temperature respectively, to complement the bias discussion of section 3.2;
- Table A1, maximum mixed layer depth and convective area for each model and each region, corresponding to the DMVs discussed in sections 3.1 and 3.2;
- Table A2, salinity and temperature of NADW and AABW for each model, briefly discussed in sections 3.1 and 3.2, and
- 575 used for the thickness computations for Figs. 5 and 6.




Figure A1. [Maximum monthly mixed layer depth in the North Atlantic over 1985-2014 for the model CanESM5: left, using the model output 'mlostst'; right, when computed from the monthly temperature and salinity. Over the entire 30-year period, the root mean square error in the Nordic seas is 305 m; in the subpolar gyre, 21 m.](#)




Figure A2. Southern Ocean reference bottom practical salinity (top left panel, top colorbar), and for each CMIP6 model, bottom practical salinity bias (model minus reference) averaged over 1985-2014. White number for each model is its RMSE over the entire Southern Ocean deeper than 1000 m. Thick black line indicates maximum mixed layer deeper than 2000 m. Thin grey line, the 2000 m isobath.




Figure A3. Southern Ocean reference bottom potential temperature (top left panel, top colorbar), and for each CMIP6 model, bottom potential temperature bias (model minus reference) averaged over 1985-2014. White number for each model is its RMSE over the entire Southern Ocean deeper than 1000 m. Thick black line indicates maximum mixed layer deeper than 2000 m. Thin grey line, the 2000 m isobath.



Figure A4. North Atlantic reference bottom practical salinity (top left panel, top colorbar), and for each CMIP6 model, bottom practical salinity bias (model minus reference) averaged over 1985–2014. White numbers for each model is its RMSE over the GIN (top) and SPG (bottom) areas, for depths over 1000 m. Thick black line indicates maximum mixed layer deeper than 1000 m; cyan dotted line in GIN, deeper than 700 m. Thin grey line, the 1000 m isobath.




Figure A5. North Atlantic reference bottom potential temperature (top left panel, top colorbar), and for each CMIP6 model, bottom potential temperature bias (model minus reference) averaged over 1985-2014. White numbers for each model is its RMSE over the GIN (top) and SPG (bottom) areas, for depths over 1000 m. Thick black line indicates maximum mixed layer deeper than 1000 m; cyan dotted line in GIN, deeper than 700 m. Thin grey line, the 1000 m isobath.

Table A1. Supplementary table version of tables 2 and 3 showing 30 year max MLD (m) and max area (in 10 000 km², which is the approximate area of a 1° cell).

model	SPG		GIN		Weddell		Amery		Ross	
	depth	area	depth	area	depth	area	depth	area	depth	area
ACCESS-CM2	2550	123	3623	53	5087	134	0	0	4428	91
ACCESS-ESM1-5	3013	108	3620	52	5328	125	0	0	4425	77
BCC-CSM2-MR	3787	216	3305	36	5334	183	3221	4	0	0
BCC-ESM1	3787	160	3787	34	5334	317	0	0	3305	4
CAMS-CSM1-0	2897	152	2810	31	5316	113	2698	3	3831	4
CESM2	2280	143	2017	22	0	0	0	0	0	0
CESM2-FV2	3102	115	2061	15	0	0	0	0	0	0
CESM2-WACCM	2392	130	2102	20	0	0	0	0	0	0
CESM2-WACCM-FV2	2872	138	1858	22	0	0	0	0	0	0
CNRM-CM6-1	3770	69	3699	59	0	0	4849	48	0	0
CNRM-ESM2-1	3930	34	3699	59	0	0	4849	81	0	0
CanESM5	1269	3	3216	69	2264	0.2	0	0	0	0
EC-Earth3	3071	55	3699	64	5306	89	4496	28	0	0
EC-Earth3-Veg	4006	72	3699	58	5374	49	4737	23	0	0
GFDL-CM4	4500	216	3500	36	6000	291	3500	2	4500	16
GFDL-ESM4	3760	148	3734	60	0	0	0	0	4214	18
GISS-E2-1-G	4008	225	3342	48	0	0	0	0	0	0
GISS-E2-1-G-CC	4007	204	3341	42	0	0	0	0	0	0
GISS-E2-1-H	3000	263	3500	90	4500	63	3898	21	3159	3
HadGEM3-GC31-LL	3826	59	3699	53	5395	21	0	0	2333	0
INM-CM5-0	1360	14	0	0	4500	111	2590	10	2886	37
IPSL-CM6A-LR	2686	20	3699	56	5036	60	0	0	2472	1
MCM-UA-1-0	4662	1	3373	1	4662	3	4662	0.4	4662	1
MIROC-ES2L	2590	41	4065	88	6240	104	0	0	5190	156
MIROC6	4740	144	4065	102	6240	234	5190	23	5190	289
MPI-ESM-1-2-HAM	3770	89	3395	70	5131	107	3760	4	4195	37
MPI-ESM1-2-HR	3388	101	3033	33	5170	94	0	0	4195	27
MPI-ESM1-2-LR	3395	47	1829	64	4872	68	3770	6	4195	18
MRI-ESM2-0	4033	135	3650	52	2394	1	2958	2	2705	2
NESM3	3292	118	1951	32	3772	50	0	0	4506	52
NorCPM1	1005	0.3	0	0	0	0	0	0	0	0
NorESM2-LM	2741	133	3614	61	5410	215	4191	16	2844	10
NorESM2-MM	2770	130	2836	82	5408	217	4297	44	3323	51
SAM0-UNICON	3843	193	3380	98	0	0	0	0	2649	13
UKESM1-0-LL	3750	52	3699	85	5037	22	2195	0.2	0	0

Table A2. For each CMIP6 model, 30 year median practical salinity S and potential temperature θ ($^{\circ}\text{C}$) of the NADW formed in the subpolar gyre (SPG) or Nordic seas (GIN), and of the AABW.

model	NADW (SPG)		NADW (GIN)		AABW	
	S	θ	S	θ	S	θ
ACCESS-CM2	35.126 \pm 0.024	5.95 \pm 0.26	34.988 \pm 0.021	0.65 \pm 0.17	34.613 \pm 0.045	-0.84 \pm 0.15
ACCESS-ESM1-5	35.328 \pm 0.050	5.89 \pm 0.32	35.176 \pm 0.031	0.46 \pm 0.26	34.581 \pm 0.005	-1.06 \pm 0.02
BCC-CSM2-MR	35.155 \pm 0.038	6.00 \pm 0.19	35.129 \pm 0.002	1.92 \pm 0.00	34.440 \pm 0.025	-0.44 \pm 0.05
BCC-ESM1	35.086 \pm 0.022	6.12 \pm 0.16	35.043 \pm 0.010	2.20 \pm 0.17	34.308 \pm 0.011	-0.91 \pm 0.09
CAMS-CSM1-0	35.104 \pm 0.004	6.04 \pm 0.14	34.909 \pm 0.001	0.77 \pm 0.01	34.423 \pm 0.011	-0.28 \pm 0.07
CESM2	35.264 \pm 0.018	5.37 \pm 0.18	35.064 \pm 0.010	1.97 \pm 0.11	34.677 \pm 0.030	-0.50 \pm 0.34
CESM2-FV2	35.242 \pm 0.027	5.17 \pm 0.22	35.023 \pm 0.005	1.84 \pm 0.10	34.713 \pm 0.014	-0.63 \pm 0.08
CESM2-WACCM	35.260 \pm 0.025	5.22 \pm 0.19	35.112 \pm 0.004	1.75 \pm 0.07	34.682 \pm 0.016	-0.31 \pm 0.15
CESM2-WACCM-FV2	35.258 \pm 0.026	5.40 \pm 0.19	35.036 \pm 0.013	2.39 \pm 0.07	34.683 \pm 0.017	-0.53 \pm 0.10
CNRM-CM6-1	34.968 \pm 0.030	3.57 \pm 0.19	34.927 \pm 0.025	0.06 \pm 0.33	34.679 \pm 0.001	0.25 \pm 0.02
CNRM-ESM2-1	35.009 \pm 0.026	4.12 \pm 0.13	34.904 \pm 0.024	0.12 \pm 0.26	34.641 \pm 0.003	1.00 \pm 0.04
CanESM5	34.981 \pm 0.008	4.77 \pm 0.07	34.799 \pm 0.017	0.54 \pm 0.10	34.655 \pm 0.015	-1.07 \pm 0.02
EC-Earth3	35.086 \pm 0.028	4.90 \pm 0.13	34.902 \pm 0.060	0.46 \pm 0.64	34.601 \pm 0.000	0.13 \pm 0.01
EC-Earth3-Veg	35.261 \pm 0.018	5.56 \pm 0.12	35.068 \pm 0.024	1.73 \pm 0.23	34.607 \pm 0.014	0.17 \pm 0.02
GFDL-CM4	35.179 \pm 0.025	5.25 \pm 0.21	34.936 \pm 0.004	1.20 \pm 0.63	34.460 \pm 0.019	-1.08 \pm 0.10
GFDL-ESM4	35.251 \pm 0.021	5.40 \pm 0.16	35.102 \pm 0.017	0.58 \pm 0.17	34.563 \pm 0.037	-0.57 \pm 0.15
GISS-E2-1-G	35.197 \pm 0.027	4.81 \pm 0.14	34.843 \pm 0.004	0.39 \pm 0.05	34.559 \pm 0.011	-0.31 \pm 0.08
GISS-E2-1-G-CC	35.193 \pm 0.039	4.80 \pm 0.18	34.840 \pm 0.006	0.57 \pm 0.05	34.551 \pm 0.010	-0.23 \pm 0.08
GISS-E2-1-H	35.010 \pm 0.024	3.96 \pm 0.31	34.719 \pm 0.008	1.47 \pm 0.15	34.788 \pm 0.033	0.96 \pm 0.02
HadGEM3-GC31-LL	35.193 \pm 0.013	4.72 \pm 0.09	34.991 \pm 0.008	0.07 \pm 0.03	34.628 \pm 0.032	-0.17 \pm 0.16
INM-CM5-0	35.310 \pm 0.007	2.97 \pm 0.21	35.512 \pm 0.003	-1.59 \pm 0.01	35.106 \pm 0.017	-0.45 \pm 0.12
IPSL-CM6A-LR	35.032 \pm 0.018	3.89 \pm 0.11	35.001 \pm 0.005	1.70 \pm 0.05	34.662 \pm 0.007	-1.25 \pm 0.01
MCM-UA-1-0	34.697 \pm 0.039	4.10 \pm 0.31	34.708 \pm 0.009	1.60 \pm 0.29	34.265 \pm 0.003	-1.23 \pm 0.10
MIROC-ES2L	34.946 \pm 0.007	5.25 \pm 0.25	34.781 \pm 0.014	0.73 \pm 0.10	34.347 \pm 0.032	1.56 \pm 0.20
MIROC6	35.081 \pm 0.004	4.86 \pm 0.03	34.953 \pm 0.015	1.55 \pm 0.07	34.419 \pm 0.018	0.22 \pm 0.13
MPI-ESM-1-2-HAM	35.195 \pm 0.013	5.94 \pm 0.13	35.128 \pm 0.003	1.50 \pm 0.04	34.581 \pm 0.012	-0.12 \pm 0.05
MPI-ESM1-2-HR	35.175 \pm 0.033	6.07 \pm 0.22	34.954 \pm 0.003	1.05 \pm 0.06	34.554 \pm 0.002	-0.35 \pm 0.04
MPI-ESM1-2-LR	35.163 \pm 0.017	6.10 \pm 0.18	35.101 \pm 0.001	1.19 \pm 0.02	34.554 \pm 0.009	-0.01 \pm 0.05
MRI-ESM2-0	35.094 \pm 0.024	4.22 \pm 0.33	35.042 \pm 0.009	2.16 \pm 0.11	34.526 \pm 0.025	-1.21 \pm 0.10
NESM3	34.894 \pm 0.013	3.96 \pm 0.65	34.896 \pm 0.001	-0.36 \pm 0.00	34.689 \pm 0.000	0.84 \pm 0.01
NorCPM1	35.227 \pm 0.011	4.49 \pm 0.08	35.276 \pm 0.000	-1.52 \pm 0.00	34.659 \pm 0.000	-0.72 \pm 0.00
NorESM2-LM	35.373 \pm 0.013	4.54 \pm 0.13	35.157 \pm 0.024	-0.31 \pm 0.16	34.543 \pm 0.115	-1.48 \pm 0.11
NorESM2-MM	35.391 \pm 0.014	4.48 \pm 0.21	35.376 \pm 0.042	-0.74 \pm 0.44	34.796 \pm 0.137	-1.57 \pm 0.07
SAM0-UNICON	35.138 \pm 0.030	4.43 \pm 0.66	35.055 \pm 0.007	0.93 \pm 0.06	34.745 \pm 0.016	-1.04 \pm 0.07
UKESM1-0-LL	35.141 \pm 0.041	4.43 \pm 0.21	34.990 \pm 0.026	0.38 \pm 0.22	34.639 \pm 0.037	-0.74 \pm 0.20
multimodel mean	35.163 \pm 0.143	4.86 \pm 0.81	35.001 \pm 0.169	0.77 \pm 0.99	34.607 \pm 0.154	-0.45 \pm 0.73

Appendix B: Transports

In this section, you will find two tables to complement section 3.3:

- Table B1 presents the AMOC and southernmost extent of NADW in each model;
- Table B2 presents the SMOC and northernmost extent of AABW in the Atlantic, Indian and Pacific oceans, in each model.

Table B1. For each CMIP6 model, 30 year median AMOC at 35°N (in Sv), and southernmost latitude (in degrees north) of the 2000 m -thick NADW layer in the Atlantic from Fig. 6.

model	AMOC	latitude
ACCESS-CM2	19.8 ± 3.5	48.5
ACCESS-ESM1-5	20.0 ± 2.1	44.5
BCC-CSM2-MR	26.1 ± 2.8	-49.5
BCC-ESM1	25.3 ± 3.7	-59.5
CAMS-CSM1-0	14.8 ± 3.5	50.5
CESM2	24.9 ± 6.5	-49.5
CESM2-FV2	25.6 ± 6.4	-50.5
CESM2-WACCM	24.7 ± 7.3	-49.5
CESM2-WACCM-FV2	24.8 ± 7.0	-50.5
CNRM-CM6-1	19.4 ± 4.9	-46.5
CNRM-ESM2-1	20.0 ± 5.1	-47.5
CanESM5	15.1 ± 5.7	60
EC-Earth3	17.8 ± 5.1	41.5
EC-Earth3-Veg	19.5 ± 5.0	31.5
GFDL-CM4	8.9 ± 10.7	-43.5
GFDL-ESM4	N/A	-48.5
GISS-E2-1-G	22.2 ± 4.8	-51.5
GISS-E2-1-G-CC	24.0 ± 4.1	-50.5
GISS-E2-1-H	16.4 ± 12.9	60
HadGEM3-GC31-LL	18.8 ± 4.0	50.5
INM-CM5-0	63.1 ± 19.8	60
IPSL-CM6A-LR	13.4 ± 5.3	60
MCM-UA-1-0	17.9 ± 2.4	8.5
MIROC-ES2L	15.0 ± 6.0	47.5
MIROC6	19.0 ± 6.2	47.5
MPI-ESM-1-2-HAM	29.6 ± 8.7	2.5
MPI-ESM1-2-HR	1.8 ± 12.8	-50.5
MPI-ESM1-2-LR	25.4 ± 7.4	-13.5
MRI-ESM2-0	18.6 ± 17.7	-48.5
NESM3	8.8 ± 4.2	19.5
NorCPM1	N/A	-49.5
NorESM2-LM	18.0 ± 7.3	60
NorESM2-MM	21.4 ± 7.6	40.5
SAM0-UNICON	24.9 ± 5.6	-50.5
UKESM1-0-LL	18.6 ± 4.5	40.5
multimodel median	19.5 ± 9.5	

Table B2. For each CMIP6 model, 30 year median southern MOC at 30°S (SMOC, in Sv) and northernmost latitude (in degrees north) of the 2000 m -thick AABW layer in each ocean from Fig. 5.

model	Atlantic		Indian		Pacific	
	SMOC	lat.	SMOC	lat.	SMOC	lat.
ACCESS-CM2	4.2 ± 1.2	-35.5	1.5 ± 3.1	16.5	3.9 ± 3.8	54.5
ACCESS-ESM1-5	3.6 ± 1.0	-34.5	1.9 ± 3.3	21.5	5.9 ± 3.5	54.5
BCC-CSM2-MR	2.7 ± 1.1	7.5	4.8 ± 9.1	16.5	6.7 ± 3.5	57.5
BCC-ESM1	4.2 ± 1.5	13.5	3.6 ± 8.3	17.5	8.2 ± 2.7	58.5
CAMS-CSM1-0	5.9 ± 1.8	8.5	1.6 ± 4.6	-12.5	2.8 ± 3.5	58.5
CESM2	2.3 ± 1.2	-38.5	2.2 ± 2.8	-49.5	3.9 ± 4.0	52.5
CESM2-FV2	2.3 ± 1.2	-40.5	2.0 ± 3.1	-51.5	3.5 ± 3.7	17.5
CESM2-WACCM	2.1 ± 1.2	-38.5	1.9 ± 3.0	-39.5	3.3 ± 3.7	53.5
CESM2-WACCM-FV2	2.7 ± 1.3	-38.5	1.6 ± 3.0	-50.5	3.2 ± 3.7	52.5
CNRM-CM6-1	1.5 ± 1.7	-26.5	3.4 ± 4.6	21.5	6.3 ± 4.8	57.5
CNRM-ESM2-1	1.8 ± 1.6	-31.5	3.1 ± 4.8	19.5	6.2 ± 4.5	60.5
CanESM5	4.0 ± 1.6	-3.5	3.7 ± 2.8	11.5	6.3 ± 3.4	57.5
EC-Earth3	3.8 ± 2.2	14.5	1.5 ± 4.5	14.5	4.9 ± 4.2	60.5
EC-Earth3-Veg	2.8 ± 2.1	-25.5	1.3 ± 4.8	25.5	4.3 ± 4.1	59.5
GFDL-CM4	3.0 ± 2.4	13.5	11.1 ± 18.2	11.5	3.2 ± 3.1	59.5
GFDL-ESM4	N/A	14.5	N/A	25.5	N/A	60.5
GISS-E2-1-G	0.4 ± 0.5	-40.5	8.7 ± 5.7	-43.5	10.1 ± 5.8	-42.5
GISS-E2-1-G-CC	0.3 ± 0.5	-41.5	8.9 ± 5.6	-37.5	10.9 ± 6.2	-44.5
GISS-E2-1-H	0.2 ± 1.6	-33.5	10.3 ± 8.3	-40.5	7.3 ± 6.7	-53.5
HadGEM3-GC31-LL	3.1 ± 1.9	-26.5	2.3 ± 3.4	25.5	7.1 ± 4.8	59.5
INM-CM5-0	3.4 ± 1.9	52.5	3.0 ± 2.8	-50	10.8 ± 3.9	-50
IPSL-CM6A-LR	3.8 ± 2.6	-2.5	2.3 ± 4.4	16.5	5.8 ± 5.4	60.5
MCM-UA-1-0	3.5 ± 0.6	-33.5	1.5 ± 1.6	21.5	3.9 ± 1.9	-23.5
MIROC-ES2L	0.3 ± 0.5	13.5	5.1 ± 4.2	17.5	12.1 ± 5.3	58.5
MIROC6	4.0 ± 1.3	14.5	5.1 ± 3.7	23.5	13.6 ± 4.0	60.5
MPI-ESM-1-2-HAM	2.9 ± 1.5	13.5	3.1 ± 5.3	-30.5	3.0 ± 3.4	60.5
MPI-ESM1-2-HR	5.8 ± 2.1	13.5	4.1 ± 4.9	-17.5	5.1 ± 3.5	60.5
MPI-ESM1-2-LR	2.9 ± 1.6	13.5	3.3 ± 4.9	-30.5	2.9 ± 3.7	60.5
MRI-ESM2-0	2.8 ± 1.2	-27.5	2.6 ± 4.0	25.5	6.6 ± 5.6	56.5
NESM3	1.4 ± 1.4	-6.5	2.3 ± 4.8	-4.5	4.4 ± 4.7	57.5
NorCPM1	N/A	-47.5	N/A	-47.5	N/A	-23.5
NorESM2-LM	1.7 ± 1.1	-39.5	3.3 ± 4.3	-28.5	10.5 ± 4.8	60.5
NorESM2-MM	1.4 ± 1.0	5.5	4.0 ± 4.8	14.5	10.6 ± 5.3	59.5
SAM0-UNICON	1.7 ± 1.1	-42.5	2.9 ± 3.8	-49.5	5.6 ± 4.5	-11.5
UKESM1-0-LL	3.4 ± 1.9	-36.5	3.0 ± 3.3	25.5	9.0 ± 4.2	59.5
multimodel median	2.8 ± 1.4		3.0 ± 2.5		5.9 ± 3.0	

Author contributions. C.H. designed and conducted the analyses, and wrote the paper.

Competing interests. The author declares no competing interest.

Acknowledgements. This work is supported by the Swedish Research Council (dnr. 2018-03859). We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modeling groups
585 (whose models are listed in Table 1 of this paper) for producing and making available their model output. [The author thanks the two anonymous reviewers whose comments greatly improved the quality of this manuscript.](#) C.H. would [also](#) like to thank Jonathan Rheinländer for the constructive discussion that inspired this [manuscript](#)~~work~~, Martin Mohrmann for the regular CMIP6 deep convection chats during its writing, and Matthew Menary for freely sharing his AMOC data (in fact, not used in this manuscript).

References

- Armour, K.: Energy budget constraints on climate sensitivity in light of inconstant climate feedbacks, *Nature Climate Change*, 7, <https://doi.org/10.1038/nclimate3278>, 2017.
- Årthun, M., Eldevik, T., and Smedsrud, L.: The Role of Atlantic Heat Transport in Future Arctic Winter Sea Ice Loss, *Journal of Climate*, 32, <https://doi.org/10.1175/JCLI-D-18-0750.1>, 2019.
- Ba, J., Keenlyside, N., Latif, M., Park, W., Ding, H., Lohmann, K., Mignot, J., Menary, M., Otterå, O., Wouters, B., and Salas y Melia, D.: A multi-model comparison of Atlantic multidecadal variability, *Climate dynamics*, 43, <https://doi.org/10.1007/s00382-014-2056-1>, 2014.
- Beadling, R., Russell, J., Stouffer, R., Mazloff, M., Talley, L., Goodman, P., Sallée, J., Hewitt, H., Hyder, P., and Pandde, A.: Representation of Southern Ocean properties across Coupled Model Intercomparison Project generations: CMIP3 to CMIP6, *Journal of Climate*, EOR, <https://doi.org/10.1175/JCLI-D-19-0970.1>, 2020.
- Behrens, E., Rickard, G., Morgenstern, O., Martin, T., Osprey, A., and Joshi, M.: Southern Ocean deep convection in global climate models: A driver for variability of subpolar gyres and Drake Passage transport on decadal timescales, *Journal of Geophysical Research: Oceans*, 121, <https://doi.org/10.1002/2015JC011286>, 2016.
- Briegleb, P., Danabasoglu, G., and Large, G.: An overflow parameterization for the ocean component of the Community Climate System Model, <https://doi.org/10.5065/D69K4863>, 2010.
- Brodeau, L. and Koenigk, T.: Extinction of the northern oceanic deep convection in an ensemble of climate model simulations of the 20th and 21st centuries, *Climate Dynamics*, 46, <https://doi.org/10.1007/s00382-015-2736-5>, 2016.
- Broecker, W. S.: *The Glacial World According to Wally*, Eldigio Press, 2 edn., 1995.
- Cabré, A., Marinov, I., and Gnanadesikan, A.: Global atmospheric teleconnections and multidecadal climate oscillations driven by Southern Ocean convection, *Journal of Climate*, 30, <https://doi.org/10.1175/JCLI-D-16-0741.1>, 2017.
- Campbell, E., Wilson, E., Moore, G., Riser, S., Brayton, C., Mazloff, M., and Talley, L.: Antarctic offshore polynyas linked to Southern Hemisphere climate anomalies, *Nature*, 570, <https://doi.org/10.1038/s41586-019-1294-0>, 2019.
- Cao, J., Wang, B., Yang, Y.-M., Ma, L., Li, J., Sun, B., Bao, Y., He, J., Zhou, X., and Wu, L.: The NUIST Earth System Model (NESM) version 3: description and preliminary evaluation, *Geoscientific Model Development*, 11, <https://doi.org/10.5194/gmd-11-2975-2018>, 2018.
- Chen, H., Morrison, A., Dufour, C., and Sarmiento, J.: Deciphering patterns and drivers of heat and carbon storage in the Southern Ocean, *Geophysical Research Letters*, 46, <https://doi.org/10.1029/2018GL080961>, 2019.
- Counillon, F., Keenlyside, N., Bethke, I., Wang, Y., Billeau, S., Shen, M., and Bentsen, M.: Flow-dependent assimilation of sea surface temperature in isopycnal coordinates with the Norwegian Climate Prediction Model, *Tellus A: Dynamic Meteorology and Oceanography*, 68, <https://doi.org/10.3402/tellusa.v68.32437>, 2016.
- Cox, P., Huntingford, C., and Williamson, M.: Emergent constraint on equilibrium climate sensitivity from global temperature variability, *Nature*, 553, <https://doi.org/10.1038/nature25450>, 2018.
- Danabasoglu, G., Lamarque, J., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., and Emmons et al., L. K.: The community earth system model version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, <https://doi.org/10.1029/2019MS001916>, 2020.
- Danek, C., Scholz, P., and Lohmann, G.: Effects of high resolution and spinup time on modeled North Atlantic circulation, *Journal of Physical Oceanography*, 49, <https://doi.org/10.1175/JPO-D-18-0141.1>, 2019.
- de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: an examination of profile data and a profile-based climatology, *Journal of Geophysical Research*, 109, <https://doi.org/10.1029/2004JC002378>, 2004.

- De Lavergne, C., Palter, J., Galbraith, E., Bernardello, R., and Marinov, I.: Cessation of deep convection in the open Southern Ocean under anthropogenic climate change, *Nature Climate Change*, 4, <https://doi.org/10.1038/nclimate2132>, 2014.
- Drucker, R., Martin, S., and Kwok, R.: Sea ice production and export from coastal polynyas in the Weddell and Ross Seas, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL048668>, 2011.
- 630 Duchez, A., Courtois, P., Harris, E., Josey, S., Kanzow, T., Marsh, R., Smeed, D., and Hirschi, J.: Potential for seasonal prediction of Atlantic sea surface temperatures using the RAPID array at 26°N, *Climate Dynamics*, 46, <https://doi.org/10.1007/s00382-015-2918-1>, 2016.
- Eyring, V., Bony, S., Meehl, G., Senior, C., Stevens, B., Stouffer, R., and Taylor, K.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 635 GEBCO Compilation Group: GEBCO 2019 Grid, <https://doi.org/10.5285/836f016a-33be-6ddc-e053-6c86abc0788e>, 2019.
- Haase, S., Matthes, K., Latif, M., and Omrani, N.: The importance of a properly represented stratosphere for northern hemisphere surface variability in the atmosphere and the ocean, *Journal of Climate*, 31, <https://doi.org/10.1175/JCLI-D-17-0520.1>, 2018.
- Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geoscientific Model Development*, 13, <https://doi.org/10.5194/gmd-13-2197-2020>, 2020.
- 640 Held, I., Guo, H., Adcroft, A., Dunne, J., Horowitz, L., Krasting, J., Shevliakova, E., Winton, M., Zhao, M., Bushuk, M., and Wittenberg, A.: Structure and performance of GFDL's CM4. 0 climate model, *Journal of Advances in Modeling Earth Systems*, 11, <https://doi.org/10.1029/2019MS001829>, 2019.
- Heuzé, C.: North Atlantic deep water formation and AMOC in CMIP5 models, *Ocean Science*, 13, <https://doi.org/10.5194/os-13-609-2017>, 2017.
- 645 Heuzé, C. and Årthun, M.: The Atlantic inflow across the Greenland-Scotland ridge in global climate models (CMIP5), *Elementa Science of the Anthropocene*, 7, <https://doi.org/10.1525/elementa.354/>, 2019.
- Heuzé, C., Heywood, K., Stevens, D., and Ridley, J.: Southern Ocean bottom water characteristics in CMIP5 models, *Geophysical Research Letters*, 40, <https://doi.org/10.1002/grl.50287>, 2013.
- 650 Heuzé, C., Heywood, K., Stevens, D., and Ridley, J.: Changes in global ocean bottom properties and volume transports in CMIP5 models under climate change scenarios, *Journal of Climates*, 28, <https://doi.org/10.1175/JCLI-D-14-00381.1>, 2015.
- Holt, J., Hyder, P., Ashworth, M., Harle, J., Hewitt, H., Liu, H., New, A., Pickles, S., Porter, A., Popova, E., and Allen, J.: Prospects for improving the representation of coastal and shelf seas in global ocean models, *Geoscientific Model Development*, 10, <https://doi.org/10.5194/gmd-10-499-2017>, 2017.
- 655 Huussen, T., Naveira-Garabato, A., Bryden, H., and McDonagh, E.: Is the deep Indian Ocean MOC sustained by breaking internal waves?, *Journal of Geophysical Research*, 117, <https://doi.org/10.1029/2012JC008236>, 2012.
- Jenkins, A.: The impact of melting ice on ocean waters, *Journal of physical oceanography*, 29, [https://doi.org/10.1175/1520-0485\(1999\)029<2370:TIOMIO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<2370:TIOMIO>2.0.CO;2), 1999.
- Johnson, G.: Quantifying Antarctic bottom water and North Atlantic deep water volumes, *Journal of Geophysical Research: Oceans*, 113, <https://doi.org/10.1029/2007JC004477>, 2008.
- 660 Killworth, P.: Deep convection in the world ocean, *Reviews of Geophysics*, 21, <https://doi.org/10.1029/RG021i001p00001>, 1983.

- Koenigk, T., Fuentes-Franco, R., Meccia, V., Gutjahr, O., Jackson, L. C., New, A. L., Ortega, P., Roberts, C., Roberts, M., Arsouze, T., Iovino, D., Moine, M., and Sein, D. V.: Deep water formation in the North Atlantic Ocean in high resolution global coupled climate models, *Ocean Science Discuss.*, <https://doi.org/10.5194/os-2020-41>, 2020.
- 665 Kuhlbrodt, T., Jones, C., Sellar, A., Storkey, D., Blockley, E., Stringer, M., Hill, R., Graham, T., Ridley, J., Blaker, A., and Calvert, D.: The low-resolution version of HadGEM3 GC3. 1: Development and evaluation for global climate, *Journal of Advances in Modeling Earth Systems*, 10, <https://doi.org/10.1029/2018MS001370>, 2018.
- Lin, P., Yu, Z., Lü, J., Ding, M., Hu, A., and Liu, H.: Two regimes of Atlantic multidecadal oscillation: cross-basin dependent or Atlantic-intrinsic, *Science Bulletin*, 64, <https://doi.org/10.1016/j.scib.2018.12.027>, 2019.
- 670 Lique, C. and Thomas, M.: Latitudinal shift of the Atlantic Meridional Overturning Circulation source regions under a warming climate, *Nature Clim Change*, 8, <https://doi.org/10.1038/s41558-018-0316-5>, 2018.
- Liu, W., Fedorov, A., and Sévellec, F.: The mechanisms of the Atlantic meridional overturning circulation slowdown induced by Arctic sea ice decline, *Journal of Climate*, 32, <https://doi.org/10.1175/JCLI-D-18-0231.1>, 2019.
- Locarnini, R., Mishonov, A., Baranova, O., Boyer, T., Zweng, M., Garcia, H., Reagan, J., Seidov, D., Weathers, K., Paver, C., and Smolyar, I.: World Ocean Atlas 2018, Volume 1: Temperature, A. Mishonov Technical Ed.; NOAA Atlas NESDIS 81, 2018.
- 675 Lozier, M., Li, F., Bacon, S., Bahr, F., Bower, A., Cunningham, S., De Jong, M., De Steur, L., Deyoung, B., Fischer, J., and Gary, S.: A sea change in our view of overturning in the subpolar North Atlantic, *Science*, 363, <https://doi.org/10.1126/science.aau6592>, 2019.
- Lumpkin, R. and Speer, K.: Global ocean meridional overturning, *Journal of Physical Oceanography*, 37, <https://doi.org/10.1175/JPO3130.1>, 2007.
- 680 Lurton, T., Balkanski, Y., Bastrikov, V., Bekki, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Contoux, C., Cozic, A., and Cugnet, D.: Implementation of the CMIP6 Forcing Data in the IPSL-CM6A-LR Model, *Journal of Advances in Modeling Earth Systems*, 12, <https://doi.org/10.1029/2019MS001940>, 2020.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., and Fast, I.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1. 2) and its response to increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, <https://doi.org/10.1029/2018MS001400>, 2019.
- 685 Meijers, A., Shuckburgh, E., Bruneau, N., Sallée, J., Bracegirdle, T., and Wang, Z.: Representation of the Antarctic Circumpolar Current in the CMIP5 climate models and future changes under warming scenarios, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/10.1029/2012JC008412>, 2012.
- Menary, M. and Wood, R.: An anatomy of the projected North Atlantic warming hole in CMIP5 models, *Climate Dynamics*, 50, <https://doi.org/10.1007/s00382-017-3793-8>, 2018.
- 690 Menary, M., Hodson, D., Robson, J., Sutton, R., Wood, R., and Hunt, J.: Exploring the impact of CMIP5 model biases on the simulation of North Atlantic decadal variability, *Geophysical Research Letters*, 42, <https://doi.org/10.1002/2015GL064360>, 2015.
- Menary, M., Robson, J., Allan, R., Booth, B., Cassou, C., Gastineau, G., Gregory, J., Hodson, D., Jones, C., Mignot, J., Ringer, M., Sutton, R., Wilcox, L., and Zhang, R.: Aerosol-forced AMOC changes in CMIP6 historical simulations, *Geophysical Research Letters*, accepted, 2020.
- 695 Menviel, L., Spence, P., Skinner, L., Tachikawa, K., Friedrich, T., Missiaen, L., and Yu, J.: Enhanced Mid-depth Southward Transport in the Northeast Atlantic at the Last Glacial Maximum Despite a Weaker AMOC, *Paleoceanography and Paleoclimatologies*, 35, <https://doi.org/10.1029/2019PA003793>, 2020.
- Mohrmann, M., Heuzé, C., and Swart, S.: Southern Ocean polynyas in CMIP6 models, *The Cryosphere*, *subm.*

- Müller, W., Jungclaus, J., Mauritsen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M., Ghosh, R., Haak, H., and Ilyina, T.: A Higher-resolution Version of the Max Planck Institute Earth System Model (MPI-ESM1. 2-HR), *Journal of Advances in Modeling Earth Systems*, 10, <https://doi.org/10.1029/2017MS001217>, 2018.
- Notz, D., Jahn, A., Holland, M., Hunke, E., Massonnet, F., Stroeve, J., Tremblay, B., and Vancoppenolle, M.: The CMIP6 Sea-Ice Model Intercomparison Project (SIMIP): understanding sea ice through climate-model simulations, *Geoscientific Model Development*, 9, <https://doi.org/10.5194/gmd-9-3427-2016>, 2016.
- Nowicki, S., Payne, T., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and Shepherd, A.: Ice sheet model intercomparison project (ISMIP6) contribution to CMIP6, *Geoscientific Model Development*, 9, <https://doi.org/10.5194/gmd-9-4521-2016>, 2016.
- Odaka, T., Banihirwe, A., Eynard-Bontemps, G., Ponte, A., Maze, G., Paul, K., Baker, J., and Abernathy, R.: *Tools and Techniques for High Performance Computing*, Springer, 2020.
- Ohshima, K., Fukamachi, Y., Williams, G., Nihashi, S., Roquet, F., Kitade, Y., Tamura, T., Hirano, D., Herraiz-Borreguero, L., Field, I., and Hindell, M.: Antarctic Bottom Water production by intense sea-ice formation in the Cape Darnley polynya, *Nature Geoscience*, 6, <https://doi.org/10.1038/ngeo1738>, 2013.
- Orsi, A.: Recycling bottom waters, *Nature Geoscience*, 3, <https://doi.org/10.1038/ngeo854>, 2010.
- Park, S., Shin, J., Kim, S., Oh, E., and Kim, Y.: Global climate simulated by the seoul national university atmosphere model version 0 with a unified convection scheme (sam0-unicon), *Journal of Climate*, 32, <https://doi.org/10.1175/JCLI-D-18-0796.1>, 2019.
- Patara, L. and Böning, C.: Abyssal ocean warming around Antarctica strengthens the Atlantic overturning circulation, *Geophysical Research Letters*, 41, <https://doi.org/10.1002/2014GL059923>, 2014.
- Roach, L., Dörr, J., Holmes, C., Massonnet, F., Blockley, E., Notz, D., Rackow, T., Raphael, M., O'Farrell, S., Bailey, D., and Bitz, C.: Antarctic sea ice area in CMIP6, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2019GL086729>, 2020.
- Rong, X. Y., Li, J., and Chen, H. M.: Introduction of CAMS-CSM model and its participation in CMIP6, *Climate Change Res*, 6, <https://doi.org/10.12006/j.issn.1673-1719.2019.186>, 2019.
- Sallée, J., Shuckburgh, E., Bruneau, N., Meijers, A., Bracegirdle, T., and Wang, Z.: Assessment of Southern Ocean mixed-layer depths in CMIP5 models: Historical bias and forcing response, *Journal of Geophysical Research: Oceans*, <https://doi.org/10.1002/jgrc.20157>, 2013.
- Séférián, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., and Sénési, S.: Evaluation of CNRM Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-Day and Future Climate, *Journal of Advances in Modeling Earth Systems*, <https://doi.org/10.1029/2019MS001791>, 2019.
- Sellar, A., Walton, J., Jones, C., Wood, R., Abraham, N., Andrejczuk, M., Andrews, M., Andrews, T., Archibald, A., de Mora, L., and Dyson, H.: Implementation of UK Earth system models for CMIP6, *Journal of Advances in Modeling Earth Systems*, <https://doi.org/10.1029/2019MS001946>, 2020.
- Shu, Q., Wang, Q., Song, Z., Qiao, F., Zhao, J., Chu, M., and Li, X.: Assessment of sea ice extent in CMIP6 with comparison to observations and CMIP5, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2020GL087965>, 2020.
- Snow, K., Hogg, A., Downes, S., Sloyan, B., Bates, M., and Griffies, S.: Sensitivity of abyssal water masses to overflow parameterisations, *Ocean Modelling*, 89, <https://doi.org/10.1016/j.ocemod.2015.03.004>, 2015.
- Swart, N., Cole, J., Kharin, V., Lazare, M., Scinocca, J., Gillett, N., Anstey, J., Arora, V., Christian, J., Hanna, S., and Jiao, Y.: The Canadian Earth System Model version 5 (CanESM5. 0.3), *Geoscientific Model Development*, 12, <https://doi.org/10.5194/gmd-2019-177>, 2019.

- Sweetman, A., Thurber, A., Smith, C., Levin, L., Mora, C., Wei, C., Gooday, A., Jones, D., Rex, M., Yasuhara, M., and Ingels, J.: Major impacts of climate change on deep-sea benthic ecosystems, *Elementa Science of the Anthropocene*, 5, <https://doi.org/10.1525/elementa.203>, 2017.
- 740 Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe, M., Saito, F., Chikira, M., Watanabe, S., Mori, M., Hirota, N., Kawatani, Y., Mochizuki, T., Yoshimura, K., Takata, K., O'ishi, R., Yamazaki, D., Suzuki, T., Kurogi, M., Kataoka, T., Watanabe, M., and Kimoto, M.: Description and basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6, *Geoscientific Model Development*, 12, <https://doi.org/10.5194/gmd-12-2727-2019>, 2019.
- Taylor, K., Stouffer, R., and Meehl, G.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 745 93, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- Tjiputra, J. F., Schwinger, J., Bentsen, M., Morée, A. L., Gao, S., Bethke, I., Heinze, C., Goris, N., Gupta, A., He, Y.-C., Olivié, D., Seland, Ø., and Schulz, M.: Ocean biogeochemistry in the Norwegian Earth System Model version 2 (NorESM2), *Geoscientific Model Development*, 13, <https://doi.org/10.5194/gmd-13-2393-2020>, 2020.
- Våge, K., Pickart, R., Thierry, V., Reverdin, G., Lee, C., Petrie, B., Agnew, T., Wong, A., and Ribergaard, M.: Surprising return of deep 750 convection to the subpolar North Atlantic Ocean in winter 2007–2008, *Nature Geoscience*, 2, <https://doi.org/10.1038/ngeo382>, 2009.
- Voldoire, A., Saint-Martin, D., Sénési, S., Decharme, B., Alias, A., Chevallier, M., and Colin et al., J.: Evaluation of CMIP6 DECK Experiments With CNRM-CM6-1, *Journal of Advances in Modeling Earth Systems*, 11, <https://doi.org/10.1029/2019MS001683>, 2019.
- Volodin, E. and Gritsun, A.: Simulation of observed climate changes in 1850–2014 with climate model INM-CM5, *Earth System Dynamics*, 9, <https://doi.org/10.5194/esd-9-1235-2018>, 2018.
- 755 Wang, Z., Wu, Y., Lin, X., Liu, C., and Xie, Z.: Impacts of open-ocean deep convection in the Weddell Sea on coastal and bottom water temperature, *Climate Dynamics*, 48, <https://doi.org/10.1007/s00382-016-3244-y>, 2017.
- Wu, T., Lu, Y., Fang, Y., Xin, X., Li, L., Li, W., Jie, W., Zhang, J., Liu, Y., Zhang, L., Zhang, F., Zhang, Y., Wu, F., Li, J., Chu, M., Wang, Z., Shi, X., Liu, X., Wei, M., Huang, A., Zhang, Y., and Liu, X.: The Beijing Climate Center Climate System Model (BCC-CSM): the main progress from CMIP5 to CMIP6, *Geoscientific Model Development*, 12, <https://doi.org/10.5194/gmd-12-1573-2019>, 2019.
- 760 Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., and Yabu, S.: The Meteorological Research Institute Earth System Model version 2.0, MRI-ESM2. 0: Description and basic evaluation of the physical component, *Journal of the Meteorological Society of Japan*, <https://doi.org/10.2151/jmsj.2019-051>, 2019.
- Zanna, L., Khatiwala, S., Gregory, J., Ison, J., and Heimbach, P.: Global reconstruction of historical ocean heat storage and transport, *Proceedings of the National Academy of Sciences*, 116, <https://doi.org/10.1073/pnas.1808838115>, 2019.
- 765 Zanowski, H., Hallberg, R., and Sarmiento, J.: Abyssal ocean warming and salinification after Weddell polynyas in the GFDL CM2G coupled climate model, *Journal of Physical Oceanography*, 45, <https://doi.org/10.1175/JPO-D-15-0109.1>, 2015.
- Zelinka, M., Myers, T., McCoy, D., Po-Chedley, S., Caldwell, P., Ceppi, P., Klein, S., and Taylor, K.: Causes of higher climate sensitivity in CMIP6 models, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2019GL085782>, 2020.
- Zickfeld, K., Solomon, S., and Gilford, D.: Centuries of thermal sea-level rise due to anthropogenic emissions of short-lived greenhouse 770 gases, *Proceedings of the National Academy of Sciences*, 114, <https://doi.org/10.1073/pnas.1612066114>, 2017.
- Ziehn, T., Lenton, A., Law, R. M., Matear, R. J., and Chamberlain, M.: The carbon cycle in the Australian Community Climate and Earth System Simulator (ACCESS-ESM1)-Part 2: Historical simulations, *Geoscientific Model Development*, 10, <https://doi.org/10.5194/gmd-10-2591-2017>, 2017.

Zweng, M., Reagan, J., Seidov, D., Boyer, T., Locarnini, R., Garcia, H., Mishonov, A., Baranova, O., Weathers, K., Paver, C., and Smolyar, I.: World Ocean Atlas 2018, Volume 2: Salinity, A. Mishonov Technical Ed.; NOAA Atlas NESDIS 82, 2018.