

Interannual Variability in contributions of the Equatorial Undercurrent (EUC) to Peruvian Upwelling source water

Gandy Maria Rosales Quintana¹, Robert Marsh², and Luis Alfredo Icochea Salas³

¹Tokyo University of Marine Science and Technology, Japan

²University of Southampton, UK

³Universidad Nacional Agraria La Molina, Peru

Correspondence: Gandy Maria Rosales Quintana (gandy.rosales@gmail.com)

Abstract. Time-varying sources of upwelling waters off the coast of northern Peruvian are analysed in a Lagrangian framework, tracking virtual particles backwards in time for 12 months. Particle trajectories are calculated with temperature, salinity and velocity fields from a hindcast spanning 1988-2007, obtained with an eddy-resolving ($1/12^\circ$) global configuration of the NEMO ocean model. At 30-m and 100-m, where coastal upwelling rates exceed 50 m per month, particles are seeded at monthly intervals in proportion to the upwelling rate. Ensemble maps of particle concentration, age, depth, temperature, salinity and density reveal that a substantial but variable fraction of the particles upwelling off Peru arrive via the Equatorial Undercurrent (EUC). Particles follow the EUC core within the depth range 125-175-m, characterised by temperatures $<17^\circ\text{C}$, salinities in the range 34.9-35.2, and densities of $\sigma_\theta = 25.5 - 26.5$. Additional inflows are via two slightly deeper branches further south from the main system, at around $\approx 3^\circ\text{S}$ and $\approx 8^\circ\text{S}$. Averaged across the hindcast, annual-mean percentages of particles upwelling at 30-m (100-m) associated with the EUC vary from 57.4% (52%) at 92°W to 19.2% (17.9%) at 165°W . Considerable interannual variability in these percentages reveal that more of the Peruvian upwelling can be traced back to the EUC during warm events, such as El Niño. In contrast, upwelling waters are of more local origin during cold events such as La Niña. Despite weaker EUC transport during El Niño, relative flattening of the equatorial thermocline brings the EUC upwelling waters much closer to the Peruvian coast than under neutral or La Niña conditions. Annually averaging EUC transport at specific longitudes, a notable negative-to-positive transition is evident during the major El Niño/La Niña events of 1997/99. On short timescales, a degree of longitudinal coherence is evident in EUC transport, with transport anomalies at 160°W evident at the Galapagos Islands (92°W) around 30-35 days later. It is concluded that the Peruvian upwelling system is subject to a variable EUC influence, on a wide range of timescales, most notably the interannual timescale of El Niño Southern Oscillation (ENSO). Identifying this variability as a driver of shifts in population and catch data for several key species, during the study period, these new findings might inform sustainable management of commercially-important fisheries off northern Peru.

Copyright statement. TEXT

1 Introduction

A key feature of the tropical Pacific circulation is the Equatorial Undercurrent (EUC) (Cromwell et al., 1954; Knauss, 1959; Lukas, 1986). The EUC originates in the western equatorial Pacific, just north of Papua New Guinea, as an eastward thermo-
cline flow in the depth range 180-280 m at 147°E. The EUC strengthens eastward (Bryden, 1985; Tsuchiya et al., 1989; Johnson
et al., 2002), reaching peak velocity and transport at around 140°W (Knauss, 1958, 1959), in a core located within $\approx 3^\circ\text{N}$ and
3°S of the Equator (Blanke and Raynaud, 1997; Johnson et al., 2002; Brown et al., 2007). Near the Galapagos Islands, the
EUC core shifts to around 0.5°S (Kessler, 2006; Karnauskas et al., 2010, 2020), before strengthening again towards the eastern
boundary (Lukas, 1986; Johnson et al., 2002), where it feeds major currents along the western South American coastlines of
Ecuador, Peru and Chile (Lukas, 1986; Karnauskas et al., 2010; Montes et al., 2010). The EUC is located at the depth of the
equatorial thermocline, which varies seasonally across the Pacific (Johnson et al., 2002), leading to seasonality in EUC core
velocity. This sub-surface flow plays a crucial role in regional climate and biogeochemistry, through substantial transport of
nutrient and carbon-rich cold water to the surface, feeding the so-called “cold tongue” upwelling region (Chavez et al., 1998;
Pennington et al., 2006; Chavez and Messié, 2009; Qin et al., 2015; Wang et al., 2019).

Around the global coastline, wind-driven upwelling results in high nutrient supply to the surface layer, enhancing primary
production where light and nutrient levels are optimal. At the eastern boundary of the Atlantic and Pacific basins in each
hemisphere, the “Big Four” upwelling systems - Benguela, California, Iberia/Canary and Chile/Peru - are the most active in
the world, accounting for approximately 12 out of 17 million metric tons of marine fish catch-year between 2000 to 2007
(representing 20% of the global taken over an area of less than 1% of the global ocean) according to FAO (on Chavez and
Messié (2009)). Of the Big Four, the Peruvian part of the Peru/Chile system presents the highest average volume of upwelled
waters (1.6 Sv), even though upwelling-favorable winds are the weakest in average (5.7 m s^{-1}) (Chavez and Messié, 2009;
Kämpf and Chapman, 2016).

Peruvian upwelling is dynamically linked to the EUC, itself a sub-surface consequence of equatorial surface flows that are
driven by easterly trade winds, pushing surface water to the west along the Equator in the South Equatorial Current and creating
a pressure head in the western Pacific. Beneath this wind-dominated surface layer, where the Coriolis effect disappears at the
Equator, an eastward pressure gradient drives water back to the east within the EUC, which shoals across the basin from around
250 m in the west to reach the surface in the east (Knauss, 1959; Johnson et al., 2002).

This equilibrium is interrupted as the trade winds weaken or reverse (specially west to 155°W) under El Niño along with
considerable flattening of the thermocline (McPhaden, 1999), seen in historical model simulations (Terada et al., 2020) and
moored buoys (Kessler and McPhaden, 1995). Associated with El Niño are changes in the quantity and properties of upwelled
waters off Peru. For instance, in 1982-83 and 1997-98 El Niño events, principal fisheries such the Peruvian Anchovy, Peruvian
hake and others collapsed, due to migration and dispersion of the biomass in the region (Ñiquen and Bouchon, 2004; Tam
et al., 2006; Wolff et al., 2007). Total pelagic fish landings decreased from 3.3 million tonnes in 1982 to 1.4 millions in 1983
and to almost zero in 1984, as catch species transitioned to Sardine (Arntz and Tarazona, 1989) and others such Jack Mackerel
and Pacific Mackerel (Ñiquen and Bouchon, 2004).

In the present study, we address the origin and nature of the upwelled waters of one of the most productive upwelling systems in the world. We specifically address the role of the EUC in Peruvian upwelling, relative to local wind-driven coastal upwelling in the zone 5-10°S. We aim to quantify absolute and relative changes in the provenance of waters upwelling off northern Peru, to establish the extent of interannual variability and links to ENSO events. We investigate how El Niño and La Niña conditions thus modulate the Peruvian upwelling system. Central to our analysis is an ocean model hindcast spanning 1988-2007. We use a combination of Eulerian and Lagrangian diagnostics, the latter to track upwelling water backwards through time on annual timescales.

In the following, Section 2 describes the methodology used along with brief specifications of the numerical model and hindcast, and details of our diagnostics. In Section 3, we present a range of results, to illustrate variability in the source of upwelling waters, and in the EUC itself. In Section 4, we summarize and discuss our findings in relation to previous studies. In conclusion, Section 5, we emphasize the consequences of El Niño and La Niña for the EUC, Peruvian upwelling, and associated marine ecosystems. Section 6 provides details of code and data availability.

2 Methodology

We first describe the model that provides the hindcast data needed for the Lagrangian analysis, EUC transport calculations, and diagnostics of coastal upwelling, which are subsequently outlined.

2.1 Model Description

We sample 5-day averages of temperature, salinity and velocity from a hindcast spanning 1988-2007, previously obtained with the Nucleus for European Modelling of the Ocean (NEMO) ocean model (Madec, 2008) in eddy-resolving global configuration (ORCA12), henceforth NEMO-ORCA12. For details of parameterization, initialisation and forcing of this hindcast, see Blaker et al. (2015). The model has a horizontal resolution at the equator of $1/12^\circ$ (9.277 km), with 75 vertical levels from the surface up to 5902 meters depth, alongside finer grid spacing near the surface (38 levels from 0 to 411 m). The region of focus in this study is the eastern tropical Pacific, extracted as the region from 170°W to 75°W and from 5°N to 15°S, for the purposes of particle trajectory calculations. An advantage of using fields from a fully global model is that remote influences on the region of interest are fully represented, rather than prescribed at the boundaries in a regional model, which can be problematic. The high resolution NEMO-ORCA12 hindcast is evaluated with observations from three moored NOAA buoys, from the surface up to 400 m depth along the Equatorial Pacific.

2.2 Lagrangian Analysis

To efficiently analyze the provenance of water upwelling off the Peruvian Coast of NEMO-ORCA12, we use the ARIANE Lagrangian code, based on the original method of Blanke and Raynaud (1997). This mass-preserving numerical Lagrangian approach has proved to be an appropriate method for studying the origin and fate of water masses in a wide range of studies (Doos, 1995; Blanke et al., 2002). Ensembles of particles are “seeded” in the coastal upwelling zone off Peru, defined here

east of 82.5°W, from 10°S to the Equator. These particles are then tracked in “backward” mode, reversing in time the analytical calculation of particle progress through grid cells, to reveal a range of pathways.

We allocate particles in proportion to upward vertical velocity where this exceeds a threshold of 50 m per month (1.9×10^{-5} m.s⁻¹). We specifically allocate 1 particle per 10 m per month of upwelling in excess of the threshold, per grid cell. For example, in the simple case that upwelling is 90 m per month, the excess upwelling rate is 40 m per month, so we allocate 4 particles to this grid cell. Initial locations were evenly located within grid cells on the 5-day ORCA12 mesh.

A variable number of particles are released at three “release depths”, model levels closest to 30-m, 50-m and 100-m, at monthly intervals throughout 1989-2007. This number ranges from 38,292 (at 30-m in 1998) up to 98,951 (at 30-m in 2005). Calculations based on initial upwelling at 50-m release depth were relatively similar to those at 30-m release depth, so we show results for 30-m and 100-m release depths only. For examples of the initial positions of particles at both release depths, in Fig. 1 we illustrate seeded particles for the cases of 1997 and 1998.

Particles were released on the last day of each month (December-January) for each year and followed back in time, to sample inter-annual changes in 3D pathways and water properties (temperature, salinity, potential density) of consequence for Peruvian upwelling. Particle data were statistically analyzed on a grid of resolution 0.5° x 0.5° to quantify particle “concentration” as a fractional particle presence in Log10 scale, dividing the number of particle occurrences passing through each grid cell by the total number of particles occurrences during the course of the year.

For an average representation of main pathways in the study region, we computed a “grand ensemble” for the entire data 1989-2007 (for the 30-m and 100-m release depths). Alongside particle concentration, we also average particle age, depth, salinity, temperature and potential density on the 0.5° x 0.5° grid, providing further context for interannual variability of inflow to the Peruvian upwelling system. Then, given the initial (total) number of upwelled particles off Peru (where upwelling exceeds 50 m per month) and the total number of particles crossing specific locations along the equatorial Pacific, we calculate a percentage recruitment from the EUC (3°N -3°S) west to east for further yearly comparisons.

2.3 Pathways and Transport

Informed by the Lagrangian analysis, we also compute EUC volume transport by integrating eastward flow between 3°N to 3°S, encompassing most particle trajectories, and in agreement with previous studies (Blanke and Raynaud, 1997; Johnson et al., 2002; Brown et al., 2007), at selected longitudes from 160°W to 85°W, every 10° from 160°W-110°W, and at 95°W, 92°W and 85°W respectively. Naturally accommodating EUC flow along shoaling isopycnals, we specifically compute transports binned in potential density, σ_θ , in the range $\sigma_\theta = 23.0 - 27.0$, at intervals of $\sigma_\theta = 0.1$, using the NEMO “CDFTOOLS” diagnostic routine “cdfsigtrp” (see <https://github.com/meom-group/CDFTOOLS>). We thus assemble time series of 5-day averaged EUC transport in density space at the selected longitudes to get time series of monthly averages and anomalies.

2.4 Diagnostics of coastal upwelling

Finally, to evaluate the contribution of local winds to the coastal upwelling, Ekman flux and Upwelling flux were calculated. We sample 5-day averages of vertical velocity east of 82.5°W longitude and between 5°S to 10°S (avoiding the near-Equator

120 zone) at 30 m, where this exceeds 50m per month. Area-integrated, we thus obtain the Upwelling flux in units of Sv. Then, to estimate the wind-driven component of coastal upwelling, we calculate zonal Ekman transport (M_x) associated with the meridional component of wind stress (τ_y) at each coastal gridpoint, per $1/12^\circ$ of latitude, as $M_x = \tau_y / f\rho$, where f is the Coriolis parameter and ρ is a representative density for seawater (1026 kg.m^{-3}). Multiplied by meridional gridcell length ($1/12 \times 111 \text{ km}$), we obtain offshore transport in units of Sv that is proportional to coastal upwelling, satisfying continuity of
125 volume. Summed in the same latitudinal range (avoiding low values of f equatorward of 5°S), we thus obtain a monthly index of wind-driven coastal upwelling by averaging the 5-daily index.

3 Results

We begin with a brief evaluation of the NEMO-ORCA12 hindcast in our study region. We then provide an overview of the Lagrangian calculations, before we focus on the pathways followed by water upwelling off Peru, and variation in the source and
130 transport of these waters. We then consider variability in the wind-driven component of upwelling, concluding with analysis of EUC transport anomalies along the Equator.

3.1 Evaluation of NEMO-ORCA12 hindcast in the equatorial Pacific

To establish whether the NEMO-ORCA12 hindcast realistically simulates the equatorial Pacific circulation, we extract vertical profiles of the monthly mean velocities from the model at longitudes along the Equator (170°W , 140°W and 110°W) corresponding with observational data from NOAA-ADCPs, as shown in Fig. 2. The principal eastward subsurface (e.g. EUC) and
135 westward surface currents (e.g. South Equatorial Current - SEC), as described in previous studies (Cromwell et al., 1954; Knauss, 1959; Lukas, 1986; Johnson et al., 2002), are clearly reproduced by NEMO/ORCA12. Shoaling of eastward-flowing core of the EUC towards the Galapagos Islands, can be well identified between 50 and 200 m depth with velocities higher than 0.5 m.s^{-1} in most of the sections shown in Fig. 2. At 140°W and 110°W , the EUC core is seen at around 50 to 150 m,
140 with values notably higher than 1 m.s^{-1} . The highest core velocity along the equator, is found at 140°W longitude in this comparison.

3.2 Upwelling rates and backtracking at annual timescales

We define the upwelling off the northern Peruvian coast where upward advection exceeds $1.9 \times 10^{-5} \text{ m.s}^{-1}$ (50 m per month), for each release depth in the water column, chosen to sample upwelling over a depth range typical of the Peruvian upwelling
145 system (Kämpf and Chapman, 2016). To illustrate vertical velocities at two of our chosen depths, monthly NEMO-ORCA12 climatologies (1988-2007) at 33 m and 100 m are presented in Figs. 3 and 4 respectively (we have used here 33m, since is the closest depth available to 30m).

Monthly upward velocities are elevated along both the equatorial upwelling region and off the Peruvian coast, with clear seasonal variability. During austral summer (December-January-February), upwelling is stronger and more widespread north
150 and south of the Equator, and slightly weaker off Peru. During austral autumn (March-April-May), equatorial upwelling is

characterised as a narrower zone (restricted equatorward of 3°), with higher values towards the east; off Peru, stronger upwelling is evident, compared to summer. During austral winter (June-July-August), strong upwelling extends across the easternmost equatorial region (140°W to Galapagos) and off Peru, when south-east trade winds are seasonally stronger.

A notable difference is evident in the 100-m analysis, with downwelling stronger around the Galapagos Islands and near the
155 Ecuadorian coast. This is also consistent with the climatology rates calculated for the Upwelling and Ekman coastal fluxes, peaking during austral autumn and winter seasons and reduced during austral summer season (when also winds are weakest), as forced by the meridional component of the wind stress along the north-central Peruvian coast.

Returning to particles released off Peru, for most years (notably 1991, 1997 and 2004 - not shown, we observe a westward origin for some particles, from the Galapagos Islands via the Equator. This is consistent with eastward shoaling of the EUC
160 (around 75 m). However, in other years, particles originate from a northern shallower region, in the vicinity of the Ecuadorian coast (e.g. 1988, 1989 and 1990 - not shown). For particles initialised at 100 m, we observe a similar pattern of inflow at greater depths (around 125 m), although with a clearer origin from the equatorial band and southern hemisphere, this can be also observed in the grand ensemble.

3.3 Mean pathways at annual timescale

165 As outlined in Section 2.2, particle concentration and mean age maps are obtained on a $0.5^\circ \times 0.5^\circ$ mesh for each year from 1989 to 2007, further averaged to obtain the “grand ensemble” results for 30-m and 100-m release depths that are shown in Fig. 5, right and left panels respectively. Tracking backwards, Figs. 5 a-f show the corresponding particle concentration, age, depth, temperature, salinity, and potential density (σ_θ) distributions, for flows feeding the upwelling source water off Peru.

We use a logarithmic scale for particle concentration, to emphasize a particularly wide range of this diagnostic (in Fig. 5a, red represents high concentration while blue colors low concentration in average). Mean particle age (5b) is expressed in days
170 back-tracked, so for end-December releases, day 0 to day 365 runs from December 31st back to January 1st.

In Fig. 5a,b (left and right), highest particle concentration and youngest age (transit time range between 0-120 days) are naturally located near the release points in both experiments. After around 2 months, particles are back-tracked to the Galapagos Islands with relative higher concentrations before further back-conveyance along the equatorial band. As particles are traced
175 westward, high concentration along $\approx 3^\circ\text{N}$ to 3°S are associated with sub-surface eastward flow of the EUC, widening slightly west of the Galapagos Islands. Particle concentration rapidly declines further south and north of 3° to the west of 110°W , although, we note patches of relatively higher concentrations forming two branches in the southeast of our region.

Inflows from 30-m release depth remain at a relatively shallow depth (above ≈ 75 m) during the first 150-180 days (Fig. 5c, left panel) described by the Isotherms of 17°C - 20°C and Isohalines between 34.9 and 35.1 along $\sigma_\theta = 25.5 - 26.0$ isopycnals
180 (Fig. 5d,e left panels). Particles released at 100-m more notably deepen after the first 60-90 days (October-September) of release, remaining above 125 m around Galapagos Island and the eastern coast. In this region, particles were characterised with temperatures in the range 15°C - 17°C and salinities around 35-35.1 along isopycnals $\sigma_\theta = 26.0 - 26.5$ (Fig. 5f right panel).

West of Galapagos, after 210 days of back-tracking in both experiments, particles lie at greater depth (below 125 m) where
185 particle concentration is the highest. Along the EUC core, between $\approx 2^\circ\text{N}$ to $\approx 2^\circ\text{S}$, temperatures fall below 17°C (Fig. 5d, left
and right panels) and salinities lie in the range 34.9 – 35.2 (Fig. 5e, left and right panels), combining for $\sigma_\theta > 26.0$ isopycnal
(Fig. 5f, left and right panels) and populating the isopycnal surface around $\sigma_\theta = 26.5$ for the deeper release depth.

As previously mentioned, two relatively deep branches (Fig. 5a, left and right panels), in agreement with previous studies
(Lukas, 1986; Johnson and Moore, 1997; Donohue et al., 2002; Montes et al., 2010; Kuntz and Schrag, 2018), are identified
190 further south of the main equatorial system. Along $\approx 3^\circ\text{S}$ and $\approx 8^\circ\text{S}$ (Fig. 5a, left and right panels), more clearly distinguished
with potential density (Fig. 5f) for the deeper release depth experiment. From both release depths, these southern branches are
associated with higher density (e.g. $> \sigma_\theta \approx 26.5$ in the case of 100-m release depth), with the southern most branch ($\approx 8^\circ\text{S}$)
carrying the densest inflow and the northern branch ($\approx 3^\circ\text{S}$) at somewhat lower density (i.e. Figure 5f left panel).

Our results suggest that these branches are likely associated with a bifurcation of the EUC, to the west of the Galapagos
195 Islands as previous studies have highlighted (Johnson et al., 2002; Karnauskas et al., 2010; Jakoboski et al., 2020). This source
of Peruvian water masses in the upwelling region is thus directly modulated by the EUC, when the latter is at its strongest; this
may be particularly the case from April to June, when EUC transports reach maxima (Table 1). We will return to this point in
the following section.

Regarding the denser southern branch ($\approx 8^\circ\text{S}$, left panels for Fig. 5d, Fig. 5e and Fig. 5f) traced by those particles released at
200 30-m, our experiments indicate that the high density is coincident with high salinity rather than low temperature (> 35.1 , 17°C
or less, and $\sigma_\theta = 25.5 - 26.5$). Somewhat different characteristics were found for particles released at 100-m (right panels for
Fig. 5c, Fig. 5d, Fig. 5e and Fig. 5f), for which temperature is a more dominant influence on density.

3.4 EUC transport across the eastern Pacific and the variable EUC contribution to Peruvian upwelling

Back-tracked trajectories from the released particles off Peru revealed that a substantial proportion of water masses are recruited
205 from the main EUC system, traveling along the equatorial Pacific for around 10,000 km (from 170°W to 80°W) before reaching
the Peruvian upwelling region. Integrated positive transports across density classes show a longitudinal-seasonal EUC flow
variability when averaging over 20 years from 1988-2007 (Table 1 and Fig. 6). In the first half of the study region between
 160°W and 135°W , EUC transport peaks from March to June with average values > 36.2 Sv. Maximum values during April-
May are almost double the annual average of 36.3 Sv. East of 135°W , EUC transport tends to peak during April-June (see Table
210 1), consistent with observational data (Johnson et al., 2002). Climatological EUC transport tends to decrease from August to
December across the whole region.

This variability underpins transports of energy and other properties from the western Pacific to the Peruvian Upwelling
region. For instance, using Lagrangian trajectories, Qin et al. (2016) found that an elevated iron injection associated with the
1997/98 El Niño showed 30% higher total dissolved iron concentrations in the eastern equatorial Pacific around 13 months later,
215 consistent with suppressed productivity responses in the central, eastern Pacific and off Peru, from models and observations
(Ryan et al., 2006; Slemons et al., 2009; Chever et al., 2015). Moreover, eastward transports along the EUC are higher in the

western part of the study region than close to Galapagos (e.g. 36.2 Sv at 160°W and 12.5 Sv at 92°W, on average), consistent with loss of transport to progressive upwelling along the Equator.

220 Seasonality is evident for most years of the hindcast, as shown in the monthly average and anomalies of Fig. 6. To emphasize seasonal cycles, we add contour lines for transports of 25 and 50 Sv in Fig. 6a. We note that the lowest values of <5 Sv coincide with major El Niño events, such 1997/98 and 1991/92, and other warm event such 2002/03 in the western region (\approx 160°W-140°W). At these times, the West Pacific Warm Pool expands across the central Pacific, the thermocline largely flattens, and the EUC quasi-disappears for a short period (McPhaden, 1999).

225 During strong La Niñas 1988-1989 and 1998/99, positive anomalies in transport coincide with restoration of an eastward-shoaling thermocline after strong El Niños the year before, associated with “peaks” of the >50 Sv isoline in Fig. 6a, displaced westward from 140°W to 160°W. West-East displacement can be also seen following isolines for anomalies (Fig. 6b) during El Niños. For instance, in late 1996 and early 1997, the warm pool with anomalies >+10 Sv was rapidly transported from 160°W to 92°W, next to the Galapagos Islands - in less than a year - before reaching the Peruvian coast. A similar transient is observed for smaller events (e.g., 2002/03 and 2006/07), although less intense than during strong El Niños. We identify other strong 230 displacements, such during 2000/01 and 2004/05, when ENSO variability is not so clearly implicated.

This EUC transport variability is consistent with our Lagrangian trajectory analysis, most clearly during major events, when the signal is clear. Fig. 7 shows yearly variations of the number of particles (a and b for 30-m and 100-m respectively) and its percentage (d and c for 30-m and 100-m respectively) after a year of tracking. In general, our results suggest that particles originating in the western basin cross the Pacific mainly within the EUC, at different intensities varying between 19.2% - 235 52.0% for 30-m (17.9% - 57.4% for 100-m) between 165°W and 92°W (see dashed lines in Fig. 7c and Fig. 7d).

We found that during strong El Niño and La Niña events, the percentage of particles feeding the eastern basin changes dramatically. Highest and lowest percentages of particles are observed during the major ENSO of 1997/98. Particles recruited by the EUC in 1997 rapidly crossed the Pacific, showing in the easternmost region (92°W) the highest record of 71.1% (80.7% for 100-m), consistent with an enhancement of volume transport in Qin et al. (2016). However, during a following cold La Niña 240 phase, in the same longitude range, smaller percentages arrived via the EUC (3.7% for 30-m and 1.7% for 100-m in Fig. 7c and Fig. 7d). This reduced contribution of the EUC to Peruvian upwelling coincides with a western displacement of transport in mid-1998, which persists into 1999 (see anomalies in Fig. 6). Following the major La Niña of 1998-99, notably into 2000, positive transport anomalies (>10Sv) prevailed across the Pacific (Fig. 6), consistent with a high percentage of particles coming from the EUC (with 68.4%-76.7% to 39.2%-38.7% at 160°W and 92°W for 30-100-m respectively, see Fig. 7).

245 Another major event categorised as a “Moderate Eastern Event” (Yu et al., 2011), unfolded over 1991/92 in the Pacific. For the 1991-92 warm event, our trajectories in percentage showed that above-average numbers of particles are recruited by the EUC close to Galapagos (up to 53.5% for 30-m and 43.7% for 100-m, in 1991 at 165°W). In the case of a 2002/03 event, we observed the same tendency for warm and cold phases of the event, although less intense compared to that of 1991/92. Conversely, for the 2006/07 “Moderate Eastern” El Niño, we can differentiate between the warm and cold phases, with higher 250 particles recruited by the EUC in 2006 at 165°W (52.7% for 30-m and 74.9% for 100-m) than during 2007 La Niña (43.1% for

30-m and 53.6% for 100-m). Evidently, the EUC contribution to Peruvian upwelling is most dominant before-during major El Niño or anomalously warm events, than cold or major La Niña conditions.

3.5 The coastal wind-driven contribution to Peruvian upwelling

Upwelling off Peru is augmented by equatorward winds that induce offshore transport in a surface layer, supplied by “rapid upwelling” in a relatively narrow coastal zone. Further offshore, “slow upwelling” is induced by negative wind stress curl that induces large-scale Ekman suction (Rykaczewski and Checkley, 2008). Variability in wind-driven upwelling has been often linked to the biological decadal variability in upwelling system such as off Peru (Rykaczewski and Checkley, 2008; Espinoza-Morriberón et al., 2019).

Here, we focus on variability in coastal upwelling associated with the meridional component of wind stress (Fig. 8a), used to compute Ekman offshore transport which we term the Ekman flux (Fig. 8b), compared to the full Upwelling flux, obtained from the vertical velocity component at 30 m (Fig. 8c). The three time series in Fig. 8 all trend towards positive anomalies after 2000, most obviously for the Ekman flux, that is a direct consequence of strengthening in the equatorward wind stress. This may be part of a wider strengthening trend in the trade wind system of the Pacific, that has been identified as a driver of the early-2000s hiatus in global warming (England et al., 2014). We further observe that, during strong and Moderate La Niña events such 1988-1999, 1998-1999 and 1993-1994, equatorward wind stress decreases drastically during austral summer with values $\leq 0.01 \text{ N}\cdot\text{m}^{-2}$, and a further minimum during the relatively warm event of summer 2000. From the turn of the millennium, anomalies are generally positive (exceeding climatology), peaking in late 2005 at $\approx 0.07 \text{ N}\cdot\text{m}^{-2}$.

During the 1997-1998 El Niño, wind stress was not completely shut down in the northern part off Peru, although presenting negative anomalies (reduction) in the first quarter or the year. This is consistent with the high percentage of particles appearing off Peru during extreme El Niños that arrived via the EUC, rather than being locally sourced via coastal Ekman dynamics. An alternative source for upwelled waters is further evidenced in the Upwelling flux (Fig. 8), where positive anomalies coincide with a developing El Niño (e.g., second half of 1997).

3.6 Longitudinal coherence of EUC transport anomalies

To the extent that EUC transport anomalies are coherent across the equatorial Pacific, variability of upwelling via the EUC may be predictable. To examine the longitudinal coherence of EUC transport anomalies, we compute correlation between EUC transport at each selected longitude (Table 2). Despite generally low values, we find significant (p-value < 0.01, 99% confidence interval) positive correlations at zero-lag. This is consistent with rapid and near-simultaneous variations in EUC transport across the eastern Pacific, although weakening of correlation with separation (in longitude) indicates that transport anomalies may advect eastward, associated with inertial terms in the momentum balance and eastward propagation of Kelvin waves.

The development of an anomalous EUC flow to the east may thus be predictable for an appropriate time lag. Experimenting with lagged cross correlations of transport at 160°W and 92°W, we obtain highly significant $R = +0.31$ (p-value < 0.01) between transport at 160°W and transport 30-35 days later at 92°W. With further investigation, beyond scope here, more skilful EUC

predictability may provide useful advance warning of substantial changes in the Peruvian upwelling system, and subsequent
285 impacts on important marine resources. A further step would be assess the variability and predictability of year-round inflows
to the Peruvian upwelling system, also beyond the scope of this investigation.

4 Summary and Discussion

With a range of diagnostics, we have evaluated variability of the Peruvian upwelling system in an eddy-resolving model
hindcast spanning 1988-2007. Wind-driven coastal upwelling is associated with alongshore winds, complemented by shoaling
290 of the Equatorial Undercurrent. Local and remote drivers of upwelling are highly variable on seasonal to interannual timescales.
Contrasting scenarios are shown in Fig. 9. In Fig. 9a, we highlight upwelling dominated by local winds, when the EUC shoals
well to the west of the Galapagos Island. In Fig. 9b, the EUC shoals more gradually to the east, reaching the coastal upwelling
zone, where weakened winds otherwise limit local upwelling.

We now consider impacts on Peruvian upwelling that are linked to variable EUC contributions. Both the 1997/98 and 2000/01
295 events had important effects in fisheries. May-June 1997 fieldwork conducted by the Peruvian Sea Institute (IMARPE), revealed
an unusual southern and deeper (> 150-200 m) migration of Peruvian Hake (*Merluccius gayi peruanus*; generally distributed
between 0°N to 8°S), easily reaching 12°S and even more southern areas along the Peruvian coast (Castillo et al., 1997).
Conversely, Hake tended to migrate northward and sometimes closer to the surface (pelagic behaviour, considering that hake
is a demersal specie), due bottom oxygen deficiency, during the La Niña of 1998/99. This is consistent with the physical-
300 biogeochemical character of the EUC. Where a southward branch of the EUC reaches the northern Peruvian coast (Fig. 5), it
refreshes the sub-surface layer with high concentrations of dissolved oxygen (Echevin et al., 2020) and other vital nutrients
(Qin et al., 2016; Espinoza-Morriberón et al., 2019).

A similar impact afflicts the second most abundant demersal species in Peruvian coastal waters, the Conger eel (*Ophichthus*
remiger). During El Niño events (December 1997 to May 1998; March-July 1992) and subsequent La Niña events (November
305 1993 to March 1994; April-August 1998), significant increased and reduced Catch per Unit Effort (CPUE) was reported,
respectively (Castillo et al., 2000; Martina, 2004). The oceanographic and fisheries evidence together suggest crucial links
between demersal species and the EUC. More recently, Martina (2018) also highlighted dramatic reduction in biomass during
2004/05, followed by significant increase in 2007. The latter increase coincides with positive EUC transport anomalies taking
place during early 2007 (see Fig. 6b).

310 During the moderate Eastern Pacific El Niño of 1991/92, an alternative transition is observed. A relatively high percentage
of upwelling particles is traced back to the EUC, amounting to 48.2% (36.5%) and 22.6% (8.7%) at 160°W and 92°W during
1991, for 30-m (100-m) release depths respectively (Fig. 7c and Fig. 7d). These enhanced percentages of EUC origin, at
160°W from 1991 to 1992, coincide with relatively high anomalies in transports (Fig. 7b). From the second half of 1991, east
of 140°W, positive anomalies $>+10Sv$ were found, while at the end of 1991 to the first half of 1992, high negative anomalies
315 were observed from 160°W to 110°W. This is consistent with eastward transport of warm pool waters and the relatively high

percentage of EUC particles found in 1992 near Galapagos (92°W), associated with deepening of the thermocline followed by relaxation-shoaling, in the eastern Pacific (Kessler and Mcphaden, 1995).

320 Related to 1991/92 El Niño, a dramatic size-age and biomass reduction of the Peruvian Hake happened. This was first attributed to an overfishing (Castillo, 1996), although observations indicated an enhanced EUC at this time. This likely led to drastic relocation of species farther south, causing high mortality for adults and leaving young species in the northern region of Peru. Plus, sub-optimal oceanographic conditions when the EUC weakened after its peaks passed. Species are often slow to recover in size-age-abundance indicators. Even after the major El Niño of 1997/98, Hake has shown signs of recovery in CPUE data as wider size structure and higher catch numbers (Ballón et al., 2008; Lassen et al., 2009a).

325 Furthermore, it has further been suggested that some species also use the EUC as a migration conveyor, such as the Pacific Giant squid (*Dosidicus gigas*). Largely absent from east Pacific coastal waters over 1950-90, catches of this species increased from the beginning of 1991 through 1992. During 1995-96, overfishing impacted this population (in the northern American coast), primary due to the high catch effort of foreign fishing vessels (Contreras Paya, 2017). Subsequent to the El Niño, of 1997/98, a denser nearshore migration of the squid, more accessible for fishing, was reported off the North American coast (CALCOFI, 2000). Changes in coastal squid populations may be linked to the equatorially and then coastally trapped Kelvin waves (e.g., Shaffer et al. (1997)), which may favour southward (northward) migration when the EUC is stronger (weaker) around the Galapagos Island, as the squid seek out for optimal conditions (Cheung W. L. et al., 2018).

330 We identify less intense events at other times, such late 2002/03 (Fig. 6 and Fig. 7), when Peruvian Hake almost collapsed off northern Peru, leading to a moratorium on fishing until March 2004 (Lassen et al., 2009b; Benites and Barriga, 2011). This followed the moderate Eastern El Niño of 2006/07, with positive transport anomalies ($>+10$ Sv) during the first half of 2006, transporting water masses to the east. Then, from the end of 2006 to early 2007, negative anomalies were observed in the west, indicating that water masses reached the east. This is consistent with higher percentage of particles in 2006 and a reduction of those in 2007 (Fig. 7) from both experiments.

340 In summary, the EUC contribution to upwelling region along the northern Peruvian coast is sometimes substantial, depending on the type of event (strong, moderate or weak), altering environmental conditions of consequence for many marine species. Changes in the EUC in the Pacific, as a first stage before El Niño happens off Peru, can have positive and negative effects in fisheries regarding the species (Icochea et al., 1989; Castillo et al., 1997; Contreras Paya, 2017; Tam et al., 2008; Taylor et al., 2008). Identifying this variability as a driver of shifts in population for several key species, might efficiently inform sustainable management of fisheries in Peru.

5 Conclusions

345 We have systematically quantified the origin of waters upwelling in one of the most productive regions in the global ocean, based on a high resolution NEMO-ORCA12 eddy-resolving model hindcast spanning 1988-2007, which includes major warm and cold events associated with El Niño and La Niña respectively. Through Lagrangian analysis of virtual particle ensembles, we identified an interannually variable fraction of upwelled water that is recruited via the Equatorial Undercurrent (EUC),

flowing between $\approx 3^{\circ}\text{N}$ to 3°S at depths ranging 125 m to 175 m. A key finding is that the northern Peruvian upwelling system
350 is sensitive to highly variable EUC inflow. Particle back-trajectories - sampling the upwelling at depths of 30-m and 100-m off
Peru - trace the EUC as far west as 170°W on an annual timescale, moving at depths, temperatures, salinities and densities that
are consistent with observations (Johnson et al., 2002). Back-trajectories further identified two relatively deep branches south
of the main equatorial system, along $\approx 3^{\circ}\text{S}$ and $\approx 8^{\circ}\text{S}$, in agreement with previous studies (Lukas, 1986; Johnson and Moore,
1997; Donohue et al., 2002; Montes et al., 2010). Only a small percentage of particles otherwise originate from latitudes
355 poleward of 3° , as far west as 170°W .

Over 1989-2007, we quantified the variable contribution of EUC waters to Peruvian upwelling. At 92°W , we identify highest
and lowest percentages of EUC-sourced particles during the strong El Niño of 1997/98 and the subsequent strong La Niña of
1998/99, respectively. The EUC is thus most influential – in relative terms – when the equatorial thermocline deepens across
the eastern Pacific, during early El Niño, allowing the EUC to extend as far as the eastern boundary and upwelling to the east of
360 Galapagos. During a strong La Niña, EUC waters conversely upwell with a strongly shoaling thermocline to reach the surface
layer in the central Pacific, with weaker EUC upwelling further to the east. In this scenario, the waters upwelling off Peru are
of local provenance (see Fig. 9).

Variable provenance of oxygenated, nutrient-rich EUC waters in the 1990s can be linked to substantial changes in Peruvian
coastal fisheries. Further variability in the 2000s is associated with a range of El Niño and La Niña events. On sub-annual
365 timescales, particles most rapidly cross the eastern Pacific with the EUC during peak transport around March and April,
consistent with previous studies (Flores et al., 2009). Correlations between 5-day averaged EUC transport at selected longitudes
in the range $92\text{-}160^{\circ}\text{W}$ indicate a high degree of longitudinal coherence, with evidence of a time lag of 30-35 days for transport
anomalies at 160°W to reach the Galapagos Islands at 92°W .

In highlighting the impact of variable EUC influences on regional biogeochemistry, ecosystems and fisheries, this study
370 provides the basis for informed analysis and prediction of an unfolding El Niño or La Niña, for a sustainable approach to
management of marine resources in the Peruvian upwelling system. A next step would be to include biogeochemical analyses,
using models and observations, to better understand the consequences of variable nutrient supply for primary productivity at
the base of the food chain.

6 Code and data availability

375 The NEMO-ORCA12 data analysed here is archived at the National Oceanography Centre, Southampton. The original ver-
sion of ARIANE software used here, available from <http://stockage.univ-brest.fr/grima/Ariane/>, was adapted at the National
Oceanography Centre and the University of Southampton. Specific trajectory data and NEMO-ORCA12 diagnostics presented
here are available from the authors, on request. Equatorial current data from the TAO/TRITON Array are available from NOAA
via <https://www.pmel.noaa.gov/gtmba/pmel-theme/pacific-ocean-tao>.

All authors have contributed equally to this paper.

Competing interests. .

The authors declare that they have no conflict of interest.

Acknowledgements. We are grateful to the European Mundus Joint Master Degree (ERASMUS MUNDUS Scholarship) and the Marine
385 Environment and Resources program (MER+) for the financial support provided during this 2-years project. We acknowledge the National
Oceanographic Centre Southampton (NOCS) for undertaking the NEMO-ORCA12 simulation (using the NEMO framework developed by a
consortium of European institutions), and Jeff Blundell at the University of Southampton, for development and installation of a local version
of the ARIANE software that is central to this study. We also thank to KAKENHI-JSPS-2020-Japan project for financially supporting this
publication.

390 References

- Arntz, W. E. and Tarazona, J.: Summary for Policymakers, in: *Climate Change 2013 - The Physical Science Basis*, edited by Intergovernmental Panel on Climate Change, vol. 52, pp. 1–30, Cambridge University Press, Cambridge, <https://doi.org/10.1017/CBO9781107415324.004>, https://www.cambridge.org/core/product/identifier/CBO9781107415324A009/type/book_part, 1989.
- 395 Ballón, M., Wosnitzer-Mendo, C., Guevara-Carrasco, R., and Bertrand, A.: The impact of overfishing and El Niño on the condition factor and reproductive success of Peruvian hake, *Merluccius gayi peruanus*, *Progress in Oceanography*, 79, 300–307, <https://doi.org/10.1016/j.pocean.2008.10.016>, <http://dx.doi.org/10.1016/j.pocean.2008.10.016>, 2008.
- Benites, C. and Barriga, E.: La Poblacion de la merluza durante el verano 2004. Crucero BIC Olaya 0401-02, Tech. rep., Instituto del Mar del Peru, Callao, <https://doi.org/ISSN 0378-7702>, 2011.
- 400 Blaker, A. T., Hirschi, J. J., McCarthy, G., Sinha, B., Taws, S., Marsh, R., Coward, A., and de Cuevas, B.: Historical analogues of the recent extreme minima observed in the Atlantic meridional overturning circulation at 26°N, *Climate Dynamics*, 44, 457–473, <https://doi.org/10.1007/s00382-014-2274-6>, 2015.
- Blanke, B. and Raynaud, S.: Kinematics of the Pacific Equatorial Undercurrent: An Eulerian and Lagrangian approach from GCM results, *Journal of Physical Oceanography*, 27, 1038–1053, [https://doi.org/10.1175/1520-0485\(1997\)027<1038:KOTPEU>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<1038:KOTPEU>2.0.CO;2), 1997.
- 405 Blanke, B., Arhan, M., Lazar, A., and Prévost, G.: A Lagrangian numerical investigation of the origins and fates of the salinity maximum water in the Atlantic, *Journal of Geophysical Research C: Oceans*, 107, 27–1, <https://doi.org/10.1029/2002jc001318>, 2002.
- Brown, J. N., Godfrey, J. S., and Fiedler, R.: A zonal momentum balance on density layers for the central and eastern equatorial Pacific, *Journal of Physical Oceanography*, 37, 1939–1955, <https://doi.org/10.1175/JPO3090.1>, 2007.
- Bryden, H. L. B. E. C.: Diagnostic Model of the Three-Dimensional Circulation in the Upper Equatorial Pacific Ocean, *Journal of Physical Oceanography*, 15, 1255–1273, 1985.
- 410 CALCOFI: California Cooperative Oceanic Fisheries Investigations, Tech. Rep. 5, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, La Jolla CA, <https://doi.org/10.1163/156851801511873>, http://www.calcofi.org/publications/calcofireports/v41/CalCOFI_Rpt_Vol_41_2000.pdf, 2000.
- Castillo, R.: Informe tecnico sobre la situacion de la merluza 1990-1995, Tech. rep., Instituto del Mar del Peru, Callao, <http://biblioimarpe.imarpe.gob.pe/bitstream/123456789/899/1/IP17.pdf>, 1996.
- 415 Castillo, R., Ganoza, F., Aliaga, A., Gutiérrez, M., and Guevara-Carrasco, R.: Distribución, concentración y biomasa de la merluza peruana en otoño 1997 con el método hidroacústico. Crucero BIC Humboldt 9705-06, Tech. rep., Instituto del Mar del Peru, Callao, <http://biblioimarpe.imarpe.gob.pe/handle/123456789/1486>, 1997.
- Castillo, R., Gomez, E., and Paredes, F.: Pesquería y biología de la Anguila común *Ophichthus pacifici* (Günther)* en el Perú, Tech. rep., Instituto del Mar del Peru, Callao, <http://biblioimarpe.imarpe.gob.pe/bitstream/123456789/1168/1/IP134.pdf>, 2000.
- 420 Chavez, F. P. and Messié, M.: A comparison of Eastern Boundary Upwelling Ecosystems, *Progress in Oceanography*, 83, 80–96, <https://doi.org/10.1016/j.pocean.2009.07.032>, <http://dx.doi.org/10.1016/j.pocean.2009.07.032>, 2009.
- Chavez, F. P., Strutton, P. G., and McPhaden, M. J.: Biological-physical coupling in the central equatorial Pacific during the onset of the 1997-98 El Niño, *Geophysical Research Letters*, 25, 3543–3546, <https://doi.org/10.1029/98GL02729>, 1998.
- 425 Cheung W. L., W., Bruggeman, J., and Butenschön, M.: Chapter 4: Projected changes in global and national potential marine fisheries catch under climate change scenarios in the twenty-first century, in: *Impacts of climate change on fisheries and aquaculture. Synthesis of current*

- knowledge, adaptation and mitigation options., edited by Barange, M., Bahri, T., Beveridge C. M., M., Cochrane L., K., Funge-Smith, S., and Poulain, F., vol. 627, chap. 4, pp. 113–138, Food and Agriculture Organization of the United Nations, Rome, 2018.
- 430 Chever, F., Rouxel, O. J., Croot, P. L., Ponzevera, E., Wuttig, K., and Auro, M.: Total dissolvable and dissolved iron isotopes in the water column of the Peru upwelling regime, *Geochimica et Cosmochimica Acta*, 162, 66–82, <https://doi.org/10.1016/j.gca.2015.04.031>, <http://dx.doi.org/10.1016/j.gca.2015.04.031>, 2015.
- 435 Contreras Paya, I.: DOCUMENTO CONSOLIDADO, Convenio de desempeño 2016. Estatus y posibilidades de explotación biológicamente sustentable de los principales recursos nacionales del 2017: Jibia, 2017, Tech. rep., Instituto de Fomento Pesquero, <https://doi.org/10.13140/RG.2.2.20164.63365>, http://www.ifop.cl/wp-content/contenidos/uploads/RepositorioIfop/InformeFinal/P-483253_jibia.pdf, 2017.
- Cromwell, T., Montgomery, R., and Stroup, E.: Equatorial Undercurrent in Pacific Ocean Revealed by New Methods, *Science*, 119, 648–649, <https://doi.org/10.1126/science.119.3097.648>, 1954.
- 440 Donohue, K. A., Firing, E., Rowe, G. D., Ishida, A., and Mitsudera, H.: Equatorial Pacific subsurface countercurrents: A model-data comparison in stream coordinates, *Journal of Physical Oceanography*, 32, 1252–1264, [https://doi.org/10.1175/1520-0485\(2002\)032<1252:EPSCAM>2.0.CO;2](https://doi.org/10.1175/1520-0485(2002)032<1252:EPSCAM>2.0.CO;2), 2002.
- Doos, K.: Interocean exchange of water masses, *Journal of Geophysical Research*, 100, 499–514, 1995.
- Echevin, V., Gévaudan, M., Espinoza-morriberon, D., Tam, J., Aumont, O., Gutierrez, D., and Colas, F.: Physical and biogeochemical impacts of RCP8.5 scenario in the Peru upwelling system, *Biogeosciences*, pp. 1–42, <https://doi.org/doi.org/10.5194/bg-2020-4>, 2020.
- 445 England, M. H., Mcgregor, S., Spence, P., Meehl, G. A., Timmermann, A., Cai, W., Gupta, A. S., Mcphaden, M. J., Purich, A., and Santoso, A.: Recent intensification of wind-driven circulation in the Pacific and the ongoing warming hiatus, *Nature Climate Change*, 4, 222–227, <https://doi.org/10.1038/nclimate2106>, 2014.
- Espinoza-Morriberón, D., Echevin, V., Colas, F., Tam, J., Gutierrez, D., Graco, M., Ledesma, J., and Quispe-Ccalluari, C.: Oxygen variability during ENSO in the Tropical South Eastern Pacific, *Frontiers in Marine Science*, 5, 1–20, <https://doi.org/10.3389/fmars.2018.00526>, 2019.
- 450 Flores, R., Tenorio, J., and Dominguez, N.: Variaciones de la Extension Sur de la Corriente Cromwell frente al Peru entre los 3 y 14°S, Tech. rep., Instituto del Mar del Peru, Callao, <http://biblioiarpe.imarpe.gob.pe/bitstream/123456789/1091/1/BOL24%281-2%29-7.pdf>, 2009.
- Icochea, L., Chipollini, A., and Ñiquen, M.: Analisis de la pesqueria de arrastre pelagica en la costa peruana durante 1983-1987 y su relacion con el medio ambiente, in: *Memorias del Simposio Internacional de los Recursos Vivos y las Pesquerias en el Pacifico Sudeste*, pp. 455–465, Instituto del Mar del Peru, Viña del Mar, https://biblioteca.imarpe.gob.pe/opac_css/index.php?lvl=notice_display&id=13778, 1989.
- 455 Jakoboski, J., Todd, R. E., Brechner Owens, W., Karnauskas, K. B., and Rudnick, D. L.: Bifurcation and upwelling of the equatorial undercurrent west of the galápagos archipelago, *Journal of Physical Oceanography*, 50, 887–905, <https://doi.org/10.1175/JPO-D-19-0110.1>, 2020.
- Johnson, G. C. and Moore, D. W.: The Pacific subsurface countercurrents and an inertial model, *Journal of Physical Oceanography*, 27, 2448–2459, [https://doi.org/10.1175/1520-0485\(1997\)027<2448:TPSCAA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<2448:TPSCAA>2.0.CO;2), 1997.
- 460 Johnson, G. C., Sloyan, B. M., Kessler, W. S., and McTaggart, K. E.: Direct measurements of upper ocean currents and water properties across the tropical Pacific during the 1990s, *Progress in Oceanography*, 52, 31–61, [https://doi.org/10.1016/S0079-6611\(02\)00021-6](https://doi.org/10.1016/S0079-6611(02)00021-6), 2002.
- Kämpf, J. and Chapman, P.: *Upwelling Systems of the World. A Scientific Journey to the Most Productive Marine Ecosystems*, Springer Nature, Switzerland, <https://doi.org/10.1007/978-3-319-42524-5>, 2016.

- Karnauskas, K. B., Murtugudde, R., and Busalacchi, A. J.: Observing the Galápagos-EUC interaction: Insights and challenges, *Journal of Physical Oceanography*, 40, 2768–2777, <https://doi.org/10.1175/2010JPO4461.1>, 2010.
- 465 Karnauskas, K. B., Jakoboski, J., Johnston, T. M., Owens, W. B., Rudnick, D. L., and Todd, R. E.: The Pacific Equatorial Undercurrent in Three Generations of Global Climate Models and Glider Observations, *Journal of Geophysical Research: Oceans*, 125, 1–12, <https://doi.org/10.1029/2020JC016609>, 2020.
- Kessler, S. and McPhaden, J.: The 1991–1993 El Niño in the central Pacific, *Deep Sea Research II*, 42, 295–333, [https://doi.org/https://doi.org/10.1016/0967-0645\(95\)00041-N](https://doi.org/https://doi.org/10.1016/0967-0645(95)00041-N), 1995.
- 470 Kessler, W. S.: The circulation of the eastern tropical Pacific: A review, *Progress in Oceanography*, 69, 181–217, <https://doi.org/10.1016/j.pocean.2006.03.009>, 2006.
- Kessler, W. S. and McPhaden, M. J.: The 1991–1993 El Niño in the central Pacific, *Deep-Sea Research Part II*, 42, 295–333, [https://doi.org/10.1016/0967-0645\(95\)00041-N](https://doi.org/10.1016/0967-0645(95)00041-N), 1995.
- 475 Knauss, J. A.: Observations of the Pacific Equatorial Undercurrent, *Nature*, pp. 601–602, <https://doi.org/https://doi.org/10.1038/182601a0>, <https://www.nature.com/articles/182601a0>, 1958.
- Knauss, J. A.: Measurements of the Cromwell current, *Deep Sea Research (1953)*, 6, 265–286, [https://doi.org/10.1016/0146-6313\(59\)90086-3](https://doi.org/10.1016/0146-6313(59)90086-3), 1959.
- Kuntz, L. B. and Schrag, D. P.: Hemispheric asymmetry in the ventilated thermocline of the tropical Pacific, *Journal of Climate*, 31, 1281–1288, <https://doi.org/10.1175/JCLI-D-17-0686.1>, 2018.
- 480 Lassen, H., Barriga, E., Palacios, J., Vargas, N., Díaz, E., and Argüelles, J.: Evaluación del estado del stock de merluza (*Merluccius gayi* peruanus Ginsburg) en el mar peruano. 2008, Tech. Rep. 1-2, Instituto del Mar del Peru, Callao, https://hdl.handle.net/20.500.12958/1016_2009a.
- Lassen, H., Barriga, E., Palacios, J., Vargas, N., Díaz, E., and Argüelles, J.: Evaluación del estado del stock de merluza (*Merluccius gayi* peruanus Ginsburg) en el mar peruano. 2008, Tech. Rep. 1-2, Instituto del Mar del Peru, Callao, 2009b.
- 485 Lukas, R.: The termination of the Equatorial Undercurrent in the eastern Pacific, *Progress in Oceanography*, 16, 63–90, [https://doi.org/10.1016/0079-6611\(86\)90007-8](https://doi.org/10.1016/0079-6611(86)90007-8), 1986.
- Madec, G.: NEMO ocean engine, Note du Pôle de modélisation de l’Institut Pierre-Simon Laplace No 27, France, 2008.
- Martina, J.: Analisis de la captura por unidad de esfuerzo (CPUE) de la anguila común (*Ophichthus pacifici*) y su relacion con la Corriente Cromwell, Ph.D. thesis, Universidad Nacional Agraria La Molina, http://ban.lamolina.edu.pe/search~S1*sp?/?cM11+M37+-+T/cm+++11+m37+t/-3%2C-1%2C0%2CB/frameset&FF=cm++++11+m37+t&1%2C1%2C, 2004.
- 490 Martina, J.: Estimación del coeficiente de capturabilidad (q) y de la biomasa de anguila común (*Ophichthus remiger*) del norte de Perú mediante análisis geoestadístico de las densidades de captura using likelihood-based geostatistical method on fish density, *Anales Científicos*, 79, 168–177, <https://doi.org/http://dx.doi.org/10.21704/ac.v79i1.1159>, ISSN2519-7398(Versión electrónica), 2018.
- 495 McPhaden, M. J.: Genesis and Evolution of the 1997–98 El Niño, *Science*, 283, 950–954, <https://doi.org/https://doi.org/10.1126/science.283.5404.950>, 1999.
- Montes, I., Colas, F., Capet, X., and Schneider, W.: On the pathways of the equatorial subsurface currents in the eastern equatorial Pacific and their contributions to the Peru-Chile Undercurrent, *Journal of Geophysical Research: Oceans*, 115, 1–16, <https://doi.org/10.1029/2009JC005710>, 2010.
- 500 Ñiquen, M. and Bouchon, M.: Impact of El Niño events on pelagic fisheries in Peruvian waters, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 51, 563–574, <https://doi.org/10.1016/j.dsr2.2004.03.001>, 2004.

- Pennington, J. T., Mahoney, K. L., Kuwahara, V. S., Kolber, D. D., Calienes, R., and Chavez, F. P.: Primary production in the eastern tropical Pacific: A review, *Progress in Oceanography*, 69, 285–317, <https://doi.org/10.1016/j.pocean.2006.03.012>, 2006.
- 505 Qin, X., Gupta, A. S., and van Sebille, E.: Variability in the origins and pathways of Pacific Equatorial Undercurrent, *Journal of Geophysical Research: Oceans*, 120, 3113–3128, <https://doi.org/10.1002/2014JC010549>, 2015.
- Qin, X., Menviel, L., Sen Gupta, A., and van Sebille, E.: Iron sources and pathways into the Pacific Equatorial Undercurrent, *Geophysical Research Letters*, 43, 9843–9851, <https://doi.org/10.1002/2016GL070501>, 2016.
- Ryan, J. P., Ueki, I., Chao, Y., Zhang, H., Polito, P. S., and Chavez, F. P.: Western Pacific modulation of large phytoplankton blooms in the central and eastern equatorial Pacific, *Journal of Geophysical Research: Biogeosciences*, 111, 1–14, <https://doi.org/10.1029/2005JG000084>,
510 2006.
- Rykaczewski, R. R. and Checkley, D. M.: Influence of ocean winds on the pelagic ecosystem in upwelling regions, *Proceedings of the National Academy of Sciences of the United States of America*, 105, 1965–1970, <https://doi.org/10.1073/pnas.0711777105>, 2008.
- Shaffer, G., Pizarro, O., Djurfeldt, L., Salinas, S., and Rutllant, J.: Circulation and low-frequency variability near the Chilean coast: Remotely forced fluctuations during the 1991–92 El Niño, *Journal of Physical Oceanography*, 27, 217–235, [https://doi.org/10.1175/1520-0485\(1997\)027<0217:CALFVN>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<0217:CALFVN>2.0.CO;2), 1997.
- 515 Slemmons, L., Gorgues, T., Aumont, O., Menkes, C., and Murray, J. W.: Biogeochemical impact of a model western iron source in the Pacific Equatorial Undercurrent, *Deep-Sea Research Part I: Oceanographic Research Papers*, 56, 2115–2128, <https://doi.org/10.1016/j.dsr.2009.08.005>, <http://dx.doi.org/10.1016/j.dsr.2009.08.005>, 2009.
- Tam, J., Purca, S., Duarte, L. O., Blaskovic, V., and Espinoza, P.: Changes in the diet of hake associated with El Niño 1997–1998 in the northern Humboldt Current ecosystem, *Advances in Geosciences*, 6, 63–67, <https://doi.org/10.5194/adgeo-6-63-2006>, 2006.
- 520 Tam, J., Taylor, M. H., Blaskovic, V., Espinoza, P., Michael Ballón, R., Díaz, E., Wosnitza-Mendo, C., Argüelles, J., Purca, S., Ayón, P., Quipuzcoa, L., Gutiérrez, D., Goya, E., Ochoa, N., and Wolff, M.: Trophic modeling of the Northern Humboldt Current Ecosystem, Part I: Comparing trophic linkages under La Niña and El Niño conditions, *Progress in Oceanography*, 79, 352–365, <https://doi.org/10.1016/j.pocean.2008.10.007>, <http://dx.doi.org/10.1016/j.pocean.2008.10.007>, 2008.
- 525 Taylor, M. H., Tam, J., Blaskovic, V., Espinoza, P., Michael Ballón, R., Wosnitza-Mendo, C., Argüelles, J., Díaz, E., Purca, S., Ochoa, N., Ayón, P., Goya, E., Gutiérrez, D., Quipuzcoa, L., and Wolff, M.: Trophic modeling of the Northern Humboldt Current Ecosystem, Part II: Elucidating ecosystem dynamics from 1995 to 2004 with a focus on the impact of ENSO, *Progress in Oceanography*, 79, 366–378, <https://doi.org/10.1016/j.pocean.2008.10.008>, <http://dx.doi.org/10.1016/j.pocean.2008.10.008>, 2008.
- Terada, M., Minobe, S., and Deutsch, C.: Mechanisms of future changes in equatorial upwelling: CMIP5 intermodel analysis, *Journal of Climate*, 33, 497–510, <https://doi.org/10.1175/JCLI-D-19-0128.1>, 2020.
- 530 Tsuchiya, M., Lukas, R., Fine, R. A., Firing, E., and Lindstrom, E.: Source waters of the Pacific Equatorial Undercurrent, *Progress in Oceanography*, 23, 101–147, [https://doi.org/10.1016/0079-6611\(89\)90012-8](https://doi.org/10.1016/0079-6611(89)90012-8), 1989.
- Wang, Q., Wang, F., Feng, J., Hu, S., Zhang, L., Jia, F., and Hu, D.: The Equatorial Undercurrent and Its Origin in the Region Between Mindanao and New Guinea, *Journal of Geophysical Research: Oceans*, 124, 2313–2330, <https://doi.org/10.1029/2018JC014842>, 2019.
- 535 Wolff, M., Taylor, M., Mendo, J., and Yamashiro, C.: A catch forecast model for the Peruvian scallop (*Argopecten purpuratus*) based on estimators of spawning stock and settlement rate, *Ecological Modelling*, 209, 333–341, <https://doi.org/10.1016/j.ecolmodel.2007.07.013>, 2007.
- Yu, J. Y., Kao, H. Y., Lee, T., and Kim, S. T.: Subsurface ocean temperature indices for Central-Pacific and Eastern-Pacific types of El Niño and La Niña events, *Theoretical and Applied Climatology*, 103, 337–344, <https://doi.org/10.1007/s00704-010-0307-6>, 2011.




Figure 1. Representation of particles “seeded” (initial positions) off Peru and Ecuador at 30-m depth (red dots) and 100-m depth (blue dots) for upwelling rates exceeding $1.9 \times 10^{-5} \text{ m s}^{-1}$ (50 m per month) in the region bounded by 82.5°W, 10°S and the Equator, in this particular case for 1997 and 1998 years. Particles were released at the end of each month December-January, for each year, over the period 1988 to 2007. Consider that initial particles here represented as blue and red dots are overlapping each-other, although, 30-m released depth is in the front while 100-m released depth is in the background respectively.




Figure 2. Vertical profiles for zonal and meridional velocity components (red and blue respectively) averaged over 1988-2007 (first top panel), and monthly vertical sections for zonal velocity component (bottom panels, contour lines highlight 0.5 m.s^{-1} and 1.0 m.s^{-1}) in the NEMO-ORCA12 hindcast and NOAA mooring at 170°W , 140°W and 110°W longitudes respectively.




Figure 3. Monthly climatology (averaged over 1988-2007) of vertical velocity at a depth of 33 m in the eastern Pacific, from NEMO-ORCA12 hindcast.




Figure 4. Monthly climatologies (averaged over 1988-2007) of vertical velocity at a depth of 100 m in the eastern Pacific, from NEMO-ORCA12 hindcast.




Figure 5. Grand ensemble of back-trajectory data (averaging the 1989-2007 ensembles), for particles back-tracked at the end of each month (December-January) at 30-m depth (left panel) and 100-m depth (right panel) respectively, binning at $0.5^\circ \times 0.5^\circ$ resolution: (a) particle concentration (calculated as a fractional particle presence in Log10 scale, by dividing the number of particle occurrence passing through each grid cell by the total number of particles occurrences during the course of the year); (b) particle age (days, backward in time); (c) particle depth (m); (d) particle temperature ($^\circ\text{C}$), (e) particle salinity (psu); (f) particle potential density (σ_θ , $\text{kg}\cdot\text{m}^{-3}$).

Monthly Climatology	160°W	150°W	140°W	130°W	120°W	110°W	95°W	92°W
January	25.3	28.9	30.7	30.1	24.9	18.4	13.5	12.1
February	28.3	27.4	25.3	23.1	21.3	17.2	13.2	12.2
March	44.2	45.0	39.1	30.8	22.1	17.1	12.6	11.7
April	53.5	54.0	49.9	43.8	33.6	26.9	22.2	18.7
May	56.1	51.3	44.0	37.6	31.0	26.1	18.3	14.6
June	45.2	42.5	37.0	32.5	28.0	21.9	15.1	13.0
July	39.2	34.7	30.0	25.1	20.6	18.5	11.3	9.5
August	31.9	29.7	27.2	25.1	21.6	17.1	11.6	10.3
September	26.1	27.4	29.1	28.3	27.1	23.5	14.0	12.4
October	26.5	27.8	29.3	29.2	27.1	23.6	16.5	13.8
November	28.5	32.6	34.2	33.0	30.5	22.9	11.8	10.4
December	29.1	33.4	35.8	34.1	31.1	23.4	13.0	11.2
Annual average	36.2	36.2	34.3	31.0	26.6	21.4	14.4	12.5

Table 1. Monthly Climatology (from 5-day averaged data from 1988 to 2007) and annual average for the EUC transports (Sv) at 160°W, 150°W, 140°W, 130°W, 120°W, 110°W, and 92°W.




Figure 6. Integrated EUC transport based on calculations at 160°W, 150°W, 140°W, 130°W, 120°W, 110°W, 95°W and 92°W (Galapagos Islands), in the latitude range 3°N-3°S where the concentration of particles backtracked is the highest: (a) monthly averages; (b) monthly anomalies.




Figure 7. Annual number (from December-January monthly releases) of particles and percentage associated to the EUC (3°N-3°S) at 160°W, 150°W, 140°W, 130°W, 120°W, 110°W, 100°W and 92°W, for 30-m releases in (a) and (b), and for 100-m releases in (c) and (d).





Figure 8. Diagnostics of upwelling in the north Peruvian Upwelling System, spanning the latitude range 5-10°S: (a) equatorward wind stress ($\text{N}\cdot\text{m}^{-2}$); (b) Ekman flux (Sv); (c) Upwelling flux (Sv). Climatologies are obtained by averaging across the 1988-2007 time series.

(a) Local winds dominate upwelling

- ✓ Weak EUC in the Pacific.
- ✓ EUC does not influence in the upwelling water off Peru much.
- ✓ Easterly winds present.
- ✓ Warm pool restricted to the western basin.



(b) EUC dominates upwelling

- ✓ Strong EUC in the Pacific.
- ✓ EUC highly influences in the upwelling water off Peru.
- ✓ Strong Westerly winds (weak or reversed easterly winds).




Figure 9. Schematic drivers and processes associated with the Peruvian upwelling system: (a) dominated by local winds; (b) dominated by EUC source waters.

R	160°W	150°W	140°W	130°W	120°W	110°W	92°W
160°W	1.00	0.79	0.53	0.27	N/S	N/S	-0.17
150°W	0.79	1.00	0.71	0.45	0.23	N/S	-0.14
140°W	0.53	0.71	1.00	0.68	0.42	0.19	-0.11
130°W	0.27	0.45	0.68	1.00	0.68	0.39	N/S
120°W	N/S	0.23	0.42	0.68	1.00	0.70	0.20
110°W	N/S	N/S	0.19	0.39	0.70	1.00	0.43
92°W	-0.17	-0.14	-0.11	N/S	0.20	0.43	1.00

Table 2. Correlation coefficient (R) matrix at 0-lag for the EUC transport at the selected longitudes, for R values with p-values <0.01 (99 % confidence interval). Not significant R values are indicated as N/S.