

Salinity as a key control on the diazotrophic community composition in the Baltic Sea

Christian Furbo Reeder¹, Ina Stoltenberg¹, Jamileh Javidpour¹, Carolin Regina Löscher^{1,2}

¹Department of Biology, University of Southern Denmark, Campusvej 55, 5230 Odense M, DK

5 ²Danish Institute for Advanced Study, University of Southern Denmark, Campusvej 55, 5230 Odense M, DK

Correspondence should be addressed to crfurbo@biology.sdu.dk

Abstract. Over the next decade, the Baltic Sea is predicted to undergo severe changes including ~~a-decreased~~ in salinity due to ~~altering~~~~-altered~~ precipitation ~~related to climate changes~~. This will likely impact the distribution and community composition of Baltic Sea ~~dinitrogen~~ (N_2 -) fixing microbes, of which especially ~~heterocytous~~ ~~heteroeystous~~-cyanobacteria are adapted to low salinities and may expand to waters with currently higher salinity, including the Danish Strait and Kattegat, while other high-salinity adapted N_2 fixers might decrease in abundance.

10

In order to explore the impact of salinity on the distribution and activity of different diazotrophic clades, we followed the natural salinity gradient from the Eastern Gotland and Bornholm Basins through the Arkona Basin to the Kiel Bight and combined N_2 fixation rate measurements with a molecular analysis of the diazotrophic community using the key functional marker gene for N_2 fixation *nifH*, as well as the key functional marker genes *anf* and *vnf*, encoding for the two alternative nitrogenases.

15

We detected N_2 fixation rates between 0.7 and 6 nmol N L⁻¹ d⁻¹, and the diazotrophic community was dominated by the cyanobacterium ~~related to~~ *Nodularia spumigena* and the small unicellular, cosmopolitan cyanobacterium UCYN-A. *Nodularia* was present in ~~gene~~ abundances between 8.07 x 10⁵ and 1.6 x 10⁷ copies L⁻¹ in waters with salinities of 10 and below, while UCYN-A reached ~~gene~~ abundances of up to 4.5 x 10⁷ copies L⁻¹ in waters with salinity above 10. Besides those two cyanobacterial diazotrophs, we found several clades of proteobacterial N_2 fixers and alternative nitrogenase genes associated with *Rhodopseudomonas palustris*, a purple non-sulfur bacterium. Based on ~~statistical-testing~~~~principal component analysis (PCA)~~, salinity was identified as the primary parameter describing the diazotrophic distribution, while pH and temperature did not have a ~~similarly~~-significant influence on the diazotrophic distribution. While this statistical analysis will need to be explored in direct experiments, it gives an indication for a future development of diazotrophy in a freshening Baltic Sea with UCYN-A retracting to more saline North Sea waters and ~~heterocytous~~ ~~heteroeystous~~-cyanobacteria expanding as salinity decreases.

20

25

30 1 Introduction

The Baltic Sea (Fig. 1) is a marginal, brackish sea characterized by a natural salinity gradient increasing from the North-East to the South-West. ~~The Baltic Sea covers an area of 415000 km² with a permanent halocline preventing vertical mixing, oxygen (O₂) depleted waters in the deeper basins and coastal systems, accompanied with the occasional accumulation of hydrogen sulfide (H₂S) and ammonium (NH₄⁺) below the chemocline (Conley et al., 2002; Hietanen et al., 2012; Lennartz et al., 2014).~~ The Baltic Sea covers an area of 415000 km² with a permanent halocline in the Baltic Sea proper, preventing vertical mixing, oxygen (O₂)-depleted waters in the deeper basins and coastal systems, accompanied with the occasional accumulation of hydrogen sulfide (H₂S) and ammonium (NH₄⁺) below the chemocline (Conley et al., 2002; Hietanen et al., 2012; Lennartz et al., 2014). It is further challenged by a high land-derived influx of phosphorous leading to a substantial internal surface water and sedimentary phosphorous load (Gustafsson et al., 2017; Stigebrandt and Andersson, 2020). Prior studies have shown that the Baltic Sea experienced a 10-fold increase in O₂-depleted areas over the last 115 years, covering an area of 12000 – 70000 km² making it one of the ocean areas most severely affected by deoxygenation (Diaz and Rosenberg, 2008; Reusch et al., 2018). Between 1871 and 2013, the Baltic Sea showed an increase in temperature of 0.1°C decade⁻¹ exceeding the average global trend of about 0.06°C decade⁻¹ (Reusch et al., 2018; Rutgersson et al., 2014) and decreased salinity (1-2 kg⁻¹) (Liblik and Lips, 2019). Nitrogen (N) deposition rates are now among the highest in marine areas (Reusch et al., 2018). The resulting eutrophication ~~affects~~ severely affects sensitive coastal areas resulting in high pelagic production, frequent events of anoxia, and decreased biodiversity (Breitburg et al., 2018; Carstensen et al., 2014; Maar et al., 2016; Reusch et al., 2018; Rutgersson et al., 2014). In this context, the Baltic Sea has been described as a "time machine" for how future oceans will respond to climate change (Reusch et al., 2018), making it an ideal environment to investigate biological fixation of dinitrogen gas (N₂ fixation) in response to such changes.

N₂ fixation is the primary external source of new N to life in the ocean and plays a crucial role for primary production, and thus carbon dioxide (CO₂) uptake from the atmosphere (Capone and Carpenter, 1982; Gruber, 2005). Specialized organisms, called diazotrophs, carry out this highly energy-demanding process. The diversity of diazotrophs is not easy to describe, as those organisms spread across the microbial tree of life, and are found in both, the bacterial and archaeal domains (Zehr et al., 1998). N₂ fixation is catalyzed by an enzyme complex, the nitrogenase, which exists in three different subtypes: the molybdenum (Mo)-iron (Fe) type referred to as Nif, the vanadium (V)-Fe type called Vnf, and a Fe-Fe type called Anf (Chisnell et al., 1988; Robson et al., 1986). The different metal cofactors are essential in the context of the redox sensitivity of the three nitrogenases, respectively. Mo is preferentially available in the presences of at least traces of O₂, contrary to V and Fe, which are more soluble and thus available under anoxic conditions (Bennett and Canfield, 2020; Bertine and Turekian, 1973; Crusius et al., 1996; Dixon and Kahn, 2004; Morford and Emerson, 1999). The differential availability of those trace metals along redox gradients might have played a role in the evolutionary development of those nitrogenase types (Anbar, 2008; Mus et al., 2019) and their distribution might change in a future ocean impacted by acidification, deoxygenation (Keeling et al., 2010; Löscher et al., 2014; Schmidtko et al., 2017; Stramma et al., 2008) and

desalination (Olofsson et al., 2020). To date, most studies focused mainly on Nif-type nitrogenases in the marine environment, while Vnf and Anf nitrogenases were often overlooked. Molecular detection of the *nifH* gene does not fully recover the diversity of *vnfH* and *anfH*, possibly resulting in an underestimation of alternative nitrogenases in the environment (Affourtit et al., 2001; Farnelid et al., 2009, 2013; Man-Aharonovich et al., 2007; Steward et al., 2004; Zehr et al., 1995). Nonetheless, the few available studies focusing on alternative nitrogenases identified them to be abundant and diverse and suggested to play a role for N₂ fixation in various ecosystems, including the upper water column of the ocean, O₂-depleted waters, and Baltic Sea sediments (Bellenger et al., 2011, 2014; Betancourt et al., 2008; Christiansen and Löscher, 2019; Farnelid et al., 2009; Loveless et al., 1999; McRose et al., 2017; Tan et al., 2009; Zehr et al., 2003; Zhang et al., 2016).

In the Baltic Sea, N₂ fixation is considered to take place in sunlit surface waters mainly and carried out by three genera of heterocystous cyanobacteria, *Aphanizomenon* sp., *Nodularia-N. spumigena* and *Anabaena* spp. (now referred to as *Dolichosperum* spp. (Klawonn et al., 2016; Wacklin et al., 2009)), all of them containing the classic Nif-nitrogenase (Janson et al., 1994; Leppanen et al., 1988), and contributed massively to N₂ fixation with rates of up to ~~82-115~~ ~~- 491-295~~ nmol N L⁻¹ d⁻¹ (Larsson et al., 2001). At the same time, cyanobacteria consume only a fraction of the N they fix with estimates in the range of ~~2-1933.33~~ ~~- 785.74-72~~ nmol N L⁻¹ d⁻¹ (Janson et al., 1994; Larsson et al., 2001; Wasmund, 1997). Using an EA-IRMS-based approach, Klawonn et al., 2016 measure N₂ fixation rates up to 300 ± 216 nmol N L⁻¹ d⁻¹ and More recent studies, based on cell-specific measurement of *Aphanizomenon* spp and *Nodularia-N. spumigena*, revealed that they release up to 35% of fixed N, translating into a substantial fraction of fixed N leaking out, available for other primary production producers (Ploug et al., 2010, 2011). In addition, low rates of N₂ fixation (0.44 nmol N L⁻¹ d⁻¹) have been described in anoxic waters of the Baltic Sea (Farnelid et al., 2013). These rates were accompanied by a highly diverse Nif-containing heterotrophic diazotroph community at and below the chemocline in anoxic NH₄⁺-rich waters (Farnelid et al., 2013). Heterotrophic diazotrophs were closely related to those in other O₂-depleted marine systems regions including the eastern tropical South (ETSP) and North Pacific (ETNP), the Arabian Sea (AS), and an anoxic basin in the Californian Bight (Fernandez et al., 2011; Hamersley et al., 2011; Jayakumar et al., 2012, 2017; Löscher et al., 2014). Non-cyanobacterial diazotrophs have been found to fix N₂ actively. However, it is still inconclusive what regulates and controls their N₂ fixation activity.

Over the last three decades, the Baltic Sea has experienced a trend of freshening in the range of 1-2 kg⁻¹ in surface waters (Liblik and Lips, 2019) or between 0.4 to 1.2 (Saraiva et al., 2019). Earlier studies suggested that salinity impacts the diazotrophic community composition and the activity of nitrogenases in pure cultures of *Azotobacter* sp. marine microbial mats and estuaries (Dicker and Smith, 1981; Marino et al., 2006; Severin et al., 2012). Surface salinity in the Baltic Sea ranges between 0 and 10 (Fig. 1) making it one of the major drivers for microbial composition and distribution in the Baltic Sea (Dupont et al., 2014; Olofsson et al., 2020; Wulff et al., 2018). Typically, *Aphanizomenon* sp. dominates the northern part (e.g. the Bothnian Sea) less saline waters (e.g., Bothnian Sea) with an optimum growth at salinity of 0-2, while *Nodularia-N. spumigena* prefers the higher saline waters in southern part (e.g. the Southern Baltic Proper) with an optimum

~~growth in salinity of salinities of 8-10 (Lehtimäki et al., 1997; Rakko and Seppälä, 2014). Additionally, the small, unicellular cyanobacterial symbiont UCYN-A has been detected in the Baltic Sea (Bentzon-Tilia et al., 2015). This cosmopolitan diazotroph has previously been shown to be abundant throughout most marine systems (Tang et al., 2019; Zehr et al., 2016), and to substantially contribute to N₂ fixation rates (Martínez-Pérez, C., Mohr, W., Löscher, 2016; Mills et al., 2020). However, the factors controlling the biogeography of UCYN-A is poorly understood. Though, latest study suggested salinity having a positive correlation to UCYN-A. These biogeography (Li et al., 2021). These observations speak for indicate potential changes in the cyanobacterial composition in future freshening events (Olofsson et al., 2020; Wasmund et al., 2011). To now explore the impact and importance of salinity for diazotrophy in the Baltic Sea and in order to complement existing datasets on N₂ fixation from high-productivity seasons, we carried out a survey including chemical profiling, and rate measurements in the low productive fall season. The focus of this study is a molecular genetic assessment of the diazotrophic community diversity and genetic activity including heterocyst and unicellular cyanobacteria, heterotroph diazotrophs and diazotrophs carrying alternative nitrogenase genes. To now explore the impact and importance of salinity for diazotrophy in the Baltic Sea, we carried out a survey including chemical profiling, rate measurements, and molecular genetic mining in a low productive fall season. In addition, we compared our data to available datasets to explore controls on Baltic Sea N₂ fixation and to be able to predict the future distribution of Baltic Sea diazotrophs and of N₂ fixation rates.~~

2 Material and Methods

2.1 Seawater sampling

Samples were collected during the cruise AL528 ~~on using~~ the German research vessel RV Alkor ~~to the Baltic Sea~~ from 17.09.2019 to 28.09.2019 along a transect through the major basins of the Baltic Sea (see Fig. 1 for a cruise plan and table S2 for overview of all stations). Specifically, samples were collected at a station in the Kiel fjord (KB06), one in the Arkona basin (H21), four stations in the Bornholm Basin (BB08, BB15, BB23, BB31) and the Eastern Gotland Basin (GB84, GB90a, GB107, GB108), respectively, at water depths between 3 to 115 meters (m) using a 10 L Niskin bottle rosette equipped with a conductivity- temperature- depth (CTD) sensor. Water for dissolved inorganic nitrogen (~~NO_x NO_x-(Nitrate (NO₃⁻)+ Nitrite (NO₂⁻))~~) was collected from the Niskin bottles, filtered through an acid-washed 0.8 µm cellulose acetate filter (Sartorius™), using a syringe and stored in 20 ml scintillation vials at -20°C until analysis. Samples for PO₄³⁻ analysis were filtered through a 200 µm mesh to avoid contamination with large zooplankton. Samples were filled in 20 ml scintillation vials and stored at -80°C until analysis. ~~NO_x DIN~~ and PO₄³⁻ samples were analyzed on a SKALAR SAN^{plus} analyzer (Thermo Fischer, Waltham, US) according to Grasshoff (1999) with a detection limit of 0.05 µmol N L⁻¹ for ~~NO_x DIN~~ and 0.03 µmol P L⁻¹ for PO₄³⁻. Nutrient samples were taken in triplicates, handled with nitrile gloves and all equipment was rinsed with MilliQ water in between stations to avoid contamination. Samples for dissolved inorganic carbon (DIC) were collected by filling triplicates of 12 mL exetainers bubble-free at stations KB06, H31, H21 and BB15. Samples were ~~fixed~~ fixed with 20µl of a saturated mercury chloride solution and stored at room temperature until analysis. DIC measurements

were carried out using a 2 mM NaHCO₃ solution as standard and as described previously (Hall and Aller, 1992). ~~Water samples for DNA extraction were collected from Niskin bottles. 0.5 to 1 L of seawater~~ From all stations and depths 0.5 to 1 L of seawater were collected from Niskin bottles and were immediately filtered onto a 0.22 µm pore size membrane filters (Millipore, Bilateralica, USA; exact seawater volumes were constantly recorded), and filters were stored at -80 ° C until DNA extraction and further analysis.

2.3 ¹⁵N₂ and ¹³C seawater incubations

N₂ and C fixation rates were determined using stable isotope labelling at KB06, H21, BB15, BB08 and GB107, at water depths between 3 and 41 m. We used the bubble addition method (Montoya et al., 1996) to ensure comparability to previous studies from the Baltic Sea. Triplicate water samples were filled from Niskin bottles into 2.4 L glass bottles until it was top-filled (Schott-Duran, Wertheim, Germany). To each incubation bottle, 1 mL ¹⁵N₂ gas (¹⁵N₂ 98%+ Lot#I-16727, Cambridge Isotope Laboratories, Inc., USA) and 1 mL H¹³CO₃ (1g 50 mL⁻¹, Sigma-Aldrich, Saint Louis Missouri US) was added through an air-tight septum and bottles were inverted to ensure mixing with final calculated concentrations of ~~was~~ 0.05% ¹⁵N₂ and 10 µg mL⁻¹ H¹³CO₃. After an incubation time of 24 h, samples were filtered onto pre-combusted GF/F filters (GE Healthcare Life Sciences, Whatman, USA). A parallel set of triplicate samples was collected from each sampling depth for determining the natural abundance of N and C isotopes in the particulate organic material. Filters were stored at -20°C until further analysis. Filters were then acidified, and ~~followed by~~ dried in an oven at 65 °C. Dried filters were analysed on an Elemental Analyzer Flash EA 1112 series (Thermo Fisher), coupled to an isotope ratio mass spectrometer (Finnigan Delta Plus XP, Thermo Fisher).

2.2 Molecular methods

For nucleic acid extraction, ~~0.22 µm pore sized millipore~~ Millipore membrane filters (Millipore, Bilateralica, USA) were flash-frozen in liquid N and subsequently crushed. DNA was purified with the MasterPure Complete DNA and RNA purification Kit (Lucigen, Wisconsin, US) according to the manufacturer's protocol, with the minor modification of using 1 mL of lysis buffer to completely cover the filter pieces. RNA was purified using Qiagen AllPrep DNA/RNA Mini Kit (Qiagen, Hilden, Germany). The remaining DNA was removed with the gDNA wipeout mix, and a cDNA library was constructed using the QuantiTect Reverse Transcription Kit (Qiagen, Hilden, Germany) with the supplied RT primer set. The nucleic acid concentration and quality were checked spectrophotometrically on a MySpec spectrofluorometer (VWR, Darmstadt, Germany).

A total of 15 DNA samples were amplified for *nifH* using nested PCR with primers and PCR conditions as described in Zani et al. (2000). *anf/vnfD* were amplified using a nested PCR with primers and conditions as described in Bellenger et al. (2014) and McRose et al. (2017). Amplicons were TOPO TA-cloned (Topo TA cloning Kit for sequencing, Thermo Fisher Scientific, Waltham, US) and Sanger-sequenced. Sequencing was carried out at the Institute of Clinical Molecular Biology

160 in Kiel, Germany. Samples for *nifH* and *anf/vnfD* amplification were the same as the ones where N₂ rate measurements were carried out (KB06, H21, BB15, BB08 and GB107 between 3 and 41 meters, see table S1).

Sequences were quality checked and trimmed with BioEdit (Hall, 1999). *nifH* sequences were trimmed to 321 bp and *vnf/anfD* to 591 bp. This resulted in 182 *nifH* and 69 *vnf/anfD* amplicon sequences. Sequences were BLASTX-searched against the NCBI database. A reference library was created, and sequences with lower than 80% identity were discarded. Sequences were aligned in MEGAX (Kumar et al., 2018), and a maximum likelihood tree was constructed using 1000 bootstraps. To explore the overall microbial diversity, 23 samples from Kiel Fjord (station KB06, depth 5 and 10 m), Arkona Basin (station H2, depth 4, 12, 42 m and station H31, depth 4 and 12 m), Bornholm Basin (Station BB23, depth 4, 48, 85 m, station BB15, depth 7, 41, 65 m and station BB08, 5 and 10 m) and East Gotland Basin (Station GB84, depth 9, 29, 110 m, station GB107, depth 8, 40 and 112 m and station GB108, depth 10 and 115 m) were sent to amplicon sequencing of the 16s rDNA V1V2 region using Illumina HiSeq technology. Depths were chosen based on the water column chemistry (halocline, oxycline, oxygen minimum zones and chlorophyll peaks). Sequencing was carried out by the Institute of Clinical Molecular Biology in Kiel, Germany. Raw reads were trimmed, quality filtered in dada2 (v1.18) (Callahan et al., 2016), and taxonomically annotated using SILVA database (v138) (Quast et al., 2013). Visualization of the cyanobacterial distribution was accessed with *phyloseq* (1.34.0) (McMurdie and Holmes, 2013).

175 Quantitative real-time PCRs were carried out targeting *nifH* clade-specifically for *Nodularia*, UCYN-A and gamma-proteobacterial diazotrophs (Gamma AO, Gamma PO) as described previously (e.g., Boström et al., 2007; Langlois et al., 2008; Loescher et al., 2014). As standards, serial dilutions of plasmids (10⁷ to 10¹ copies) containing the target *nifH* genes were used. Samples, standards, and non-template controls were run in duplicates on a Biorad qPCR machine (Biorad, Hercules, USA), and reactions were considered uncontaminated if no amplification could be detected in non-template controls after 40 qPCR cycles. To further ensure that no DNA contamination was present in our cDNA, we ran additional qPCR reactions of RNA samples. No amplification was observed. Sequences were submitted to GenBank with accession numbers MZ063808-MZ063873 and MZ063874-MZ064057 and 16S rDNA with accession number SAMN20309768-SAMN20309783.

2.4 Statistical Methods

185 Principle component analyses (PCA) on qPCR and environmental data were performed in R using the *vegan* package (Oksanen et al., 2020; R Foundation for Statistical Computing, 2017). The *simper* function (from *vegan* package) was used to identify parameters with the highest impact on diazotroph abundance. A principal component analysis (PCA) was subsequently performed using *prcomp* and visualized with the *factoextra* package (Kassambara and Mundt, 2020). To maximize the amount of explained variance of the dataset, ~~Included in the~~ figure ~~are both included both~~ component one
190 versus two (Fig. 8A), and one versus three (Fig. 8B).

3 Results and Discussion

3.1 Hydrochemistry

During the cruise, surface waters above 40 m water depth exhibited the typical Baltic Sea salinity gradient with salinities decreasing from South-West to North-East (10 – 24 in the Kiel and Arkona Basin and 7-9 in the Bornholm and Eastern Gotland Basins), accompanied by lower temperatures of 13-14°C in the Bornholm and Eastern Gotland Basin compared to slightly higher temperatures of 15-16°C in the Kiel Fjord and Arkona Basin (Fig. 2). A thermocline was observed at 45 m in the Bornholm Basin and 65 m in the Eastern Gotland Basin. The coastal station we sampled in the Arkona basin showed a similar stratification pattern (Fig. 2) with a salinity of 10 on the surface and 17 below 30 m water depth. This stratification pattern is typical for the region and time of the year and could also be observed from annual data from the International Council for the Exploration of the Sea (ICES Copenhagen, 2020) (Fig. 1, B). Surface waters were well-oxygenated with O₂ concentrations above 156.31 μmol L⁻¹ in the top 40 m of the water column. Low O₂ water masses with concentrations below 15 μmol L⁻¹ were only observed in the Bornholm basin-Basin in waters deeper than 65 m (Fig. 2).

Compared to the South-western part, the DIC concentrations were lower in the upper water column of the Eastern Gotland and Bornholm Basin with concentrations of around 1.6 mmol L⁻¹ and increased towards the south-western part of the cruise track. An opposite trend was visible in nutrient distribution with NO_x (nitrite + nitrate) and phosphate (PO₄³⁻) decreasing from the North-Eastern part of the cruise track towards the South-Western part with surface water concentrations of 9.04 μmol L⁻¹ for NO_x and 2.98 μmol L⁻¹ for PO₄³⁻ in surface waters of the Eastern Gotland Basin and 0.27 μmol L⁻¹ for NO_x and 0.18 μmol L⁻¹ for PO₄³⁻ in the Kiel Fjord (Fig. 2). The observed nutrient concentrations were in the range of the Helsinki commission (HELCOM) dataset containing data from 2015 to 2020 and the HELCOM core indicator report (Helcom, 2018, Fig S2). The detected somewhat higher nutrient concentrations in the Bornholm and Eastern Gotland Basins could result from a decaying phytoplankton bloom releasing nutrients including ammonia, decreased microbial activity or increased eutrophication. ~~The detected somewhat higher nutrient concentrations in the Bornholm and Eastern Gotland Basins could result from a decaying phytoplankton bloom, decreased microbial activity or increased eutrophication.~~ High chlorophyll concentrations in the Bornholm basin (3 μM Chl a in August and 2 μM Chl a in September), derived from the HELCOM dataset (from same areas as this cruise), indeed give evidence of a decaying phytoplankton bloom in the Bornholm Basin in September 2019 (Fig. S3). These values are in range with the HELCOM indicator report with an average of 4 μM Chl a from June to September, and a previous study presenting a range between 2 and 6 μM Chl a (Helcom, 2018; Suikkanen et al., 2010).

Our dataset indicates an N:P ratio well below the Redfield ratio of 16:1 in surface waters along the cruise track, with excess P classically supposed to promote N₂ fixation. However, NO_x DIN_ is was still available to the phytoplankton and microbial community as supported by a positive intercept of the trendline, which might be considered unfavourable for N₂ fixation (Fig. 3). Remaining NO_x concentrations in the surface waters above 40 m thus speak for a limitation of primary production by either micronutrients, temperature, or light availability and less for a limitation by N or PO₄³⁻ (Arrigo, 2005;

Saito et al., 2008). Nitrogen repletion is supported by POC: PON ratios of 3.6 to 4.94 in the Kiel Fjord and Arkona Basin and 2.92 to 6.15 in the Bornholm and Eastern Gotland Basins (Fig. 3), which are close to the Redfield ratio (Redfield, 1934). Altogether, nutrient concentrations do not strongly support a niche for N₂ fixation.

3.2 Nitrogen fixation

While no clear niche for N₂ fixation could be identified, we still detected N₂ fixation rates in our samples from water depths between 3 and 41 m (Fig. 4). N₂ fixation rates were between 0 to 6 ± 4.73 nmol N L⁻¹ d⁻¹ in the Arkona Basin, 0.7 ± 0.97 to 27.3 ± 38.21 nmol N L⁻¹ d⁻¹ in Bornholm Basin and 0.67 ± 0.51 to 2 ± 0.4 nmol N L⁻¹ d⁻¹ in the Eastern Gotland Basin (Fig. 4). In Kiel Fjord, N₂ fixation rates were below the detection limit (0.03 nmol N L⁻¹ d⁻¹) of our method. One outlier was observed in the Bornholm Basin at a water depth of 5 m, with an N₂ fixation rate of 27.3 nmol N L⁻¹ d⁻¹. This high rate was only detected in one out of our three replicates and was not included in our further analysis. Without this outlier, a conservative estimate of N₂ fixation in the Bornholm Basin would be between 0.7 ± 0.97 and 5.27 ± 1.37 nmol N L⁻¹ d⁻¹. Though these rates are at the lower end of previous observations from the Baltic Sea (~~Table S3~~Table S3), they are comparable to a previous study from the same region, but not the same season, with N₂ fixation rates of 7.6 ± 1.76 nmol N L⁻¹ d⁻¹ from July and August (Farnelid et al., 2013). N₂ fixation activity is supported by the δ¹⁵N -PON in our control samples which was in the range of control samples was found to be 0.3 to 2 ‰ in samples from surface waters above 40 m from the Eastern Gotland and Bornholm Basins (Fig. 4). Values between -2 to +2 ‰ are considered indicative for N₂ fixation (Dähnke and Thamdrup, 2013; Delwiche and Steyn, 1970).

Seasonal variations might explain the observed lower N₂ fixation rates if compared to previous studies. N₂ fixation rates are generally described sensitive to low temperatures (Brauer et al., 2013; Church et al., 2009; Englund and Meyerson, 1974; Moisaner et al., 2010; Stal, 2009) with the energetic costs of N₂ fixation sharply rising at temperatures below 21 °C (Brauer et al., 2013). During our cruise, surface temperatures were between 12 °C and 16 °C and would most likely play a role in the low rates observed. Yet, growth of *N. ~~odularia~~ spumigena* has been demonstrated at 4 °C suggesting a relevance during wintertime in the Baltic Sea (Olofsson et al., 2019). Additionally, lower light intensities during fall and winter ~~lower light intensities~~ could directly influence the abundance of cyanobacterial diazotrophs in the Baltic Sea (Dera and Woźniak, 2010; Staal et al., 2002).

N₂ fixation rates were accompanied by low C fixation rates compared to rates in the micromolar range as typically detected in the Baltic Sea (e.g., (Klawonn et al., 2016)) ~~low C fixation rates~~ (Fig. 4), which were, however, increasing over depth, with an average C fixation of 29.16 ± 10.89 nmol C L⁻¹ d⁻¹ in Kiel Fjord, 42.62 ± 7.86 nmol C L⁻¹ d⁻¹ the Arkona Basin, 40.52 ± 13.16 nmol C L⁻¹ d⁻¹ in Bornholm Basin and 58.52 ± 15.42 nmol C L⁻¹ d⁻¹ in Eastern Gotland Basin. Assuming Redfield stoichiometry, N₂ fixation sustained between 8.8 and 108% with an average of 43% of C fixation during our cruise. These rates are comparable with the contribution of N₂ fixation to C fixation in other oceanic regions, including the Atlantic (20-25%) and Pacific Ocean (1-4%), the Mediterranean Sea (8%), and the Bay of Bengal (<1%) (Church et al.,

2009; Dore et al., 2002; Fonseca-Batista et al., 2019; Rahav et al., 2013; Saxena et al., 2020; Tang et al., 2019) (Table 21). The high contribution to primary production indicates that N₂ fixation may promote or extend primary production into the fall season if only at low rates. ~~Notably, the high contribution might be bias due to~~ Other sources, however, might include anthropogenic land or riverine input.

260 3.3 Diazotrophic community composition

We assessed the diazotrophic diversity based on *nifH* and *vnf/anfD* amplicons (Fig. 5 and 6). Sequences belonging to *Nodularia-N. spumigena* (19%) and UCYN-A (26%) were dominant in the sequence pool. While *Nodularia* clades are classically abundant in the Baltic Sea waters (Boström et al., 2007; Farnelid et al., 2013; Klawonn et al., 2016; Larsson et al., 2001), only one previous report of UCYN-A from the Baltic Sea exists, where it had been detected in waters of the Danish Strait and the Great Belt (Bentzon-Tilia et al., 2015). Moreover, we detected *Pseudanabaena*-like sequences representing 8 % of the sequence pool consistent with previous studies (e.g. Acinas et al., 2009; Farnelid et al., 2013; Klawonn et al., 2016; Stal et al., 2003). Only 1.5 % of the sequences belonged to *Aphanizomenon sp.*, a diazotroph known to dominate N₂ fixation early in the season with salinity at 5-6 (Klawonn et al., 2016; Svedén et al., 2015). The low number found in this study might be seasonally related, be adapted to waters with extremely low salinity (0-2) as present in the Northern part of the Baltic Sea (Lehtimäki et al., 1997; Rakko and Seppälä, 2014) and below the salinity observed during our cruise (>8).

270 Notably, based on our *nifH* sequence analysis, we did not identify, *Anabaena-Dolichospermum spp. lemmermannii*, one of the common N₂ fixing cyanobacteria for the Baltic Sea (Bentzon-Tilia et al., 2015; Boström et al., 2007; Farnelid et al., 2013). However, sequences related to *A. lemmermannii* were recovered from our 16S rDNA amplicon sequencing (Fig. S1 ~~in~~ supplementary). This discrepancy might speak for a bias against their specific *nifH* genes by our approach, or the clades we found in the 16S rDNA dataset might lack a *nifH* gene.

We detected *Nodularia*-like sequences (primer/probe constructed against *Nodularia-N. spumigena* and environmental *Nodularia* clusters, see Boström et al., 2007), in abundances of 3.37 x 10⁵ to 4.06 x 10⁷ copies L⁻¹ at 3-5 m water depth and 8.64 x 10⁴ – 4.46 x 10⁶ copies L⁻¹ at water depths of 20-41 m. *Nodularia*-specific *nifH* transcript abundance decreased over depth from 4.20 x 10⁵ (5 m) to 3.5 x 10² (20 m) transcripts L⁻¹ (Fig. 7). *Nodularia* followed the temperature-salinity gradient with the highest abundance (up to 1.6 x 10⁷ copies L⁻¹) in waters with salinity at 10 and below (the Bornholm and Eastern Gotland basin) and lowest abundances (up 1.46 x 10⁶ copies L⁻¹) in waters with a salinity above 10 (Kiel Fjord and the Arkona basin). *Nodularia*-specific *nifH* gene and transcript abundances were in the same range as previously described (10⁶ gene copies L⁻¹ and 10⁵ transcripts L⁻¹ (Farnelid et al., 2013)). UCYN-A was identified based on both *nifH* and 16S rDNA sequences and quantified using cluster-specific qPCR. UCYN-A was present in abundances of 3.73 x 10³ – 9.97 x 10⁷ copies L⁻¹, similar to the one available previous study where the abundance of *nifH* specific for UCYN-A peaked in September (approximately 10⁶ copies L⁻¹) (Bentzon-Tilia et al., 2015) (Fig. 7). The same study showed *nifH* expression (roughly 10³ transcript L⁻¹) of UCYN-A in the Baltic Sea surface waters (1 m). These transcript abundances are comparable

with those obtained from our study with transcript abundances between 2.98×10^2 and 3.73×10^2 transcripts L^{-1} at 5 m water depth and up to 3.71×10^5 transcripts L^{-1} at a water depth of 20 m (Fig. 7). Notably, the above-mentioned study took place in the Danish Strait, and thus in more saline waters (salinity of 13-17).

Contrary to *Nodularia*, UCYN-A reached highest abundances (up to 4.52×10^7 copies L^{-1}) in higher saline waters impacted by the North Sea (the Kiel Fjord and Arkona Basin) and the lowest abundance (up to 6.29×10^5 copies L^{-1}) in low saline waters (Fig. 7, Supplementary Figure S1). UCYN-A is increasingly found throughout the oceans, often playing an important role in N_2 fixation (e.g. Martínez-Pérez, C., Mohr, W., Löscher, 2016; Tang and Cassar, 2019). Over the recent years, UCYN-A has been shown to have a crucial part in N_2 fixation and has been identified all over the globe, from polar to tropical regions and tolerating between ~~12°C~~ 4°C and 30°C (Gradoville et al., 2020; Harding et al., 2018; Martínez-Pérez, C., Mohr, W., Löscher, 2016; Mills et al., 2020; Short and Zehr, 2007; Tang and Cassar, 2019). Despite UCYN-A exposing highest abundances in Baltic Sea waters with higher salinity, we found UCYN-A present and active across the salinity gradient.

Besides those cyanobacterial diazotrophs, we detected several clades of proteobacteria related to *Rhodopseudomonas palustris* (4%), *Pelobacter sp* (9%) and gamma proteobacteria of the Gamma-AO clade (3%, Langlois et al., 2008). We quantified gamma-proteobacterial *nifH* in abundances between 1.78×10^4 - 1.82×10^5 copies L^{-1} (Gamma AO) and their transcripts in abundances of 2.89×10^3 - 3.91×10^3 transcripts L^{-1} . Despite lack of clones clustering with Gamma PO, the group could be detected in abundances between 8.21×10^2 - 1.37×10^5 copies L^{-1} and their transcripts in abundances between 3.04×10^3 - 8.05×10^3 transcripts L^{-1} , indicating a potential contribution to N_2 fixation of those clades (Fig. 7). Diazotrophic proteobacteria are common in the ocean, however, their role for N_2 fixation is not fully understood (Benavides et al., 2018; Chen et al., 2019; Turk-Kubo et al., 2014). Diazotrophs related to Cluster III were dominated by *Desulfovibrio*-like (1.5%), *Opitutaceae*-like (6%) and *Clostridium*-like sequences (1.5%). However, both *nifH* gene and transcript abundances were below the detection limit of our qPCR. Altogether, the gene and transcript abundances of cyanobacterial diazotrophs were three to four orders of magnitude higher than those of non-cyanobacterial clusters suggesting that heterotrophic microbes may have only played a minor role for N_2 fixation in Baltic Sea surface waters during our cruise.

In the pool of alternative nitrogenase sequences of the *anfD* type, we identified the purple non-sulfur bacterium *Rhodopseudomonas palustris*. We could not recover any *vnfD* sequences, which might be an adaptation to low vanadium concentrations in the Baltic Sea (Bauer et al., 2017). Purple non-sulfur bacteria (e.g. *Rhodopseudomonas palustris*) typically require anoxic conditions to fix N_2 (Masepohl and Hallenbeck, 2010), at the sample locations monitored during our cruise. The role of alternative nitrogenases in the environment is poorly understood, partly due to a lack of data and detection systems. Our data shows however the presence of alternative nitrogenases in the Baltic Sea surface water, which may sustain N_2 fixation under certain conditions, for example in anaerobic micro-niches (Bertagnolli and Stewart, 2018; Farnelid et al., 2019; Paerl and Prufert, 1987; Pelve et al., 2017), including sinking particles (Chakraborty et al., 2021; Pedersen et al., 2018).

3.4 Factors impacting the diazotrophic distribution

To identify parameters impacting on the distribution of diazotrophs during our cruise, we carried out a *simper* analysis. Interestingly, pH and temperature were less important for describing the diazotrophic distribution. The PCA suggested that depth and salinity explained the distribution of both the diazotrophic community and N₂ fixation best (Fig. 8). Moreover, it revealed a correlation of N₂ fixation and the *Nodularia* abundance and less of a correlation between N₂ fixation and UCYN-A or other diazotroph clusters (Fig. 8). A decreasing salinity would thus not only promote the distribution of *Nodularia*-like diazotrophs, and decrease UCYN-A abundances, but also promote N₂ fixation rates in such a future scenario. Intuitively, it is of little surprise that our data (Fig. 8) and previous studies indicate salinity to be a key control for cyanobacterial distribution and N₂ fixation in the Baltic Sea (Dupont et al., 2014; Olofsson et al., 2020; Rakko and Seppälä, 2014). A recent study did also suggest a positive correlation between UCYN-A and salinity (Li et al., 2021). ~~However, a~~ While a previous study based on a compiled dataset (1979-2017) and modelling approaches indicated that salinity does not affect the biovolumes of the filamentous *Nodularia-N. spumigena* but rather species-interactions (Karlberg and Wulff, 2013; Olofsson et al., 2020). Our study cannot evaluate such interactions and their impact of N₂ fixation; however, our data do point towards *Nodularia* expanding into low salinity waters in the future thus complementing UCYN-A and possibly increasing N₂ fixation rates in those waters.

Our analysis did not show temperature and pH to be major descriptors of N₂ fixation during this cruise, consistent with previous findings with elevated ~~carbon dioxide~~ CO₂ concentrations not causing any response in N₂ fixation in the Baltic Sea (Olofsson et al., 2019; Paul et al., 2016; Wulff et al., 2018). Moreover, a very recent study showed that ocean acidification has an impact on the diazotroph community composition and ecology and can decrease bulk N₂ fixation rates in the subtropical Atlantic Ocean ~~Moreover, a very recent study showed that ocean acidification has an impact on the diazotroph community composition and can decrease N₂ fixation rates in the subtropical Atlantic Ocean~~ (Singh et al., 2021). Most likely, the contradicting results are due to other factors obscuring the stimulation by N₂ fixation (Karlberg and Wulff, 2013; Wannicke et al., 2018). Besides pH, temperature has also been suggested as a control on diazotrophy, either directly or indirectly, by impacting on O₂ solubility (Stal, 2009).

Besides, increasing nutrient loads in the Baltic Sea might further stimulate primary production leading to increased anoxia and PO₄³⁻ release from sediments (Ingall and Jahnke, 1997), which will, in turn, fuel N₂ fixation and primary production (Canfield, 2006; Saraiva et al., 2019). Filamentous cyanobacteria benefit from excess PO₄³⁻ (Olofsson et al., 2016), thus, excess PO₄³⁻ resulting from progressive deoxygenation might stimulate blooms of *Nodularia* (Degerholm et al., 2006; Schoffelen et al., 2018).

In case of a potential future freshening of the upper water column (Liblik and Lips, 2019) in combination with increased PO₄³⁻ availability both through land-derived influx and through phosphorous mobilization via bottom-water anoxia (Gustafsson et al., 2017; Ingall and Jahnke, 1997; Stigebrandt and Andersson, 2020; Vahtera et al., 2007), N₂ fixation by e.g. *Nodularia* will likely increase in the region covered by our cruise. ~~In case of a future freshening of the upper water column~~

(Liblik and Lips, 2019), our data points towards N₂ fixation by cyanobacteria such as *Nodularia* will increase. This could
355 lead to a scenario with increased bloom formations and expansion of such large ~~heterocytous heteroeyst~~ cyanobacteria across
the Baltic Sea, as previously observed (Kahru et al., 1994; Kahru and Elmgren, 2014). An increased organic matter load
would possibly lead to enhanced respiration by heterotrophic microbes promoting the further expansion of O₂ depleted
waters. This could lead to increased denitrification resulting in N-loss and emission of N₂O, ~~fueling~~ fuelling global warming,
an increase in euxinic events as already observed in coastal Baltic waters (Breitburg et al., 2018; Carstensen et al., 2014;
360 Lennartz et al., 2014). Further, blooms of *Nodularia* could lead to an increased load of Baltic Sea waters with the toxin
Nodularin (Sivonen et al., 1989), which can harm Baltic Sea biota and poison animals (Main et al., 1977; Nehring, 1993) and
ultimately humans by impacting e.g. fishery industry (Karjalainen et al., 2007).

4 Conclusion

During our cruise, we explored the diazotrophic community composition and N₂ fixation rates along the natural salinity
365 gradient in the South Baltic Sea in the fall season. N₂ fixation rates were detectable but low for the Baltic Sea and sustained
by a diazotrophic community dominated by the ~~heterocytous heteroeystous~~ cyanobacterium *Nodularia* and the small
unicellular cyanobacterium UCYN-A. The two different types of cyanobacteria occupied two different niches defined by
salinity ranges, respectively, with *Nodularia* dominating in lower saline waters (8-9) and UCYN-A in high-salinity waters
(>10). Statistical analysis revealed that N₂ fixation is quantitatively mainly driven by *Nodularia* clades and that both,
370 *Nodularia* abundances and N₂ fixation rates, are best explained by salinity. In the context of a predicted freshening of the
Baltic Sea, the habitat of *Nodularia*-like ~~heterocytous heteroeystous~~ cyanobacteria would extend towards the South-Western
part of the Baltic Sea, possibly replacing the community of UCYN-A and increasing N₂ fixation in those waters. Enhanced
N₂ fixation might facilitate primary productivity, and organic matter export to waters below the euphotic zone and could thus
have severe impacts on the Baltic Sea biogeochemistry including increased respiration, O₂ and N loss.

375 Author contribution

C. Reeder and C. Löscher designed the experiments and C. Reeder carried out the experimental work. J. Javidpour and I.
Stoltenberg designed the sampling strategy during the expedition and contributed essential datasets. C. Reeder and C.
Löscher prepared the manuscript with contributions from all the co-authors.

Competing interest

380 The authors declare that they have no conflict of interest.

Acknowledgement

We thank the captain and crew and the chief scientist J. Siling of Alkor 528 for their support during the cruise. We thank E. Laursen and R. Orloff Holm for their technical assistance. Funding for this study was received from the Villum Foundation (Grants #16518 to D. Canfield and #29411 to C. Löscher) and the European Union's Horizon 2020 research and innovation program (Grant agreement no. 774499 to J. Javidpour) for funding the GoJelly project. We thank Dr. Kuosa and an anonymous reviewer for reviewing the paper. Further support was received from the Nordcee labs at SDU and the GEOMAR- Helmholtz Centre for Ocean Research Kiel by providing access to their research vessel and facilities.

References

- 390 Acinas, S. G., Haverkamp, T. H. A., Huisman, J. and Stal, L. J.: Phenotypic and genetic diversification of *Pseudanabaena* spp. (cyanobacteria), *ISME J.*, 3(1), 31–46, doi:10.1038/ismej.2008.78, 2009.
- Affourtit, J., Zehr, J. P. and Paerl, H. W.: Distribution of nitrogen-fixing microorganisms along the Neuse River Estuary, North Carolina, *Microb. Ecol.*, 41(2), 114–123, doi:10.1007/s002480000090, 2001.
- Anbar, A. D.: Oceans: Elements and evolution, *Science* (80-.), 322(5907), 1481–1483, doi:10.1126/science.1163100, 2008.
- Arrigo, K. R.: Marine microorganisms and global nutrient cycles, *Nature*, 438(7064), 122, doi:10.1038/nature04265, 2005.
- 395 Bauer, S., Blomqvist, S. and Ingri, J.: Distribution of dissolved and suspended particulate molybdenum, vanadium, and tungsten in the Baltic Sea, *Mar. Chem.*, 196(August), 135–147, doi:10.1016/j.marchem.2017.08.010, 2017.
- Bellenger, J. P., Wichard, T., Xu, Y. and Kraepiel, A. M. L.: Essential metals for nitrogen fixation in a free-living N₂-fixing bacterium: Chelation, homeostasis and high use efficiency, *Environ. Microbiol.*, 13(6), 1395–1411, doi:10.1111/j.1462-2920.2011.02440.x, 2011.
- 400 Bellenger, J. P., Xu, Y., Zhang, X., Morel, F. M. M. and Kraepiel, A. M. L.: Possible contribution of alternative nitrogenases to nitrogen fixation by asymbiotic N₂-fixing bacteria in soils, *Soil Biol. Biochem.*, 69, 413–420, doi:10.1016/j.soilbio.2013.11.015, 2014.
- Benavides, M., Martias, C., Elifantz, H., Berman-Frank, I., Dupouy, C. and Bonnet, S.: Dissolved organic matter influences N₂ fixation in the New Caledonian lagoon (western tropical South Pacific), *Front. Mar. Sci.*, 5(MAR), 1–12, doi:10.3389/fmars.2018.00089, 2018.
- 405 Bennett, W. W. and Canfield, D. E.: Redox-sensitive trace metals as paleoredox proxies: A review and analysis of data from modern sediments, *Earth-Science Rev.*, 204(March), 103175, doi:10.1016/j.earscirev.2020.103175, 2020.
- Bentzon-Tilia, M., Traving, S. J., Mantikci, M., Knudsen-Leerbeck, H., Hansen, J. L. S., Markager, S. and Riemann, L.: Significant N₂ fixation by heterotrophs, photoheterotrophs and heterocystous cyanobacteria in two temperate estuaries, *ISME J.*, 9(2), 273–285, doi:10.1038/ismej.2014.119, 2015.
- 410 Bertagnolli, A. D. and Stewart, F. J.: Microbial niches in marine oxygen minimum zones, *Nat. Rev. Microbiol.*, 16(12), 723–729, doi:10.1038/s41579-018-0087-z, 2018.
- Bertine, K. K. and Turekian, K. K.: Molybdenum in marine deposits, *Geochim. Cosmochim. Acta*, 37(6), 1415–1434, doi:10.1016/0016-7037(73)90080-X, 1973.

- 415 Betancourt, D. A., Loveless, T. M., Brown, J. W. and Bishop, P. E.: Characterization of diazotrophs containing Mo-independent nitrogenases, isolated from diverse natural environments, *Appl. Environ. Microbiol.*, 74(11), 3471–3480, doi:10.1128/AEM.02694-07, 2008.
- Boström, K. H., Riemann, L., Zweifel, U. L. and Hagström, Å.: *Nodularia* sp. *nifH* gene transcripts in the Baltic Sea proper, *J. Plankton Res.*, 29(4), 391–399, doi:10.1093/plankt/fbm019, 2007.
- 420 Brauer, V. S., Stomp, M., Rosso, C., Van Beusekom, S. A. M., Emmerich, B., Stal, L. J. and Huisman, J.: Low temperature delays timing and enhances the cost of nitrogen fixation in the unicellular cyanobacterium *Cyanothece*, *ISME J.*, 7(11), 2105–2115, doi:10.1038/ismej.2013.103, 2013.
- Breitburg, D., Levin, L. A., Oschlies, A., Grégoire, M., Chavez, F. P., Conley, D. J., Garçon, V., Gilbert, D., Gutiérrez, D., Isensee, K., Jacinto, G. S., Limburg, K. E., Montes, I., Naqvi, S. W. A., Pitcher, G. C., Rabalais, N. N., Roman, M. R., Rose, K. A., Seibel, B. A., Telszewski, M., Yasuhara, M. and Zhang, J.: Declining oxygen in the global ocean and coastal waters, *Science* (80-.), 359(6371), doi:10.1126/science.aam7240, 2018.
- 425 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A. and Holmes, S. P.: DADA2: High-resolution sample inference from Illumina amplicon data, *Nat. Methods*, 13(7), 581–583, doi:10.1038/nmeth.3869, 2016.
- Canfield, D. E.: Models of oxic respiration, denitrification and sulfate reduction in zones of coastal upwelling, *Geochim. Cosmochim. Acta*, 70(23 SPEC. ISS.), 5753–5765, doi:10.1016/j.gca.2006.07.023, 2006.
- 430 Capone, D. and Carpenter, E. J.: Nitrogen Fixation in the Marine Environment, , 4565(4565), 1140–1142 [online] Available from: <http://www.jstor.org/stable/1689643>, 1982.
- Carstensen, J., Andersen, J. H., Gustafsson, B. G. and Conley, D. J.: Deoxygenation of the baltic sea during the last century, *Proc. Natl. Acad. Sci. U. S. A.*, 111(15), 5628–5633, doi:10.1073/pnas.1323156111, 2014.
- 435 Chakraborty, S., Andersen, K. H., Follows, M. J. and Riemann, L.: Quantifying nitrogen fixation by heterotrophic bacteria in sinking marine particles, *Nat. Commun.*, doi:10.1038/s41467-021-23875-6, 2021.
- Chen, M., Lu, Y., Jiao, N., Tian, J., Kao, S. J. and Zhang, Y.: Biogeographic drivers of diazotrophs in the western Pacific Ocean, *Limnol. Oceanogr.*, 64(3), 1403–1421, doi:10.1002/lno.11123, 2019.
- 440 Chisnell, J. R., Premakumar, R. and Bishop, P. E.: Purification of a second alternative nitrogenase from a *nifHDK* deletion strain of *Azotobacter vinelandii*, *J. Bacteriol.*, 170(1), 27–33, doi:10.1128/jb.170.1.27-33.1988, 1988.
- Christiansen, C. F. and Löscher, C. R.: Facets of diazotrophy in the OMZ off Peru revisited- what we couldn't see from a single marker gene approach, , doi:https://doi.org/10.1101/558072, 2019.
- Church, M. J., Mahaffey, C., Letelier, R. M., Lukas, R., Zehr, J. P. and Karl, D. M.: Physical forcing of nitrogen fixation and diazotroph community structure in the North Pacific subtropical gyre, *Global Biogeochem. Cycles*, 23(2), doi:10.1029/2008GB003418, 2009.
- 445 Conley, D. J., Humborg, C., Rahm, L., Savchuk, O. P. and Wulff, F.: Hypoxia in the baltic sea and basin-scale changes in phosphorus biogeochemistry, *Environ. Sci. Technol.*, 36(24), 5315–5320, doi:10.1021/es025763w, 2002.
- Crusius, J., Calvert, S., Pedersen, T. and Sage, D.: Rhenium and molybdenum enrichments in sediments as indicators of oxic, suboxic and sulfidic conditions of deposition, *Earth Planet. Sci. Lett.*, 145(1–4), 65–78, doi:10.1016/s0012-821x(96)00204-x, 1996.
- 450 Dähnke, K. and Thamdrup, B.: Nitrogen isotope dynamics and fractionation during sedimentary denitrification in Boknis

- Eck, Baltic Sea, Biogeosciences Discuss., 10(1), 681–709, doi:10.5194/bgd-10-681-2013, 2013.
- 455 Degerholm, J., Gundersen, K., Bergman, B. and Söderbäck, E.: Phosphorus-limited growth dynamics in two Baltic Sea cyanobacteria, *Nodularia* sp. and *Aphanizomenon* sp., FEMS Microbiol. Ecol., 58(3), 323–332, doi:10.1111/j.1574-6941.2006.00180.x, 2006.
- Delwiche, C. C. and Steyn, P. L.: Nitrogen Isotope Fractionation in Soils and Microbial Reactions, Environ. Sci. Technol., 4(11), 929–935, doi:10.1021/es60046a004, 1970.
- Dera, J. and Woźniak, B.: Solar radiation in the baltic sea, Oceanologia, 52(4), 533–582, doi:10.5697/oc.52-4.533, 2010.
- 460 Diaz, R. J. and Rosenberg, R.: Spreading Dead Zones and Consequences for Marine Ecosystems, , 321(5891), 926–929, 2008.
- Dicker, H. J. and Smith, D. W.: Effects of Salinity on Acetylene Reduction (Nitrogen Fixation) and Respiration in a Marine Azotobacter, Appl. Environ. Microbiol., 42(4), 740–744, doi:10.1128/aem.42.4.740-744.1981, 1981.
- Dixon, R. and Kahn, D.: Genetic regulation of biological nitrogen fixation, Nat. Rev. Microbiol., 2(8), 621–631, doi:10.1038/nrmicro954, 2004.
- 465 Dore, J. E., Brum, J. R., Tupas, L. M. and Karl, D. M.: Seasonal and interannual variability in sources of nitrogen supporting export in the oligotrophic subtropical North Pacific Ocean, Limnol. Oceanogr., 47(6), 1595–1607, doi:10.4319/lo.2002.47.6.1595, 2002.
- 470 Dupont, C. L., Larsson, J., Yooseph, S., Ininbergs, K., Goll, J., Asplund-Samuelsson, J., McCrow, J. P., Celepli, N., Allen, L. Z., Ekman, M., Lucas, A. J., Hagström, Å., Thiagarajan, M., Brindefalk, B., Richter, A. R., Andersson, A. F., Tenney, A., Lundin, D., Tovchigrechko, A., Nylander, J. A. A., Brami, D., Badger, J. H., Allen, A. E., Rusch, D. B., Hoffman, J., Norrby, E., Friedman, R., Pinhassi, J., Venter, J. C. and Bergman, B.: Functional tradeoffs underpin salinity-driven divergence in microbial community composition, PLoS One, 9(2), doi:10.1371/journal.pone.0089549, 2014.
- Englund, B. . and Meyerson, H.: In situ Measurement of Nitrogen Fixation at Low Temperatures, , 25(3), 283–287, 1974.
- 475 Farnelid, H., Öberg, T. and Riemann, L.: Identity and dynamics of putative N₂-fixing picoplankton in the Baltic Sea proper suggest complex patterns of regulation, Environ. Microbiol. Rep., 1(2), 145–154, doi:10.1111/j.1758-2229.2009.00021.x, 2009.
- Farnelid, H., Bentzon-Tilia, M., Andersson, A. F., Bertilsson, S., Jost, G., Labrenz, M., Jürgens, K. and Riemann, L.: Active nitrogen-fixing heterotrophic bacteria at and below the chemocline of the central Baltic Sea, ISME J., 7(7), 1413–1423, doi:10.1038/ismej.2013.26, 2013.
- 480 Farnelid, H., Turk-Kubo, K., Ploug, H., Ossolinski, J. E., Collins, J. R., Van Mooy, B. A. S. and Zehr, J. P.: Diverse diazotrophs are present on sinking particles in the North Pacific Subtropical Gyre, ISME J., 13(1), 170–182, doi:10.1038/s41396-018-0259-x, 2019.
- Fernandez, C., Farías, L. and Ulloa, O.: Nitrogen fixation in denitrified marine waters, PLoS One, 6(6), doi:10.1371/journal.pone.0020539, 2011.
- 485 Fonseca-Batista, D., Li, X., Riou, V., Michotey, V., Deman, F., Fripiat, F., Guasco, S., Brion, N., Lemaitre, N., Tonnard, M., Gallinari, M., Planquette, H., Planchon, F., Sarthou, G., Elskens, M., Laroche, J., Chou, L. and Dehairs, F.: Evidence of high N₂ fixation rates in the temperate northeast Atlantic, Biogeosciences, 16(5), 999–1017, doi:10.5194/bg-16-999-2019, 2019.
- Gradoville, M. R., Farnelid, H., White, A. E., Turk-kubo, K. A., Stewart, B., Ribalet, F., Ferrón, S., Pinedo-gonzalez, P.,

- 490 Armbrust, E. V., Karl, D. M., John, S. and Zehr, J. P.: Latitudinal constraints on the abundance and activity of the cyanobacterium UCYN-A and other marine diazotrophs in the North Pacific, , 1–18, doi:10.1002/lno.11423, 2020.
- Grasshoff, G., Kremling, K. and Erhardt, M.: *Methods of seawater analysis*, Wiley VCH, Weinheim, 3ed, 1999.
- Gruber, N.: A bigger nitrogen fix, *Nature*, 436(7052), 786–787, doi:10.1038/436786a, 2005.
- Gustafsson, E., Savchuk, O. P., Gustafsson, B. G. and Müller-Karulis, B.: Key processes in the coupled carbon, nitrogen, and phosphorus cycling of the Baltic Sea, *Biogeochemistry*, 134(3), 301–317, doi:10.1007/s10533-017-0361-6, 2017.
- 495 Hall, P. O. J. and Aller, R. C.: Rapid, small-volume, flow injection analysis for SCO₂, and NH₄⁺ in marine and freshwaters, *Limnol. Oceanogr.*, 37(5), 1113–1119, doi:10.4319/lo.1992.37.5.1113, 1992.
- Hall, T. A.: *BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows*, 1999.
- 500 Hamersley, M. R., Turk, K. A., Leinweber, A., Gruber, N., Zehr, J. P., Gunderson, T. and Capone, D. G.: Nitrogen fixation within the water column associated with two hypoxic basins in the Southern California Bight, *Aquat. Microb. Ecol.*, 63(2), 193–205, doi:10.3354/ame01494, 2011.
- Harding, K., Turk-Kubo, K. A., Sipler, R. E., Mills, M. M., Bronk, D. A. and Zehr, J. P.: Symbiotic unicellular cyanobacteria fix nitrogen in the Arctic Ocean, *Proc. Natl. Acad. Sci. U. S. A.*, 115(52), 13371–13375, doi:10.1073/pnas.1813658115, 2018.
- Helcom: HELCOM. [online] Available from: <https://helcom.fi/baltic-sea-trends/indicators/>, 2018.
- 505 Hietanen, S., Jääntti, H., Buizert, C., Jürgens, K., Labrenz, M., Voss, M. and Kuparinen, J.: Hypoxia and nitrogen processing in the Baltic Sea water column, *Limnol. Oceanogr.*, 57(1), 325–337, doi:10.4319/lo.2012.57.1.0325, 2012.
- ICES Copenhagen: International Council for the Exploration of the Sea (ICES) - Dataset on Ocean Hydrography, [online] Available from: <https://ocean.ices.dk/>, 2020.
- 510 Ingall, E. and Jahnke, R.: Influence of water-column anoxia on the elemental fractionation of carbon and phosphorus during sediment diagenesis, *Mar. Geol.*, 139(1–4), 219–229, doi:10.1016/S0025-3227(96)00112-0, 1997.
- Janson, S., Carpenter, E. J. and Bergman, B.: Fine structure and immunolocalisation of proteins in *Aphanizomenon* sp. from the Baltic Sea Fine structure and immunolocalisation of proteins in *Aphanizomenon* sp. from the Baltic Sea, , 0262, doi:10.1080/09670269400650651, 1994.
- 515 Jayakumar, A., Al-Rshaidat, M. M. D., Ward, B. B. and Mulholland, M. R.: Diversity, distribution, and expression of diazotroph nifH genes in oxygen-deficient waters of the Arabian Sea, *FEMS Microbiol. Ecol.*, 82(3), 597–606, doi:10.1111/j.1574-6941.2012.01430.x, 2012.
- Jayakumar, A., Chang, B. X., Widner, B., Bernhardt, P., Mulholland, M. R. and Ward, B. B.: Biological nitrogen fixation in the oxygen-minimum region of the eastern tropical North Pacific ocean, *ISME J.*, 11(10), 2356–2367, doi:10.1038/ismej.2017.97, 2017.
- 520 Kahru, M. and Elmgren, R.: Multidecadal time series of satellite-detected accumulations of cyanobacteria in the Baltic Sea, *Biogeosciences*, 11(13), 3619–3633, doi:10.5194/bg-11-3619-2014, 2014.
- Kahru, M., Horstmann, U. and Rud, O.: Satellite detection of increased cyanobacteria blooms in the Baltic Sea: Natural fluctuation or ecosystem change?, *Ambio*, 23(8), 469–472, doi:10.2307/4314262, 1994.
- Karjalainen, M., Engström-Öst, J., Korpinen, S., Peltonen, H., Pääkkönen, J. P., Rönkkönen, S., Suikkanen, S. and Viitasalo,

- 525 M.: Ecosystem consequences of cyanobacteria in the northern Baltic Sea, *Ambio*, 36(2–3), 195–202, doi:10.1579/0044-7447(2007)36[195:ECOCIT]2.0.CO;2, 2007.
- Karlberg, M. and Wulff, A.: Impact of temperature and species interaction on filamentous cyanobacteria may be more important than salinity and increased pCO₂ levels, *Mar. Biol.*, 160(8), 2063–2072, doi:10.1007/s00227-012-2078-3, 2013.
- Kassambara, A. and Mundt, F.: *factoextra: Extract and Visualize the Results of Multivariate Data Analyses*, 2020.
- 530 Keeling, R. F., Körtzinger, A. and Gruber, N.: Ocean Deoxygenation in a Warming World, *Ann. Rev. Mar. Sci.*, 2(1), 199–229, doi:10.1146/annurev.marine.010908.163855, 2010.
- Klawonn, I., Nahar, N., Walve, J., Andersson, B., Olofsson, M., Svedén, J. B., Littmann, S., Whitehouse, M. J., Kuypers, M. M. M. and Ploug, H.: Cell-specific nitrogen- and carbon-fixation of cyanobacteria in a temperate marine system (Baltic Sea), *Environ. Microbiol.*, 18(12), 4596–4609, doi:10.1111/1462-2920.13557, 2016.
- 535 Kumar, S., Stecher, G., Li, M., Knyaz, C. and Tamura, K.: MEGA X: Molecular evolutionary genetics analysis across computing platforms, *Mol. Biol. Evol.*, 35(6), 1547–1549, doi:10.1093/molbev/msy096, 2018.
- Langlois, R. J., Hümmer, D. and LaRoche, J.: Abundances and distributions of the dominant nifH phylotypes in the Northern Atlantic Ocean, *Appl. Environ. Microbiol.*, 74(6), 1922–1931, doi:10.1128/AEM.01720-07, 2008.
- Larsson, U., Hajdu, S., Walve, J. and Elmgren, R.: Baltic Sea Nitrogen Fixation Estimated from the Summer Increase in Upper Mixed Layer., 2001.
- 540 Lehtimäki, J., Lehtimäki, L., Moisander, P., Kaarina Sivonen, † and Kononen, K.: Growth, Nitrogen Fixation, and Nodularin Production by Two Baltic Sea Cyanobacteria., 1997.
- Lennartz, S. T., Lehmann, A., Herrford, J., Malien, F., Hansen, H. P., Biester, H. and Bange, H. W.: Long-term trends at the Boknis Eck time series station (Baltic Sea), 1957-2013: Does climate change counteract the decline in eutrophication?, *Biogeosciences*, 11(22), 6323–6339, doi:10.5194/bg-11-6323-2014, 2014.
- 545 Leppanen, J., Niemi, A. and Rinne, I.: Nitrogen fixation of Cyanobacteria (blue-green algae) and the nitrogen cycle of the Baltic Sea, *Symbiosis (Philadelphia, PA)*, 6(1), 181–194, 1988.
- Li, L., Wu, C., Huang, D., Ding, C., Wei, Y. and Sun, J.: Integrating Stochastic and Deterministic Process in the Biogeography of N₂-Fixing Cyanobacterium Candidatus *Atelocyanobacterium Thalassa*, *Front. Microbiol.*, 12(October), 1–15, doi:10.3389/fmicb.2021.654646, 2021.
- 550 Liblik, T. and Lips, U.: Stratification has strengthened in the baltic sea – an analysis of 35 years of observational data, *Front. Earth Sci.*, 7(July), 1–15, doi:10.3389/feart.2019.00174, 2019.
- Löscher, C. R., Großkopf, T., Desai, F. D., Gill, D., Schunck, H., Croot, P. L., Schlosser, C., Neulinger, S. C., Pinnow, N., Lavik, G., Kuypers, M. M. M., Laroche, J. and Schmitz, R. a: Facets of diazotrophy in the oxygen minimum zone waters off Peru., *ISME J.*, 1–13, doi:10.1038/ismej.2014.71, 2014.
- 555 Loveless, T. M., Saah, J. R. and Bishop, P. E.: Isolation of nitrogen-fixing bacteria containing molybdenum-independent nitrogenases from natural environments, *Appl. Environ. Microbiol.*, 65(9), 4223–4226, doi:10.1128/aem.65.9.4223-4226.1999, 1999.
- Maar, M., Markager, S., Madsen, K. S., Windolf, J., Lyngsgaard, M. M., Andersen, H. E. and Møller, E. F.: The importance of local versus external nutrient loads for Chl a and primary production in the Western Baltic Sea, *Ecol. Modell.*, 320, 258–272, doi:10.1016/j.ecolmodel.2015.09.023, 2016.
- 560

- Main, D. C., Berry, P. H., Peet, R. L. and Robertson, J. P.: Sheep mortalities associated with the blue green alga *Nodularia spumigena*, 1977.
- 565 Man-Aharonovich, D., Kress, N., Zeev, E. B., Berman-Frank, I. and Béjà, O.: Molecular ecology of *nifH* genes and transcripts in the eastern Mediterranean Sea, *Environ. Microbiol.*, 9(9), 2354–2363, doi:10.1111/j.1462-2920.2007.01353.x, 2007.
- Marino, R., Chan, F., Howarth, R. W., Pace, M. L. and Likens, G. E.: Ecological constraints on planktonic nitrogen fixation in saline estuaries. I. Nutrient and trophic controls, *Mar. Ecol. Prog. Ser.*, 309(March), 25–39, doi:10.3354/meps309025, 2006.
- 570 Martínez-Pérez, C., Mohr, W., Löscher, C. et al: The small unicellular diazotrophic symbiont, UCYN-A, is a key player in the marine nitrogen cycle, *Nat Microbiol* 1, 16163, 2016.
- Masepohl, B. and Hallenbeck, P. C.: Nitrogen and Molybdenum Control of Nitrogen Fixation in the Phototrophic Bacterium *Rhodobacter capsulatus*, in *Adv. Exp. Med. Biol.*, 675, pp. 49–70., 2010.
- 575 McMurdie, P. J. and Holmes, S.: Phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data, *PLoS One*, 8(4), doi:10.1371/journal.pone.0061217, 2013.
- McRose, D. L., Zhang, X., Kraepiel, A. M. L. and Morel, F. M. M.: Diversity and activity of alternative nitrogenases in sequenced genomes and coastal environments, *Front. Microbiol.*, 8(FEB), 1–13, doi:10.3389/fmicb.2017.00267, 2017.
- Mills, M. M., Turk-Kubo, K. A., van Dijken, G. L., Henke, B. A., Harding, K., Wilson, S. T., Arrigo, K. R. and Zehr, J. P.: Unusual marine cyanobacteria/haptophyte symbiosis relies on N₂ fixation even in N-rich environments, *ISME J.*, (3), 580 doi:10.1038/s41396-020-0691-6, 2020.
- Moisander, P. H., Beinart, R. A., Hewson, I., White, A. E., Johnson, K. S., Carlson, C. A., Montoya, J. P., Zehr, J. P., Moisander, P. H., Beinart, R. A., White, A. E., Johnson, K. S., Carlson, C. A., Montoya, J. P. and Zehr, J. P.: Unicellular Cyanobacterial Distributions Broaden the Oceanic N₂ Fixation Domain, , 327(5972), 1512–1514, 2010.
- 585 Montoya, J. P., Voss, M., Ka, P. and Capone, D. G.: A Simple, High-Precision, High-Sensitivity Tracer Assay for N₂ Fixation ", , 62(3), 986–993, 1996.
- Morford, J. L. and Emerson, S.: The geochemistry of redox sensitive trace metals in sediments, *Geochim. Cosmochim. Acta*, 63(11–12), 1735–1750, doi:10.1016/S0016-7037(99)00126-X, 1999.
- Mus, F., Colman, D. R., Peters, J. W. and Boyd, E. S.: Geobiological feedbacks, oxygen, and the evolution of nitrogenase, *Free Radic. Biol. Med.*, 140(October 2018), 250–259, doi:10.1016/j.freeradbiomed.2019.01.050, 2019.
- 590 Nehring, S.: Mortality of dogs associated with a mass development of *Nodularia spumigena* (Cyanophyceae) in a brackish lake at the German North Sea coast, *J. Plankton Res.*, 15(7), 1993.
- Oksanen, F. J., Blanchet, G., Friendly, M., Roeland Kindt, P., Legendre, D., McGlenn, P. R., Minchin, R. B. O., Simpson, G. L., Peter Solymos, M., Stevens, H. H., Szoecs, E. and Wagner, H.: *vegan: Community Ecology Package*, 2020.
- 595 Olofsson, M., Egardt, J., Singh, A. and Ploug, H.: Inorganic phosphorus enrichments in Baltic Sea water have large effects on growth, carbon fixation, and N₂ fixation by *Nodularia spumigena*, *Aquat. Microb. Ecol.*, 2016.
- Olofsson, M., Torstensson, A., Karlberg, M., Steinhoff, F. S., Dinasquet, J., Riemann, L., Chierici, M. and Wulff, A.: Limited response of a spring bloom community inoculated with filamentous cyanobacteria to elevated temperature and pCO₂, *Bot. Mar.*, 62(1), 3–16, doi:10.1515/bot-2018-0005, 2019.

- 600 Olofsson, M., Suikkanen, S., Kobos, J., Wasmund, N. and Karlson, B.: Basin-specific changes in filamentous cyanobacteria community composition across four decades in the Baltic Sea, *Harmful Algae*, 91(October 2019), 101685, doi:10.1016/j.hal.2019.101685, 2020.
- Paerl, H. W. and Prufert, L. E.: Oxygen-Poor Microzones as Potential Sites of Microbial N₂ Fixation in Nitrogen-Depleted Aerobic Marine Waters, *Appl. Environ. Microbiol.*, 53(5), 1078–1087, doi:10.1128/aem.53.5.1078-1087.1987, 1987.
- 605 Paul, A. J., Achterberg, E. P., Bach, L. T., Boxhammer, T., Czerny, J., Haunost, M., Schulz, K. G., Stuhr, A. and Riebesell, U.: No observed effect of ocean acidification on nitrogen biogeochemistry in a summer Baltic Sea plankton community, *Biogeosciences*, 13(13), 3901–3913, doi:10.5194/bg-13-3901-2016, 2016.
- Pedersen, J. N., Bombar, D., Paerl, R. W. and Riemann, L.: Diazotrophs and N₂-Fixation associated with particles in coastal estuarine waters, *Front. Microbiol.*, 9(NOV), 1–11, doi:10.3389/fmicb.2018.02759, 2018.
- 610 Pelve, E. A., Fontanez, K. M. and DeLong, E. F.: Bacterial succession on sinking particles in the ocean's interior, *Front. Microbiol.*, 8(NOV), 1–15, doi:10.3389/fmicb.2017.02269, 2017.
- Ploug, H., Musat, N., Adam, B., Moraru, C. L., Lavik, G., Vagner, T., Bergman, B. and Kuypers, M. M. M.: Carbon and nitrogen fluxes associated with the cyanobacterium *Aphanizomenon* sp. in the Baltic Sea, *ISME J.*, 4(9), 1215–1223, doi:10.1038/ismej.2010.53, 2010.
- 615 Ploug, H., Adam, B., Musat, N., Kalvelage, T., Lavik, G., Wolf-Gladrow, D. and Kuypers, M. M. M.: Carbon, nitrogen and O₂ fluxes associated with the cyanobacterium *Nodularia spumigena* in the Baltic Sea, *ISME J.*, 5(9), 1549–1558, doi:10.1038/ismej.2011.20, 2011.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools, *Nucleic Acids Res.*, 41(D1), 590–596, doi:10.1093/nar/gks1219, 2013.
- 620 R Foundation for Statistical Computing: R: A language and environment for statistical computing, [online] Available from: <https://www.r-project.org/>, 2017.
- Rahav, E., Herut, B., Levi, A., Mulholland, M. R. and Berman-Frank, I.: Springtime contribution of dinitrogen fixation to primary production across the Mediterranean Sea, *Ocean Sci.*, 9(3), 489–498, doi:10.5194/os-9-489-2013, 2013.
- 625 Rakko, A. and Seppälä, J.: Effect of salinity on the growth rate and nutrient stoichiometry of two Baltic Sea filamentous cyanobacterial species, *Est. J. Ecol.*, 63(2), 55–70, doi:10.3176/eco.2014.2.01, 2014.
- Redfield, A. C.: Redfield 1934.pdf, Univ. Press Liverpool, James Johnstone Meml. Vol., 176–192 [online] Available from: <http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:On+the+proportions+of+organic+derivatives+in+sea+water+and+their+relation+to+the+composition+of+plankton#0>, 1934.
- 630 Reusch, T. B. H., Dierking, J., Andersson, H. C., Bonsdorff, E., Carstensen, J., Casini, M., Czajkowski, M., Hasler, B., Hinsby, K., Hyytiäinen, K., Johannesson, K., Jomaa, S., Jormalainen, V., Kuosa, H., Kurland, S., Laikre, L., Mackenzie, B. R., Margonski, P., Melzner, F., Oesterwind, D., Ojaveer, H., Refsgaard, J. C., Sandström, A., Schwarz, G., Tonderski, K., Winder, M. and Zandersen, M.: The Baltic Sea as a time machine for the future coastal ocean. [online] Available from: <http://advances.sciencemag.org/>, 2018.
- 635 Robson, R. L., Eady, R. R., Richardson, T. H., Miller, R. W., Hawkins, M. and Postgate, J. R.: The alternative nitrogenase of *Azotobacter chroococcum* is a vanadium enzyme, *Nature*, 1986.
- Rutgersson, A., Jaagus, J., Schenk, F. and Stendel, M.: Observed changes and variability of atmospheric parameters in the

- Baltic Sea region during the last 200 years, *Clim. Res.*, 61(2), 177–190, doi:10.3354/cr01244, 2014.
- Saito, M. A., Tyler, G. J. and Ritt, J. T.: Some Thoughts on the Concept of Colimitation : Three Definitions and the Importance of Bioavailability, , 53(1), 276–290, 2008.
- 640 Saraiva, S., Markus Meier, H. E., Andersson, H., Höglund, A., Dieterich, C., Gröger, M., Hordoir, R. and Eilola, K.: Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates, *Clim. Dyn.*, 52(5–6), 3369–3387, doi:10.1007/s00382-018-4330-0, 2019.
- Saxena, H., Sahoo, D., Khan, M. A., Kumar, S., Sudheer, A. K. and Singh, A.: Dinitrogen fixation rates in the Bay of Bengal during summer monsoon, *Environ. Res. Commun.*, 2(5), 051007, doi:10.1088/2515-7620/ab89fa, 2020.
- 645 Schmidtko, S., Stramma, L. and Visbeck, M.: Decline in global oceanic oxygen content during the past five decades, *Nature*, 542(7641), 335–339, doi:10.1038/nature21399, 2017.
- Schoffelen, N. J., Mohr, W., Ferdelman, T. G., Littmann, S., Duerschlag, J., Zubkov, M. V., Ploug, H. and Kuypers, M. M. M.: Single-cell imaging of phosphorus uptake shows that key harmful algae rely on different phosphorus sources for growth, *Sci. Rep.*, 8(1), 1–14, doi:10.1038/s41598-018-35310-w, 2018.
- 650 Severin, I., Confurius-Guns, V. and Stal, L. J.: Effect of salinity on nitrogenase activity and composition of the active diazotrophic community in intertidal microbial mats, *Arch. Microbiol.*, 194(6), 483–491, doi:10.1007/s00203-011-0787-5, 2012.
- Short, S. M. and Zehr, J. P.: Nitrogenase gene expression in the Chesapeake Bay Estuary, *Environ. Microbiol.*, 9(6), 1591–1596, doi:10.1111/j.1462-2920.2007.01258.x, 2007.
- 655 Singh, A., Bach, L. T., Löscher, C. R., Paul, A. J., Ojha, N. and Riebesell, U.: Impact of increasing carbon dioxide on dinitrogen and carbon fixation rates under oligotrophic conditions and simulated upwelling, *Limnol. Oceanogr.*, 1–13, doi:10.1002/lno.11795, 2021.
- Sivonen, K., Kononen, K., Carmichael, W. W., Dahlem, A. M., Rinehart, K. L., Kiviranta, J. and Niemela, S. I.: Occurrence of the hepatotoxic cyanobacterium *Nodularia spumigena* in the Baltic Sea and structure of the toxin., *Appl. Environ. Microbiol.*, 55(8), 1990–1995, doi:10.1128/aem.55.8.1990-1995.1989, 1989.
- 660 Staal, M., Te Lintel Hekkert, S., Herman, P. and Stal, L. J.: Comparison of models describing light dependence of N₂ fixation in heterocystous cyanobacteria, *Appl. Environ. Microbiol.*, 68(9), 4679–4683, doi:10.1128/AEM.68.9.4679-4683.2002, 2002.
- Stal, L. J.: Is the distribution of nitrogen-fixing cyanobacteria in the oceans related to temperature?: Minireview, *Environ. Microbiol.*, 11(7), 1632–1645, doi:10.1111/j.1758-2229.2009.00016.x, 2009.
- Stal, L. J., Albertano, P., Bergman, B., Von Bröckel, K., Gallon, J. R., Hayes, P. K., Sivonen, K. and Walsby, A. E.: BASIC: Baltic Sea cyanobacteria. An investigation of the structure and dynamics of water blooms of cyanobacteria in the Baltic Sea - Responses to a changing environment, *Cont. Shelf Res.*, 23(17–19), 1695–1714, doi:10.1016/j.csr.2003.06.001, 2003.
- 670 Steward, G. F., Jenkins, B. D., Ward, B. B., Zehr, J. P. and Short, S. M.: Fingerprinting diazotroph communities in the Chesapeake Bay by using a DNA microarray, *Appl. Environ. Microbiol.*, 70(3), 1767, doi:10.1128/AEM.70.3.1455, 2004.
- Stigebrandt, A. and Andersson, A.: The Eutrophication of the Baltic Sea has been Boosted and Perpetuated by a Major Internal Phosphorus Source, *Front. Mar. Sci.*, 7(November), doi:10.3389/fmars.2020.572994, 2020.
- Stramma, L., Johnson, G. C., Sprintall, J. and Mohrholz, V.: Expanding Oxygen-Minimum Zones in the Tropical Oceans,

- Source Sci. New Ser., 320(5876), 655–658 [online] Available from: <http://www.jstor.org/stable/20055146> (Accessed 26 February 2018), 2008.
- 675 Suikkanen, S., Kaartokallio, H., Hällfors, S., Huttunen, M. and Laamanen, M.: Life cycle strategies of bloom-forming, filamentous cyanobacteria in the Baltic Sea, *Deep. Res. Part II Top. Stud. Oceanogr.*, 57(3–4), 199–209, doi:10.1016/j.dsr2.2009.09.014, 2010.
- 680 Svedén, J. B., Adam, B., Walve, J., Nahar, N., Musat, N., Lavik, G., Whitehouse, M. J., Kuypers, M. M. M. and Ploug, H.: High cell-specific rates of nitrogen and carbon fixation by the cyanobacterium *Aphanizomenon* sp. at low temperatures in the Baltic Sea, *FEMS Microbiol. Ecol.*, 91(12), 1–11, doi:10.1093/femsec/fiv131, 2015.
- Tan, J. W., Thong, K. L., Arumugam, N. D., Cheah, W. L., Lai, Y. W., Chua, K. H., Rahim, R. A. and Vikineswary, S.: Development of a PCR assay for the detection of *nifH* and *nifD* genes in indigenous photosynthetic bacteria, *Int. J. Hydrogen Energy*, 34(17), 7538–7541, doi:10.1016/j.ijhydene.2009.04.029, 2009.
- 685 Tang, W. and Cassar, N.: Data-Driven Modeling of the Distribution of Diazotrophs in the Global Ocean, *Geophys. Res. Lett.*, 46(21), 12258–12269, doi:10.1029/2019GL084376, 2019.
- Tang, W., Wang, S., Fonseca-Batista, D., Dehairs, F., Gifford, S., Gonzalez, A. G., Gallinari, M., Planquette, H., Sarthou, G. and Cassar, N.: Revisiting the distribution of oceanic N₂ fixation and estimating diazotrophic contribution to marine production, *Nat. Commun.*, 10(1), doi:10.1038/s41467-019-08640-0, 2019.
- 690 Turk-Kubo, K. A., Karamchandani, M., Capone, D. G. and Zehr, J. P.: The paradox of marine heterotrophic nitrogen fixation: Abundances of heterotrophic diazotrophs do not account for nitrogen fixation rates in the Eastern Tropical South Pacific, *Environ. Microbiol.*, 16(10), 3095–3114, doi:10.1111/1462-2920.12346, 2014.
- Vahtera, E., Conley, D. J., Gustafsson, B. G., Kuosa, H., Pitkänen, H., Savchuk, O. P., Tamminen, T., Viitasalo, M., Voss, M., Wasmund, N. and Wulff, F.: Internal ecosystem feedbacks enhance nitrogen-fixing cyanobacteria blooms and complicate management in the Baltic Sea, *Ambio*, 36, 186–194, doi:10.1579/0044-7447(2007)36[186:IEFENC]2.0.CO;2, 2007.
- 695 Wacklin, P., Hoffmann, L. and Komárek, J.: Nomenclatural validation of the genetically revised cyanobacterial genus *Dolichospermum* (Ralfs ex Bornet et Flahault) comb. nova, *Fottea*, 9(1), 59–64, doi:10.5507/fot.2009.005, 2009.
- 700 Wannicke, N., Frey, C., Law, C. S. and Voss, M.: The response of the marine nitrogen cycle to ocean acidification, *Glob. Chang. Biol.*, 24(11), 5031–5043, doi:10.1111/gcb.14424, 2018.
- Wasmund, N.: Occurrence of cyanobacterial blooms in the baltic sea in relation to environmental conditions, *Int. Rev. der Gesamten Hydrobiol.*, 82(2), 169–174, doi:10.1002/iroh.19970820205, 1997.
- Wasmund, N., Tuimala, J., Suikkanen, S., Vandepitte, L. and Kraberg, A.: Long-term trends in phytoplankton composition in the western and central Baltic Sea, *J. Mar. Syst.*, 87(2), 145–159, doi:10.1016/j.jmarsys.2011.03.010, 2011.
- 705 Wulff, A., Karlberg, M., Olofsson, M., Torstensson, A., Riemann, L., Steinhoff, F. S., Mohlin, M., Ekstrand, N. and Chierici, M.: Ocean acidification and desalination: climate-driven change in a Baltic Sea summer microplanktonic community, *Mar. Biol.*, 165(4), 1–15, doi:10.1007/s00227-018-3321-3, 2018.
- 710 Zani, S., Mellon, M. T., Collier, J. L. and Zehr, J. P.: Expression of *nifH* Genes in Natural Microbial Assemblages in Lake George, New York, Detected by Reverse Transcriptase PCR, *Appl. Environ. Microbiol.*, 66(7), 3119 [online] Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC92123/> (Accessed 21 April 2019), 2000.
- Zehr, J. P., Mellon, M., Braun, S., Litaker, W., Steppe, T. and Paerl, H. W.: Diversity of heterotrophic nitrogen fixation

genes in a marine cyanobacterial mat, *Appl. Environ. Microbiol.*, 61(7), 2527–2532, doi:10.1128/aem.61.7.2527-2532.1995, 1995.

715 Zehr, J. P., Mellon, M. T. and Zani, S.: New nitrogen-fixing microorganisms detected in oligotrophic oceans by amplification of Nitrogenase (*nifH*) genes., *Appl. Environ. Microbiol.*, 64(9), 3444–50 [online] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9726895> (Accessed 21 April 2019), 1998.

Zehr, J. P., Jenkins, B. D., Short, S. M. and Steward, G. F.: Nitrogenase gene diversity and microbial community structure: A cross-system comparison, *Environ. Microbiol.*, 5(7), 539–554, doi:10.1046/j.1462-2920.2003.00451.x, 2003.

720 Zehr, J. P., Shilova, I. N., Farnelid, H. M., Muñoz-Maríncarmen, M. D. C. and Turk-Kubo, K. A.: Unusual marine unicellular symbiosis with the nitrogen-fixing cyanobacterium UCYN-A, *Nat. Microbiol.*, 2, doi:10.1038/nmicrobiol.2016.214, 2016.

Zhang, X., McRose, D. L., Darnajoux, R., Bellenger, J. P., Morel, F. M. M. and Kraepiel, A. M. L.: Alternative nitrogenase activity in the environment and nitrogen cycle implications, *Biogeochemistry*, 127(2–3), 189–198, doi:10.1007/s10533-016-0188-6, 2016.

725

730

735




Figure 1. Overview of cruise track and stations from this study (A) Distribution of O₂, salinity (sal) in surface water (5 m) and the deepest sample at the station (last) in the Baltic Sea. Stations from where N₂ fixation were obtained are coloured red. Data for figure B were obtained through ICES oceanography spanning from 2019-08-01 to 2019-10-30 (ICES Copenhagen, 2020).





Figure 2 Distribution of surface water O₂ ($\mu\text{mol L}^{-1}$), temperature (temp) ($^{\circ}\text{C}$), salinity (sal), dissolved inorganic carbon (DIC) (mmol L^{-1}) and NO_x (nitrate + nitrite) and PO₄³⁻ in $\mu\text{mol L}^{-1}$.



750

Figure 4-3 N:P (A) and C:N (B) ratios from station KB06, BB23, BB15, BB08 and GB107. The black lines depict the N:P (16:1) and C:N (106:16) ratios after Redfield. The red lines indicate the trendlines of our data. An N:P ratio below the canonical Redfield ratio of 16:1 suggests a niche for N₂ fixation. However, the positive intercept speaks for DIN still being available, which might be considered as unfavorable for N₂ fixation (A). A C:N ratio below Redfield might suggest a slight deficiency in N (B), this might result from degrading organic matter in the POM pool.

755




Figure 5-4 C (A) and N₂ (B) fixation rates [nmol L⁻¹ d⁻¹]. Similar in all stations (five stations covering three depths), C fixation increased over depth, followed by a decrease at a water depth of 41 m. Higher rates were observed in the Bornholm and Eastern Gotland Basins compared to the Kiel and Arkona Basins. N₂ fixation rates increased slightly over depth, followed by a decrease at 41 m water depth in the Bornholm Basin. (C) δ¹⁵N [‰] from station KB06, H21, BB08, BB15 and GB107. The red line indicates a δ¹⁵N of 2 ‰. Values below 2 ‰ speak for previous or current N₂ fixation (Dähnke and Thamdrup, 2013).

760





Figure 5 Maximum likelihood tree of *nifH* amplicons (321 bp) identified by Sanger sequencing. Clusters identified are denoted as grey triangles while identified individual sequences are denoted as 'Seq + NCBI submission ID'. Cluster I cyanobacteria are shown in green, Cluster I proteobacteria are shown in blue, and Cluster III diazotrophs are shown in orange.



770 **Figure 6** Maximum likelihood tree of *vnf/anfD* amplicons (591 bp) identified by Sanger sequencing. All recovered sequences clustered with *Rhodospseudomonas palustris* (grey triangle). *anfD* sequences are represented in brown, *vnfD* sequences are represented in green. A *nifD* Sequence has been chosen as the outgroup.

775

780




Figure 7 Cluster-specific *nifH* (A) gene, (B) transcript abundances and (C) transcript per gene copy of *Nodularia*, *UCYN-A* and *Gamma-PO/AO*. In general, both cyanobacterial groups are decreasing in *nifH* gene abundance over depth at all stations. Transcript abundances of *Nodularia-nifH* were higher in surface waters (5 m), contrary to *UCYN-A*, which showed an opposite trend. *Gamma-PO/AO* gene and transcripts abundance generally increased over depth.




Figure 8 Principal component analysis (PCA) of components one and two (A), one and three (B). Positive correlations were identified between salinity and UCYN-A, while *Nodularia* and salinity are negatively correlated. Moreover, N₂ fixation correlated positively with *Nodularia*. Stations are color-coded. Each circle denoted one sampling point, bigger circles indicate the center for each station.

800

Table 1 Contribution of N₂ fixation to primary production (PP) in different ocean basins. Values represent mean values for the respective regions.

Location	PP Contribution (%)	Source
North east Atlantic	25	(Fonseca-Batista et al., 2019; Tang et al., 2019)
West Northeast Atlantic	20	
Mediterranean Sea	8	(Church et al., 2009; Dore et al., 2002; Fonseca-Batista et al., 2019; Rahav et al., 2013; Saxena et al., 2020)
North Pacific Ocean	1-4	
Bay of Bengal	<1	
Baltic Sea	8-25	This study

805